

Synthesis, and Fluorescence Properties of Coumarin and Benzocoumarin Derivatives Conjugated Pyrimidine Scaffolds for Biological Imaging Applications

Najim A. Al-Masoudi¹ · Niran J. Al-Salihi¹ · Yossra A. Marich¹ · Timo Markus²

Received: 10 June 2015 / Accepted: 28 September 2015
© Springer Science+Business Media New York 2015

Abstract Series of coumarin and 5,6-benzomcomarin substituted pyrimidine derivatives **11–15** and **22–25** were synthesized, aiming to develop new imaging fluorescent agents. Analogously, treatment of 4-chloropyrimidine analog **16** with coumarin 3-carbohyrazide **5** under MWI condition followed by boiling with NH₄OAc in HOAc furnished coumarin-1,2,4-triazolo-pyrimidine analog **18**. The fluorescence property was investigated spectrophotometrically in MeOH with Rhodamine 6G as standard dye. All the compounds showed emission in the region between 331 and 495 nm. The quantum yield of all the compounds were found to be weak, except methyl benzocoumarin 3-carboxylate **22** which showed ($\Phi_F=0.98$) in comparison to Rhodamine 6G as standard ($\Phi_F=0.95$).

Keywords Coumarins · Fluorescence · Pyrimidines · UV-visible absorption

Introduction

Coumarin and its derivatives are one of the important classes of heterocyclic compounds which occur in many

natural products with pharmacological activity [1–6]. Coumarin compounds are known to possess a wide range of biological activities such as antibacterial [7], anticancer [8, 9], anticoagulants [10], anti-HIV protease inhibitors [11], anti-HIV integrases [12, 13], serine protease inhibitors [14], inhibitors of steroid 5 α -reductase [15], and

NO synthase inhibitors [16]. Geiparvarin **1** (Fig. 1), a naturally occurring product bearing the coumarin residue, has been shown to possess a significant inhibitory activity against a variety of cancer cell lines [17]. Due to efficient light emission properties, the coumarin derivatives are of the most importance as significant organic fluorescent materials [18] for biochemical and biological imaging applications [19]. Lee et al. [20] have reported novel coumarin-based fluorogenic probe bearing the 2-picolyl as a fluorescent chemosensor with high selectivity and suitable affinity in biological systems toward Cu²⁺, meanwhile Rajisha et al. [21] have synthesized new benzocoumarin-oxadiazolyls as strong blue-green fluorescent brighteners research with good bathochromic shifts. Other coumarin derivatives are used as fluorescent imaging agents such as 7-amino-4-methyl coumarin-3-acetic acid (AMCA) **2** [22] in fluorohistochemical examination of human kidney glomeruli and 3-(4-aminophenyl)-2H-chromen-2-one (CMC) **3** (Fig. 1) can be used for in situ fluorescent imaging of myelin in the vertebrate nervous system [23].

Fluorescence Imaging (FI) is one of the most popular imaging modes in biomedical sciences for the visualisation of cells and tissues both in vitro and in vivo [24]. Despite that a number of fluorescent imaging agents have been reported, we synthesized here new coumarin and benzocoumarin derivatives conjugated pyrimidines scaffolds with study of their fluorescence properties, aiming to develop new imaging agents can be used in living cells.


Electronic supplementary material The online version of this article (doi:10.1007/s10895-015-1677-z) contains supplementary material, which is available to authorized users.

✉ Najim A. Al-Masoudi
najim.al-masoudi@gmx.de

¹ Department of Chemistry, College of Science, University of Basrah, Basrah 61001, Iraq

² Department of Chemistry, University of Konstanz, P.O. Box 5560, 78457 Konstanz, Germany

Fig. 1 Some potential coumarin derivatives



Experimental Details

Physical Measurements

Melting points are uncorrected and were measured with a Büchi melting point apparatus B-545 (Büchi Labortechnik AG, Switzerland). NMR data were obtained on 400 and 600 MHz (^1H) and 100 and 150.91 MHz (^{13}C) spectrometers (Avance III, Bruker, Germany), respectively, with TMS as internal standard and on the δ scale in ppm. Heteronuclear assignments were verified by ^1H , ^{13}C HMBC and ^1H , ^{13}C HSQC NMR experiments. Microanalytical data were obtained with a Vario EL (Shimadzu, Japan). Analytical silica gel TLC plates 60 F₂₅₄ were purchased from Merck. Microwave-assisted reactions were carried out in a CEM Focused Microwave Synthesis System (100–450 W). All reagents were obtained from commercial suppliers and were used without further purification.

General Procedure for the Preparation of Coumarinyl-Pyrimidine Analogues (**11**–**15**)

To a stirred solution of coumarin-3-carbohydrazide (**5**) (100 mg, 0.50 mmol) in DMF (10 mL) and substituted arylazopyrimidines **6**–**10** (0.50 mmol) were heated in an oil bath at 90–100 °C for 4–5 h. The mixture was evaporated to dryness and the residue was co-evaporated with silica gel (1.0 g) in MeOH then poured on a short SiO₂ column (5.0 g). Elution, in gradient, with MeOH (0–10 % v:v) and CHCl₃ as eluent afforded the desired products.

N'-(2,6-Diamino-5-((4-chlorophenyl)diazenyl)pyrimidin-4-yl)-2-oxo-2H-chromene-3-carbohydrazide (**11**)

From 2,6-diamino-4-chloro-5-*p*-chlorophenylazopyrimidine (**6**) (141 mg). Yield: 120 mg (53 %), m.p. 203–210 °C R_f =0.65. ^1H NMR (DMSO- d_6): δ 11.14 (br s., 1H, NH), 9.26 (br s., 1H, NH), 9.01 (s, 1H, H_{coum.}-4), 8.18 (br s., 2H, NH₂), 7.82 (dd, 1H, $J_{5,7}$ =1.4 Hz, $J_{5,6}$ =7.7 Hz, H-5), 7.70 (dt, 1H, $J_{6,7}$ = $J_{7,8}$ =7.8 Hz, H-7), 7.57 (d, 2H, $J_{2,3'}$ =8.0 Hz, H_{arom.}-2'+H_{arom.}-6'), 7.41 (d, 2H, $J_{5,6'}$ =8.0 Hz, H_{arom.}-3'+H_{arom.}-5'), 7.00 (m, 2H, H_{coum.}-6+H_{coum.}-8), 6.81 (d, 2H, J =4.5 Hz, NH₂). ^{13}C NMR (DMSO- d_6): δ 165.1 (C₉=O), 163.2 (C_{pyrimid.}-4), 161.7 (C_{pyrimid.}-2), 159.1 (C_{coum.}-2), 157.0 (C_{pyrimid.}-6), 151.1 (C_{coum.}-8a), 133.9 (C-Cl), 131.1 (C_{arom.}-2'+C_{arom.}-6'), 129.8 (C_{arom.}-3'+C_{arom.}-5'), 129.0 (C_{coum.}-5+C_{coum.}-7), 128.6 (C_{arom.}-1'), 123.5 (C_{coum.}-6), 119.2 (C_{coum.}-4), 118.7 (C_{coum.}-

4a), 117.0 (C_{coum.}-8), 113.1 (C_{coum.}-3), 102.4 (C_{pyrimid.}-5). Anal. Calcd for C₂₀H₁₅ClN₈O₃ (450.84): C, 53.28; H, 3.35; N, 24.85. Found: C, 53.01; H, 3.28; N, 24.77 %.

N'-(2,6-Diamino-5-((4-bromophenyl)diazenyl)pyrimidin-4-yl)-2-oxo-2H-chromene-3-carbohydrazide (**12**)

From 2,6-diamino-4-chloro-5-*p*-bromophenylazopyrimidine (**7**) (164 mg). Yield: 153 mg (62 %), m. p. 222–223 °C (dec.). R_f =0.72. ^1H NMR (DMSO- d_6): δ 11.11 (br s., 1H, NH), 9.40 (br s., 1H, NH), 9.00 (s, 1H, H_{coum.}-4), 8.35 (m, 3H, NH₂+H-5), 7.98 (dt, 1H, $J_{5,7}$ =1.5 Hz, $J_{7,8}$ = $J_{6,7}$ =7.7 Hz, H_{coum.}-7), 7.68 (d, 2H, $J_{2,3'}$ =7.9 Hz, H_{arom.}-2'+H_{arom.}-6'), 7.41 (m, 3H, H_{arom.}-3'+H_{arom.}-5'+H_{coum.}-6), 6.97 (m, 3H, NH₂+H_{coum.}-8). ^{13}C NMR (DMSO- d_6): δ 165.1 (C₉=O), 163.2 (C_{pyrimid.}-4), 161.7 (C_{pyrimid.}-2), 159.1 (C_{coum.}-2), 156.4 (C_{pyrimid.}-6), 151.8 (C_{coum.}-8a), 133.7 (C-Br), 132.7 (C_{arom.}-2'+C_{arom.}-6'), 131.3 (C_{coum.}-5+C_{coum.}-7), 123.8 (C_{coum.}-6), 120.1 (C_{coum.}-4), 119.1 (C_{coum.}-4a), 118.7 (C_{coum.}-8), 112.5 (C_{coum.}-3), 102.7 (C_{pyrimid.}-5'). Anal. Calcd for C₂₀H₁₅BrN₈O₃ (495.29): C, 48.50; H, 3.05; N, 22.62. Found: C, 48.31; H, 2.98; N, 22.40 %.

N'-(2,6-Diamino-5-((4-nitrophenyl)diazenyl)pyrimidin-4-yl)-2-oxo-2H-chromene-3-carbohydrazide (**13**)

From 2,6-diamino-4-chloro-5-*p*-nitrophenylazopyrimidine (**8**) (164 mg). Yield: 138 mg (60 %), m.p. 198–201 °C (dec.). R_f =0.79. ^1H NMR (DMSO- d_6): δ 11.11 (br s., 1H, NH), 9.40 (br s., 1H, NH), 9.00 (s, 1H, H_{coum.}-4), 8.48 (br s., 2H, NH₂), 8.36 (d, 2H, $J_{5,6'}$ =8.9 Hz, H_{arom.}-3'+H_{arom.}-5'), 7.98 (m, 2H, H_{coum.}-5+H_{coum.}-7), 7.69 (d, 2H, $J_{2,3'}$ =8.9 Hz, H_{arom.}-2'+H_{arom.}-6'), 7.40 (m, 2H, H_{coum.}-6+H_{coum.}-8), 6.98 (d, 2H, J =5.9 Hz, NH₂). ^{13}C NMR (DMSO- d_6): δ 163.6 (C₉=O), 161.3 (C_{pyrimid.}-4'), 159.4 (C_{pyrimid.}-2'), 158.5 (C_{coum.}-2), 156.2 (C_{pyrimid.}-6), 155.7 (C_{coum.}-8a), 146.5 (C-NO₂), 133.3 (C_{arom.}-1'), 130.7 (C_{arom.}-2'+C_{arom.}-6'), 126.9 (C_{coum.}-5+C_{coum.}-7), 124.9 (C_{coum.}-6), 122.0 (C_{arom.}-3'+C_{arom.}-5'), 119.5 (C_{coum.}-4), 118.1 (C_{coum.}-4a), 116.4 (C_{coum.}-8), 112.5 (C_{coum.}-3), 103.0 (C_{pyrimid.}-5). Anal. Calcd for C₂₀H₁₅BrN₉O₅ (461.39): C, 52.06; H, 3.28; N, 27.32. Found: C, 51.89; H, 3.17; N, 27.08 %.

Methyl 4-((2,6-diamino-6-(2-oxo-2H-chromene-3-carbonyl)hydrazinyl)pyrimidin-5-yl) diazenyl)benzoate (**14**)

From methyl 4-((2,6-diamino-4-chloropyrimidin-5-yl)diazanyl)benzoate (**9**) (153 mg). Yield: 52 mg (22 %),

m.p. 212–216 °C. R_f =0.60. ^1H NMR (DMSO- d_6): δ 11.12 (br s., 1H, NH), 9.00 (s, 2H, NH+H-4), 8.00 (d, 1H, J =8.7 Hz, $\text{H}_{\text{arom.}-2'}+\text{H}_{\text{arom.}-6'}$), 7.69 (m, 2H, $\text{H}_{\text{coulm.}-5}+\text{H}_{\text{coulm.}-7}$), 7.43 (d, 2H, J =8.7 Hz, $\text{H}_{\text{arom.}-3'}+\text{H}_{\text{arom.}-5'}$), 7.00–6.95 (m, 4H, $\text{NH}_2+\text{H}_{\text{coulm.}-6}+\text{H}_{\text{coulm.}-8}$), 3.90 (s, 3H, CO_2Me). ^{13}C NMR (DMSO- d_6): δ 164.1 (CO_2Me), 163.3 ($\text{C}_9=\text{O}$), 162.6 ($\text{C}_{\text{pyrimid.}-4}$), 159.6 ($\text{C}_{\text{pyrimid.}-2}$), 158.5 ($\text{C}_{\text{coulm.}-2}$), 155.5 ($\text{C}_{\text{pyrimid.}-6}$), 152.8 ($\text{C}_{\text{coulm.}-8a}$), 133.1 ($\text{C}_{\text{arom.}-1'}$), 130.7 ($\text{C}_{\text{arom.}-4'}$), 129.4 ($\text{C}_{\text{arom.}-3'}+\text{C}_{\text{arom.}-5'}$), 127.8 ($\text{C}_{\text{coulm.}-5}+\text{C}_{\text{coulm.}-7}+\text{C}_{\text{arom.}-2'}+\text{C}_{\text{arom.}-6'}$), 124.2 ($\text{C}_{\text{coulm.}-6}$), 119.5 ($\text{C}_{\text{coulm.}-4}$), 118.1 ($\text{C}_{\text{coulm.}-4a}$), 116.4 ($\text{C}_{\text{coulm.}-8}$), 113.1 ($\text{C}_{\text{coulm.}-3}$), 99.9 ($\text{C}_{\text{pyrimid.}-5}$), 51.6 (CO_2Me). Anal. Calcd for $\text{C}_{22}\text{H}_{18}\text{N}_8\text{O}_5$ (474.43): C, 55.70; H, 3.82; N, 23.62. Found: C, 55.49; H, 3.70; N, 23.40 %.

N-(4-((2,4-Diamino-6-(2-oxo-2H-chromene-3-carbonyl)hydrazinyl)pyrimidin-5-yl)diazenyl)phenyl)acetamide (**15**)

From 2,6-diamino-4-chloro-5-*p*-acetamidophenylazopyrimidine (**8**)**10** (153 mg). Yield: 88 mg (37 %), m.p. 288 °C (dec.). R_f =0.62. ^1H NMR (DMSO- d_6): δ 11.11 (br s., 1H, NH), 9.00 (s, 1H, $\text{H}_{\text{coulm.}-4}$), 7.70 (m, 6H, $\text{NH}_2+\text{H}_{\text{arom.}-2'}+\text{H}_{\text{arom.}-6'}+\text{H}_{\text{coulm.}-5}+\text{H}_{\text{coulm.}-7}$), 7.42 (d, 2H, $J_{3,4}=8.5$ Hz, $\text{H}_{\text{arom.}-3'}+\text{H}_{\text{arom.}-5'}$), 7.39 (m, 2H, $\text{H}_{\text{coulm.}-6}+\text{H}_{\text{coulm.}-8}$), 7.11 (s, 1H, NHMe), 6.97 (d, 2H, J =6.0 Hz, NH_2), 2.32 (s, 3H, NHMe). ^{13}C NMR (DMSO- d_6): δ 169.0 (CONH), 165.0 ($\text{C}_9=\text{O}$), 162.6 ($\text{C}_{\text{pyrimid.}-4}$), 160.1 ($\text{C}_{\text{pyrimid.}-2}$), 158.5 ($\text{C}_{\text{coulm.}-2}$), 156.5 ($\text{C}_{\text{pyrimid.}-6}$), 153.1 ($\text{C}_{\text{coulm.}-8a}$), 137.8 ($\text{C}_{\text{arom.}-4'}$), 133.1 ($\text{C}_{\text{arom.}-2'}+\text{C}_{\text{arom.}-6'}$), 130.7 ($\text{C}_{\text{arom.}-1'}$), 127.7 ($\text{C}_{\text{coulm.}-5}+\text{C}_{\text{coulm.}-7}$), 123.6 ($\text{C}_{\text{coulm.}-6}$), 119.5 ($\text{C}_{\text{coulm.}-4}+\text{C}_{\text{arom.}-3'}+\text{C}_{\text{arom.}-5'}$), 118.1 ($\text{C}_{\text{coulm.}-4a}$), 116.4 ($\text{C}_{\text{coulm.}-8}$), 113.9 ($\text{C}_{\text{coulm.}-3}$), 99.4 ($\text{C}_{\text{pyrimid.}-5}$), 25.0 (CONHMe). Anal. Calcd for ($\text{C}_{22}\text{H}_{19}\text{N}_9\text{O}_4$) (473.44): C, 55.81; H, 4.05; N, 26.63. Found: C, 54.61; H, 3.93; N, 26.4 %.

3-(5,7-Diamino[1,2,4]triazolo[4,3-*c*]pyrimidin-3-yl)-2H-chromene-2-one (**18**)

A mixture of 2,6-diamino-4-chloropyrimidine (**16**) (200 mg, 1.39 mmol) and **5** (283 mg, 1.39 mmol) was irradiated under microwave at 180 °C for 25 min. The crude mixture was heated under reflux in glacial HOAc (10 ml) for 4 h. After cooling, water was added the precipitate was filtered, washed with cold EtOH and dried. Recrystallization from EtOH afforded **18** (176 mg, 43 %), m.p. 210 °C (dec.), R_f =0.17. ^1H NMR (DMSO- d_6): δ 9.72 (br s., 2H, NH_2), 8.04 (s, 1H, $\text{H}_{\text{coulm.}-4}$), 7.73 (dd, 1H, $J_{5,6}=8.3$ Hz, $J_{5,7}=2.3$ Hz, $\text{H}_{\text{coulm.}-5}$), 7.59–7.32 (m, 3H, $\text{H}_{\text{coulm.}-6}+\text{H}_{\text{coulm.}-7}+\text{H}_{\text{coulm.}-8}$), 6.32 (br s., 2H, NH_2), 5.71 (s, 1H, $\text{H}_{\text{pyrimid.}-5}$). ^{13}C NMR (DMSO- d_6): δ 168.2 ($\text{C}_{\text{pyrimid.}-2'}$), 164.9 ($\text{C}_{\text{pyrimid.}-6'}$), 159.3 ($\text{C}_{\text{coulm.}-2}$), 152.8 ($\text{C}_{\text{coulm.}-8a}$), 149.3 ($\text{C}_{\text{pyrimid.}-4'}$), 146.1 ($\text{C}_{\text{triazol-3}}$), 144.7 ($\text{C}_{\text{coulm.}-4}$), 130.4 ($\text{C}_{\text{coulm.}-3}$), 128.3 ($\text{C}_{\text{coulm.}-7}$), 126.5 ($\text{C}_{\text{coulm.}-5}$), 124.7 ($\text{C}_{\text{coulm.}-6}$), 120.4 ($\text{C}_{\text{coulm.}-4a}$), 115.8 ($\text{C}_{\text{coulm.}-8}$), 93.7 ($\text{C}_{\text{pyrimid.}-5}$). Anal. Calcd for $\text{C}_{14}\text{H}_{10}\text{N}_6\text{O}_2$ (294.27): C, 57.14; H, 3.33; N, 28.56. Found: C, 56.83; H, 3.33; N, 28.29 %.

Methyl 3-oxo-3H-benzochromene-2-carboxylate (**22**)

A mixture of 2-hydroxynaphthaldehyde (**19**) (500 mg, 2.91 mmol) and dimethylmalonate (**20**) (385 mg, 2.91 mmol) in the presence of piperidine (2.0 mL) was heated under reflux for 1 h. After cooling, the reaction mixture was acidified with HCl and cooled. The precipitate was filtered, washed with small amount of cold water, the product was dried and recrystallized from MeOH to give **22** (593 mg, 85 %), m.p. 148 °C, R_f =0.69. ^1H NMR (DMSO- d_6): δ 9.45 (s, 1H, H-4), 8.63 (dd, 1H, $J_{5,6}=7.6$ Hz, $J_{5,7}=1.5$ Hz, H-5), 8.36 (dd, 1H, $J_{7,8}=7.6$ Hz, $J_{6,8}=1.5$ Hz, H-8), 8.11 (m, 1H, H-6), 7.82 (m, 1H, H-7), 7.70 (d, 1H, $J_{9,10}=8.0$ Hz, H-9), 7.64 (d, 1H, $J_{9,10}=8.0$ Hz, H-10), 3.90 (s, 3H, CO_2Me). ^{13}C NMR (DMSO- d_6): δ 163.9 (CO_2Me), 155.8 (C-2), 145.2 (C-10a), 136.7 (C-4), 129.6 (C-9+C-5a), 129.5 (C-8), 127.0 (C-6), 122.8 (C-7+C-8a), 120.3 (C-5), 117.0 (C-4a), 115.3 (C-10), 115.3 (C-4a), 100.0 (C-3), 53.0 (OMe). Anal. Calcd for $\text{C}_{15}\text{H}_{10}\text{O}_4$ (254.24): C, 70.86; H, 3.96. Found: C, 70.66; H, 3.89 %.

Ethyl 3-imino-3H-benzochromene-2-carboxylate (**23**)

To a stirred solution of **19** (361 mg, 2.01 mmol) in EtOH (10 ml) were added ethyl cyanoacetate (**21**) (227 mg, 2.01 mmol), NH_4OAc (200 mg) and glacial HOAc (1.0 ml). The mixture was heated under reflux for 1 h, and then poured onto crushed ice. The precipitate was filtered, dried and recrystallized from MeOH to give **23** (456 mg, 81 %), m.p. 202–205 °C (dec.), R_f =0.32. ^1H NMR (DMSO- d_6): δ 9.29 (s, 1H, H-4), 9.07 (s, 1H, NH), 8.66 (d, 1H, J =8.5 Hz, H-5), 8.34 (d, 1H, J =9.1 Hz, H-8), 8.12 (d, 1H, $J_{6,7}=8.1$ Hz, H-6), 7.82 (m, 1H, H-7), 7.81 (d, 1H, $J_{9,10}=8.0$ Hz, H-9), 7.68 (m, 2H, H-9+H-10), 4.23 (q, 2H, J =7.8 Hz, CH_2CH_3), 1.27 (t, 3H, CH_2CH_3). ^{13}C NMR (DMSO- d_6): δ 167.1 (CO_2Et), 165.0 (C=NH), 154.3 (C-10a), 135.6 (C-8), 130.6 (C-5a), 129.5, 129.4 (C-4+C-6), 127.1 (C-9), 123.1 (C-5), 122.1 (C-7+C-8a), 118.8 (C-4a), 117.0 (C-10), 112.6 (C-3), 60.5 (CH_2), 14.9 (CH_3). Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{NO}_3$ (267.28): C, 71.90; H, 4.91; N, 5.24. Found: C, 71.74; H, 7.80; N, 5.07 %.

3-Oxo-3H-benzochromene-2-carbohydrazide (**24**)

A solution of **22** (100 mg, 0.37 mmol) in EtOH (7 mL) and hydrazine hydrate (2 mL) was heated under reflux for 3 h. After cooling, the solution was poured into crushed ice with stirring. The separated solid was filtered, washed with water, dried and recrystallized from EtOH to give **24** (75 mg, 80 %), as a yellow solid, m. p. 260–263 °C (Lit. [25], m.p. 260–262 °C),

N'-(2,6-Diamino-5-((4-bromophenyl)diazenyl)pyrimidin-4-yl)-3-oxo-3H-benzochromene-2-carbohydrazide (**25**)

To a stirred solution of **24** (67 mg, 0.30 mmol) in DMF (5 mL) was added the azopyrimidine **7** (98 mg, 0.30 mmol) and the mixture was heated at 100 °C for 3 h. After cooling, the residue was evaporated to dryness and the residue was purified on a short SiO₂ column (5.0 g). Elution, in gradient, with MeOH (0–10 %) and CHCl₃ as eluent afforded **25** (95 mg, 58 %), m.p. 279–282 °C (dec.). ¹H NMR (DMSO-*d*₆): δ 10.02 (br s., 2H, NH₂), 9.26 (br s., 1H, NH), 8.78 (d, 1H, *J*_{9,10}=8.8 Hz, H_{coum.}-10), 8.00 (d, 1H, *J*_{7,8}=8.0 Hz, H_{coum.}-8), 7.89 (d, 1H, *J*_{5,6}=8.0 Hz, H-5), 7.82 (d, 1H, *J*_{9,10}=8.8 Hz, H_{coum.}-9), 7.55 (d, 2H, *J*_{2',3'}=8.0 Hz, H_{arom.}-2'+H_{arom.}-6'), 7.50 (m, 2H, H_{coum.}-6+H_{coum.}-7), 7.38 (d, 2H, *J*_{5',6'}=8.0 Hz, H_{arom.}-3'+H_{arom.}-5'), 7.25 (s, 1H, H_{coum.}-4), 6.97 (d, 2H, *J*=5.0 Hz, NH₂). ¹³C NMR (DMSO-*d*₆): δ 164.9 (CONH), 162.1 (C_{pyrimid.}-4), 161.2 (C_{pyrimid.}-2), 157.8 (C_{coum.2}=O), 156.1 (C_{pyrimid.}-6), 137.2 (C_{coum.}-3), 134.5 (C-8a), 132.8 (C-4a), 132.3, 131.0, 130.3, 129.4, 128.4, 128.0, 127.3, 127.0, 125.2, 124.3, 123.1 (C_{coum.}+C_{arom.}), 120.1 (C_{coum.}-4), 102.1 (C_{pyrimid.}-5). Anal. Calcd for C₂₄H₁₇BrN₈O₃ (545.35): C, 52.86; H, 3.14; N, 20.55. Found: C, 52.66; H, 3.03; N, 20.36 %.

Results and Discussion

Chemistry

Treatment of methyl coumarin-3-carboxylate **4** with hydrazine hydrate afforded the carbohydrazide analog **5**. 4-Chloro-azopyrimidine derivatives **6–10** have been selected as starting materials for the synthesis of the target compounds **11–15**, since the electron withdrawing group, azo residue would facilitated replacement of the chloro group at C-4 by


the carbohydrazide scaffolds. Thus, treatment of **6–10** with **5** in hot DMF for 4–5 h furnished, after chromatographic purification, **11–15** in 22–62 % yield (Scheme 1).

The structures of **11–15** were determined by ¹H, ¹³C and 2D NMR spectroscopy. In the ¹H NMR spectra, H-4 of the coumarin ring were resonated as singlets at δ=9.01–9.00 ppm, meanwhile H-5 - H-8 together with the aromatic protons conjugated pyrimidine ring and other substituents were fully analysed. In the ¹³C NMR spectra of **11–15**, the carbonyl carbon atom (C₉=O) appeared at the regions δ=165.2–163.3 ppm, while C-4 of the pyrimidine ring resonated at δ=163.3–161.3 ppm. The resonances at the regions δ=161.7–159.4, 157.0–155.7 and 103–99.4 ppm were assigned to carbon atoms 2, 6 and 5 of the pyrimidine backbone, respectively, whereas the signals at the regions δ=159.1–158.5 and 155.7–151 ppm were attributed to carbon atoms 2, and 8a of the coumarin scaffold, respectively. Other coumarin and aromatic carbon atoms as well as the substituents were fully analysed (*c.f.* Experimental section).


Next, treatment of **16** with carbohydrazide **5** under microwave irradiation (400 Watt, 180 °C) for 20 min afforded the crud product, 4-coumarinyl-pyrimidine carbohydrazide derivative **17** which has been used directly, without purification. Boiling of the crude **17** with glacial HOAc for 4 h furnished, after chromatographic purification, the cyclized triazolo analog **18** (43 %) (Scheme 2).

The structure of **18** was determined from its ¹H and ¹³C NMR spectra. In the ¹H NMR spectrum, H-4 of the coumarin ring appeared as a singlet at δ=8.04 ppm, while H-5 resonated as a doublet of doublets at δ=7.73 ppm (*J*_{5,6}=8.3 Hz, *J*_{5,7}=2.3 Hz). H-5 of the pyrimidine backbone appeared as a singlet δ=7.71 ppm, whereas the other coumarin protons were analysed. The ¹³C NMR spectrum showed four signals at the lower fields, δ=168.2, 164.9, 159.3 and 152.8 ppm, were assigned to C-2', C-6' of the pyrimidine ring and C-2, 8a of coumarin moiety, respectively. The resonances at δ=149.3, 146.1 and 144.7 ppm were attributed C-4' of the pyrimidine

Scheme 1 Conditions and reagents: (i) NH₂NH₂·H₂O, EtOH, reflux, 6 h; (ii) DMF, 90–100 °C, 4–5 h



Scheme 2 Synthesis of 3-(5,7-diamino[1,2,4-]triazolo[4,3-*c*]pyrimidin-3-yl)-2*H*-chromene-2-one (**18**)



ring, C-3 of the triazole and C-4 of coumarin moiety, respectively. C-5 of the pyrimidine ring resonated at $\delta=93.7$ ppm.

Our work was modified by selecting **19** as a precursor for the synthesis of new benzocoumarinyl-pyrimidine, aiming to examine its fluorescence property in comparison to the coumarin analogs **11–15**. Thus, treatment of **19** with dimethylmalonate **20** in the presence of piperidine afforded, after purification, the benzocoumarin 3-methyl ester **22** (85 %). Alternatively, treatment of **19** with ethyl cyanoacetate **21** in the presence of NH_4OAc and HOAc in boiling EtOH furnished the 3-iminoethyl benzocoumarin ester **23** (81 %) which has been structurally confirmed by the ^1H and ^{13}C NMR spectra. Treatment of **22** with hydrazine hydrate afforded the carbohydrazide derivative **24** in 30 % yield. Compound **25** was obtained in 58 % yield by condensation of **24** with the 4-bromo-azopyrimidine derivative **7** in hot DMF (Scheme 3).

The structures of **22–25** were assigned by their ^1H and ^{13}C NMR spectra, where the protons and carbon atoms of the coumarin moiety showed almost similar pattern. The singlets at $\delta=9.24$, 9.29, 8.87 and 7.25 ppm were assigned for H-4 of the coumarin ring, respectively. The other aromatic, NH and NH_2 protons were fully analysed (*c.f.* Experimental section). In the ^{13}C NMR spectrum of **23**, CO_2Et and $\text{C}=\text{NH}$ carbon

atoms resonated at $\delta=167.1$ and 165.0 ppm, respectively. Further, the resonances at $\delta=164.9$ and 162.1 ppm were assigned for CONH of coumarin and C-4' of pyrimidine backbone, respectively. The rest of benzocoumarin and aromatic carbon atoms were fully assigned (*cf.* Experimental section).

All the synthesized compounds were further analysed by the HSQC [26] and HMBC [27] NMR spectroscopy.

UV–Vis Absorption and Fluorescence Spectra

The UV-visible absorption and fluorescence spectra of the synthesized coumarin analogs **4**, **5**, **11–15**, **18** and **22–25** in MeOH were obtained and are presented in Table 1. The absorption and fluorescence maxima of some of the synthesized analogs showed good bathochromic shifts with wavelength of maximum absorption in the UV or visible region ($\lambda_{\text{ex.}}$) (315–405 nm). Wavelengths of maximum emission for the compounds ($\lambda_{\text{em.}}$) (322–483 nm) were observed in MeOH at room temperature. Although, compounds **4**, **12**, **22** and **23** showed low quantum yield values (Φ_{F}) of 0.02, 0.01 and 0.15 nm, respectively, but exhibited fluorescence spectral properties with Stoke's shift of 89, 87, 72 and 85 nm, respectively (Fig. 2). However, **22**

Scheme 3 Conditions and reagents: (i) **20**, piperidine, EtOH, reflux, 1 h; (ii) **21**, NH_4OAc , HOAc , EtOH, reflux, 1 h; (iii) $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$, EtOH, reflux, 3 h; (iv) **7**, DMF, 100 °C, 3 h




Table 1 UV-vis, fluorescence data, and quantum yields (Φ_F) of compounds some coumarin and benzocoumarin analogs

Compd.	λ_{ex} [nm]	$\log \epsilon$ [mM] ⁻¹ [cm] ⁻¹	λ_{em} [nm]	Φ_F	Stoke's shift
4	331	4.04	331	0.02	89
5	315	4.03	322	<0.01	7
11	365	4.41	385	<0.01	20
12	365	4.78	483	0.01	87
13	405	4.43	405	<0.01-	10
14	[355]	[4.61]	355	<0.01-	8
15	[355]	[4.42]	360	<0.01-	5
18	293	4.56	300	<0.01-	7
22	371	3.90	443	0.98	72
23	363	3.81	448	0.15	85
24	360	4.09	373	0.01	13
25	379	4.27	391	0.12	12

Annotations: Ref.: Rh6G in EtOH (Φ_F : 0.95) at room temperature; Equipment and software: Perkin Elmer LS 50, Cary UV/vis 50; SpekWin 1.71.3, FL WinLab. []: shoulder


showed a very exciting fluorescence result in MeOH with a quantum yield (Φ_F =0.98) with stoke's shift of 72 nm (Fig. 2), which characterized by a brightly intense blue emission light at λ_{em} =443 nm, in comparison for those of Rh6G (Φ_F =0.95). These result encouraged us to proceed further in our current fluorescence imaging (FI) for the visualisation of cells and tissues both in vitro and in vivo as well as in coronary angiographic applications, with full biochemical study of **22**.

In conclusion, the replacement of 3-ester group in the coumarin of benzocoumarin backbone by hydrazide group (e.g., **5** and **24**) or hydrazide-pyrimidine moiety (e.g., **11–15** and **25**) would reduced the fluorescence property unless there is an electron donating group at the aromatic ring of the coumarin moiety. Such argument is in agreement with Mahadevan et al. results [28].

Biological Imaging Application

Organic fluorescent dyes are routinely used as contrast agents in immunohistochemical staining protocols. Owing to their low molecular weight, high quantum yield, relatively low cost and the existence of well-developed conjugation protocols, they represent an ideal optical label for molecular imaging probes [29]. Additionally, fluorescent imaging is a relatively recent imaging method and thus still developing in many ways, and typically performed by exogenous delivery of a fluorescent probe (e.g., an organic dye) that interacts with the target or by direct imaging of an endogenously expressed fluorescent protein. Signal generation is achieved by exciting the fluorescent probe or protein at a given light wavelength (λ) and detecting light emission at another λ with a charge-coupled device camera built into either a planar (fluorescence reflectance imaging) or a tomographic (fluorescence molecular tomography) imaging system [30, 31]. Most of the above imaging fluorescence agents exhibited several side effects in vivo (especially by human treatment), including the allergy, toxicity, solubility and radioactivity. Update ¹²⁵I and sodium diatrizoate (sodium 3-(acetylamino)-2,4,6-triiodobenzoate) are used in the coronary arterials angiography [32].

Our small molecule, the benzocoumarin ester **22** has been selected for a primary fluorescence imaging application by using an experimental animal model (rabbit, weight~3.2 kg) induced intravenously by **22** with a concentration of 10^{-3} M in a mixture of MeOH-H₂O (1:9 v:v) solution, following Johnston et al. method [33]. The X-ray contrast media for coronary angiography showed some encouraging peripheral vascular angiographic imaging pictures. The application is under further investigation to overcome the solubility problem of **22** in vivo by introducing a more polar group such hydroxyl group.

**Fig. 2** UV-visible and fluorescence spectra of compounds **12** and **22** in MeOH

Conclusion

A new series of coumarin and 5,6-benzocoumarin substituted pyrimidine derivatives **11–15** and **22–25** were synthesized from treatment of coumarin and benzocoumarin 3-carbohydrazide with 4-chloro-5-azoarylpyrimidines analogs. Further, the coumarin-1,2,4-triazolo-pyrimidine analog **18** was prepared from 4-chloropyrimidine analog **16** with coumarin 3-carbohydrazide **5** under MWI condition, aiming to develop new imaging fluorescence agents. The new analogs were investigated spectrophotometrically in MeOH for their fluorescence properties, which showed emission in the region between 331 and 495 nm. Interestingly, methyl benzocoumarin 3-carboxylate (**22**) showed a higher quantum yield ($\Phi_F=0.98$) in comparison for those of Rhodamine 6G as standard ($\Phi_F=0.95$). The primary biological fluorescence imaging application of **22** in an experimental animal model showed some peripheral vascular angiographic imaging pictures by using a fluorescence emission with some pharmacokinetic studies.

Acknowledgments We thank Professor Surg. Abdul-Bari A. Alfaris of College of Veterinary Medicine, Department of Surgery, University of Basrah (Iraq) for the primary biological imaging fluorescence experiments. Miss A. Friemel of Chemistry Department, Konstanz University, Germany is highly acknowledged for the 2D-NMR experiments.

References

- Ito C, Itoigawa M, Katsuno S, Omura M, Tokuda H, Nishino H, Furukawa H (2000) Chemical constituents of *Clausena excavata*: isolation and structure elucidation of novel furanone-coumarins with inhibitory effects for tumor-promotion. *J Nat Prod* 63:1218–1224
- Abyshev AZ, D'Yachuk GI, Semenov EV, Pukhov MP (1993) Synthesis and biological activity of derivatives of benzopyran-2-one. *Pharm Chem J* 27:766–770
- Kontogiorgis CA, Hadjipavlou-Litina DJ (2004) Synthesis and biological evaluation of # novel coumarin derivatives with a 7-azomethine linkage. *Bioorg Med Chem Lett* 14:611–614
- Abyshev AZ, Alekseev AT, Platonov VG, Byrkin IA (1996) Synthesis and antiviral activity of benzopyran-2-one derivatives. *Pharm Chem J* 30:441–444
- Teran C, Santana L, Uriarte E, Fall Y, Lena L, Unelius L, Tolf B-R (1998) Phenyl- piperazine derivatives with strong affinity for 5HT_{1A}, D_{2A} and D₃ receptors. *Bioorg Med Chem Lett* 8:3567–3570
- Al-Soud YA, Al-Sauodoni HH, Amajaour HAS, Salih KSM, Mubarak MS, Al-Masoudi NA, Jaber IH (2008) Synthesis, characterization and anti-HIV activities of new coumarin derivatives. *Z Naturforsch* 63b:83–89
- Kayser O, Kolodziej H (1997) Antibacterial activity of extracts and constituents of *Pelargonium sidoides* and *Pelargonium reniforme*. *Planta Med* 63:508–510
- Wang CJ, Hsieh YJ, Chu CY, Lin YL, Tseng TH (2002) Inhibition of cell cycle progression in human leukemia HL-60 cells by esculetin. *Cancer Lett* 183:163–186
- Itoigawa M, Ito C, Tan T-WT, Kuchide M, Tokuda H, Nishino H, Furukawa H (2000) Cancer chemopreventive agents, 4-phenylcoumarins from *Calophyllum inophyllum*. *Cancer Lett* 169:15–19
- Cravotto G, Nano GM, Palmisano G, Tagliapietra S (2001) An asymmetric approach to coumarin anticoagulants via hetero-Diels-Alder cycloaddition. *Tetrahedron: Asym* 12:707–709
- Al-Soud YA, Al-Masoudi IA, Saeed B, Beifuß U, Al-Masoudi NA (2006) Synthesis of new 1H-1,2,4-triazolylcoumarins and their antitumor and anti-HIV activities. *Chem Heterocycl Comp* 42:667–676
- Kirkiacharian S, Thuy DT, Sicsic S, Bakhchinian R, Kurkjian R, Tonnaire R (2002) Structure-activity relationships of some 3-substituted-4-hydroxycoumarins as HIV-1 protease inhibitors. *Farmaco* 57:703–708
- Yamaguchi T, Fukuda T, Ishibashi F, Iwao M (2006) The first total synthesis of lamellarin α 20-sulfate, a selective inhibitor of HIV-1 integrase. *Tetrahedron Lett* 47:3755–3757
- Pochet L, Frederic R, Masereel B (2004) Coumarin and isocoumarin as serin protease inhibitors. *Curr Pharm Design* 10:3781–3796
- Fan G-J, Mar W, Park MK, Choi EW, Kim K, Kim S (2001) A novel class of inhibitors for steroids 5 α -reductase: synthesis and evaluation of umbelliferone derivatives. *Bioorg Med Chem Lett* 11:2361–2363
- Murakami A, Gao G, Omura M, Yano M, Ito C, Furukawa H, Takahashi D, Koshimizu K, Ohigashi H (2000) 1,1-Dimethylallylcoumarins potently suppress both lipopolysaccharide and interferon- γ -induced nitric oxide generation in mouse macrophage RAW 264.7 cells. *Bioorg Med Chem Lett* 10:59–62
- Jerris JP, Smith AB (1981) Synthesis of geiparvarin: a novel anti-tumor agent. *J Org Chem* 46:577–585
- Christie RM, Lui C-H (2000) Studies of fluorescent dyes: part 2. an investigation of the synthesis and electronic spectral properties of substituted 3-(2-benzimidazolyl) coumarins. *Dyes Pigments* 47:79–89
- Zhao Y, Zheng Q, Dakin K, Xu K, Martinez ML, Li W-H (2004) New caged coumarin fluorophores with extraordinary uncaging cross sections suitable for biological imaging applications. *J Am Chem Soc* 126:4653–4663
- Jung HS, Kwon PS, Lee JW, Kim JI, Hong CS, Kim JW, Yan S, Lee JY, Lee JH, Joo J, Kim JS (2009) Coumarin-derived Cu²⁺-selective fluorescence sensor: synthesis, mechanisms, and applications in living cells. *J Am Chem Soc* 131:2008–2012
- Rajesh G, Kumar HK, Naik HSB, Mahadevan KM (2011) Synthesis of new benzocoumarinyl oxadiazolyls as strong blue-green fluorescent brighteners. *South Afr J Chem* 64:88–94
- Khalfan H, Abuknesha R, Rand-Weaver M, Price RG, Robinson D (1986) Aminomethyl coumarin acetic acid: a new fluorescence labelling agent proteins. *Histochem J* 19:497–509
- Wang C, Popescu DC, Wu C, Zhu J, Macklin W, Wang Y (2010) In situ fluorescence imaging of myelination. *J Histochem Cytochem* 58:611–621
- Wu Q, Merchant FA, Castleman KR (2008) Microscope image processing. Academic, New York
- Bogdal D (1998) Coumarins: fast synthesis by Knoevenagel condensation under microwave irradiation. *J Chem Res (S)*: 468–469
- Chiyomi M, Toshinobu M, Yasuko K, Kenichiro T, Hideyuki Y, Hitoshi N, Masatoshi Y, Akira T (2005) Improvement of fluorescence characteristics of coumarins: syntheses and fluorescence properties of 6-methoxycoumarin and benzocoumarin derivatives as novel fluorophores emitting in the longer wavelength region and their application to analytical reagents. *Chem Pharm Bull* 53:750–758

27. Davis AL, Keeler J, Laue ED, Moskau D (1992) Experiments for recording pure- absorption heteronuclear correlation spectra using pulsed field gradients. *J Magn Reson* 98:207–216
28. Willker W, Leibfritz D, Kerssebaum R, Bermel W (1993) Gradient selection in inverse heteronuclear correlation spectroscopy. *Magn Reson Chem* 31:287–292
29. Harishkumar HN, Mahadevan KM, Masagalli JN, Chandrashekarappa KKH (2012) Synthesis and fluorescence study of phenylcoumarin/cyanophenylbenzocoumarin-3- carboxylates. *Organ Commun* 5:196–208
30. Hellebust A, Richards-Kotum R (2012) Advances in molecular imaging: targeted optical contrast agents for cancer diagnostics. *Nanomedicine (Lond)* 7:429–445
31. Ntziachristos V (2006) Fluorescence molecular imaging. *Annu Rev Biomed Eng* 8:1–33
32. Rao J, Dragulescu-Andrasi A, Yao H (2007) Fluorescence imaging in vivo: recent advances. *Curr Opin Biotech* 18:17–25
33. Hoppe JO, Archer S (1960) X-ray contrast media for cardiovascular angiography. *Angiology* 11:244–254