

Article

OF WOLLONGON

### Direct Synthesis of Cyclopropanes from *gem*-Dialkyl Groups through Double C–H Activation

Antonin Clemenceau, Pierre Thesmar, Maxime Gicquel, Alexandre Le Flohic, and Olivier Baudoin

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.0c05887 • Publication Date (Web): 12 Aug 2020

Downloaded from pubs.acs.org on August 12, 2020

### **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# Direct Synthesis of Cyclopropanes from *gem*-Dialkyl Groups through Double C–H Activation

Antonin Clemenceau,<sup>†</sup> Pierre Thesmar,<sup>†</sup> Maxime Gicquel,<sup>‡</sup> Alexandre Le Flohic,<sup>‡</sup> and Olivier Baudoin<sup>†\*</sup>

<sup>†</sup>University of Basel, Department of Chemistry, St. Johanns-Ring 19, CH-4056 Basel, Switzerland.

<sup>‡</sup>Oril Industrie, 13 rue Auguste Desgenétais, CS60125, 76210 Bolbec, France.

Supporting Information Placeholder

**ABSTRACT:** Cyclopropanes are important structural motifs found in numerous bioactive molecules, and a number of methods are available for their synthesis. However, one of the simplest cyclopropanation reactions involving the intramolecular coupling of two C-H bonds on gem-dialkyl groups has remained an elusive transformation. We demonstrate herein that this reaction is accessible using aryl bromide or triflate precursors and the 1,4-Pd shift mechanism. The use of pivalate as the base was found to be crucial to divert the mechanistic pathway toward the cyclopropane instead of the previously obtained benzocyclobutene product. Stoichiometric mechanistic studies allowed the identification of aryl- and alkylpalladium pivalates, which are in equilibrium via a five-membered palladacycle. With pivalate, a second  $C(sp^3)$ -H activation leading to the four-membered palladacycle intermediate and the cyclopropane product is favored. A catalytic reaction was developed and showed a broad scope for the generation of diverse arylcyclopropanes, including valuable bicyclo[3.1.0] systems. This method was applied to a concise synthesis of lemborexant, a recently approved anti-insomnia drug.

### INTRODUCTION

Cyclopropanes are widespread structural motifs in natural products and bioactive molecules (see Figure 1 for examples).<sup>1</sup> Indeed, the cyclopropane ring can improve the pharmacological properties of active substances by reducing their lipophilicity, increasing their metabolic stability, or modifying their acidobasicity.<sup>2</sup> Although a great number of methods have been developed to synthesize cyclopropanes,<sup>3</sup> the cycloaddition of alkenes and carbenes or carbenoids largely prevails for applications to complex molecules.<sup>4</sup> In this context, one of the simplest ways to construct cyclopropanes, through C–C bond formation from two geminal alkyl groups (Scheme 1a), has remained an elusive transformation to this date. In 2006, Yu and coworkers reported the synthesis of cyclopropanes via a two-step sequence involving palladium(II)-catalyzed, oxazoline-directed  $C(sp^3)$ –H diiodination of *gem*dimethyl groups followed by radical cyclization. (Scheme 1b).<sup>5</sup> Besides, cyclopropanes have been generated through domino reactions including olefin carbopalladation and  $C(sp^3)$ –H activation, leading to putative 4-membered palladacyclic intermediates (Scheme 1c).<sup>6,7</sup>



**Figure 1.** Examples of drugs or drug candidates containing an arylcyclopropane motif relevant to this study.

The current work reports the first single-step synthesis of cyclopropanes via coupling between two geminal methyl groups or one methyl and one activated methylene group (Scheme 1d). This reaction is based on the propension of the arylpalladium halide, generated in the initial oxidative addition step, to undergo 1,4-Pd shift to generate a  $\sigma$ -alkylpalladium intermediate.<sup>8</sup> Since seminal observations by Dyker,9 such alkylpalladium intermediates have been exploited in various transformations:  $\beta$ -H elimination to generate olefins,<sup>10</sup> trapping with boronic acids<sup>11</sup> and anilines,<sup>12</sup> or dearomatizing carbopalladation. intramolecular furnishing spiroannulation products.<sup>13</sup> Recently, we showed that such organopalladium species are also able to perform the cleavage of an activated  $C(sp^3)$ -H bond (i. e., benzylic and adjacent to a carbonyl group) to generate oxygen and nitrogen heterocycles.<sup>14</sup> The

extension of this mechanism to the formation of cyclopropanes is reported herein.

### **RESULTS AND DISCUSSION**

**Initial Work and Mechanistic Study.** The origin of this work stems from our initial studies on the generation of benzocyclobutenes (BCBs) by  $Pd^{0}$ -catalyzed  $C(sp^{3})$ -H activation.<sup>15</sup> Recent reinvestigations pointed to a spectacular influence of the base on the reaction outcome (Scheme 2).

Scheme 1. Synthesis of Cyclopropanes by Pd-Catalyzed C(sp<sup>3</sup>)–H Activation

a) elusive transformation

Me

 $\begin{array}{c} H \\ H \\ H \\ H \\ H^{3} \\ H^{2} \\ H^{2}$ 

Pd<sup>0</sup> cat.

b) cyclopropanes via double C(sp3)-H iodination and radical cyclization



c) cyclopropanes via alkene carbopalladation and  $C(\ensuremath{\mathsf{sp}}^3)\ensuremath{-}\ensuremath{\mathsf{H}}$  activation





**X** = **Br**, **OTf**  $R^2$ ,  $R^3$  = H or CH<sub>2</sub>EWG (EWG = electron-withdrawing group)

As initially reported, using potassium carbonate as the base in the intramolecular C–H arylation of substrate **1a** 

under the shown conditions provided BCB **2a** as the sole observable C–H activation product (entry 1), together with the proto-dehalogenated byproduct (not shown). Replacing carbonate with pivalate led to a complete change of selectivity, with cyclopropane **3a** becoming the sole C–H activation product (entry 2). The ligand also had a noticeable effect on the selectivity, with the bulky P(*t*-Bu)<sub>3</sub> favoring the BCB product, as previously reported (entry 3).<sup>15</sup>

# Scheme 2. Influence of the Base on the Formation of Benzocyclobutene vs. Cyclopropane



 $^{a}$  <sup>1</sup>H NMR yield using CH<sub>2</sub>Br<sub>2</sub> as internal reference.  $^{b}$  Using Pd<sub>2</sub>dba<sub>3</sub> (5 mol%)/P(*t*-Bu)<sub>3</sub> (20 mol%) instead of Pd(PPh<sub>3</sub>)<sub>4</sub>.

Following these initial observations, and inspired from a study by Martin and co-workers,<sup>16</sup> we decided to reinvestigate the reaction mechanism by following each step through stoichiometric experiments (see the Supporting Information for details). A tentative mechanism, represented in qualitative energy profile format, is displayed in Figure 2. The oxidative addition of prototypical aryl bromide **1b** to Pd(PPh<sub>3</sub>)<sub>4</sub> gives rise to complex **A**, which undergoes bromide substitution with pivalate, producing complex **B**<sup>1</sup>.



Figure 2. Qualitative energy profile of the reaction of substrate 1b based on experimental observations. The relative stabilities of  $B^2$  and  $D^1$ ,  $C^1$  and  $E^1$ , and 2b and 3b were calculated by DFT (see the Supporting Information for details). L =

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29 30 31

60

 $PPh_3$ . TS = transition state; OA = oxidative addition; Sub = substitution; CH = C-H activation; PROT = protonation; RE = reductive elimination.

The latter is in equilibrium with complex  $B^2$ , containing one phosphine ligand and pivalate coordinated in the  $\kappa^2$  mode,<sup>17c</sup> which undergoes basemediated  $C(sp^3)$ -H activation via the concerted metalation-deprotonation mechanism<sup>15,17</sup> to give fivemembered palladacycle  $C^1$ . Reductive elimination from  $C^1$  provides the BCB product **2b**. Alternatively, protonation of  $C^1$  with pivalic acid<sup>15,17e</sup> furnishes alkylpalladium complex  $D^1$ . The sequence  $B^1 \rightarrow C^1 \rightarrow$  $D^1$  results in an overall 1,4-Pd shift.<sup>8,14</sup> A second baseinduced  $C(sp^3)$ -H activation from  $D^1$  leads to the rare four-membered palladacycle  $E^{1,6,7,18}$  already suggested by Martin,<sup>16</sup> which reductively eliminates to generate cyclopropane **3b**.

The oxidative addition of the bulky aryl bromide **1b** was followed by <sup>1</sup>H and <sup>31</sup>P NMR and was found to be slow using PPh<sub>3</sub> as the ligand, with only 26% of complex **A** formed at 80 °C after 16 h, and 72% at 120 °C together with 20% remaining starting material (Scheme 3a). Complex **A** was independently synthesized in 88% yield from **1b** and Pd(PPh<sub>3</sub>)<sub>4</sub> (toluene, 120 °C), and its three-dimensional structure was confirmed by Xray diffraction analysis (Scheme 3). Consistent with results displayed in Scheme 2, heating **A** to 100 °C in the presence of K<sub>2</sub>CO<sub>3</sub> furnished BCB **2b** in 65% yield (Scheme 3b). The substitution step was then analyzed by

mixing complex A with KOPiv (Scheme 3b). This step was relatively easy, with  $\sigma$ -arylpalladium pivalate **B**<sup>1</sup> being slowly formed at room temperature (25% after 16 h, not shown) and in 82% yield at 60 °C. Raising the temperature to 80 and 100 °C led to the formation of cyclopropane **3b** at the expense of complex  $B^1$ . This result shows that the formation of the cyclopropane product is faster than the oxidative addition, and therefore that the latter is the slowest step of the reaction. The reaction of the known  $\sigma$ -alkylpalladium complex (PhCMe<sub>2</sub>CH<sub>2</sub>)Pd(COD)Cl  $D^2$  (COD = cyclooctadiene)<sup>19</sup> in the presence of KOPiv and PPh<sub>3</sub> in toluene at room temperature led to alkylpalladium pivalate  $D^1$  in 88% yield (Scheme S1, see Figure 2 or Scheme 3 for the structure of  $D^1$ ). In contrast, at 60 °C arylpalladium pivalate  $B^1$  was formed from  $D^2$ , thus showing that  $\mathbf{B}^1$  is more stable than  $\mathbf{D}^1$ . The thermal decomposition of  $B^1$  was next analyzed at various temperatures (Scheme 3c). At 80 °C, both D<sup>1</sup> and cyclopropane 3b were formed, and the proportion of cyclopropane markedly increased at 100 °C. The reactivity of alkylpalladium pivalate  $D^1$  was also analyzed (Scheme 3d). At 80 °C, a new complex, which was assigned to arylpalladium species  $B^2$  containing one PPh<sub>3</sub> ligand and the  $\kappa^2$ -coordinated pivalate, was formed together with degradation products.

### Scheme 3. Stepwise Study of the Reaction Mechanism through Stoichiometric Experiments<sup>a</sup>



<sup>*a*</sup> <sup>1</sup>H NMR yields using CH<sub>2</sub>Br<sub>2</sub> as internal reference. All reactions were performed in C<sub>6</sub>D<sub>6</sub> for 16 h. <sup>*b*</sup> Thermal ellipsoids at 50% probability, H atoms omitted for clarity. COD = 1,5-cyclooctadiene.

Adding PPh<sub>3</sub> (1 equiv) led to a cleaner transformation, with both mono- and bis-ligated complexes  $B^1-B^2$  and cyclopropane 3b being formed in comparable amounts. Heating this mixture to 100 °C furnished a higher proportion of  $\mathbf{B}^2$  and cyclopropane **3b**, and upon addition of 1 equiv of KOPiv the cyclopropane was formed exclusively. These results show that aryl- and alkylpivalate complexes  $B^1$ ,  $B^2$  and  $D^1$  are in equilibrium from 80 °C and are all competent intermediates en route to cyclopropane **3b**. Finally, since we did not observe the formation of five-membered palladacycles such as  $C^1$  (Figure 2) during the above studies, likely because they are higher-energy intermediates, we decided to prepare the known<sup>20</sup> fivemembered COD-stabilized palladacycle  $C^2$  and study its reactivity (Scheme 3e). In the presence of KOPiv, PivOH and PPh<sub>3</sub> at room temperature, complex  $D^1$  was formed as major product, together with minor amounts of  $B^1$ , thereby showing that the protonation with PivOH is kinetically favored on the aromatic vs. the alkyl ligand. Heating to 80 °C inversed the ratio between D<sup>1</sup> and  $\mathbf{B}^1$ , thus confirming the higher thermodynamic stability of  $B^1$ , and produced a minor amount of cyclopropane 3b. At 100 °C, only the latter was formed (61% yield). In contrast, heating palladacycle  $C^2$  to 80 °C in the presence of PPh<sub>3</sub> only provided BCB 2b in good yield.<sup>16</sup>

Taken together, these experimental observations allow to deduce the following relative stabilities for

organopalladium complexes:  $\mathbf{B}^1 > \mathbf{B}^2 > \mathbf{D}^1 > \mathbf{C}^1$ . In addition, DFT calculations indicated that the fourmembered palladacycle  $E^1$  is less stable than the fivemembered isomer  $C^1$  by ca. 7 kcal mol<sup>-1</sup>, hence establishing the stability order  $\mathbf{B}^1 > \mathbf{B}^2 > \mathbf{D}^1 > \mathbf{C}^1 > \mathbf{E}^1$ . Furthermore, cyclopropane **3b** was found to be more stable than BCB **2b** by 1.7 kcal mol<sup>-1</sup>. Figure 2 provides a qualitative overview of the reaction profile summarizing all experimental results and calculations. The oxidative addition is the highest kinetic barrier, which arises from the fact that the relatively electronically neutral PPh<sub>3</sub> was employed as ligand. In the presence of carbonate and the absence of pivalate, reductive elimination from five-membered palladacycle C<sup>1</sup> occurs preferentially to give BCB product 2b. In contrast. pivalate promotes the formation of cyclopropane **3b** via  $C(sp^3)$ -H activation of a second producing the four-membered methyl group palladacycle intermediate  $E^1$ . Both bicarbonate<sup>21</sup> and pivalic acid are able to open up palladacycle  $C^1$  to give alkylpalladium intermediates, and therefore the unique ability of pivalate to promote the second  $C(sp^3)$ -H activation event leading to cyclopropanation remains unclear, and will be the object of later studies.

**Reaction Optimization and Scope.** Taking advantage of the above mechanistic study, a catalytic version of this transformation was successfully developped (see the Supporting Information for details). Unsubstituted cycloproprane **3b** was obtained from both aryl bromide

1

2

3

4

5

6

7

8 9

10

11 12

13

14

15

16

17

18

19

20

21

22

23

24

25 26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

60

and triflate precursors in good yield using 10 mol%  $Pd(PPh_3)_4$  and 2 equiv potassium pivalate in toluene at 120 °C (Scheme 4a). However, this volatile product was best isolated from the triflate precursor in 72% yield. For functionalized cyclopropanes, a temperature of 140 °C was essential to ensure complete conversion. Besides, the addition of DMSO (5% vol.) was found to diminish the formation of the proto-dehalogenated side-product, presumably by increasing the solubility of pivalate.

With these standard conditions in hand, the versatility  $C(sp^3)-H$ this double activation-based of cyclopropanation was inspected. Substrates bearing an ester (3a, 3c), amide (3d), protected amine (3e) or protected alcohol (3i, 3m) on the quaternary benzylic carbon worked well under the optimized conditions (72-85% yields). Of note, the reaction producing cyclopropane **3a** was successfully scaled up to 1 mmol using o-xylene as the solvent for practical reasons. In addition, substrate **1f** bearing a substituted linear alkyl chain provided a mixture of the desired cyclopropane 3f (50% yield) and olefin 4f arising from methylene C-H activation and  $\beta$ -H elimination, consistent with previous work.<sup>10a,17e</sup> Reactants containing various electrondonating groups on the aromatic ring such as dimethoxy (3g-i), methylenedioxy (3j) or TIPS-protected phenol (3k) gave the desired cyclopropanes in good yields (65-81%). Likewise, electron-withdrawing groups such as fluoro (31, 3m, 3o), nitro (3n) and trifluoromethyl (3p) were found to be compatible. The position of the substituent on the aromatic ring (meta or para) did not have a significant influence on the reaction outcome. In addition, the pyridinylcyclopropane 3q was generated in 70% yield, hence indicating the compatibility of heteroarenes. Moreover, substrates functionalized in *ortho* position to the quaternary carbon (1r) or to the bromine atom (1s, 1t) provided the corresponding cyclopropane products 3r-t in good yields (69-78%). Remarkably, a twofold cyclopropanation reaction was successfully carried out, allowing formation of biscyclopropane 3u in 92% yield via two consecutive cyclopropanations including four C(sp<sup>3</sup>)–H activation steps.

As expected, substrates bearing benzyl instead of methyl groups did not undergo C(sp<sup>3</sup>)-H activation and cyclopropanation, but rather direct  $C(sp^2)$ -H arylation, as illustrated with the 6-membered ring product 5 (Scheme 4b). In contrast, in the case of substrates bearing a quaternary carbon with one methyl and one methylene adjacent to an ester or nitrile group, the construction of trisubstituted cyclopropanes was possible. First, cyclopropanes 3aa and 3ab containing two esters were synthesized in average yields and good diastereoselectivity. probably arising from thermodynamic control. The positional isomers 3ac and **3ad** containing nitrile groups were also produced, albeit with a reduced diastereoselectivity. Finally, the trisubstituted cyclopropane 3ae was generated in 86% combined yield and with a weak, but opposite diastereoselectivity compared to 3ab.



<sup>*a*</sup> Reactions were performed on a 0.2 mmol scale. The aryl bromide precursor was employed unless otherwise noted. Yields refer to the isolated product or mixture of diastereoisomers. Diastereomeric ratios (d.r.) were determined by <sup>1</sup>H NMR of the crude mixture. Relative configurations were determined by NOESY NMR. <sup>*b*</sup> Performed at 120 °C without DMSO. <sup>*c*</sup> <sup>1</sup>H NMR yield using CH<sub>2</sub>Br<sub>2</sub> as internal reference. <sup>*d*</sup> Performed on a 1 mmol scale using *o*-xylene as the solvent.

Molecules containing an aryl-substituted [3.1.0] bicyclic system show interesting biological activities,<sup>2</sup> but only a few methods have been developped for their synthesis<sup>22-26</sup> The access to these attractive scaffolds was investigated by applying our newly developped Pdcatalyzed double C(sp<sup>3</sup>)-H activation reaction (Scheme 5). Starting from lactone 6a, the cyclopropane-fused product 7a was formed in low yield and the major product was the butenolide 8a resulting from  $\beta$ hydrogen elimination (Scheme 5a). A similar result was obtained from lactam 6b. To supress the olefin formation, the reaction was conducted with  $\gamma$ -lactones and lactams 9 bearing the quaternary carbon in  $\beta$ position to the carbonyl group (Scheme 5b). Gratifyingly, this modification led to the cyclopropanefused lactones 10a-b and lactam 10c in 65-79% yields. To further increase the complexity, a more substituted substrate bearing a CH<sub>2</sub>CO<sub>2</sub>Me instead of the CH<sub>3</sub> group was employed. A separable 3:1 diastereomeric mixture of fused  $\gamma$ -lactams was obtained in 79% combined yield, with the major diastereoisomer **10c** having the aryl and ester groups in *cis* relationship. This example shows that our methodology can also be employed to couple two activated methylene groups for the synthesis of tetrasubstituted cyclopropanes.

Lemborexant 11 (Dayvigo<sup>®</sup>, Figure 1) is a dual antagonist of the orexin OX<sub>1</sub> and OX<sub>2</sub> receptors, which was recently approved for the treatment of insomnia.<sup>27</sup> The application of the current method to a short synthesis of this bioactive target is depicted in Scheme 6. Esterification of commercially available aryl bromide 12 and sequential alkylation and allylation led to methyl ester 13. The latter was subjected to ester reduction and oxidative cleavage of the alkene, which directly provided γ-butyrolactone **9e**.<sup>28</sup> The kev cyclopropanation of 9e was performed under the optimized conditions, to furnish the fused lactone 10e in good yield (79%). Direct amidation of the latter with aminopyridine **14** and LiHMDS according to Szostak and co-workers,<sup>29</sup> followed by Mitsunobu reaction with pyrimidinol **16** furnished racemic lemborexant **11** in 28% overall yield over 8 steps.

### Scheme 5. Synthesis of Cyclopropane-Fused Lactones and Lactams

a) with an  $\alpha$ -quaternary carbon



<sup>*a*</sup> Conditions: Pd(PPh<sub>3</sub>)<sub>4</sub> (10 mol%), KOPiv (2 equiv), toluene/DMSO (95:5), 140 °C, 16 h. <sup>*b*</sup> NMR yield. <sup>*c*</sup> Yield of the isolated product. <sup>*d*</sup> Yield of the isolated minor diastereoisomer. <sup>*e*</sup> Ratio determined by <sup>1</sup>H NMR of the crude mixture, relative configuration determined by NOESY NMR.

### CONCLUSIONS

The elusive intramolecular cyclopropanation reaction cleaving two C-H bonds and forging a C-C bond has been developed by taking advantage of the 1,4-Pd shift mechanism. Stoichiometric studies indicated that oxidative addition is the rate-limiting step with the employed substrates and catalyst. The characterization of intermediate Pd complexes allowed the delineation of a mechanistic pathway including  $\sigma$ -aryl- and  $\sigma$ alkylpalladium pivalates, which are in equilibrium via the five-membered diorganopalladacycle. With pivalate as the base and  $PPh_3$  as the ligand, the reductive elimination from this palladacycle is disfavored and the activation of a second C(sp<sup>3</sup>)-H bond is favored, leading to a high-energy four-membered palladacycle which reductively eliminates to give the cyclopropane product. A catalytic version was developed, which enabled the generation of an array of arylcyclopropanes in good yields, including fused ring systems, from simple aryl bromide or triflate precursors. The generation of fused cyclopropanes was applied to a straightforward synthesis of the anti-insomnia drug lemborexant.

# Scheme 6. Application to the Synthesis of Lemborexant<sup>a</sup>



<sup>*a*</sup> Conditions: 1. MeOH,  $H_2SO_4$  (0.1 equiv), 50 °C; 2. LDA (1.2 equiv), MeI (1.05 equiv), -78 to 25 °C; 3. LiHMDS (3 equiv), allyl bromide (5 equiv), THF, 0 to 25 °C (72% for 3 steps); 4. DIBAL-H (2.1 equiv), Et<sub>2</sub>O, -78 to 25 °C (82%); 5. RuCl<sub>3</sub> (5 mol%), NaIO<sub>4</sub> (6 equiv), MeCN/H<sub>2</sub>O 10:1, 25 °C; 6. Pd(PPh<sub>3</sub>)<sub>4</sub> (10 mol%), KOPiv (2 equiv), *o*-xylene/DMSO 95:5, 140 °C; 7. LiHMDS (3 equiv), **14** (3 equiv), toluene, 25 °C (89%); 8. **16** (1.5 equiv), PPh<sub>3</sub> (1.5 equiv), DIAD (1.5 equiv), THF, 25 °C (68%).

This work further illustrates that tuning the catalyst and base in Pd-catalyzed C–H activation reactions may allow to divert mechanistic pathways toward the formation of high value-added products.

### ASSOCIATED CONTENT

### **Supporting Information**

The Supporting Information is available free of charge on the ACS Publications website at DOI: Full optimization tables, crystallographic, procedural and spectral data.

#### **AUTHOR INFORMATION**

#### **Corresponding Author**

\*olivier.baudoin@unibas.ch

### ORCID

Olivier Baudoin: 0000-0002-0847-8493 Antonin Clemenceau: 0000-0002-2024-2061

### Notes

The authors declare no competing financial interest.

### ACKNOWLEDGMENT

This work was financially supported by Oril Industrie, affiliated to Les Laboratoires Servier, and the University of Basel. We thank Dr. Lucile Vaysse-Ludot, Dr. R. Tamion and J. Fournier, ORIL Industrie, for fruitful discussions and constant support, Dr. D. Häussinger, University of Basel, for NMR experiments, and Dr. M. Pfeffer, University of Basel, for MS analyses.

### REFERENCES

- (1) Faust, R. Fascinating Natural and Artificial Cyclopropane Architectures. *Angew. Chem. Int. Ed.* **2001**, *40*, 2251–2253.
- (2) (a) Talele, T. T. The "Cyclopropyl Fragment" is a Versatile Player that Frequently Appears in Preclinical/Clinical Drug Molecules. J. Med. Chem. 2016, 59, 8712–8756. (b) Časar, Z. Synthetic Approaches to Contemporary Drugs that Contain the Cyclopropyl Moiety. Synthesis 2020, 52, 1315–1345.
- (3) (a) Roy, M.-N.; Lindsay, V. N. G.; Charette, A. B. Cyclopropanation Reactions. In *Science of Synthesis*, De Vries, J. G.; Molander, G. A.; Evans, P. A., Eds.; Thieme: Stuttgart, 2011; Vol. 1, pp 731–817. (b) Kulinkovich, O. G. *Cyclopropanes in Organic Synthesis*, John Wiley & Sons, Inc.: Hoboken, 2015.
- (4) (a) Chen, D. Y.-K.; Pouwerb, R. H.; Richard, J.-A. Recent Advances in the Total Synthesis of Cyclopropane-Containing Natural Products. *Chem. Soc. Rev.* 2012, *41*, 4631–4642. (b) Ebner, C.; Carreira, E. M. Cyclopropanation Strategies in Recent Total Syntheses. *Chem. Rev.* 2017, *117*, 11651–11679.
- (5) Giri, R.; Wasa, M.; Breazzano, S. P.; Yu, J.-Q. Converting gem-Dimethyl Groups into Cyclopropanes via Pd-Catalyzed Sequential C-H Activation and Radical Cyclization. Org. Lett. 2006, 8, 5685–5688.
- (6) (a) Liron, F.: Knochel, P. Reactivity of Stable Neopentyl-Pd Intermediates in the Absence of Nucleophile. Tetrahedron Lett. 2007, 48, 4943-4946. (b) Kim, H. S.; Gowrisankar, S.; Kim, S. H.; Kim, J. N. Synthesis of 6-Oxacyclopropa[a]indene Derivatives Starting from Baylis-Hillman Adducts via Pd-Mediated C(sp3)-H Activation. Tetrahedron Lett. 2008, 49, 3858-3861. (c) Huang, Q.; Larock, R. C. Synthesis of Cyclopropanes by Pd-Catalyzed Activation of Alkyl C-H Bonds. Tetrahedron Lett. 2009, 50, 7235-7238. (d) Mao, J.; Zhang, S.-Q.; Shi, B.-F.; Bao, W. Palladium(0)-Catalyzed Cyclopropanation of Benzyl Bromides via C(sp3)-H Bond Activation. Chem. Commun. 2014, 50, 3692-3694. (e) Du, W.; Gu, Q.; Li, Z.; Yang, D. Palladium(II)-Catalyzed Intramolecular Tandem Aminoalkylation via Divergent C(sp<sup>3</sup>)-H Functionalization. J. Am. Chem. Soc. 2015, 137, 1130-1135. (f) Chung, D. S.; Lee, J. S.; Ryu, H.; Park, J.; Kim, H.; Lee, J. H.; Kim, U B.; Lee, W. K.; Baik, M.-H.; Lee, S.-g. Palladium-Catalyzed Divergent Cyclopropanation by Regioselective Solvent-Driven C(sp<sup>3</sup>)-H Bond Activation. Angew. Chem. Int. Ed. 2018, 57, 15460-15464.
- (7) For the construction of cyclopropanes via C–C activation and C(sp<sup>3</sup>)–H activation: Cao, J.; Chen, L.; Sun, F.-N.; Sun, Y.-L.; Jiang, K.-Z.; Yang, K.-F.; Xu, Z.; Xu, L.-W. Pd-Catalyzed Enantioselective Ring Opening/Cross-Coupling and Cylopropanation of Cyclobutanones. *Angew. Chem. Int. Ed.* **2019**, *58*, 897–901.
  - (8) (a) Ma, S.; Gu, Z. 1,4-Migration of Rhodium and Palladium in Catalytic Organometallic Reactions. *Angew. Chem. Int. Ed.* 2005, 44, 7512–7517. (b) Shi, F.; Larock, R. C. Remote C–H Activation via Through-Space Palladium and Rhodium Migrations. *Top. Curr. Chem.* 2010, 292, 123–164. (c)

Rahim, A.; Feng, J.; Gu, Z. 1,4-Migration of Transition Metals in Organic Synthesis. *Chin. J. Chem.* **2019**, *37*, 929– 945.

- (9) (a) Dyker, G. Palladium-Catalyzed C-H Activation of Methoxy Groups: A Facile Synthesis of Substituted 6H-Dibenzo[b,d]pyrans. Angew. Chem. Int. Ed. 1992, 31, 1023– 1025. (b) Dyker, G. Palladium-Catalyzed C-H Activation of tert-Butyl Groups: A Simple Synthesis of 1,2-Dihydrocyclobutabenzene Derivatives. Angew. Chem. Int. Ed. 1994, 33, 103–105. (c) Dyker, G. Palladium-Catalyzed C-H Activation at Methoxy Groups: Regiochemistry of the Domino Coupling Process. Chem. Ber. 1994, 127, 739–742.
- (10) (a) Baudoin, O.; Herrbach, A.; Guéritte, F. The Palladium-Catalyzed C-H Activation of Benzylic gem-Dialkyl Groups. Angew. Chem. Int. Ed. 2003, 42, 5736-5740. (b) Hitce, J.; Retailleau, P.; Baudoin, O. Palladium-Catalyzed Intramolecular C(sp<sup>3</sup>)-H Functionalization: Catalyst Development and Synthetic Applications. Chem. Eur. J. 2007, 13, 792-799. (c) Motti, E.; Catellani, M. Catalytic Dehydrogenation of o-Alkylated or o-Alkoxylated Iodoarenes with Concomitant Hydrogenolysis. Adv. Synth. Catal. 2008, 350, 565-569. (d) Bheeter, C. B.; Jin, R.; Bera, J. K.; Dixneuf, P. H.; Doucet, H. Palladium-Catalysed Dehydrogenative sp<sup>3</sup> C-H Bonds Functionalisation into Alkenes: A Direct Access to N-Alkenylbenzenesulfonamides. Adv. Synth. Catal. 2014, 356, 119-124.
- (11) Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. Catalysts for Suzuki–Miyaura Coupling Processes: Scope and Studies of the Effect of Ligand Structure. J. Am. Chem. Soc. 2005, 127, 4685–4696.
- (12) Pan, J.; Su, M.; Buchwald, S. L. Palladium(0)-Catalyzed Intermolecular Amination of Unactivated C<sub>sp3</sub>-H Bonds. *Angew. Chem. Int. Ed.* 2011, 50, 8647–8651.
- (13) Tan, B.; Bai, L.; Ding, P.; Liu, J.; Wang, Y.; Luan, X. Palladium-Catalyzed Intermolecular [4+1] Spiroannulation by C(sp<sup>3</sup>)-H Activation and Naphthol Dearomatization. *Angew. Chem. Int. Ed.* 2019, *58*, 1474–1478.
- (14) Rocaboy, R.; Anastasiou, I.; Baudoin, O. Redox-Neutral Coupling between Two C(sp<sup>3</sup>)-H Bonds Enabled by 1,4-Palladium Shift for the Synthesis of Fused Heterocycles. *Angew. Chem. Int. Ed.* 2019, 58, 14625–14628.
- (15) Chaumontet, M.; Piccardi, R.; Audic, N.; Hitce, J.; Peglion, J.-L.; Clot, E.; Baudoin, O. Synthesis of Benzocyclobutenes by Palladium-Catalyzed C–H Activation of Methyl Groups: Method and Mechanistic Study. J. Am. Chem. Soc. 2008, 130, 15157–15166.
- (16) Gutiérrez-Bonet, Á.; Juliá-Hernández, F.; de Luis, B.; Martin, R. Pd-Catalyzed C(sp<sup>3</sup>)–H Functionalization/Carbenoid Insertion: All Carbon Quaternary Centers via Multiple C–C Bond Formation. J. Am. Chem. Soc. 2016, 138, 6384–6387.
- (17) (a) Lafrance, M.; Gorelsky, S. I.; Fagnou, K. High-Yielding Palladium-Catalyzed Intramolecular Alkane Arylation: Reaction Development and Mechanistic Studies. J. Am. Chem. Soc. 2007, 129, 14570-14571. (b) Kefalidis, C. E.; Baudoin, O.; Clot, E. DFT Study of the Mechanism of Benzocyclobutene Formation by Palladium-Catalysed C(sp3)-H Activation: Role of the Nature of the Base and the Phosphine. Dalton Trans. 2010, 39, 10528-10535. (c) Rousseaux, S.; Gorelsky, S. I.; Chung, B. K. W.; Fagnou, K. Investigation of the Mechanism of C(sp<sup>3</sup>)-H Bond Cleavage in Pd(0)-Catalyzed Intramolecular Alkane Arylation Adjacent to Amides and Sulfonamides. J. Am. Chem. Soc. 2010, 132, 10692-10705. (d) Rousseaux, R.; Davi, M.; Sofack-Kreutzer, J.; Pierre, C.; Kefalidis, C. E.; Clot, E.; Fagnou, K.; Baudoin, O. Intramolecular Palladium-Catalyzed Alkane C-H Arylation from Aryl Chlorides. J. Am. Chem. Soc. 2010, 132, 10706-10716. (e) Kefalidis, C. E.; Davi, M.; Holstein, P. M.; Clot, E.; Baudoin, O. Mechanistic Study on the Selectivity of Olefin versus Cyclobutene Formation by Palladium(0)-

58 59

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

60

Catalyzed Intramolecular C(sp<sup>3</sup>)–H Activation. J. Org. Chem. 2014, 79, 11903–11910.

- (18) (a) McNally, A.; Haffemayer, B.; Collins, B. S. L.; Gaunt, M. J. Palladium-Catalysed C–H Activation of Aliphatic Amines to Give Strained Nitrogen Heterocycles. *Nature* 2014, *510*, 129–133. (b) He, C.; Gaunt, M. J. Ligand-Enabled Catalytic C–H Arylation of Aliphatic Amines by a Four-Membered-Ring Cyclopalladation Pathway. *Angew. Chem., Int. Ed.* 2015, *54*, 15840-15844. (c) Su, B.; Bunescu, A.; Qiu, Y.; Zuend, S. J.; Ernst, M.; Hartwig, J. F. Palladium-Catalyzed Oxidation of β-C(sp<sup>3</sup>)–H Bonds of Primary Alkylamines through a Rare Four-Membered Palladacycle Intermediate. *J. Am. Chem. Soc.* 2020, *142*, 7912–7919.
- (19) Gutiérrez, E.; Nicasio, M. C.; Paneque,; M.; Ruiz, C.; Salazar, V. Synthesis and Reactivity of New Palladium Alkyl Complexes Containing PMe<sub>3</sub> Ligands: Insertion Reactions and Formation of Bis(pyrazolyl)borate Derivatives J. Organomet. Chem. **1997**, 549, 167–176.
- (20) Campora, J.; López, J. A.; Palma, P.; del Rio, D.; Carmona, E.; Valerga, P.; Graiff, C.; Tiripicchio, A. Synthesis and Insertion Reactions of the Cyclometalated Palladium–Alkyl Complexes Pd(CH<sub>2</sub>CMe<sub>2</sub>-o-C<sub>6</sub>H<sub>4</sub>)L<sub>2</sub>. Observation of a Pentacoordinated Intermediate in the Insertion of SO<sub>2</sub>. *Inorg. Chem.* **2001**, 40, 4116–4126.
- (21) The opening of five-membered palladacycle C<sup>1</sup> under carbonate-mediated conditions was indirectly observed through the formation BCB isomers and deuterium incorporation on the aryl ring.<sup>15</sup>
- (22) 2-Aryl cyclopropane-fused γ-lactones: (a) Milewska, M. J.; Gdaniec, M.; Polonski, T. Synthesis, Stereochemistry, and Chiroptical Spectra of Cyclopropyl Lactones and Thionolactones. *Tetrahedron Asymmetry* 1996, 7, 3169–3180.
  (b) Muraoka, O.; Tanabe, G.; Momose, T. Accentuation of the Di-p-Methane Reactivity by Central Carbon Substitution in the 4-(Phenylmethyl)-2(5H)-furanone. *Heterocycles* 1990, *31*,1589–1592.

- (23) 1-Aryl cyclopropane-fused γ-lactones: (a) Shuto, S.; Ono, S.; Hase, Y.; Kamiyama, N.; Matsuda, A. Synthesis of (+)- and (-)-milnaciprans and their conformationally restricted analogs. *Tetrahedron Lett.* **1996**, *37*, 641–644. (b) Doyle, M. P.; Hu, W. A New Enantioselective Synthesis of Milnacipran and an Analogue by Catalytic Asymmetric Cyclopropanation. *Adv. Synth. Catal.* **2001**, *343*, 299–302.
- (24) 1-Aryl cyclopropane-fused γ-lactams: Wang, K.-B.; Ran, R.-Q.; Xiu, S.-D.; Li, C.-Y. Synthesis of 3-Azabicyclo[3.1.0]hexan-2-one Derivatives via Gold-Catalyzed Oxidative Cyclopropanation of N-Allylynamides. *Org. Lett.* 2013, 15, 2374–2377.
- (25) Pedroni, J.; Cramer, N. Enantioselective C-H Functionalization-Addition Sequence Delivers Densely Substituted 3-Azabicyclo[3.1.0]hexanes. J. Am. Chem. Soc. 2017, 139, 12398–12401.
- (26) Zhang, Z.-Z.; Han, Y.-Q.; Zhan, B.-B.; Wang, S.; Shi, B.-F. Synthesis of Bicyclo[n.1.0]alkanes by a Cobalt-Catalyzed Multiple C(sp<sup>3</sup>)–H Activation Strategy. *Angew. Chem. Int. Ed.* **2017**, *56*, 13145–13149.
- (27) Yoshida, Y.; Naoe, Y.; Terauchi, T.; Ozaki, F.; Doko, T.; Takemura, A.; Tanaka, T.; Sorimachi, K.; Beuckmann, C. T.; Suzuki, M.; Ueno, T.; Ozaki, S.; Yonaga, M. Discovery of (1*R*,2*S*)-2-{[(2,4-Dimethylpyrimidin-5-yl)oxy]methyl}-2-(3fluorophenyl)-*N*-(5-fluoropyridin-2-yl)cyclopropanecarboxam ide (E2006): A Potent and Efficacious Oral Orexin Receptor Antagonist. J. Med. Chem. 2015, 58, 4648–4664.
- (28) Gao, R.; Fan, R.; Canney, D. J. Synthesis of  $\beta$ , $\beta$ -Disubstituted  $\gamma$ -Butyrolactones by Chemoselective Oxidation of 1,4-Diols and  $\gamma$ -Hydroxy Olefins with RuCl<sub>3</sub>/NaIO<sub>4</sub>. *Synlett* **2015**, *26*, 661–665.
- (29) Li, G.; Ji, C.-L.; Hong, X.; Szostak, M. Highly Chemoselective, Transition-Metal-Free Transamidation of Unactivated Amides and Direct Amidation of Alkyl Esters by N–C/O–C Cleavage. J. Am. Chem. Soc. 2019, 141, 11161– 11172.



TOC graphic