


This article was downloaded by: [University of Edinburgh]

On: 09 June 2013, At: 07:14

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lncn20>

Synthesis and Antimicrobial Activity of Some S- β -D-Glucosides of 4-Mercaptopyrimidine

A. H. Moustafa^a, H. A. Morsy^a, M. G. Assy^a & A. Z. Haikal^a

^a Department of Chemistry, Faculty of Science, Zagazig University, Zagazig, Egypt

Published online: 04 Sep 2009.

To cite this article: A. H. Moustafa, H. A. Morsy, M. G. Assy & A. Z. Haikal (2009): Synthesis and Antimicrobial Activity of Some S- β -D-Glucosides of 4-Mercaptopyrimidine, *Nucleosides, Nucleotides and Nucleic Acids*, 28:9, 835-845

To link to this article: <http://dx.doi.org/10.1080/15257770903172639>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS AND ANTIMICROBIAL ACTIVITY OF SOME S- β -D-GLUCOSIDES OF 4-MERCAPTOPYRIMIDINE

A. H. Moustafa, H. A. Morsy, M. G. Assy, and A. Z. Haikal

Department of Chemistry, Faculty of Science, Zagazig University, Zagazig, Egypt

□ 5-Acetyl-2-aryl-6-methyl-4-(2,3,4,6-tetra-O-acetyl- β -D-glucopyranosylmercapto)pyrimidines **3a–c** were obtained by the reaction of 5-acetyl-2-aryl-6-methyl-pyrimidine thiol **1a–c** with 2,3,4,6-tetra-O-acetyl- α -D-glucopyranosyl bromide (**2**) in aq. KOH/acetone. The reaction of **1a–c** with peracetylated galactose **5** and peracetylated ribose **8** under MW irradiation gave 5-acetyl-2-aryl-6-methyl-4-(2,3,4,6-tetra-O-acetyl- β -D-galactopyranosylmercapto)pyrimidine **6a–c** and 5-acetyl-2-aryl-6-methyl-4-(2,3,5-tri-O-acetyl- β -D-ribofuranosylmercapto)pyrimidines **9a–c**. The deprotection of **3a–c**, **6a–c**, and **9a–c** in the presence of methanol and TEA/H₂O yielded the deprotected products **4a–c**, **7a–c**, and **10a–c**. The structures of the compounds were confirmed by using IR, ¹H, ¹³C spectra and microanalysis. Selected members of these compounds were screened for antimicrobial activity.

Keywords Glycosides; microwave; antimicrobial activity


INTRODUCTION

The pyrimidines and their nucleosides are interesting classes in medicinal and agricultural chemistry. Some pyrimidine nucleosides are known as antiviral,^[1–3] antibacterials,^[4,5] antitumor,^[6] antipyretic,^[7] antiinflammatory drugs,^[8] pesticides,^[9] and plant growth regulators.^[10]

Glycosylthio-heterocycles have attracted much attention because of their abilities to function as biological inhibitors, inducers, and ligands for affinity of chromatography of carbohydrate-process enzymes and proteins.^[11] The utility of microwave (MW) as a nonconventional energy source has advantages such as significant rate enhancements of reactions, higher product yields, and greater selectivity of organic reactions.^[12–14] The combination of solvent-free reaction conditions and microwave irradiation has also been investigated.^[15–17]

Received 13 May 2009; accepted 30 June 2009.

Address correspondence to Ahmed H. Moustafa, Chemistry Department, Faculty of Science, Zagazig University, Zagazig, Egypt. E-mail: ah.hu.mostafa@yahoo.com



SCHEME 1


RESULTS AND DISCUSSION

The syntheses of 5-acetyl-6-methyl-2-arylpyrimidine-4-thiols **1a-c** have been achieved as reported in literature.^[18] Reaction of **1a-c** with glycosyl bromide **2** in the presence of aqueous potassium hydroxide/acetone^[4,5] gave 5-acetyl-6-methyl-2-aryl-4-(2,3,4,6-tetra-*O*-acetyl- β -D-glycopyranosylthio)pyrimidines (**3a-c**) (Scheme 1). Irradiation of **1a-c** with **5** in the presence of silica gel as a solid support in MW oven^[4,19] for 10 minutes gave the corresponding thiogalactosides **6a-c** in 40–45% yield (Scheme 1).

The structure of compounds **3a-c** was confirmed on the basis of their elemental analysis and spectral data. Their IR spectra showed a band at $1765\text{--}1745\text{ cm}^{-1}$ for the acetoxy carbonyl groups in addition to a band at 1695 cm^{-1} due to the carbonyl of 5-acetylpyrimidine.

The ^1H NMR spectrum of **3a** showed the presence of signals at δ 1.53, 1.96, 2.00, and 2.02 ppm characteristic for 4 CH_3 of OAc groups, in addition to 2 CH_3 at δ 2.36, 2.58 ppm and a doublet at δ 6.23 ppm for anomeric proton with $J_{1',2'} = 8.97\text{ Hz}$, which confirmed the presence of the β -configuration. The ^{13}C NMR spectrum of **3a** showed peak at δ 80.1 ppm, characteristic for anomeric carbon, and absence of $\text{C}=\text{S}$ group, which indicates the formation of S-glycoside not N-glycoside.

The ^1H NMR spectrum of **3b** and **3c** showed similar results. Also the glycosides **6a-c** showed spectral characteristic conforming their



SCHEME 2

β-configuration. On the other hand, reaction of **1a-c** with **8** under MW irradiation in the presence of silica gel as a solid support^[4,20] for 8 minutes gave the corresponding ribosides **9a-c** in 40–45% yield (Scheme 2).

The ¹H NMR spectrum of **9a** showed signals at δ 2.06, 2.07, and 2.15 ppm for 3 CH₃ of OAc groups, and doublet at δ 6.46 ppm for anomeric proton with coupling constant $J_{1',2'} = 2.80$ Hz indicating the presence of β-configuration. Similarly **9b** and **9c** gave similar results. Deprotection of the previous prepared S-glucosides **3a-c**, **6a-c**, and **9a-c** using triethylamine in presence of methanol and few drops of water^[4,5,20] gave the deprotected glycosides **4a-c**, **7a-c**, and **10a-c** (Schemes 1 and 2).

The ¹H, ¹³C NMR spectra for **4a-c** showed the absence of the four signals of the 4 OAc groups and presence of signals at δ 4.53, 5.11, 5.24, and 5.51 ppm for 4 OH groups of sugar moieties, in addition of doublet at δ 5.61, 5.68, and 6.26 ppm for the anomeric protons of **4a-c**. The IR spectra of **4a-c** showed the appearance of the broad band at 3417 cm⁻¹ for 4 OH groups. Similar results were obtained for **7a-c** and **10a-c**.

Antimicrobial Activity

Compound **3c** showed higher activity against *B. subtilis* as gram positive and *E. coli* as gram negative bacteria, while **3a,b** and **6b** showed only moderate activity. Compounds **3** and **9a** showed activity towards fungi (e.g., *Penicillium sp.* and *aspergillus sp.*). Compounds **6a,c** and **9b,c** did not show any activity against the tested microorganisms (Table 1).

EXPERIMENTAL

All melting points are uncorrected and were measured using an Electrothermal IA 9100 apparatus. Thin layer chromatography (TLC) was performed on Merck Silica Gel 60F₂₅₄ with detection by UV light and charring with H₂SO₄/EtOH (5:95); The IR spectra (KBr disc) were recorded on a Pye Unicam Sp-3-300 and Shimadzu FTIR 8101 PC infrared

TABLE 1 Antimicrobial activity of tested compounds (inhibition zones [mm], minimum inhibitory concentration [$\mu\text{g/mL}$])

Compound no.	<i>E. Coli</i>	<i>B. Subtilis</i>	<i>Penicillium Sp.</i>	<i>Aspergillus Sp.</i>
3a	—	15	15	10
3b	—	10	—	—
3c	10	25	—	—
6a	—	—	—	—
6b	—	5	—	—
6c	—	—	—	—
9a	—	10	10	8
9b	—	—	—	—
9c	—	—	—	—

spectrophotometer. The ultraviolet (UV) spectra were recorded by UV-160A, UV-visible recording spectrometer Shimadzu using EtOH 95% as a solvent. The ^1H and ^{13}C NMR spectra were recorded on a Varian Mercury VX-NMR 300 & 100 MHz spectrometer. Chemical shifts are expressed on the $\delta(\text{ppm})$ scale using TMS as the standard and coupling-constant values are given in Hz. Elemental analyses were determined on a Perkin Elmer 240.

General Methods for Preparation of Glucosides

Method A

To a solution of pyrimidinethiol **1a–c** (0.01 mol) in aq. KOH [0.01 mol in distilled water (6 mL)] was added a solution of 2,3,4,6-tetra-*O*-acetyl- α -**D**-glucopyranosyl bromide (**2**) (0.011 mol) in acetone (30 mL). The reaction mixture was stirred at room temperature and followed by *tlc* until the reaction was finished (4.5–5 hours), the reaction mixture was evaporated under reduced pressure and the residue was washed with distilled water to remove potassium bromide. The product was dried and crystallized from the appropriate solvent.

Method B

A mixture of pyrimidinethiol **1a–c** (0.001 mol) and (0.001 mol) of per acetylated galactose (**5**) or ribose (**8**) was dissolved in methylene chloride, then 1 gm of silica gel (200–400 mesh) was added, the solvent was removed by evaporation, and then the dried residue was transferred into a glass beaker and irradiated for (8–10 minutes) in a domestic microwave oven. The product was extracted with methylene chloride, evaporated to dryness, and crystallized from appropriate solvent.

5-Acetyl-6-methyl-2-phenyl-4-(2,3,4,6-tetra-*O*-acetyl- β -D**-glucopyranosyl-thio)pyrimidine (3a).** *Method A*: yield 26%. As colorless crystals from ethanol; m.p. 105–107°C; λ_{max} = 300 nm. ^1H NMR (DMSO- d_6 , 300 MHz):

δ 1.53, 1.96, 2.00, 2.02 (4s, 12 H, 4 CH₃CO), 2.36 (s, 3 H, CH₃ pyrimidine), 2.58 (s, 3 H, CH₃CO pyrimidine), 4.00 (m, 1 H, H-5'), 4.13 (dd, 1 H, J_{6'',5'} = 2.10, J_{6'',6'} = 12.32 Hz, H-6''), 4.26 (dd, 1 H, J_{6',5'} = 5.96, J_{6',6''} = 12.3 Hz, H-6'), 5.12 (t, 1 H, J_{4',3'} = 8.78, J_{4',5'} = 9.13 Hz, H-4'), 5.31 (t, 1 H, J_{2',1'} = 7.70, J_{2',3'} = 9.20 Hz, H-2'), 5.63 (t, 1 H, J_{3',2'} = 9.20, J_{3',4'} = 8.78 Hz, H-3'), 6.23 (d, 1 H, J_{1',2'} = 8.97 Hz, H-1'), 7.56 (m, 3 H, Ar-H), 8.58 (m, 2 H, Ar-H). ¹³C NMR (DMSO-d₆, 100 MHz): δ 20.8, 21.1, 21.2, 21.4, 23.8, 32.2 (6 CH₃), 62.8 (C-6'), 69.2 (C-2'), 69.8 (C-4'), 73.8 (C-3'), 75.9 (C-5'), 80.1 (C-1'), 129.4, 129.6, 130.9, 132.4, 136.7, 162.54, 163.1, 163.6, 170.1, 170.4, 170.6 and 202.8 (Ar-C, 2 C=N and 5 C=O). Anal. Calcd for C₂₇H₃₀N₂O₁₀S (574.6): C, 56.44; H, 5.26; N, 4.88. Found: C, 56.68; H, 5.31; N, 5.03.

5-Acetyl-6-methyl-2-(4-methoxyphenyl)-4-(2,3,4,6-tetra-O-acetyl-β-D-glucopyranosylthio)pyrimidine (3b). *Method A:* yield 24%. As colorless crystals from ethanol; m.p. 100–103°C; $\lambda_{\max}$ = 301.4 nm. ¹H NMR (DMSO-d₆, 300 MHz): δ 1.54, 1.95, 2.00, 2.02 (4s, 12 H, 4 CH₃CO), 2.34 (s, 3 H, CH₃ pyrimidine), 2.59 (s, 3 H, CH₃CO pyrimidine), 3.89 (s, 3 H, OCH₃), 3.98 (m, 1 H, H-5'), 4.12 (dd, 1 H, J_{6'',5'} = 2.10, J_{6'',6'} = 12.3 Hz, H-6''), 4.26 (dd, 1 H, J_{6',5'} = 5.96, J_{6',6''} = 12.3 Hz, H-6'), 5.12 (t, 1 H, J_{4',3'} = 8.97, J_{4',5'} = 9.13 Hz, H-4'), 5.31 (t, 1 H, J_{2',1'} = 7.90, J_{2',3'} = 9.22 Hz, H-2'), 5.62 (t, 1 H, J_{3',2'} = 9.22, J_{3',4'} = 8.78 Hz, H-3'), 6.24 (d, 1 H, J_{1',2'} = 8.87 Hz, H-1'), 7.10 (d, 2 H, J = 9.0 Ar-H), 8.40 (d, 2 H, J = 9.0 Hz, Ar-H). ¹³C NMR (DMSO-d₆, 100 MHz): δ 20.8, 21.1, 21.2, 21.4, 23.9, 32.3 (6 CH₃), 56.2 (OCH₃), 62.8 (C-6'), 69.2 (C-2'), 69.8 (C-4'), 73.8 (C-3'), 75.9 (C-5'), 80.0 (C-1'), 129.2, 130.0, 131.3, 131.7, 162.3, 163.0, 163.1, 163.6, 170.1, 170.2, 170.4, 170.6 and 202.7 (Ar-C, 2 C=N and 5 C=O). Anal. Calcd for C₂₈H₃₂N₂O₁₁S (604.63): C, 55.62; H, 5.33; N, 4.63. Found: C, 55.50; H, 5.68; N, 4.60.

5-Acetyl-2-(4-chlorophenyl)-6-methyl-4-(2,3,4,6-tetra-O-acetyl-β-D-glucopyranosylthio)pyrimidine (3c). *Method A:* yield 34%. As colorless crystals from ethanol; m.p. 104–106°C; $\lambda_{\max}$ = 300 nm. IR (KBr): 1749 (C=O, esters) cm⁻¹, 1697 (C=O, acetyl) cm⁻¹. ¹H NMR (DMSO-d₆, 300 MHz): δ 1.53, 1.95, 2.00, 2.02 (4s, 12 H, 4 CH₃CO), 2.45 (s, 3 H, CH₃ pyrimidine), 2.55 (s, 3 H, CH₃CO pyrimidine), 3.98 (m, 1 H, H-5'), 4.12 (dd, 1 H, J_{6'',5'} = 2.10, J_{6'',6'} = 12.3 Hz, H-6''), 4.27 (dd, 1 H, J_{6',5'} = 5.96, J_{6',6''} = 12.3 Hz, H-6'), 4.92 (t, 1 H, J_{4',3'} = 8.97, J_{4',5'} = 9.13 Hz, H-4'), 5.08 (t, 1 H, J_{2',1'} = 7.90, J_{2',3'} = 9.22 Hz, H-2'), 5.67 (t, 1 H, J_{3',2'} = 9.22, J_{3',4'} = 8.78 Hz, H-3'), 6.21 (d, 1 H, J_{1',2'} = 7.89 Hz, H-1'), 7.59 (d, 2 H, J = 8.90 Ar-H), 8.49 (d, 2 H, J = 8.90 Ar-H). ¹³C NMR (DMSO-d₆, 100 MHz): δ 20.2, 20.3, 20.4, 22.8, 22.9, 31.3 (6 CH₃), 61.9 (C-6'), 68.2 (C-2'), 68.9 (C-4'), 72.9 (C-3'), 74.9 (C-5'), 79.2 (C-1'), 128.8, 128.9, 130.3, 130.4, 134.7, 136.5, 160.6, 162.7, 162.9, 169.4, 169.5, 169.6 and 201.9 (Ar-C, 2 C=N and 5 C=O). Anal. Calcd for C₂₇H₂₉ClN₂O₁₀S (609.04): C, 53.25; H, 4.80; N, 6.40. Found: C, 53.12; H, 4.57; N, 6.21.

5-Acetyl-6-methyl-2-phenyl-4-(β -D-glucopyranosylthio)pyrimidine (4a).

General method for deprotection of acetyl groups: Triethylamine (1ml) was added to 10 ml a solution of glucosides or ribosides (0.001mol) in (MeOH/H₂O). The mixture was stirred overnight at room temperature; it was evaporated under reduced pressure and co-evaporated with MeOH until the excess of Et₃N was removed. The residue was then crystallized from the proper solvent to give the desired product with yield >85%. Colourless crystals from ethanol; m.p. 115–117°C; IR (KBr): 3417 (br. 4 OH) cm⁻¹ and 1689 (C=O) cm⁻¹. ¹H NMR (DMSO-d₆/D₂O, 300 MHz): δ 2.46 (s, 3 H, CH₃ pyrimidine), 2.60 (s, 3 H, CH₃CO pyrimidine), 3.24–3.85 (m, 6 H, H-6', H-6'', H-5', H-4', H-3' and H-2'), 5.69 (d, 1 H, J_{1',2'} = 8.97 Hz, H-1'), 7.45 (m, 3 H, Ar-H), 8.42 (m, 2 H, Ar-H). ¹³C NMR (DMSO-d₆, 300 MHz): δ 22.5, 31.4 (2 CH₃), 61.6 (C-6'), 69.6 (C-2'), 71.7 (C-4'), 78.5 (C-3'), 81.8 (C-5'), 83.0 (C-1'), 128.2, 128.8, 128.9, 130.3, 131.4, 136.3, 161.6, 163.7 and 202.6 (Ar-C, 2 C=N and C=O). Anal. Calcd for C₁₉H₂₂N₂O₆S (406.45): C, 56.15; H, 5.46; N, 6.89. Found: C, 55.91; H, 5.55; N, 6.93.

5-Acetyl-6-methyl-2-(4-methoxyphenyl)-4-(β -D-glucopyranosylthio)

pyrimidine (4b). As for 4a, colorless crystals from ethanol; m.p. 110–111°C; IR (KBr): 3415 (br. 4 OH) cm⁻¹ and 1690 (C=O) cm⁻¹. ¹H NMR (DMSO-d₆/D₂O, 300 MHz): δ 2.45 (s, 3 H, CH₃ pyrimidine), 2.62 (s, 3 H, CH₃CO pyrimidine), 3.23–3.85 (m, 6 H, H-6', H-6'', H-5', H-4', H-3' and H-2'), 3.89 (s, 3 H, OCH₃), 6.21 (d, 1 H, J_{1',2'} = 8.90 Hz, H-1'), 7.10 (d, 2 H, J = 8.62 Hz, Ar-H), 8.40 (d, 2 H, J = 8.62 Hz, Ar-H). Anal. Calcd for C₂₀H₂₄N₂O₇S (436.48): C, 55.03; H, 5.54; N, 6.42. Found: C, 55.23; H, 5.44; N, 6.50.

5-Acetyl-2-(4-chlorophenyl)-6-methyl-4-(β -D-glucopyranosylthio)pyrimidine (4c).

As for 4a, colorless crystals from ethanol; m.p. 114–116°C; IR (KBr): 3417 (br. 4 OH) cm⁻¹ and 1700 (C=O) cm⁻¹. ¹H NMR (DMSO-d₆, 300 MHz): δ 2.46 (s, 3 H, CH₃ pyrimidine), 2.59 (s, 3 H, CH₃CO pyrimidine), 3.13–3.46 (m, 6 H, H-6', H-6'', H-5', H-4', H-3' and H-2'), 4.53 (t, 1 H, J = 3.56 Hz, OH-6'), 5.11 (d, J = 4.34 Hz, 1 H, OH-4'), 5.24 (d, 1 H, J = 5.24 Hz, OH-3'), 5.51 (d, 1 H, J = 4.96 Hz, OH-2'), 6.24 (d, 1 H, J_{1',2'} = 8.89 Hz, H-1'), 7.60 (d, 2 H, J = 8.91 Hz, Ar-H), 8.39 (d, 2 H, J = 8.91 Hz, Ar-H). ¹³C NMR (DMSO-d₆, 100 MHz): δ 22.4, 31.3 (2 CH₃), 60.7 (C-6'), 69.6 (C-2'), 71.6 (C-4'), 78.4 (C-3'), 81.9 (C-5'), 83.1 (C-1'), 128.9, 130.0, 130.4, 135.2, 136.3, 160.6, 161.7, 163.9 and 202.5 (Ar-C, 2 C=N and C=O). Anal. Calcd for C₁₉H₂₁ClN₂O₆S (440.90): C, 51.76; H, 4.80; N, 6.35. Found: C, 51.42; H, 4.36; N, 6.17.

5-Acetyl-6-methyl-2-phenyl-4-(2,3,4,6-tetra-O-acetyl- β -D-galactopyranosylthio)pyrimidine (6a).

Method B: yield 40%. As colorless crystals from ethanol; m.p. 102–103°C; $\lambda_{\max}$ = 303.2 nm. ¹H NMR (DMSO-d₆, 300 MHz): δ 1.50, 1.95, 2.03, 2.13 (4s, 12 H, 4 CH₃CO), 2.46 (s, 3 H, CH₃ pyrimidine), 2.59 (s, 3 H, CH₃CO pyrimidine), 3.95 (m, 1 H, H-5'), 4.12 (dd, 1 H, J_{6',5'} = 6.50, J_{6',6''} = 12.0 Hz, H-6'), 4.25 (dd, 1 H, J_{6',5'} = 5.60, J_{6',6''} = 12.0 Hz, H-6''),

5.21 (t, 1 H, $J_{3',2'} = 10.0$, $J_{3',4'} = 3.50$ Hz, H-3'), 5.41 (t, 1 H, $J_{2',1'} = 8.20$, $J_{2',3'} = 10.0$ Hz, H-2'), 5.62 (t, 1 H, $J_{4',3'} = 3.50$, $J_{4',5'} = 1.00$ Hz, H-4'), 6.26 (d, 1 H, $J_{1',2'} = 8.20$ Hz, H-1'), 7.57 (m, 3 H, Ar-H), 8.56 (d, 2 H, Ar-H). ^{13}C NMR (DMSO- d_6 , 100 MHz): δ 20.1, 20.3, 20.4, 22.7, 22.9, 31.3 (6 CH_3), 62.0 (C-6'), 69.1 (C-2'), 68.9 (C-4'), 73.8 (C-3'), 75.9 (C-5'), 80.1 (C-1'), 129.3, 129.6, 130.8, 132.3, 136.7, 162.5, 163.1, 163.5, 170.2, 170.3, 170.4, 170.6 and 202.6 (Ar-C, 2 C=N and 5 C=O). Anal. Calcd for $\text{C}_{27}\text{H}_{30}\text{N}_2\text{O}_{10}\text{S}$ (574.6): C, 56.44; H, 5.26; N, 4.88. Found: C, 56.53; H, 5.33; N, 5.23.

5-Acetyl-6-methyl-2-(4-methoxyphenyl)-4-(2,3,4,6-tetra-O-acetyl-β-D-galactopyranosylthio)pyrimidine (6b). *Method B:* yield 41%. As colorless crystals from ethanol; m.p. 98–99°C; $\lambda_{\text{max}} = 302.8$ nm. ^1H NMR (DMSO- d_6 , 300 MHz): δ 1.51, 1.59, 2.03, 2.13 (4s, 12 H, 4 CH_3CO), 2.49 (s, 3 H, CH_3 pyrimidine), 2.57 (s, 3 H, CH_3CO pyrimidine), 3.85 (s, 3 H, OCH_3), 3.95 (m, 1H, H-5'), 4.13 (dd, 1 H, $J_{6',5'} = 6.50$, $J_{6',6''} = 12.0$ Hz, H-6''), 4.26 (dd, 1 H, $J_{6',5'} = 5.60$, $J_{6',6''} = 12.0$ Hz, H-6'), 5.22 (t, 1 H, $J_{3',2'} = 10.0$, $J_{3',4'} = 3.50$ Hz, H-3'), 5.41 (t, 1 H, $J_{2',1'} = 8.20$, $J_{2',3'} = 10.0$ Hz, H-2'), 5.62 (t, 1 H, $J_{4',3'} = 3.50$, $J_{4',5'} = 1.00$ Hz, H-4'), 6.26 (d, 1 H, $J_{1',2'} = 8.22$ Hz, H-1'), 7.08 (d, 2 H, $J = 8.63$ Hz, Ar-H), 8.48 (d, 2 H, $J = 8.63$ Hz, Ar-H). ^{13}C NMR (DMSO- d_6 , 100 MHz): δ 20.1, 20.2, 20.4, 22.6, 22.8, 31.2 (6 CH_3), 56.2 (OCH_3), 61.8 (C-6'), 66.4 (C-2'), 70.9 (C-4'), 72.4 (C-3'), 74.4 (C-5'), 79.6 (C-1'), 128.3, 129.2, 130.4, 131.7, 161.4, 162.1, 162.7, 169.4, 169.5, 169.7, 169.8, 170.0 and 201.8 (Ar-C, 2 C=N and 5 C=O). Anal. Calcd for $\text{C}_{28}\text{H}_{32}\text{N}_2\text{O}_{11}\text{S}$ (604.63): C, 55.62; H, 5.33; N, 4.63. Found: C, 55.51; H, 5.48; N, 4.60.

5-Acetyl-2-(4-chlorophenyl)-6-methyl-4-(2,3,4,6-tetra-O-acetyl-β-D-galactopyranosylthio)pyrimidine (6c). *Method B:* yield 45%. As colorless crystals from ethanol; m.p. 106–109°C; $\lambda_{\text{max}} = 301.4$ nm. ^1H NMR (DMSO- d_6 , 300 MHz): δ 1.49, 1.58, 2.03, 2.14 (4s, 12 H, 4 CH_3CO), 2.48 (s, 3 H, CH_3 pyrimidine), 2.59 (s, 3 H, CH_3CO pyrimidine), 3.98 (m, 1 H, H-5'), 4.14 (dd, 1 H, $J_{6',5'} = 6.50$, $J_{6',6''} = 12.0$ Hz, H-6''), 4.27 (dd, 1 H, $J_{6',5'} = 5.60$, $J_{6',6''} = 12.0$ Hz, H-6'), 5.23 (t, 1 H, $J_{3',2'} = 10.0$, $J_{3',4'} = 3.50$ Hz, H-3'), 5.42 (t, 1 H, $J_{2',1'} = 8.20$, $J_{2',3'} = 10.0$ Hz, H-2'), 5.65 (t, 1 H, $J_{4',3'} = 3.50$, $J_{4',5'} = 1.00$ Hz, H-4'), 6.36 (d, 1 H, $J_{1',2'} = 8.22$ Hz, H-1'), 7.53 (d, 2 H, $J = 8.90$ Hz, Ar-H), 8.49 (d, 2 H, $J = 8.90$ Hz, Ar-H). ^{13}C NMR (DMSO- d_6 , 100 MHz): δ 20.1, 20.2, 20.4, 22.6, 22.8, 31.1 (6 CH_3), 61.6 (C-6'), 66.8 (C-2'), 69.6 (C-4'), 70.5 (C-3'), 70.6 (C-5'), 79.3 (C-1'), 128.5, 130.1, 134.4, 134.5, 160.3, 162.1, 162.6, 168.5, 168.9, 169.2, 169.3, 169.5 and 201.6 (Ar-C, 2 C=N and 5 C=O). Anal. Calcd for $\text{C}_{27}\text{H}_{29}\text{ClN}_2\text{O}_{10}\text{S}$ (609.04): C, 53.25; H, 4.80; N, 6.40. Found: C, 53.52; H, 4.78; N, 6.33.

5-Acetyl-6-methyl-2-phenyl-4-(β-D-galactopyranosylthio) pyrimidine (7a). As for **4a**, colorless crystals from ethanol; m.p. 118–119°C; IR (KBr): 3417 (br 4 OH) cm^{-1} and 1690 (C=O) cm^{-1} . ^1H NMR (DMSO- $d_6/\text{D}_2\text{O}$, 300 MHz): δ 2.45 (s, 3 H, CH_3 pyrimidine), 2.59 (s, 3 H, CH_3CO pyrimidine), 3.23–3.84 (m, 6 H, H-6', H-6'', H-5', H-4', H-3' and H-2'), 6.21 (d, 1 H, $J_{1',2'} =$

8.40 Hz, H-1'), 7.57 (m, 3 H, Ar-H), 8.53 (m, 2 H, Ar-H). Anal. Calcd for $C_{19}H_{22}N_2O_6S$ (406.45): C, 56.15; H, 5.46; N, 6.89. Found: C, 55.90; H, 5.45; N, 6.93.

5-Acetyl-6-methyl-2-(4-methoxyphenyl)-4-(β -D-galactopyranosylthio)pyrimidine (7b). As for **4a**, colorless crystals from ethanol; m.p. 112–114°C; IR (KBr): 3415 (br 4 OH) cm^{-1} and 1695 (C=O) cm^{-1} . 1H NMR (DMSO- d_6 /D $_2$ O, 300 MHz): δ 2.45 (s, 3 H, CH_3 pyrimidine), 2.59 (s, 3 H, CH_3CO pyrimidine), 3.24–3.75 (m, 6 H, H-6', H-6'', H-5', H-4', H-3' and H-2'), 3.85 (s, 3 H, OCH $_3$), 6.23 (d, 1H, $J_{1',2'} = 8.51$ Hz, H-1'), 7.52 (d, 2 H, $J = 8.90$ Hz, Ar-H), 8.48 (d, 2 H, $J = 8.90$ Hz, Ar-H). Anal. Calcd for $C_{20}H_{24}N_2O_7S$ (436.48): C, 55.03; H, 5.54; N, 6.42. Found: C, 55.43; H, 5.44; N, 6.52.

5-Acetyl-2-(4-chlorophenyl)-6-methyl-4-(β -D-galactopyranosylthio)pyrimidine (7c). As for **4a**, colorless crystals from ethanol; m.p. 119–120°C; IR (KBr): 3418 (br. 4 OH) cm^{-1} and 1692 (C=O) cm^{-1} . 1H NMR (DMSO- d_6 /D $_2$ O, 300 MHz): δ 2.45 (s, 3 H, CH_3 pyrimidine), 2.58 (s, 3 H, CH_3CO pyrimidine), 3.35–3.85 (m, 6 H, H-6', H-6'', H-5', H-4', H-3' and H-2'), 6.22 (d, 1 H, $J_{1',2'} = 8.45$ Hz, H-1'), 7.53 (d, 2 H, $J = 8.90$ Hz, Ar-H), 8.48 (d, 2 H, $J = 8.90$ Hz, Ar-H). Anal. Calcd for $C_{19}H_{21}ClN_2O_6S$ (440.90): C, 51.76; H, 4.80; N, 6.35. Found: C, 51.44; H, 4.32; N, 6.27.

5-Acetyl-6-methyl-2-phenyl-4-(2,3,5-tri-O-acetyl- β -D-ribofuranosylthio)pyrimidine (9a). *Method B*: yield 41%; as colorless crystals from methanol; m.p. 111–113°C; $\lambda_{max} = 300$ nm. 1H NMR (DMSO- d_6 , 300 MHz): δ 2.06, 2.07, 2.15 (3s, 9 H, 3 CH_3CO), 2.43 (s, 3 H, CH_3 pyrimidine), 2.59 (s, 3 H, CH_3CO pyrimidine), 4.10 (dd, 1 H, $J_{5',4'} = 3.40$, $J_{5',5''} = 11.6$ Hz, H-5'), 4.16 (dd, 2 H, $J_{5'',4'} = 3.60$, $J_{5'',5'} = 11.6$ Hz, H-5''), 4.41 (m, 1 H, H-4'), 5.54 (t, 1 H, $J_{3',2'} = 2.71$, $J_{3',4'} = 6.62$ Hz, H-3'), 5.63 (dd, 1 H, $J_{2',1'} = 2.60$, $J_{2',3'} = 2.80$ Hz, H-2'), 6.46 (d, 1 H, $J_{1',2'} = 2.80$ Hz, H-1'), 7.45–8.40 (m, 5 H, Ar-H). Anal. Calcd for $C_{24}H_{26}N_2O_8S$ (502.54): C, 57.36; H, 5.52; N, 5.57. Found: C, 57.14; H, 5.11; N, 5.75.

5-Acetyl-6-methyl-2-(4-methoxyphenyl)-4-(2,3,5-tri-O-acetyl- β -D-ribofuranosylthio)pyrimidine (9b). *Method B*: yield 36%, as colorless crystals from methanol ; m.p. 107–109°C; UV; $\lambda_{max} = 299$ nm. 1H NMR (DMSO- d_6 , 300 MHz): δ 2.05, 2.07, 2.16 (3s, 9 H, 3 CH_3CO), 2.51 (s, 3 H, CH_3CO pyrimidine), 2.62 (s, 3 H, CH_3 pyrimidine), 3.34 (s, 3 H, OCH $_3$), 4.10 (dd, 1 H, $J_{5',4'} = 3.40$, $J_{5',5''} = 11.6$ Hz, H-5'), 4.14 (dd, 1 H, $J_{5'',4'} = 3.60$, $J_{5'',5'} = 11.6$ Hz, H-5''), 4.41 (m, 1 H, H-4'), 5.43 (t, 1 H, $J_{3',2'} = 2.71$, $J_{3',4'} = 6.62$ Hz, H-3'), 5.56 (dd, 1 H, $J_{2',1'} = 2.60$, $J_{2',3'} = 2.80$ Hz, H-2'), 6.43 (d, 1 H, $J_{1',2'} = 2.75$ Hz, H-1') 7.61 (d, 2 H, $J = 8.90$ Hz, Ar-H), 8.31 (d, 2 H, $J = 8.90$ Hz, Ar-H). Anal. Calcd for $C_{25}H_{28}N_2O_9S$ (532.56): C, 56.38; H, 5.30; N, 5.26. Found: C, 56.10; H, 5.32; N, 5.19.

5-Acetyl-2-(4-chlorophenyl)-6-methyl-4-(2',3',5'-tri-O-acetyl- β -D-ribofuranosylthio)pyrimidine (9c). *Method B*: yield 40%, as colorless crystals from methanol ; m.p. 105–107°C; UV; $\lambda_{max} = 299$ nm. 1H NMR

(DMSO- d_6 , 300 MHz): δ 2.05, 2.07, 2.16 (3s, 9 H, 3 CH_3CO), 2.62 (s, 3 H, CH_3CO pyrimidine), 2.51 (s, 3 H, CH_3 pyrimidine), 4.06 (dd, 1 H, $J_{5',4'} = 3.40$, $J_{5',5''} = 11.6$ Hz, H-5'), 4.12 (dd, 1 H, $J_{5'',4'} = 3.60$, $J_{5'',5'} = 11.6$ Hz, H-5''), 4.45 (m, 1 H, H-4'), 5.42 (t, 1 H, $J_{3',2'} = 2.71$, $J_{3',4'} = 6.62$ Hz, H-3'), 5.55 (dd, 1 H, $J_{2',1'} = 2.60$, $J_{2',3'} = 2.80$ Hz, H-2'), 6.42 (d, 1 H, $J_{1',2'} = 2.60$ Hz, H-1'), 7.60 (d, 2 H, $J = 8.80$ Hz, Ar-H), 8.34 (d, 2 H, $J = 8.80$ Hz, Ar-H). ^{13}C NMR (DMSO- d_6 , 100 MHz): δ 20.7, 20.8, 21.0, 23.4, 31.8 (5 CH_3), 62.7 (C-5'), 70.5 (C-3'), 75.6 (C-2'), 79.6 (C-4'), 85.4 (C-1'), 129.3, 130.2, 135.2, 137.0, 160.9, 163.3, 164.1, 166.5, 169.7, 169.8, 170.3 and 202.1 (Ar-C, 2 C=N and 4 C=O). Anal. Calcd for $\text{C}_{24}\text{H}_{25}\text{ClN}_2\text{O}_8\text{S}$ (536.98): C, 53.68; H, 4.69; N, 5.22. Found: C, 53.61; H, 4.81; N, 4.93.

5-Acetyl-6-methyl-2-phenyl-4-(β -D-ribofuranosylthio)pyrimidine (10a).

As for **4a**, colorless crystals from ethanol; m.p. 124–126°C. IR (KBr): 3419 (3 OH) cm^{-1} , 1692 (C=O) cm^{-1} . ^1H NMR (DMSO- $d_6/\text{D}_2\text{O}$, 300 MHz): δ 2.41 (s, 3 H, CH_3 pyrimidine), 2.54 (s, 3 H, CH_3CO pyrimidine), 3.46–3.73 (m, 2 H, H-5' 5 Prime;), 3.97 (m, 1 H, H-4'), 4.12 (dd, 1 H, $J_{3',4'} = 6.60$, $J_{3',2'} = 2.70$ Hz, H-3'), 4.23 (dd, 1 H, $J_{2',1'} = 2.60$, $J_{2',3'} = 2.70$ Hz, H-2'), 6.21 (d, 1 H, $J_{1',2'} = 2.80$ Hz, H-1'), 7.53–8.37 (m, 5 H, Ar-H). Anal. Calcd for $\text{C}_{189}\text{H}_{20}\text{N}_2\text{O}_5\text{S}$ (376.43): C, 57.43; H, 5.36; N, 7.44. Found: C, 57.40; H, 4.98; N, 7.47.

5-Acetyl-6-methyl-2-(4-methoxyphenyl)-4-(β -D-ribofuranosylthio) pyrimidine (10b). As for **4a**, colorless crystals from ethanol; m.p. 117–119°C; IR (KBr): 3420 (3 OH) cm^{-1} , 1695 (C=O) cm^{-1} . ^1H NMR (DMSO- $d_6/\text{D}_2\text{O}$, 300 MHz): δ 2.45 (s, 3 H, CH_3 pyrimidine), 2.54 (s, 3 H, CH_3CO pyrimidine), 3.41–3.83 (m, 2 H, H-5' H-5 Prime;), 3.89 (s, 3 H, OCH_3), 3.96 (m, 1 H, H-4'), 4.10 (dd, 1 H, $J_{3',4'} = 6.59$, $J_{3',2'} = 2.70$ Hz, H-3'), 4.20 (t, 1 H, $J_{2',1'} = 2.60$, $J_{2',3'} = 2.70$ Hz, H-2'), 6.22 (d, 1 H, $J_{1',2'} = 2.8$ Hz, H-1'), 7.54 (d, 2 H, $J = 8.90$ Hz, Ar-H), 8.36 (d, 2 H, $J = 8.90$ Hz, Ar-H). ^{13}C NMR (DMSO- d_6 , 100 MHz): δ 20.0, 31.0 (2 CH_3), 56.3 (OCH_3), 61.5 (C-5'), 70.4 (C-3'), 75.3 (C-2'), 75.8 (C-4'), 74.0 (C-1'), 120.9, 121.3, 128.7, 129.8, 136.5, 136.9, 161.9, 169.9, and 201.8 (Ar-C, 2 C=N and C=O). Anal. Calcd for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_6\text{S}$ (406.45): C, 56.15; H, 5.46; N, 6.89. Found: C, 56.43; H, 5.32; N, 6.70.

5-Acetyl-2-(4-chlorophenyl)-6-methyl-4-(β -D-ribofuranosylthio)pyrimidine (10c). As for **4a**, colourless crystals from ethanol; m.p. 123–125°C; IR (KBr): 3422 (OH) cm^{-1} , 1690 (C=O) cm^{-1} . ^1H NMR (DMSO- $d_6/\text{D}_2\text{O}$, 300 MHz): δ 2.43 (s, 3 H, CH_3 pyrimidine), 2.57 (s, 3 H, CH_3CO pyrimidine), 3.42–3.83 (m, 2 H, H-5' 5Prime;), 3.95 (m, 1 H, H-4'), 4.12 (dd, 1 H, $J_{3',4'} = 6.58$, $J_{3',2'} = 2.75$ Hz, H-3'), 4.20 (t, 1 H, $J_{2',1'} = 2.60$, $J_{2',3'} = 2.70$ Hz, H-2'), 6.23 (d, 1 H, $J_{1',2'} = 2.80$ Hz, H-1'), 7.25 (d, 2 H, $J = 8.98$ Hz, Ar-H), 8.05 (d, 2 H, $J = 8.98$ Hz, Ar-H). Anal. Calcd for $\text{C}_{18}\text{H}_{19}\text{ClN}_2\text{O}_5\text{S}$ (410.87): C, 52.62; H, 4.66; N, 6.82. Found: C, 52.34; H, 4.54; N, 6.76.

Antibacterial Screening

The antimicrobial activities of some newly synthesized compounds were screened for their antibacterial activity against three species of bacteria, namely (*E. coli*) as Gram positive, (*B. Subtilis*) as Gram negativeve and (*Penicillium Sp.*) and (*Asperagillus Sp.*) as fungi, using saturated disks according to the a dapted method.^[21] The tested compounds were dissolved in chloroform to get a solution of 1 mg/mL concentration. The inhibition zones were measured in (mm) at the end of an incubation period of 48 hours at 37°C. Chloroform showed no inhibition zones. The fungi cultures were maintained on dextrose agar medium.

REFERENCES

1. Abdel-Rahman, A.H.; Abdel-Maged, A.E.S.; Goda, A.E.S.; Zeid, I.F.; El-Ashry, E.S.H. Synthesis and anti-HBV activity of thiouracils linked via S and N-1 to the 5-Position of methyl β -D-ribofuranoside. *Nucleosides, Nucleotides Nucleic Acids* **2003**, 22, 2027–2038.
2. Abdel-Rahman, A.H.; El-Etrawy, A.S.H.; Abdel-Maged, A.E.S.; Zeid, I.F.; El-Ashry, E.S.H. Synthesis and antiviral evaluation of novel 2,3-dihydroxypropyl nucleosides from 2- and 4-thiouracils. *Nucleosides, Nucleotides Nucleic Acids* **2008**, 27, 1257–1271.
3. Lednicer, D.; Mitscher, L.A. *The Organic Chemistry of Drug Synthesis*, Pergamon Press; New York, 1980.
4. El-Sayed, H.A.; Moustafa, A.H.; Haikal, A.Z.; Abdou, I.M.; El-Ashry, E.S.H. Synthesis and evaluation of antimicrobial activity of some pyrimidine glycosides. *Nucleosides, Nucleotides Nucleic Acids* **2008**, 27, 1061–1071.
5. Moustafa, A.H.; Saad, H.A.; Shehab, W.S.; El-Mobayed, M.M. Synthesis of some new pyrimidine derivatives of expected antimicrobial activity. *Phosphorus, Sulfer Silicon* **2008**, 183, 115–135.
6. Maria, T.C.; Cenzo, C.; Valentina, L.; Valentina, O. Synthesis and in vitro antitumoral activity of new hydrazinopyrimidine-5-carbonitrile derivatives. *Bioorg. Med. Chem.* **2006**, 14, 366–372.
7. Smith, P.A.S.; Kan, R.O. Cyclization of isothiocyanates as a route to phthalic and homophthalic acid derivatives. *J. Org. Chem.* **1964**, 29, 2261–2268.
8. Nega, S.; Aionso, J.; Diaz, A.; Junquere, F. Synthesis of 3-substituted-4-phenyl-2-thioxo-1,2,3,4,5,6,7,8-octahydrobenzo[4,5] thieno [2,3-d]pyrimidines. *J. Heterocycl. Chem.* **1990**, 27, 269–273.
9. Tetsuo, S.; Mikio, T.; Hidetoshi, H.; Daijirom, H.; Akira, I. *Jpn. Kokai Tokyo JP*, 62, 132, 884 (1987); C. A., 107 p. 198350h (1987).
10. Shishoo, C.J.; Devani, M.B.; Ullas, G.V.; Ananthan, S.; Bhadti, V.S. Studies on the Synthesis of 2-(2-Arylviny)thieno[2,3-d]pyrimidines and 5-(2-Arylviny) triazolothieno[3,2-e]pyrimidines. *J. Heterocycl. Chem.* **1985**, 22, 825–832.
11. El Ashry, E.S.H.; Awad, L.F.; Atta, A.I. Synthesis and role of glycosylthio heterocycles in carbohydrate chemistry. *Tetrahedron* **2006**, 62, 2943–2998.
12. Li, Z.; Wang, Z.-Y.; Zhu, W.; Xing, Y.-L.; Zhao, Y.-L. CeCl₃·7H₂O-KI-Catalyzed, environmentally friendly synthesis of N,N-disubstituted ureas in water under microwave irradiation. *Synth. Comm.* **2005**, 35, 2325–2331.
13. Santagada, V.; Frecentese, F.; Perissutti, E.; Cirillo, D.; Terracciano, S.; Caliendo, G. A suitable 1,2,4-oxadiazoles synthesis by microwave radiation. *Bioorg. Med. Chem. Lett.* **2004**, 14, 4491–4493.
14. a) Galema, S.A. Microwave chemistry. *Chem. Soc. Rev.* **1997**, 26, 233–238. b) Mingos, D.; Whittaker, M.P.A. In *Microwave Dielectric Heating Effects in Chemical Synthesis in Chemistry Under Extreme or Non Classical Conditions*, eds. R.V. Eldik; C.D. Hubbark, John Wiley, New York, 1997, p. 479. c) Gabriel, C.; Gabriel, S.; Grant, E.H.; Halstead, B.S.J.; Mingos, D.M.P. Dielectric parameters relevant to microwave dielectric heating. *Chem. Soc. Rev.* **1998**, 27, 213–223. d) Perrux, L.; Loupy, A. A tentative rationalization of microwave effects in organic synthesis according to the reaction medium, and mechanistic considerations. *Tetrahedron* **2001**, *Tetrahedron, Tetrahedron*. e) Hamelin, J.; Bazureau, J.P.; Texier-Boullet, F. In *Microwave in Organic Synthesis*, ed. A. Loupy; Wiley-VCH, New York, 2002, p. 253. f) El Ashry, E.S.H.; Ramadan, E.; Kassem, A.A.; Haggag, M. Microwave irradiation for accelerating

- organic reactions. Part 1: Three, four and five membered heterocycles. *Adv. Heterocycl. Chem.* **2005**, 88, 1–110. g) El Ashry, E.S.H.; Kassem, A.A.; Ramadan, E. Microwave irradiation for accelerating organic reactions. Part 2: Six, seven fused and spiro heterocyclic ring Systems. *Adv. Heterocycl. Chem.* **2006**, 90, 1–127. h) El Ashry, E.S.H.; Kassem, A.A. Account of microwave irradiation for accelerating organic reactions. *Arkovic.* **2006**, ix, 1–15.
15. a) Devi, I.; Bhuyan, P.J. An expedient method for the synthesis of 6-substituted uracils under microwave irradiation in a solvent-free medium. *Tetrahedron Lett.* **2005**, 46, 5727–5729. b) Dhakshinamoorthy, A.; Pitchumani, K. Facile clay-induced fischer indole synthesis: a new approach to synthesis of 1,2,3,4-tetrahydrocarbazole and indoles. *Appl. Catal. A* **2005**, 292, 305–311.
 16. Regnier, T.; Lavastre, O. General and environmentally friendly synthesis of heterocyclic multidentate-molecules based on microwave-assisted heating protocol. *Tetrahedron* **2006**, 62, 155–159.
 17. Yadav, L.D.S.; Yadav, B.S.; Dubey, S. Mineral supported syntheses of benzoxazine-2-thiones under microwave irradiation. *Tetrahedron* **2004**, 60, 131–136.
 18. Assy, M.G. Some reaction with ethyl 2-aryl-4-mercapto-6-ethylpyrimidine-5-carboxylates. *Sulfur Lett.* **1990**, 11, 75–82.
 19. Andrzejewska, M.; Kaminski, J.; Kazmierczuk, Z. Microwave induced synthesis of ribonucleosides on solid support. *Nucleosides Nucleotides Nucleic Acids* **2002**, 21, 73–78.
 20. Haikal, A.Z.; El-Ashry, E.S.H.; Banoub, J.H. Synthesis and structural characterization of 1-(D-glycosyloxy)phthalazines. *J. Carbohydr. Res.* **2003**, 338, 2291–2299.
 21. Gupta, R.; Saxena, R.K.; Chatarvedi, P.; Virdi, J.S.J. *Appl. Bacterial.* **1995**, 78, 378–383.