View Article Online View Journal



# Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. O. Terent'ev, M. Y. Sharipov, I. Krylov, D. V. Gaidarenko and G. Nikishin, *Org. Biomol. Chem.*, 2014, DOI:



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/obc

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxx

### ARTICLE TYPE

## Manganese triacetate as an efficient catalyst for bisperoxidation of styrenes

Alexander O. Terent'ev,\*<sup>*a,b*</sup> Mikhail Yu. Sharipov,<sup>*a,b*</sup> Igor B. Krylov,<sup>*a*</sup> Darya V. Gaidarenko,<sup>*b*</sup> Gennady I. Nikishin<sup>*a*</sup>

s Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX DOI: 10.1039/b000000x

A method was developed for the bisperoxidation of styrenes with *tert*-butyl hydroperoxide in the presence of a catalytic amount of manganese(III) acetate. It was shown that compounds of manganese in oxidation states 2, 4, and 7 also catalyze this reaction. Target [1,2-bis(*tert*-butylperoxy)ethyl]arenes were <sup>10</sup> synthesized in yields from 46 to 75%.

#### Introduction

Due to the ready availability and high efficiency, organic peroxides have been widely used for more than half a century as radical polymerization initiators in the industrial synthesis, for

- <sup>15</sup> example, of such polymers as polystyrene, polyvinyl chloride, polyacrylates, and high-pressure polyethylene. Organic peroxides are also used for the crosslinking of silicone, acrylonitrile-butadiene, and fluorinated rubbers. A wide variety of commercial monomers and their compositions demands a wide assortment of <sup>20</sup> peroxides, the most well-known of which are organic
- hydroperoxides, dialkyl peroxides, diacyl peroxides, peroxy esters, peroxydicarbonates, peroxycarbonates, peracetals, cyclic triperoxides, and geminal bishydroperoxides.<sup>1,2</sup>
- In the past decades, organic peroxides have got attention from <sup>25</sup> researchers in the fields of medicinal chemistry and pharmacology because these compounds, particularly ozonides and tetraoxanes, were found to exhibit marked antimalarial,<sup>3</sup> anthelmintic,<sup>4</sup> and antitumor activities.<sup>5</sup> Peroxides, particularly derivatives of lower aldehydes and ketones, are still attracting
- <sup>30</sup> interest also as energy-rich compounds. For instance, the explosive power of triacetone triperoxide is comparable with that of trinitrotoluene.<sup>6</sup>

Due to the tendency of peroxides to decompose, their high sensitivity to reducing agents and variable valence metal ions,

<sup>35</sup> and the ease of a large range of transformations accompanied by the O-O bond cleavage, the search for general approaches to the synthesis of peroxides and the selective synthesis of peroxides of the specified structure are difficult problems.

The transition metal (Cu, Mn, Co)/hydroperoxide system was 40 used for the first time by Kharasch in the synthesis of peroxides

- from alkenes, ketones, and tertiary amines more than six decades ago.<sup>7</sup> Since that time, the research on and application of this peroxidation method were documented in a number of publications. The formation of peroxides was observed in various
- <sup>45</sup> reactions of hydroperoxides catalyzed by metal salts and their complexes: copper,<sup>8</sup> cobalt,<sup>9</sup> and iron,<sup>10,11</sup> dinuclear nickel complexes,<sup>12</sup> palladium acetate,<sup>13</sup> ruthenium(II)-bipyridine

anchored montmorillonite,<sup>14</sup> and metalloporphyrins.<sup>15</sup> Peroxides were synthesized in high yields by the peroxidation of amines,

<sup>50</sup> amides, and lactams catalyzed by ruthenium salts.<sup>16</sup> Examples of the synthesis of such compounds are limited by the tendency of peroxides to decompose in the presence of variable valence metals.

In the present study, we developed a new approach to the <sup>55</sup> synthesis of vicinal bis(*tert*-butyl)peroxides from styrenes and *tert*-butyl hydroperoxide using manganese(III) acetate. Manganese(III) acetate is a strong oxidizer. This compound proved to be an efficient reagent in redox and free radical reactions accompanied by the C-C,<sup>17</sup> C-O,<sup>18</sup> and C-N<sup>19</sup> bond

<sup>60</sup> formation. The combination of two oxidizing agents,  $Mn(OAc)_3$ and TBHP, imparts new properties to the system. These properties were used to oxidize alkenes at the allylic position to form enones.<sup>20</sup>

The known methods for the synthesis of vicinal bisperoxides from styrenes and TBHP are based on the use of palladium acetate,<sup>13</sup> ruthenium(II) bipyridine anchored montmorillonite,<sup>14</sup> dinuclear nickel complexes,<sup>12</sup> and Mn(III) porphyrins.<sup>15</sup> The drawback of the former two approaches, although they provide good yields (65-85% and 30-69%, respectively), is the use of

- <sup>70</sup> expensive catalysts. In the third approach, the difficult-to-prepare catalyst is employed and, besides, the target bisperoxide is produced in a rather low yield (26%). The reactions with the use of iron(III) or manganese(III) porphyrins (Fe(III)TDCIPP-OAc or Mn(III)TDCIPP-OAc) give the target products with low yields.
- <sup>75</sup> In the present study, we found that styrenes are selectively peroxidized with high conversion in the reaction with TBHP and manganese(III) acetate to form vicinal bisperoxides ([1,2-bis(*tert*butylperoxy)ethyl]arenes).

#### **Results and discussion**

<sup>80</sup> Vicinal bisperoxides **2a-i** (Scheme 1) were synthesized from styrenes **1a-i** containing substituents of different nature.



a: R = H; R' = H; b: R = H; R' = Me; c: R = 4-Me; R' = H; d: R = t-Bu; R' = H; e: R = 4-OMe; R' = H; f: R = 2-OMe; R' = H; g: R = H; R' = Ph; h: R = 2,3-(CH)<sub>4</sub>; R' = H; i: R = Cl; R' = H

Scheme 1 Synthesis of vicinal bis(*tert*-butyl)peroxides 2a-i from styrenes 1a-i and *tert*-butyl hydroperoxide

In order to optimize the reaction conditions, we studied the <sup>5</sup> peroxidation of styrene **1a** to form [1,2-bis(*tert*-butylperoxy)ethyl]benzene **2a** by varying the reagent ratio and the nature of the catalyst. The best results were achieved with  $Mn(OAc)_3$  (Table 1).

 Table 1 Synthesis of [1,2-bis(*tert*-butylperoxy)ethyl]benzene 2a from

 10 styrene 1a and *tert*-butyl hydroperoxide using compounds of manganese as catalyst



Published on 26 November 2014. Downloaded by University of Washington on 02/12/2014 07:10:08.

D a			X: 11 6 9 . 0/
Run"	Molar ratio	Reaction time, n;	Yield of $2a$ , %
	<b>1a</b> : TBHP : catalyst	(solvent)	
Mn(OAc) <sub>3</sub> •2H <sub>2</sub> O			
1	1:4:3	48; (CH <sub>3</sub> CN)	29
2	1:4:0.2	48; (CH <sub>3</sub> CN)	60
3	1:3:2	48; (CH <sub>3</sub> CN)	42
4	1:3:0.5	48; (CH <sub>3</sub> CN)	58
5	1:3:0.4	48; (CH <sub>3</sub> CN)	70
6	1:3:0.2	48; (CH <sub>3</sub> CN)	73
$7^b$	1:3:0.2	1; (CH <sub>3</sub> CN)	46
8	1:3:0.1	1; (CH <sub>3</sub> CN)	7
9	1:3:0.1	48; (CH <sub>3</sub> CN)	75
10	1:3:0.05	48; (CH <sub>3</sub> CN)	53
11	1:2:0.1	48; (CH <sub>3</sub> CN)	60
12	1:3:0.1	48; (CH <sub>2</sub> Cl <sub>2</sub> )	16
13	1:3:0.1	48; (Benzene)	43
14	1:3:0.1	48; (AcOH)	37
15	1:3:0.1	48; (Acetone)	57
MnO <sub>2</sub> , KMnO <sub>4</sub> , Mn(OAc) <sub>2</sub> •4H <sub>2</sub> O, MnCl <sub>2</sub> •4H <sub>2</sub> O			
16	1:3:0.1 MnO <sub>2</sub>	48; (CH <sub>3</sub> CN)	43
17	1:3:0.3 MnO <sub>2</sub>	48; (CH <sub>3</sub> CN)	57
18	1:3:0.4 KMnO4	48; (CH <sub>3</sub> CN)	10
19	1:3:0.1 Mn(OAc) <sub>2</sub>	48; (CH <sub>3</sub> CN)	19
20	$1:3:0.4 \text{ MnCl}_2$	72; (CH <sub>3</sub> CN)	8

<sup>*a*</sup> **General procedure:** t-BuOOH and a catalyst were sequentially added with vigorous stirring to a solution of styrene **1a** (0.5 g, 4.85 mmol) in <sup>15</sup> CH<sub>3</sub>CN, CH<sub>2</sub>Cl<sub>2</sub>, benzene, AcOH, acetone (10 mL) at 20-25 °C. 2-

Phenyloxirane was obtained in all experiments in yields approximately from 1 to 5 %. <sup>b</sup> 78-80 °C.

The conditions of the synthesis of peroxide **2a** were optimized <sup>20</sup> mainly with the use of Mn(OAc)<sub>3</sub> (runs 1-15), which gave the highest yields of the product, as well as employing MnO<sub>2</sub>, KMnO<sub>4</sub>, Mn(OAc)<sub>2</sub>, and MnCl<sub>2</sub> (runs 16-20). In runs 1-15, the reagent ratio was varied from 2 to 4 moles of TBHP per mole of styrene **1a** and 0.05-3 moles of the catalyst

<sup>25</sup> 1a. All experiments, except for run 7, were carried out at 20-25 °C. In run 7, at 78-80 °C the reaction time significantly decreased from 48 h to 1 h; however, this led to a substantial decrease in the yield of bisperoxide **2a**. The optimum TBHP/styrene **1a** molar ratio was 3; Mn(OAc)<sub>3</sub>/styrene **1a** ratio was 0.1-0.4. Application of other solvents CH<sub>2</sub>Cl<sub>2</sub>, benzene, AcOH, and acetone (runs 12-15) does not permit to increase the yield of **2a**. Supposing that manganese ions in different oxidation states are involved in the formation of bisperoxide **2a**, we carried out experiments 16-20 with the use of MnO<sub>2</sub>, KMnO<sub>4</sub>, Mn(OAc)<sub>2</sub>, and MnCl<sub>2</sub>. Despite the difference in the oxidation states of manganese in the compounds used, target bisperoxide **2a** was obtained in these

experiments in yields from 8 to 57 %. In the next step of the study, using the reaction conditions of one of the best experiments (run 9, see Table 1), we performed the 40 synthesis of a number of bisperoxidated ethylarenes **2b-i** (Table 2).



The introduction of a substituent in the aromatic ring or in the <sup>45</sup> vinyl group (at the alpha position) of styrene **1a** has only a slight effect on the formation of bisperoxides. Under the conditions of run 9, bisperoxides **2b-i** were prepared from styrenes **1b-i** in moderate or good yields, which are, however, somewhat lower compared to **2a** (Table 2). These results suggest that other <sup>50</sup> structurally similar styrenes can also be used in the synthesis of the corresponding bisperoxides.

#### - Proposed mechanism of the synthesis of bisperoxides

Based on the results of peroxidation using compounds of manganese in different oxidation states and the literature data on <sup>55</sup> oxidation reactions involving manganese salts,<sup>7,15,20</sup> we proposed the mechanism of the peroxidation (Scheme 2).

In the first step, the reaction of t-BuOOH with Mn(OAc)<sub>3</sub> can proceed *via* two pathways (**a** and **b**). In the reaction *via* the pathway **a**, Mn(OAc)<sub>3</sub> directly oxidizes t-BuOOH to form the 600 peroxide radical t-BuOO•. In the reaction *via* the pathway **b**, the acetic acid moiety is replaced with *tert*-butyl hydroperoxide to form Mn<sup>III</sup>(OAc)<sub>2</sub>OO-t-Bu, which fragments on two particles (arrows **c** and **e**).<sup>7,20</sup> The radical t-BuO• rapidly abstracts hydrogen (step **d**) from t-BuOOH to give t-BuOO•.<sup>21</sup> In the 65 pathway **f**, Mn<sup>IV</sup>(OAc)<sub>2</sub>=O is produced from Mn<sup>III</sup>(OAc)<sub>2</sub>OO-tPublished on 26 November 2014. Downloaded by University of Washington on 02/12/2014 07:10:08.

Bu followed by the reaction of this intermediate with t-BuOOH (similarly to  $Mn(OAc)_3$ ) to form  $O=Mn^{IV}(OAc)OO$ -t-Bu.<sup>15</sup>



Scheme 2 Proposed mechanism of the peroxidation of styrene 1a with tert-butyl hydroperoxide using compounds of manganese

Based on the results of the experiment with MnO<sub>2</sub> (runs 16 and 17, Table 1), in which **2a** was obtained in good yield, it can be supposed that O=Mn<sup>IV</sup>(OAc)OO-t-Bu initiates a new peroxidation cycle. The radicals t-BuOO• react with styrene **1a** <sup>10</sup> (step **g**) to give the stabilized benzyl radical, which either recombines with t-BuOO• or is oxidized with Mn<sup>III</sup>(OAc)<sub>2</sub>OO-t-Bu accompanied by the transfer of the ligand OO-t-Bu (step **h**) to give the target bisperoxide **2a**.<sup>7,15,20</sup> Indirectly existence of benzyl radical is confirmed by the isolation of phenyloxirane which is <sup>15</sup> the product of intramolecular radical substitution of t-BuO•.

compound  $Mn(OAc)_2$  that is formed in the steps **a** and **i** is oxidized with t-BuOOH (step **j**) to form  $Mn(OAc)_3$ . This is how the step-by-step transformation of manganese derivatives in various oxidation states occurs in the catalytic cycle of <sup>20</sup> peroxidation of the C=C bond.

#### Conclusions

It was shown that manganese salts in oxidation states 2, 3, 4, and 7 catalyze the peroxidation of styrenes with *tert*-butyl hydroperoxide. A method was proposed for the synthesis of [1,2-

<sup>25</sup> bis(*tert*-butylperoxy)ethyl]arenes from readily available and cheap starting reagents. Despite a large number of elementary steps in this reaction, the peroxidation is accomplished in moderate or good yields (up to 75%). The synthesized compounds and the method for their preparation may be applied <sup>30</sup> for the production of radical polymerization initiators of <sup>30</sup>

unsaturated monomers.

#### Experimental

Caution: Although we have encountered no difficulties in working with peroxides, precautions, such as the use of <sup>35</sup> shields, fume hoods, and the avoidance of heating and shaking, should be taken whenever possible.

NMR spectra were recorded on a commercial instrument (300.13 MHz for <sup>1</sup>H, 75.48 MHz for <sup>13</sup>C) in CDCl<sub>3</sub>. High resolution mass spectra (HRMS) were measured using electrospray ionization

<sup>40</sup> (ESI).<sup>22</sup> The measurements were performed in positive ion mode (interface capillary voltage 4500 V); the mass ratio was from m/z 50 to 3000 Da; external/internal calibration was done with Electrospray Calibrant Solution. A syringe injection was used for solutions in MeCN (flow rate 3  $\mu$ L/min). Nitrogen was applied as <sup>45</sup> a dry gas; the interface temperature was set at 180 °C.

The TLC analysis was carried out on standard silica gel chromatography plates. Chromatography was performed on silica gel (0.060-0.200 mm, 60 A, CAS 7631-86-9).

Hexane, CH<sub>3</sub>CN, benzene, acetone, AcOH, CH<sub>2</sub>Cl<sub>2</sub>, ethyl acetate <sup>50</sup> (EA), Mn(OAc)<sub>3</sub>•2H<sub>2</sub>O, Mn(OAc)<sub>2</sub>•4H<sub>2</sub>O, MnCl<sub>2</sub>•4H<sub>2</sub>O, MnO<sub>2</sub>, and KMnO<sub>4</sub> were purchased from Acros.

#### Experiments for Table 1 (runs 1-11).

In a 15 ml round bottom flask, styrene **1a** (0.5 g, 4.85 mmol) was  $_{55}$  dissolved in CH<sub>3</sub>CN (10 mL). Then a 70% aqueous t-BuOOH

solution (1.25-2.50 g, 9.7-19.4 mmol, 2-4 mol per mole of 1a)and  $Mn(OAc)_3 \bullet 2H_2O$  (0.065-5.2 g, ; 0.243-14.55 mmol, 0.05-3 mol per mole of 1a) were successively added with stirring. The heterogeneous reaction mixture was stirred at 20-25°C (runs 1-6

- <sup>5</sup> and 8-11) for 1 h (run 8) or 48 h (runs 1-6 and 9-11) or at 78-80°C for 1 h (run 7). The reaction mixture was cooled to room temperature (run 7). Then the reaction mixture was extracted with hexane (4×20 mL). The combined organic extracts were concentrated under water aspirator pressure. 1,2-Bis(*tert*-10 butylperoxy)ethyl]benzene 2a was isolated by column chromatography on silica gel using ethyl acetate/hexane (5:95, v/v) as the eluent. 2-Phenyloxirane was obtained in all experiments in yields approximately from 1 to 5 %. In the experiment 1 2-phenyloxirane was isolated by column 15 chromatography on silica gel using EA/hexane (5:95, v/v) as the
- eluent. The yield of 2-phenyloxirane was 5% (0.029 g, 0.24 mmol)

#### Experiments for Table 1 (runs 12, 13).

Published on 26 November 2014. Downloaded by University of Washington on 02/12/2014 07:10:08

<sup>20</sup> In a 15 ml round bottom flask, styrene **1a** (0.5 g, 4.85 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> or benzene (10 mL). Then a 70% aqueous t-BuOOH solution (1.87 g, 14.55 mmol; 3 mol per mole of **1a**) and Mn(OAc)<sub>3</sub>•2H<sub>2</sub>O (0.13 g, 0.485 mmol; 0.1 mol per mole of **1a**) were successively added with stirring. The heterogeneous
<sup>25</sup> reaction mixture was stirred at 20-25°C for 48 h. Then the solution was filtered off from manganese salts. The organic phase was concentrated under water aspirator pressure. 1,2-Bis(*tert*-butylperoxy)ethyl]benzene **2a** was isolated by column chromatography on silica gel using EA/hexane (5:95, v/v) as the <sup>30</sup> eluent.

#### Experiments for Table 1 (run 14, 15).

- In a 15 ml round bottom flask, styrene **1a** (0.5 g, 4.85 mmol) was dissolved in AcOH or acetone (10 mL). Then a 70% aqueous t-<sup>35</sup> BuOOH solution (1.87 g, 14.55 mmol; 3 mol per mole of **1a**) and Mn(OAc)<sub>3</sub>•2H<sub>2</sub>O (0.13 g, 0.485 mmol; 0.1 mol per mole of **1a**) were successively added with stirring. The practically homogeneous reaction mixture was stirred at 20-25°C for 48 h. Then CHCl<sub>3</sub> (10 mL) and a solution of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>·5 H<sub>2</sub>O (200 mg)
- <sup>40</sup> in H<sub>2</sub>O (20 mL) were added, the mixture was shaken, the organic layer was separated, and the aqueous layer was extracted with CHCl<sub>3</sub> (2×10 mL). Combined organic extracts were successively washed with a saturated aqueous NaHCO<sub>3</sub> solution (15 mL) and H<sub>2</sub>O (20 mL), and dried under MgSO<sub>4</sub>. The organic phase was
- <sup>45</sup> concentrated under water aspirator pressure. 1,2-Bis(*tert*butylperoxy)ethyl]benzene **2a** was isolated by column chromatography on silica gel using EA/hexane (5:95, v/v) as the eluent.

#### 50 Experiments for Table 1 (runs 16, 17, 19).

- In a 15 ml round bottom flask, styrene **1a** (0.5 g, 4.85 mmol) was dissolved in CH<sub>3</sub>CN (10 mL). Then a 70% aqueous t-BuOOH solution (1.87 g, 14.55 mmol, 3 mol per mole of **1a**) and the catalyst  $Mn(OAc)_2 \bullet 4H_2O$  or  $MnO_2$  (0.485-1.455 mmol, 0.1-0.3
- <sup>55</sup> mol per mole of **1a**) were successively added with stirring. The heterogeneous reaction mixture was stirred at 20-25°C for 48 h. Compound **2a** was isolated as described above (in runs 1-11).

#### Experiments for Table 1 (runs 18, 20).

- <sup>60</sup> In a 15 ml round bottom flask, styrene **1a** (0.5 g, 4.85 mmol) was dissolved in CH<sub>3</sub>CN (10 mL). Then a 70% aqueous t-BuOOH solution (1.87 g, 14.55 mmol, 3 mol per mole of **1a**) and the catalyst MnCl<sub>2</sub>•4H<sub>2</sub>O or KMnO<sub>4</sub>, (1.94 mmol, 0.4 mol per mole of **1a**) were successively added with stirring. In the course of the <sup>65</sup> reaction homogeneous mixture transform to heterogeneous. The mixture was stirred at 20-25°C for 48 h (in run 20 for 72 h).
- mixture was stirred at 20-25°C for 48 h (in run 20 for 72 h). Compound **2a** was isolated as described above (in run 14,15).

#### Experiments for Table 2. Synthesis of bisperoxides 2b-i.

<sup>70</sup> In a 15 ml round bottom flask, compound **1b-1i** (0.573-0.873 g, 4.85 mmol) was dissolved in CH<sub>3</sub>CN (10 mL). Then a 70% aqueous t-BuOOH solution (1.87 g, 14.55 mmol; 3 mol per mole of **1b-1i**) and Mn(OAc)<sub>3</sub>•2H<sub>2</sub>O (0.13 g, 0.485 mmol; 0.1 mol per mole of **1b-1i**) were successively added with stirring. The <sup>75</sup> heterogeneous reaction mixture was stirred at 20-25°C for 48 h. Compounds **2b-i** were isolated as described for **2a**.

#### [1,2-Bis(tert-butylperoxy)ethyl]benzene, 2a<sup>12,15</sup>

Colourless oil;  $R_f = 0.77$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR <sup>80</sup> (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.26 (s, 9H, t-BuOO), 1.27 (s, 9H, t-BuOO), 4.09 (dd, J = 11.5, 4.0 Hz, 1H, CH<sub>2</sub>), 4.25 (dd, J = 11.5, 7.6 Hz, 1H, CH<sub>2</sub>), 5.23 (dd, J = 7.6, 4.0 Hz, 1H, CH), 7.25-7.43 (m, 5H, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  26.3, 26.4, 76.8, 80.4, 80.5, 83.3, 127.0, 127.9, 128.2, 138.6; Anal. calcd. for s<sup>5</sup> C<sub>16</sub>H<sub>26</sub>O<sub>4</sub>: C 68.06, H 9.28; found: C 68.10, H 9.22; HRMS (ESI) calcd. for C<sub>16</sub>H<sub>26</sub>NaO<sub>4</sub> [M+Na<sup>+</sup>]: 305.1723, found: 305.1720.

#### [1,2-Bis(tert-butylperoxy)-1-methylethyl]benzene, 2b<sup>14,15</sup>

Colourless oil;  $R_f = 0.82$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR

<sup>90</sup> (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.21 (s, 9H, t-BuOO), 1.26 (s, 9H, t-BuOO), 1.67 (s, 3H, CH<sub>3</sub>), 4.25 (m, 2H, CH<sub>2</sub>O), 7.24-7.55 (m, 5H, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  21.9, 26.3, 26.6, 79.2, 79.4, 80.5, 82.9, 126.1, 127.1, 127.8, 143.0; Anal. calcd. for C<sub>17</sub>H<sub>28</sub>O<sub>4</sub>: C 68.89, H 9.52; found: C 69.12, H 9.74; HRMS (ESI) <sup>95</sup> calcd. for C<sub>17</sub>H<sub>28</sub>NaO<sub>4</sub> [M+Na<sup>+</sup>]: 319.1880, found: 319.1876.

#### 1-[1,2-Bis(tert-butylperoxy)ethyl]-4-methylbenzene, 2c

Colourless oil;  $R_f = 0.75$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.25 (s, 9H, t-BuOO), 1.26 (s, 9H, t-<sup>100</sup> BuOO), 2.36 (s, 3H, CH<sub>3</sub>), 4.08 (dd, J = 11.4, 4.0 Hz, 1H, CH<sub>2</sub>), 4.25 (dd, J = 11.4, 7.7 Hz, 1H, CH<sub>2</sub>), 5.19 (dd, J = 7.7, 4.0 Hz, 1H, CH), 7.17 (d, J = 8.1 Hz, ArH), 7.27 (d, J = 8.1 Hz, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  21.2, 26.3, 26.4, 76.8, 80.4, 80.6, 83.2, 127.0, 128.9, 135.5, 137.6; Anal. calcd. for C<sub>17</sub>H<sub>28</sub>O<sub>4</sub>: <sup>105</sup> C 68.89, H 9.52; found: C 69.24, H 9.60; HRMS (ESI) calcd. for C<sub>17</sub>H<sub>32</sub>NO<sub>4</sub> [M+NH<sub>4</sub><sup>+</sup>]: 314.2326, found: 314.2324.

#### 1-[1,2-Bis(tert-butylperoxy)ethyl]-4-tert-butylbenzene, 2d

Colourless oil; R<sub>f</sub> = 0.8 (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR <sup>110</sup> (300.13 MHz, CDCl<sub>3</sub>): δ 1.25 (s, 9H, t-BuOO), 1.27 (s, 9H, t-BuOO), 1.32 (s, 9H, t-BuOO), 4.09 (dd, *J* = 11.4, 4.0 Hz, 1H, CH<sub>2</sub>), 4.25 (dd, *J* = 11.4, 7.7 Hz, 1H, CH<sub>2</sub>), 5.20 (dd, *J* = 7.7, 4.0 Hz, 1H), 7.30 (d, *J* = 8.1 Hz, ArH), 7.37 (d, *J* = 8.1 Hz, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>): δ 26.3, 26.4, 31.3, 34.5, 76.9, 80.4, <sup>115</sup> 80.6, 83.2, 125.2, 126.7, 135.5, 150.7; Anal. calcd. for C<sub>20</sub>H<sub>34</sub>O<sub>4</sub>: C 70.97, H 10.12; found: C 70.99, H 10.15; HRMS (ESI) calcd. for C<sub>20</sub>H<sub>34</sub>NaO<sub>4</sub> [M+Na<sup>+</sup>]: 361.2349, found: 361.2335.

#### 1-[1,2-Bis(tert-butylperoxy)ethyl]-4-methoxybenzene, 2e

Colourless oil;  $R_f = 0.51$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR <sup>5</sup> (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.24 (s, 9H, t-BuOO), 1.25 (s, 9H, t-BuOO), 3.81 (s, 3H, CH<sub>3</sub>O), 4.08 (dd, *J* = 11.4, 4.3 Hz, 1H, CH<sub>2</sub>), 4.26 (dd, *J* = 11.4, 7.5 Hz, 1H, CH<sub>2</sub>), 5.15 (dd, *J* = 7.5, 4.3 Hz, 1H, CH), 6.89 (d, *J* = 8.6 Hz, 2H, ArH), 7.29 (d, *J* = 8.6 Hz, 2H, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  26.3, 26.4, 55.2, 76.7, <sup>10</sup> 80.4, 80.5, 82.8, 113.7, 128.4, 130.7, 159.4; Anal. calcd. for C<sub>17</sub>H<sub>28</sub>O<sub>5</sub>: C 65.36, H 9.03; found: C 65.41, H 9.07; HRMS (ESI) calcd. for C<sub>17</sub>H<sub>32</sub>NO<sub>5</sub> [M+NH<sub>4</sub><sup>+</sup>]: 330.2275, found: 330.2275.

#### 1-[1,2-Bis(tert-butylperoxy)ethyl]-2-methoxybenzene, 2f

<sup>15</sup> Colourless oil;  $R_f = 0.55$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.26 (s, 9H, t-BuOO), 1.28 (s, 9H, t-BuOO), 3.83 (s, 3H, OCH<sub>3</sub>), 4.08 (dd, J = 11.1, 7.9 Hz, 1H, CH<sub>2</sub>), 4.17 (dd, J = 11.1, 3.1 Hz, 1H, CH<sub>2</sub>), 5.60 (dd, J = 11.1, 7.9 Hz, 1H, CH), 6.84-6.89 (m, 1H, ArH), 6.94-7.01 (m, 1H, ArH), 7.23-<sup>20</sup> 7.29 (m, 1H, ArH), 7.44-7.48 (m, 1H, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  26.3, 26.4, 55.3, 76.0, 78.8, 80.4, 80.6, 110.4, 120.4, 126.7, 127.8, 128.7, 156.7; Anal. calcd. for C<sub>17</sub>H<sub>28</sub>O<sub>5</sub>: C 65.36, H 9.03; found: C 65.31, H 9.11; HRMS (ESI) calcd. for C<sub>17</sub>H<sub>32</sub>NO<sub>5</sub> [M+NH<sub>4</sub><sup>+</sup>]: 330.2275, found: 330.2277.

#### 25

#### [1,2-Bis(*tert*-butylperoxy)-1-phenylethyl]benzene, 2g<sup>14</sup>

Colourless oil;  $R_f = 0.71$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.11 (s, 9H, t-BuOO), 1.23 (s, 9H, t-BuOO), 4.83 (s, 2H, CH<sub>2</sub>), 7.23-7.35 (m, 10H, ArH); <sup>13</sup>C NMR <sup>30</sup> (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  26.3, 26.6, 77.6, 79.6, 80.5, 85.7, 127.3, 127.5, 127.8, 142.1; Anal. calcd. for C<sub>22</sub>H<sub>30</sub>O<sub>4</sub>: C 73.71, H 8.44; found: C 73.69, H 8.50; HRMS (ESI) calcd. for C<sub>22</sub>H<sub>30</sub>NaO<sub>4</sub> [M+Na<sup>+</sup>]: 381.2036, found: 381.2031.

#### 35 1-[1,2-Bis(tert-butylperoxy)ethyl]naphthalene, 2h

Colourless oil;  $R_f = 0.77$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR (300.13 MHz, CDCl<sub>3</sub>):  $\delta = 1.31$  (s, 9H, t-BuOO), 1.34 (s, 9H, t-BuOO), 4.22-4.33 (m, 2H, CH<sub>2</sub>), 6.08-6.15 (m, 1H, CH), 7.46-7.59 (m, 3H, ArH), 7.69-7.76 (m, 1H, ArH), 7.78-7.93 (m, 2H, 40 ArH), 8.18-8.24 (m, 1H, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$ 

- 26.3, 26.5, 77.0, 80.5, 80.7, 80.8, 123.2, 124.4, 125.4, 126.0, 128.2, 128.8, 130.9, 133.8, 134.2; Anal. calcd. for  $C_{20}H_{28}O_4$ : C 72.26, H 8.49; found: C 72.38, H 8.30; HRMS (ESI) calcd. for  $C_{20}H_{28}NaO_4$  [M+Na<sup>+</sup>]: 355.1880, found: 355.1869.
- 45

#### 1-[1,2-Bis(tert-butylperoxy)ethyl]-4-chlorobenzene, 2i

Colourless oil;  $R_f = 0.69$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR (300.13 MHz, CDCl<sub>3</sub>):  $\delta$  1.23 (s, 9H, t-BuOO), 1.24 (s, 9H, t-BuOO), 4.03 (dd, J = 11.7, 4.2 Hz, 1H, CH<sub>2</sub>), 4.19 (dd, J = 11.7,

<sup>50</sup> 7.5 Hz, 1H, CH<sub>2</sub>), 5.19 (dd, J = 7.5, 4.2 Hz, 1H, CH), 7.26-7.35 (m, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  26.2, 26.3, 76.4, 80.5, 80.7, 82.6, 128.4, 128.5, 133.6, 137.2; anal. calcd. for C<sub>16</sub>H<sub>25</sub>ClO<sub>4</sub>: C 60.66, H 7.95, Cl 11.19, found: C 60.44, H 7.62, Cl 11.20; HRMS (ESI) calcd. for C<sub>16</sub>H<sub>29</sub>ClNO<sub>4</sub> [M+NH<sub>4</sub><sup>+</sup>]: <sup>55</sup> 334.1780, found: 334.1774.

#### 2-Phenyloxirane

Colourless oil;  $R_f = 0.39$  (ethyl acetate/hexane = 5:95); <sup>1</sup>H NMR

(300.13 MHz, CDCl<sub>3</sub>):  $\delta$  2.82 (dd, J = 5.50, 2.57 Hz, 1H, CH<sub>2</sub>), <sup>60</sup> 3.16 (dd, J = 5.50, 4.04 Hz, 1H, CH<sub>2</sub>), 3.88 (dd, J = 4.04, 2.57 Hz, 1H, CH), 7.23-7.45 (m, ArH); <sup>13</sup>C NMR (75.48 MHz, CDCl<sub>3</sub>):  $\delta$  51.1, 52.2, 125.4, 128.1, 128.4, 137.6.

#### Acknowledgment

This work was supported by the Russian Science Foundation <sup>65</sup> (Grant 14-23-00150). We thank to Prof. Sergey V. Popkov for useful discussions.

#### Notes and references

- <sup>a</sup> N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of
- 70 Sciences, Leninsky Prospect 47, Moscow, 119991 (Russian Federation) Fax: (+7)499-1355328
  - E-mail: terentev@ioc.ac.ru
  - <sup>b</sup> D.I. Mendeleev University of Chemical Technology of Russia, 9
  - Miusskaya square, Moscow, 125047 (Russian Federation)
- 75 † Electronic Supplementary Information (ESI) available: analytical data for peroxides 2a-2i. See DOI: 10.1039/b000000x/
  - (a) Ed. W. Ando, Organic peroxides, N.Y., Wiley, 1992; (b) Ed. S. Patai, The chemistry of peroxides, N.Y., Wiley, 1983; (c) Ed. V. Adam, Peroxide chemistry, N.Y., Wiley-VCH, 2000; (d) E.T. Denisov, T.G. Denisova, T.S. Pokidova, Handbook of Free Radical Initiators, John Wiley & Sons, Inc., 2005. (ISBN: 9780471207535).
  - (a) A. Bunge, H.-J. Hamann, J. Liebscher, Tetrahedron Lett., 2009, 50, 524-526; (b) B. Das, B. Veeranjaneyulu, M. Krishnaiah, P. Balasubramanyam, J. Mol. Catal. A, 2008, 284, 116-119; (c) P. Ghorai, P. H. Dussault, Org. Lett., 2008, 10, 4577-4579; (d) A. O. Terent'ev, M. M. Platonov, Yu. N. Ogibin, G. I. Nikishin, Synth. Commun., 2007, 37, 1281-1287; (e) A. O. Terent'ev, M. M. Platonov, A. V. Kutkin, Central Europ. J. Chem., 2006, 4, 207-215; (f) D. Azarifar, K. Khosravi, F. Soleimanei, Synthesis, 2009, 15, 2553-2556; (g) A. G. Griesbeck, D. Blunk, T. El-Idreesy, A. Raabe, Angew. Chem. Int. Ed., 2007, 46, 8883-8886; (h) A. O. Terent'ev, M. M. Platonov, I. B. Krylov, V. V. Chernyshev, G. I. Nikishin, Org. Biomol Chem., 2008, 6, 4435-4441; (i) A. O. Terent'ev, A. V. Kutkin, N. A. Troizky, Yu. N. Ogibin, G. I. Nikishin, Synthesis, 2005, 13, 2215-2219; (j) M. Ravi, D. Anand, R. Maurya, P. Chauhan, N. K. Naikade, S. K. Shukla, P. P. Yadav, Synlett, 2013, 24, 173-176; (k) A. Bartoschek, T. T. El-Idreesy, A. G. Griesbeck, L.-O. Hoeinck, J. Lex, C. Miara, J. M. Neudoerfl, Synthesis, 2005, 14, 2433-2444; (1) B.A. Solaja, N. Terzic, G. Pocsfalvi, L. Gerena, B. Tinant, D. Opsenica, W. K. Milhous, J. Med. Chem., 2002, 45, 3331-3336.
  - 3 (a) C. W. Jefford, Curr. Top. Med. Chem., 2012, 12, 373–399; (b) R. D. Slack, A. M. Jacobine, G. H. Posner, MedChemComm., 2012, 3, 281–297; (c) D. M. Opsenica, B. A. Šolaja, J. Serb. Chem. Soc., 2009, 74, 1155–1193; (d) P. Ghorai, P. H. Dussault, C. Hu, Org. Lett., 2008, 10, 2401–2404; (e) Y. Dong, Mini-Rev. Med.Chem., 2002, 2, 113–123; (f) J. L. Vennerstrom, H.-N. Fu, W. Y. Ellis, A. L. Ager, J. K. Wood, S. L. Andersen, L. Gerena, W. K. Milhous, J. Med. Chem., 1992, 35, 3023–3027; (g) N. Yadav, C. Sharma, S. K. Awasthi, RSC Adv., 2014, 4, 5469–5498; (h) A. O. Terent'ev, D. A. Borisov, I. A. Yaremenko, Chem. Heterocycl. Compd., 2013, 19, 7605–7619; (j) Y. Li, H.-D. Hao, S. Wittlin, Y. Wu, Chem. Asian. J., 2012, 7, 1881–1886.
  - J. Boissier, J. Portela, V. Pradines, F. Coslédan, A. Robert, B. Meunier, C. R. Chim., 2012, 15, 75–78; (b) J. Keiser, V. Veneziano, L. Rinaldi, L. Mezzino, U. Duthaler, G. Cringoli, Res. Vet. Sci., 2010, 88, 107–110; (c) X. Shuhua, M. Tanner, E. K. N'Goran, J. Utzinger, J. Chollet, R. Bergquist, C. Minggang, Z. Jiang, Acta Trop., 2002, 82,

175–181; (d) K. Ingram, I. A. Yaremenko, I. B. Krylov, L. Hofer, A. O. Terent'ev, J. Keiser, J. Med. Chem., 2012, **55**, 8700–8711.

- 5 (a) N. Kumar, M. Sharma, D. S. Rawat, Curr. Med. Chem., 2011, 18, 3889–3928; (b) Ž. Žižak, Z. Juranić, D. Opsenica, B. A. Šolaja, Investigational New Drugs, 2009, 27, 432–439; (c) N. Terzić, D. Opsenica, D. Milić, B. Tinant, K. S. Smith, W. K. Milhous, B. A. Šolaja, J. Med. Chem., 2007, 50, 5118–5127; (d) M. Jung, H. Kim, K. Lee, M. Park, Mini-Rev. Med. Chem., 2003, 3, 159–165; (e) J. Kim, E. J. Park, Curr. Med. Chem. Anticancer Agents, 2002, 2, 485–537.
- 6 (a) R. Matyáš, P. Jiří, Primary explosives, Germany: Springer, 2013; (b)
  J. Chen, W. Wu, A. J. McNeil. Chem. Commun., 2012, 48, 7310–7312; (c) F. Dubnikova, R. Kosloff, J. Almog, Y. Zeiri, R. Boese, H. Itzhaky, A. Alt, E. Keinan, J. Am. Chem. Soc., 2005, 127, 1146–1159; (d) C. Denekamp, L. Gottlieb, T. Tamiri, A. Tsoglin, R. Shilav, M. Kapon, Org. Lett., 2005, 7, 2461–2464; (e) R. Matyáš, R. Jirásko, A. Lyčka, J. Pachmáň, Propellants, Explosives, Pyrotechnics., 2011, 36, 219–224; (f) H. Östmark, S. Wallin, H. G. Ang, Propellants, Explosives, Pyrotechnics., 2011, 36, 219–224; (f) H. Östmark, S. Wallin, H. G. Ang, Propellants, Explosives, Pyrotechnics., 2012, 37, 12–23; (g) A. O. Terent'ev, M. M. Platonov, E. J. Sonneveld, R. Peschar, V. V. Chernyshev, Z. A. Starikova, G. I. Nikishin, J. Org. Chem., 2007, 72, 7237–7243; (h) A. O. Terent'ev, I. A. Yaremenko, V. A. Vil', I. K. Moiseev, S. A. Kon'kov, V. M. Dembitsky, D. O. Levitsky, G. I. Nikishin, Org. Biomol. Chem., 2013, 11, 2613–2623.
- 7 (a) M. S. Kharasch, P. Pauson, W. Nudenberg, J. Org. Chem., 1953, 18, 322–327; (b) M. S. Kharasch, A. Fono, J. Org. Chem., 1958, 23, 324–325; (c) M. S. Kharasch, A. Fono, J. Org. Chem., 1959, 24, 72–78; (d) M. S. Kharasch and G. Sosnovsky, Tetrahedron, 1958, 3, 105-112.
- G. B. Shul'pin, J. Gradinaru, Y. N. Kozlov, Org. Biomol. Chem., 2003, 1, 3611–3617; (b) S. Araneo, F. Fontana, F. Minisci, F. Recupero, A. Serri, J. Chem. Soc., Chem. Commun., 1995, 1399– 1400; (c) A. Bravo, H.-R. Bjørsvik, F. Fontana, L. Liguori, F. Minisci, J. Org. Chem., 1997, 62, 3849–3857; (d) M. B. Meder, L. H. Gade, Eur. J. Inorg. Chem., 2004, 2004, 2716–2722; (e) Z. Li, C.-J. Li, J. Am. Chem. Soc., 2005, 127, 6968–6969; (f) T. Punniyamurthy, L. Rout, Coord. Chem. Rev., 2008, 252, 134–154; (g) A. O. Terent'ev, D. A. Borisov, I. A. Yaremenko, V. V. Chernyshev, G. I. Nikishin, J. Org. Chem., 2010, 75, 5065–5071; (h) A. O. Terent'ev, D. A. Borisov, V. V. Semenov, V. V. Chernyshev, V. M. Dembitsky, G. I. Nikishin, Synthesis, 2011, 2011, 2091–2100.
- 9 (a) W. Treibs, G. Pellmann, *Chem. Ber.*, 1954, **87**, 1201–1205; (b) L. Saussine, E. Brazi, A. Robine, H. Mimoun, J. Fischer, R. Weiss, *J. Am. Chem. Soc.*, 1985, **107**, 3534–3540.
- 10 (a) R. A. Leising, R. E. Norman, L. Que Jr., *Inorg. Chem.*, 1990, 29, 2553–2555; (b) R. A. Leising, Y. Zang, L. Que Jr., *J. Am. Chem. Soc.*, 1991, 113, 8555–8557; (c) I. W. C. E. Arends, K. U. Ingold, D. D. M. Wayner, *J. Am. Chem. Soc.*, 1995, 117, 4710–4711; (d) W. Liu, Y. Li, K. Liu, Z. Li, *J. Am. Chem. Soc.*, 2011, 133, 10756–10759; (e) K. Liu, Y. Li, X. Zheng, W. Liu, Z. Li, *Tetrahedron*, 2012, 68, 10333–10337.
- 11 F. Minisci, F. Fontana, S. Araneo, F. Recupero, J. Chem. Soc., Chem. Commun., 1994, 1823–1824.
- 12 M. T. Rispens, O. J. Gelling, A. M. de Vries, A. Meetsma, F. van Bolhuis, B. L. Feringa, *Tetrahedron*, 1996, **52**, 3521–3546.
- 13 (a) J. Q. Yu, E. J. Corey, Org. Lett., 2002, 4, 2727–2730; (b) G. An,
   W. Zhou, G. Zhang, H. Sun, J. Han, Y. Pan, Org. Lett. 2010, 12, 4482–4485.
- 14 T. Nishimura, T. Onoue, K. Ohe, J. I. Tateiwa, S. Uemura, *Tetrahedron Lett.*, 1998, **39**, 4359–4362
- 15 F. Minisci, F. Fontana, S. Araneo, F. Recupero, S. Banfi, S. Quici, J. Am. Chem. Soc., 1995, 117, 226–232.
- 16 (a) S.-I. Murahashi, D. Zhang, Chem. Soc. Rev., 2008, 37, 1490–1501;
  (b) S.-I. Murahashi, T. Naota, T. Kuwabara, T. Saito, H. Kumobayashi, S. Akutagawa, J. Am. Chem. Soc., 1990, 112, 7820–7822;
  (c) S.-I. Murahashi, T. Naota, K. Yonemura, J. Am. Chem. Soc., 1988, 110, 8256–8258.

- (a) M. G. Vinogradov, P. A. Direy, G. I. Nikishin, J. Org. Chem. USSR, 1977, 13, 2498–2504; (b) A. Nakatani, K. Hirano, T. Satoh, M. Miura, J. Org. Chem., 2014, 79, 1377–1385; (c) K. Hattori, A. Ziadi, K. Itamiab, J. Yamaguchi, Chem. Commun., 2014, 50, 4105– 4107; (d) M. Tobisu, K. Koh, T. Furukawa, N. Chatani, Angew. Chem. Int. Ed., 2012, 51, 11363–11366. (e) G. Bar, A. F. Parsons, C. B. Thomas, Org. Biomol. Chem., 2003, 1, 373–380; (f) C. Tanyeli, D. Özdemirhan, Tetrahedron Lett., 2003, 44, 7311–7313; (g) C. Gardrat, Bull. Soc. Chim. Belges, 1984, 93, 897–902; (h) A. Citterio, R. Santi, T. Fiorani, S. Strologo, J. Org. Chem., 1989, 54, 2703–2712; (i) Z. Cong, T. Miki, O. Urakawa, H. Nishino. J. Org. Chem. 2009, 74, 3978–3981; (j) Y. Ito, Y. Tomiyasu, T. Kawanabe, K. Uemura, Y. Ushimizu, H. Nishino, Org. Biomol. Chem., 2011, 9, 1491–1507; (k) O. Alagöz, M. Yılmaz, A. T. Pekel, C. Graiff, R. Maggi, RSC Adv., 2014, 4, 14644–14654.
- 18 (a) T. Tsubusaki, H. Nishino, Tetrahedron, 2009, 65, 3745-3752; (b) A. S. Demir, O. Reis, C. Igdir, Tetrahedron, 2004, 60, 3427-3432; (c) A. O. Terent'ev, I. B. Krylov, V. P. Timofeev, Z. A. Starikova, V. M. Merkulova, A. I. Ilovaisky, G. I. Nikishin, Adv. Synth. Catal., 2013, 355, 2375-2390; (d) I. B. Krylov, A. O. Terent'ev, V. P. Timofeev, B. N. Shelimov, R. A. Novikov, V. M. Merkulova, G. I. Nikishin, Adv. Synth. Catal., 2014, 356, 2266-2280; (e) V. P. Ananikov, L. L. Khemchyan, Yu. V. Ivanova, V. I. Bukhtiyarov, A. M. Sorokin, I. P. Prosvirin, S. Z. Vatsadze, A. V. Medved'ko, V. N. Nuriev, A. D. Dilman, V. V. Levin, I. V. Koptyug, K. V. Kovtunov, V. V. Zhivonitko, V. A. Likholobov, A. V. Romanenko, P. A. Simonov, V. G. Nenajdenko, O. I. Shmatova, V. M. Muzalevskiy, M. S. Nechaev, A. F. Asachenko, O. S. Morozov, P. B. Dzhevakov, S. N. Osipov, D. V. Vorobyeva, M. A. Topchiy, M. A. Zotova, S. A. Ponomarenko, O. V. Borshchev, Y. N. Luponosov, A. A. Rempel, A. A. Valeeva, A. Yu. Stakheev, O. V. Turova, I. S. Mashkovsky, S. V. Sysolyatin, V. V. Malykhin, G. A. Bukhtiyarova, A. O. Terent'ev, I. B. Krylov, Russ. Chem. Rev., 2014, 83, 885-985; (f) T. Rahman, A. Haque, H. Igarashi, H. Nishino. Molecules 2011, 16, 9562-9581; (g) C.-Y. Qian, H. Nishino, K. Kurosawa. J. Heterocyclic Chem., 1993, 30, 209-216.
- 19 (a) W. E. Fristad, T. A. Brandvold, J. R. Peterson, S. R. Thompson, J. Org. Chem., 1985, 50, 3647–3649; (b) B. B. Snider, J. R. Duvall, Org. Lett., 2004, 6, 1265–1268.
- 20 (a) T. K. M. Shing, Y. Y. Yeung, P. L. Su, Org. Lett., 2006, 8, 3149–3151; (b) F. D. J. Cortez, D. Lapointe, A. M. Hamlin, E. M. Simmons, R. Sarpong, Tetrahedron, 2013, 69, 5665-5676.
- 21 H. Paul, R. D. Small Jr, J. C. Scaiano, J. Am. Chem. Soc., 1978, 100, 4520–4527.
- 22 P. A. Belyakov, V. I. Kadentsev, A. O. Chizhov, N. G. Kolotyrkina, A. S. Shashkov, V. P. Ananikov, *Mendeleev Commun.*, 2010, 20, 125–131.