

Stereocontrolled Formal Synthesis of
($\pm$)-Platensimycin


Jun-ichi Matsuo,* Kosuke Takeuchi, and Hiroyuki Ishibashi

Division of Pharmaceutical Sciences, Graduate School of Natural Science and
Technology, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

jimatsuo@p.kanazawa-u.ac.jp

Received July 12, 2008

ABSTRACT




The caged structure of platensimycin, known as Nicolaou's key intermediate for total synthesis of platensimycin, was synthesized stereoselectively by using the following key steps: (i) diastereoselective Diels–Alder reaction between γ -benzoyloxy enone and *tert*-butylidimethylsiloxydiene, (ii) formation of a dihydropyran ring by intramolecular catalytic oxypalladation, and (iii) transannular radical cyclization of monothioacetal with tributyltin hydride and AIBN.

Platensimycin (**1**), which was isolated from *Streptomyces platensis* by Merck's research group in 2006, has potent, broad-spectrum Gram-positive antibacterial activity and exhibits no cross-resistance to antibiotic-resistant bacteria, including methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant enterococci (VRE), and linezolid-resistant and macrolide-resistant pathogens.¹ Platensimycin (**1**) selectively inhibits FabF, an elongation condensing enzyme of fatty-acid biosynthesis, which is highly conserved among key pathogens. In addition to the novelty in the biological activity of **1**, the characteristic caged structure of **1** has attracted the interest of organic chemists, and several research groups have reported total and formal synthesis of **1**.^{2,3}

We planned a stereocontrolled synthesis of **2**, a key intermediate in Nicolaou's total synthesis of **1**,^{2a} as shown

in Scheme 1. We expected that the C12–C15 bond of **2**

Scheme 1. Retrosynthesis of **2**


(1) (a) Wang, J.; Soisson, S. M.; Young, K.; Shoop, W.; Kodali, S.; Galgoci, A.; Painter, R.; Parthasarathy, G.; Tang, Y. S.; Cummings, R.; Ha, S.; Dorso, K.; Motyl, M.; Jayasuriya, H.; Ondeyka, J.; Herath, K.; Zhang, C.; Hernandez, L.; Allocco, J.; Basilio, A.; Tormo, J. R.; Genilloud, O.; Vicente, F.; Pelaez, F.; Colwell, L.; Lee, S. H.; Michael, B.; Felcetto, T.; Gill, C.; Silver, L. L.; Hermes, J. D.; Bartizal, S. B.; Barrett, J.; Schmatz, D.; Becker, J. W.; Cully, D.; Singh, S. B. *Nature* **2006**, *441*, 358. (b) Singh, S. B.; Jayasuriya, H.; Ondeyka, J. G.; Herath, K. B.; Zhang, C.; Zink, D. L.; Tsou, N. N.; Ball, R. G.; Basilio, A.; Genilloud, O.; Diez, M. T.; Vincente, F.; Pelaez, F.; Young, K.; Wang, J. *J. Am. Chem. Soc.* **2006**, *128*, 11916.

would be formed by transannular radical cyclization⁴ of α -alkoxy radical **3** and that a six-membered cyclic ether ring of **3** would be constructed by palladium-catalyzed intramolecular alkenyl ether formation from **5** to **4**. We

envisioned that compound **5** would be accessible by diastereoselective Diels–Alder reaction of enone **6** and siloxydiene **7**.⁶ The key feature of our plan is that all stereocenters in **2** are controlled by the stereochemistry presented in **6**. Thus, stereochemistries at C8 and C9 are controlled by the diastereoselective Diels–Alder reaction, while those at C12 and C15 are determined inevitably by the transannular cyclization.

O-TBS and *O*-benzoyl-protected enones **12a,b** were prepared from 1,3-cyclohexadiene (**8**) as shown in Scheme 2. Allylboration of **8** with allyldibromoborane⁷ gave **9** in

Scheme 2. Preparation of Enone **12a,b**



71% yield after oxidative workup. The homoallylic double bond of **9** was selectively oxidized with VO(acac)₂ and *tert*-butyl hydroperoxide⁸ to afford epoxide **10** in 89% yield. Oxidation of the hydroxy group of **10** with Dess–Martin periodinane⁹ gave the corresponding β,γ -epoxy ketone and successive isomerization to γ -hydroxy enone **11** took place

readily in the presence of silica gel.¹⁰ Other oxidation methods such as Swern oxidation¹¹ and PDC¹² gave inferior results because of the lability of formed β,γ -epoxy ketone under the reaction conditions. Protection of the hydroxy group of **11** with TBSCl and benzoyl chloride gave **12a** and **12b**, respectively.

Next, the Diels–Alder reaction of thus-prepared dienophiles **12a,b** with siloxydienes **13a,b** was investigated (Table 1).¹³ The Diels–Alder reaction between TBS-protected enone **12a** and trimethylsiloxydiene **13a** in the presence of pyridine and 2,6-di-*tert*-butyl-4-methylphenol (BHT) at 200 °C (in toluene, sealed tube) gave the corresponding Diels–Alder adducts (**14a** and **15a**) in 32% yield with low diastereoselectivity (**14a**:**15a** = 4:1) after acidic treatment (entry 1). The yield and selectivity were improved by employing benzoyl-protected enone **12b** instead of **12a**, presumably due to the lower LUMO level of **12b** (entry 2). After screening various reaction conditions, it was found that the Diels–Alder reaction between **12b** and *tert*-butyldimethylsiloxydiene **13b** at 180 °C without any solvents gave the desired Diels–Alder adduct **14b** and its diastereomer **15b** in 83% combined yield in a ratio of **14b**:**15b** = 11:1 after treatment with trifluoroacetic acid (entry 3). This further improvement of chemical yield and diastereoselectivity might be explained by better thermal stability of **13b** than that of **13a** and by effective steric repulsion between the benzoyloxy group of **12b** and TBS group of **13b** in the transition state of the Diels–Alder reaction. The mixture of two diastereomers (**14b** and **15b**) was employed in the next step because of difficulty in separation at this stage.

Selective protection of the less-hindered carbonyl group by Noyori's procedure¹⁴ followed by hydrolysis of the benzoyl group and separation of diastereomers gave **16** in 80% yield as a single diastereomer. Catalytic oxypalladation of **16** by using a catalytic amount of palladium(II) chloride and copper(II) acetate under oxygen atmosphere in dimethylacetamide¹⁵ gave **17a** and **17b** in 73% combined yield (**17a**:**17b** = 10:1).¹⁶ The mixture of isomers was employed in the following steps.

To form a carbon–carbon bond between C12 and C15 by radical cyclization, we needed to prepare alkene **19** from ketone **17**. Reduction of the keto group of **17** followed by dehydration of the resulting alcohol via mesylate did not give **19**. It was found that vinyl triflate **18**, which was prepared from **17** with KHMDS and *N*-(5-chloro-2-pyridyl)triflimide,¹⁷

(2) (a) Nicolaou, K. C.; Li, A.; Edmonds, D. J. *Angew. Chem., Int. Ed.* **2006**, *45*, 7086. (b) Nicolaou, K. C.; Edmonds, D. J.; Li, A.; Tria, G. S. *Angew. Chem., Int. Ed.* **2007**, *46*, 3942. (c) Li, P.; Payette, J. N.; Yamamoto, H. J. *Am. Chem. Soc.* **2007**, *129*, 9534. (d) Lalic, G.; Corey, E. J. *Org. Lett.* **2007**, *9*, 4921. (e) Nicolaou, K. C.; Pappo, D.; Tsung, K. Y.; Gibe, R.; Chen, D. Y.-K. *Angew. Chem., Int. Ed.* **2008**, *47*, 944. (f) Kim, C. H.; Jang, K. P.; Choi, S. Y.; Chung, Y. K.; Lee, E. *Angew. Chem., Int. Ed.* **2008**, *47*, 4009. (g) Zou, Y.; Chen, C.-H.; Taylor, C. D.; Foxman, B. M.; Snider, B. B. *Org. Lett.* **2007**, *9*, 1825. (h) Nicolaou, K. C.; Tang, Y.; Wang, J. *Chem. Commun.* **2007**, 1922. (i) Tiefenbacher, K.; Mulzer, J. *Angew. Chem., Int. Ed.* **2007**, *46*, 8074. (j) Tiefenbacher, K.; Mulzer, J. *Angew. Chem., Int. Ed.* **2008**, *47*, 2548.

(3) Total synthesis of platencin, a platensimycin analogue: (a) Nicolaou, K. C.; Tria, G. S.; Edmonds, D. J. *Angew. Chem., Int. Ed.* **2008**, *47*, 1780. (b) Hayashida, J.; Rawal, V. H. *Angew. Chem., Int. Ed.* **2008**, *47*, 4373. (c) Tiefenbacher, K.; Mulzer, J. *Angew. Chem., Int. Ed.* **2008**, *47*, 6199. (d) Yun, S. Y.; Zheng, J.-C.; Lee, D. *Angew. Chem., Int. Ed.* **2008**, *47*, 6201.

(4) Transannular cyclization of cyclooct-4-enyl radical: (a) Dowbenko, R. J. *Am. Chem. Soc.* **1964**, *86*, 946. (b) Dowbenko, R. *Tetrahedron* **1964**, *20*, 1843. (c) Friedman, L. J. *Am. Chem. Soc.* **1964**, *86*, 1885. (d) Gancarz, R. A.; Kice, J. J. *J. Org. Chem.* **1981**, *46*, 4899. (e) Winkler, J. D.; Sridar, V. J. *Am. Chem. Soc.* **1986**, *108*, 1708.

(5) (a) Rawal, V. H.; Singh, S. P.; Dufour, C.; Michoud, C. J. *Org. Chem.* **1991**, *56*, 5245. (b) Mesmaeker, A. D.; Waldner, A.; Hoffmann, P.; Mindt, T.; Hug, P.; Winkler, T. *Synlett* **1990**, 687.

(6) Danishefsky, S.; Kitahara, T. J. *Am. Chem. Soc.* **1974**, *96*, 7807.

(7) Frantz, D. E.; Singleton, D. A. *Org. Lett.* **1999**, *1*, 485.

(8) Sharpless, K. B.; Michaelson, R. C. *J. Am. Chem. Soc.* **1973**, *95*, 6136.

(9) Dess, D. B.; Martin, J. C. J. *Am. Chem. Soc.* **1991**, *113*, 7277.

(10) Isomerization of β,γ -epoxycyclohexanones: Delay, F.; Ohloff, G. *Helv. Chim. Acta* **1979**, *62*, 2168.

(11) Mancuso, A. J.; Swern, D. *Synthesis* **1981**, 165.

(12) (a) Corey, E. J.; Schmidt, G. *Tetrahedron Lett.* **1979**, 399. (b) Herscovici, J.; Egron, M.-J.; Antonakis, K. J. *Chem. Soc., Perkin Trans. 1* **1982**, 1967.

(13) Some examples for the Diels–Alder reaction between 2-alkyl-2-cyclohexen-1-ones and **13a**: (a) Danishefsky, S.; Kitahara, T. *J. Org. Chem.* **1975**, *40*, 538. (b) Inokuchi, T.; Okano, M.; Miyamoto, T.; Madon, H. B.; Takagi, M. *Synlett* **2000**, 1549.

(14) Tsunoda, T.; Suzuki, M.; Noyori, R. *Tetrahedron Lett.* **1980**, *21*, 1357.

(15) (a) Smith, A. B.; Cho, Y. S.; Friestad, G. K. *Tetrahedron Lett.* **1998**, *39*, 8765. (b) Hosokawa, T.; Hirata, M.; Murahashi, S.-I.; Sonoda, A. *Tetrahedron Lett.* **1976**, 1821.

(16) Semmelhack, M. F.; Kim, C. R.; Dobler, W.; Meier, M. *Tetrahedron Lett.* **1989**, *30*, 4925.

Table 1. The Diels–Alder Reaction Between **12a,b** and **13a,b**


12a: R = TBS **13a:** R' = TMS
12b: R = Bz **13b:** R = TBS

14a,b **15a,b**
14a: R = TBS **15a:** R = TBS
14b: R = Bz **15b:** R = Bz

entry	enone	diene (equiv)	conditions	yield (%) ^a	14:15
1	12a	13a (4.0)	(1) py (1.0 equiv), BHT ^b (cat.), toluene, 200 °C, 34 h (2) 0.01 N aq HCl	32	14a:15a = 4:1
2	12b	13a (4.0)	(1) py (1.0 equiv), BHT ^b (cat.), toluene, 180 °C, 25 h (2) 0.01 N aq HCl	60	14b:15b = 8:1
3	12b	13b (2.5)	(1) neat, 180 °C (2) TFA, CH ₂ Cl ₂	83	14b:15b = 11:1

^a Combined yield of **14** and **15**. ^b 2,6-Di-*tert*-butyl-4-methylphenol.


was reduced to **19** in good yield by using a palladium catalyst, formic acid, and tributylamine (Scheme 3).¹⁸

Scheme 3. Preparation of **19** by Palladium-Catalyzed Alkenyl Ether Formation

Reduction of vinyl triflate **18** by the use of tributyltin hydride and palladium(0)¹⁹ gave alkene **19** in only 37% yield.

Monothioacetal **20**, a precursor for radical cyclization, was prepared by reaction of alkenyl ether **19** and benzenethiol at 50 °C without any solvents (Scheme 4).²⁰ Transannular radical cyclization of monothioacetal **20**²¹ was accomplished by using tributyltin hydride in the presence of AIBN in refluxing toluene. Subsequent deprotection of the acetal group gave **2** in 57% yield in two steps along with compound **22**

(18% yield), which was formed by reduction of α-alkoxy radical intermediate with tributyltin hydride followed by acid

Scheme 4. Transannular Radical Cyclization of **20** to **2**

hydrolysis. It was also observed by TLC analysis that elimination of the phenylthio group of **20** to **19** proceeded in refluxing toluene. Tributyltin hydride was found to be a better hydrogen source in this transannular radical cyclization than tris(trimethylsilyl)silane and triphenyltin hydride. At the outset, we tried to prepare the phenylselenoacetal for radical cyclization.²² However, the preparation of the phenylselenoacetal by the reaction of **19** with benzeneselenol²³ gave compound **21**, which might be formed by reductive cleavage of a tentatively introduced benzeneselenenyl group.²⁴

In summary, we have established a stereocontrolled synthesis of the platensimycin core **2** with the following key findings. (1) Diastereoselective Diels–Alder reaction proceeded efficiently by the combination of enone **12b** and siloxydiene **13b**. The steric and electronic factors of the benzoyl group of **12b** and TBS group of **13b** significantly affected the Diels–Alder reaction. (2) Intramolecular alkenyl ether formation to construct a dihydropyran ring was performed by oxypalladation with a catalytic amount of

(17) (a) Comins, D. L.; Dehghani, A.; Foti, C. J.; Joseph, S. P. *Org. Synth.* **1997**, 74, 77. (b) Comins, D. L.; Dehghani, A. *Tetrahedron Lett.* **1992**, 33, 6299.

(18) Cacchi, S.; Morera, E.; Ortar, G. *Tetrahedron Lett.* **1984**, 25, 4821.

(19) Scott, W. J.; Crisp, G. T.; Stille, J. K. *J. Am. Chem. Soc.* **1984**, 106, 4630.

(20) When the addition of benzenethiol to the alkenyl ether moiety was carried out by employing a catalytic amount of PPTS, deprotection of the acetal group of **19** also took place.

(21) Reductive radical cyclization of monothioacetal: (a) Yadav, V. K.; Fallis, A. G. *Tetrahedron Lett.* **1988**, 29, 897. (b) Sato, K.; Sasaki, M. *Org. Lett.* **2005**, 7, 2441.

(22) Reaction of 1-alkoxyalkyl radicals from selenoacetal: (a) Rawal, V. H.; Singh, S. P.; Dufour, C.; Michoud, C. *J. Org. Chem.* **1991**, 56, 5245. (b) Mesmaeker, A. D.; Waldner, A.; Hoffmann, P.; Mindt, T.; Hug, P.; Winkler, T. *Synlett* **1990**, 687. (c) Nishiyama, Y.; Yamamoto, H.; Nakata, S.; Ishii, Y. *Chem. Lett.* **1993**, 841.

(23) Braga, A. L.; Silveira, C. C.; Zeni, G.; Severo Filho, W. A.; Stefani, H. A. *J. Chem. Res. (S)* **1996**, 206.

(24) Lee, J. S.; Fuchs, P. L. *Org. Lett.* **2003**, 5, 2247.

palladium(II) chloride and copper(II) acetate under oxygen atmosphere. (3) The caged structure was constructed by transannular radical cyclization of conformationally rigid monothioacetal **20** by using tributyltin hydride and AIBN in refluxing toluene with minimization of the elimination of benzenethiol from **20** to **19** and reduction of **20** to **21**. This convergent synthesis of **2** will be useful for the synthesis of platensimycin analogues which might improve the pharmacokinetic properties²⁵ of **1**.

Acknowledgment. This study was financially supported by the Uehara Memorial Foundation and a Grant-in-Aid for

Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

Supporting Information Available: Experimental procedures and spectroscopic data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL801584R

(25) Singh, S. B.; Herath, K. B.; Wang, J.; Tsou, N.; Ball, R. G. *Tetrahedron Lett.* **2007**, 48, 5429.