

# MedChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: T. V. Volkova, I. V. Terekhova, O. Silyukov, A. N. Proshin, A. Bauer-Brandl and G. L. Perlovich, *Med. Chem. Commun.*, 2016, DOI: 10.1039/C6MD00545D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



## ARTICLE

# Towards the rational design of novel drugs based on solubility, partitioning/distribution, biomimetic permeability and biological activity exemplified by 1,2,4-thiadiazole derivatives<sup>†</sup>

T. V. Volkova<sup>a</sup>, I. V. Terekhova<sup>a</sup>, O. I. Silyukov<sup>a</sup>, A. N. Proshin<sup>b</sup>, A. Bauer-Brandl<sup>c</sup>, G. L. Perlovich<sup>a,\*</sup>

Received 00th January 20xx

Accepted 00th January 20xx



DOI: 10.1039/x0xx00000x

www.rsc.org/



Novel 1,2,4-thiadiazole derivatives as potent neuroprotectors were synthesized and identified. Their ability to inhibit the glutamate stimulated  $\text{Ca}^{2+}$  uptake was investigated. The solubility of thiadiazoles was measured in buffer solution (pH 7.4) at 298 K. The distribution coefficients in 1-octanol/buffer (pH 7.4) and 1-hexane/buffer (pH 7.4) immiscible phases as model systems imitating gastrointestinal tract epithelium and blood-brain barrier were determined. Permeation experiments across new Permeapad<sup>TM</sup> barrier using Franz diffusion cells were conducted and the apparent permeability coefficients were obtained. Influence of the compounds structure on the physicochemical properties determining the bioavailability of drug-like substances was revealed. Solubility-permeability interplay has been assessed to evaluate potential bioavailability of the compounds studied.

## 1. Introduction

In the opinion of leading experts, Alzheimer's disease is one of the most frequent and progressive condition among elderly people, comparable by occurrence with cardiac and cerebral infarctions [1]. As the etiology of Alzheimer's disease (Alzheimer type dementia) has not been found yet, there is no rational standard therapy of the disease. Currently, worldwide investigations in the sphere of neurodegenerative drugs design are performing mainly in the following directions: invention of neuroprotectors efficacious in terms of pathogenesis of the disease and capable to block a pathogenic mechanism, particularly,

effective multitarget drugs [2]; searching for new targets participating in the course of the disease. Moreover, considering business purposes, repositioning the well-known formulations to novel application is very attractive. During the creation of new neuroprotective agents fast screening of a large number of structures which exhibit biological activity at the receptors comes forward. Investigation of the drug delivery processes (transport properties) is an important stage in the creation of bioavailable drugs. Therefore, obtaining the screening parameters characterizing solubility, partitioning, and membrane permeability is an essential part of drug design.

The 1,2,4-thiadiazole derivatives are well known to have the biologically active groups to specific targets of different pathologies. Many of them reveal a wide spectrum of biological activity including antioxidant and neuroprotective ones [3-6]. Analysis of the influence of 1,2,4-thiadiazole derivatives on the glutamate-induced calcium ions uptake into synaptosomes of rat brain cortex is often used for primary evaluation of the biological activity of newly synthesized drug compounds [7]. The glutamate receptors in the central nervous system are known to play an important role both in regular neurophysiological processes and pathogenesis of a series of neurodegenerative diseases [8]. In this connection, the evaluation of a compound's

<sup>a</sup> Department of Physical Chemistry of Drugs, Krestov's Institute of Solution Chemistry, Russian Academy of Sciences, 153045 Akademicheskaya str. 1, Ivanovo, Russia. E-mail: glp@isc-ras.ru

<sup>b</sup> Institute of Physiologically Active Compounds, Russian Academy of Sciences, 142432, Chernogolovka, Russia.

<sup>c</sup> Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, 5230 Odense M, Denmark

<sup>†</sup>The authors declare no competing interests.

Electronic Supplementary Information (ESI) available: Dependence of content of the ionized forms of the molecules on the pH of buffer solution of compound **2** (Fig. 1SI); Relationships of the permeability coefficients with the distribution coefficients (Fig. 2SI); Relationship of the permeability coefficient with physicochemical descriptor HYBOT (Fig. 3SI). This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ARTICLE

## Journal Name

ability to affect the calcium channels mediated by the glutamate receptors allows to assess their common biological potential as possible neuroprotectors (in the case of inhibiting) or activators of cognition functions (in the case of activation). Members of a newly discovered class of compounds, ampakines, boost the activity of glutamate and thus make it easier to form memories. The cognition enhancers work on the neural processes that underlie such mental activities as attention, perception, learning, memory, language, planning and decision-making, usually by altering the balance of the chemical neurotransmitters involved in these processes. At present, nootropics and some neuronotrophic agents represent the available approaches to symptomatic treatment of Alzheimer's disease [9]. Attempts to improve cognitive function in patients with brain disorders have become the focus of intensive research efforts [10]. The task of developing guidelines for creating effective modulators of brain cognitive functions based on the derivatives of 1,2,4-thiadiazole is relevant from the perspective of the creation of new class of effective drugs for the treatment of Alzheimer's disease.

Lipophilicity is a property that has a major effect on *in vivo* processes of absorption, distribution, metabolism, excretion, and toxicity (ADMET) properties as well as pharmacological activity [11]. The partition coefficient is accepted by most scientists to be one of the most relevant lipophilicity descriptors applied in pharmaceutical research [12]. It can be measured by a rather simple experiment or calculated. Lipophilicity has been correlated to many other properties, such as solubility, permeability, metabolism, toxicity, protein binding, and distribution. 1-Octanol is the preferred organic solvent for phase-distribution investigations or partition coefficient determination of drugs. The 1-octanol/water partition coefficient is reported in drug handbooks and is commonly used in various types of analysis of drug properties [13]. The  $\log D^{7.4}$  ( $\log D^{oct/buf}$  in the present study) of a compound stands for its distribution coefficient at pH 7.4, and it is considered to be a property of utmost importance because of its high physiological relevance and its resemblance to real biological partitions in the body [14]. As it has been emphasized in [15], the implication of  $\log D^{7.4}$  for drug development leads to a general rule for optimal intestinal absorption by passive diffusion permeability after oral dosing. Drug substances should have a moderate  $\log D^{7.4}$ . Namely, a good balance of permeability and solubility exists when  $\log D^{7.4} = 0 - 3$ . At that, an optimum range for CNS (central nervous system) narrows to  $\log D^{7.4} = 1 - 3$ . Since all drug compounds for the prevention of Alzheimer's disease need to overcome the blood-brain barrier (BBB), the analysis of drug interactions with these barriers is an important objective in the design of effective drugs. As a rule, the water/alkane systems (usually 1-hexane/water distribution coefficients  $\log D^{hex/w}$ ) are used for BBB simulation. Selecting the alkane phase is dictated by the fact that it is a good description of non-specific interactions

occurring between the drug molecules and the rather tight and lipophilic BBB environment. A parameter describing the difference between the distribution coefficients in 1-octanol/water and 1-hexane/water phases is often used to account for specific interactions (donor-acceptor, hydrogen bonding) to drug transport properties [16].


In the present study we propose the synthesis of new 1,2,4-thiadiazole derivatives with different substitution of the phenyl ring and replacement of the phenyl ring by cyclopropyl- and isoxazole-one. The aim focuses at investigating the biological activity of the synthesized 1,2,4-thiadiazoles and revealing correlations between the structures, solubilities, lipophilicities and membrane permeabilities of the respective substances. The present work is a continuation of our previous attempts to study the passive transport properties of 1,2,4-thiadiazoles [17-23] and the relationship between the biological activity and transport characteristics of drugs and drug-like compounds [23,24].

## 2. Materials and methods

### 2.1. Synthesis

#### 2.1.1. General Procedure for Preparing Novel 1,2,4-thiadiazoles

Synthetic approaches to novel 1,2,4-thiadiazoles were carried out according to Scheme 1.



**Scheme 1.** Synthesis of the compounds studied.

5-*N*-Monoaminosubstituted 5-amino-3-(2-aminopropyl)-[1,2,4]thiadiazoles (f) were synthesized in the following matter:

A solution of 3-amino-5-methylisoxazole (a) (0.01 M) in 10 mL of acetonitrile was added drop wise to a solution of isothiocyanate (0.01 M) in 10 mL of acetonitrile under stirring. At that, the obtained isoxazole thiourea (b) in situ

transformed to [1,2,4]thiadiazole ring according to Boulton-Katritsky [25]. Upon completion of the dropping the mixture was brought to boiling and left at room temperature until the precipitation of 5-amino-3-(2-oxopropyl)-[1,2,4]thiadiazole (c) occurs. The precipitate was filtered off. If the crystallization was not taking place the reaction mixture was evaporated and the resultant oil was triturated in diethyl ether.

The resulting 5-amino-3-(2-oxopropyl)-[1,2,4]thiadiazole (c) (0.01 M) and primary amine (0.01 M) was dissolved in 10 ml of methanol and left one day at room temperature. The precipitate was filtered off. According to NMR spectroscopy, the imine (d) forming during the reaction was rearranged to the corresponding vinylamine 5-amino-3-(2-amino-1-propenyl)-[1,2,4]thiadiazole (e) which is introduced into the reduction reaction.

The resulting 5-amino-3-(2-amino-1-propenyl)-[1,2,4]thiadiazole (e) (0.01 M) was suspended in 30 mL of

methanol, heated to 50 °C. Then sodium borohydride (0.01 M) was added portionwise under vigorous stirring. Under the addition of sodium borohydride the reaction was passed and the precipitate was dissolved. Upon the completion of the reaction, methanol was evaporated, 50 mL of methylene chloride was added and washed with water (2 x 50 mL), and then an organic layer was separated, and dried over sodium sulfate. The drier was filtered off, and the filtrate was evaporated to obtain 5-amino-3-(2-hydroxypropyl)-[1,2,4]thiadiazole (f).

### 2.1.2. Chemical Procedures

5-*N*-Monoaminosubstituted 5-amino-3-(2-aminopropyl)-[1,2,4]thiadiazoles were synthesized by standard and accessible methods in the Institute of Physiologically Active Compounds of the Russian Academy of Sciences. The compounds synthesized are presented in Table 1.

**Table 1** Structures of studied compounds

Group I		Group II					
No	R-	No	R-				
1		7		13		18	
2		8		14		19	
3		9		15		20	
4		10		16		21	
5		11		17		22	
6		12					

*N*-[1-[5-(3,4-Dichlorophenylamino)-[1,2,4]thiadiazol-3-yl]-propan-2-yl]-(2,2,6,6-tetramethylpiperidine-4-yl)amine (**1**).

White powder. Yield 67%. Calcd, %: C 54.29, H 6.61, N 15.83 C<sub>20</sub>H<sub>29</sub>Cl<sub>2</sub>N<sub>5</sub>S. Found, %: C 54.67, H 6.66, N 15.48. NMR <sup>1</sup>H [200 MHz] δ: 0.79 (2H, dt, J = 12.0, 21.2 Hz, C(3)HH, C(5)HH), 1.08 and 1.12 (both s, 6 H, 2 CH<sub>3</sub>), 1.15 (3H, d, J = 6.6 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.84 (2H, dd, J = 3.3, 12.6 Hz, C(3)HH, C(5)HH), 2.84 (1H, dd, J = 6.0, 14.2 Hz, CHHCHMe), 2.92 (1H, dd, J = 4.0, 14.2 Hz, CHHCHMe), 3.00 (1H, m, C(4)H), 3.42 (1H, m, CH<sub>2</sub>CHMe), 7.17 (1H, dd, J = 2.2, 8.6 Hz, H<sub>arom</sub>), 7.45 (1H, d, J = 8.6 Hz, H<sub>arom</sub>), 7.51 (1H, d, J = 2.2 Hz, H<sub>arom</sub>).

*2*-[5-(4-Methyl-3-chlorophenylamino)-[1,2,4]thiadiazol-3-yl]-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**2**).

White powder. Yield 77%. Calcd, %: C 59.77, H 7.64, N 16.59 C<sub>21</sub>H<sub>32</sub>ClN<sub>5</sub>S. Found, %: 59.87, H 7.34, N 16.44. NMR <sup>1</sup>H [200 MHz] δ: 0.65 (1H, t, J = 12.0 Hz, CHH), 0.75 (1H, t, J = 12.0 Hz, CHH), 1.00 (6H, s, (CH<sub>3</sub>)<sub>2</sub>), 1.06 (3H, d, J = 6.3 Hz, CCH<sub>3</sub>), 1.10 (6H, s, (CH<sub>3</sub>)<sub>2</sub>), 1.71 (2H, dd, J = 3.5, 12.3 Hz, 2CH), 2.31 (3H, s, Ph-CH<sub>3</sub>), 2.71 (2H, dd, J = 6.3, 14.2 Hz, CH<sub>2</sub>), 2.89 (1H, tt, J = 3.5, 11.6 Hz, NCH), 3.32 (1H, m, NCH), 7.17 (1H, d, J = 8.4 Hz, H<sub>arom</sub>), 7.29 (1H, dd, J = 2.2, 8.4 Hz, H<sub>arom</sub>), 7.71 (1H, d, J = 2.2 Hz, H<sub>arom</sub>).

*2*-[5-(4-Fluoro-3-chlorophenylamino)-[1,2,4]thiadiazol-3-yl]-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**3**).

White powder. Yield 67%. Calcd, %: C 56.39, H 6.86, N 16.44 C<sub>20</sub>H<sub>29</sub>ClFN<sub>5</sub>S. Found, %: C 56.26, H 6.81, N 16.55. NMR <sup>1</sup>H [200 MHz] δ: 0.77 (2H, dt, J = 12.1, 21.0 Hz, C(3)HH, C(5)HH), 1.06 and 1.14 (12H, c and c, 4xCH<sub>3</sub>), 1.15 (3H, d, J = 6.6 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.81 (2H, dd, J = 3.4, 12.6 Hz, C(3)HH, C(5)HH), 2.83 (1H, dd, J = 6.0, 14.0 Hz, CHHCHMe), 2.97 (1H, dd, J = 4.1, 14.0 Hz, CHHCHMe), 3.00 (1H, m, C(4)H), 3.40 (1H, m, CH<sub>2</sub>CHMe), 7.16 (2H, m, H<sub>arom</sub>), 7.45 (1H, d, J = 8.0 Hz, H<sub>arom</sub>).

*2*-[5-(2-Methyl-5-chlorophenylamino)-[1,2,4]thiadiazol-3-yl]-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**4**).

White powder. Yield 72%. Calcd, %: C 59.77, H 7.64, N 16.59 C<sub>21</sub>H<sub>32</sub>ClN<sub>5</sub>S. Found, %: C 59.80, H 7.30, N 16.49. C<sub>21</sub>H<sub>32</sub>ClN<sub>5</sub>S. Found, %: 59.87, H 7.34, N 16.44. NMR <sup>1</sup>H [200 MHz] δ: 0.71 (1H, t, J = 12.0 Hz, CHH), 0.85 (1H, t, J = 12.0 Hz, CHH), 1.12 (6H, s, (CH<sub>3</sub>)<sub>2</sub>), 1.14 (3H, d, J = 6.2 Hz, CCH<sub>3</sub>), 1.19 (6H, s, (CH<sub>3</sub>)<sub>2</sub>), 1.85 (2H, dd, J = 3.5, 12.3 Hz, 2CH), 2.32 (3H, s, Ph-CH<sub>3</sub>), 2.831 (2H, m, CH<sub>2</sub>), 2.99 (1H, tt, J = 3.5, 11.6 Hz, NCH), 3.41 (1H, m, NCH), 7.17 (2H, m, H<sub>arom</sub>), 7.52 (1H, s, H<sub>arom</sub>).

*2*-[5-(4-Chlorophenylamino)-[1,2,4]thiadiazol-3-yl]-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**5**).

White powder. Yield 70%. Calcd, %: C 58.88, H 7.41, N 17.16 C<sub>20</sub>H<sub>30</sub>ClN<sub>5</sub>S. Found, %: C 58.80, H 7.40, N 17.34. NMR <sup>1</sup>H [200 MHz] δ: 0.86 (2H, m, C(3)HH, C(5)HH), 1.14 and 1.20 (12H, s and s, 4xCH<sub>3</sub>), 1.27 (3H, d, J = 6.65 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.87 (2H, d, J = 12.5 Hz, C(3)HH, C(5)HH), 2.84

(2H, d, J = 6.5 Hz, CH<sub>2</sub>CHMe), 3.03 (1H, m, C(4)H), 3.45 (1H, m, CH<sub>2</sub>CHMe), 7.40 and 7.50 (4H, d and d, J = 8.4 Hz, H<sub>arom</sub>). [1-Methyl-2-(5-*p*-tolylamino)-[1,2,4]thiadiazol-3-yl]-ethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**6**).

Light crystals. Yield 69%. Calcd, %: C 65.08, H 8.58, N 18.07. C<sub>21</sub>H<sub>33</sub>N<sub>5</sub>S. Found, %: C 65.01, H 8.52, N 18.09. NMR <sup>1</sup>H [200 MHz] δ: 0.71 (2H, m, C(3)HH, C(5)HH), 1.02, 1.05 (6H, s + s, 2xCH<sub>3</sub>), 1.07 (3H, d, J = 6.5 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.12, 1.14 (6H, s + s, 2xCH<sub>3</sub>), 1.75 (2H, dd, J = 3.1, 12.5 Hz, C(3)HH, C(5)HH), 2.73 (2H, dd, J = 2.3, 6.5 Hz, CH<sub>2</sub>CHMe), 2.91 (1H, m, C(4)H), 3.32 (1H, m, CH<sub>2</sub>CHMe), 7.09 and 7.38 (4H, d + d, J = 8.6 Hz, H<sub>arom</sub>), 10.56 (1H, br.s, ArNH).

*2*-[5-(3-Methylthiophenylamino)-[1,2,4]thiadiazol-3-yl]-ethyl-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**7**).

White powder. Yield 70%. Calcd, %: C 60.10, H 7.93, N 16.69 C<sub>21</sub>H<sub>33</sub>N<sub>5</sub>S<sub>2</sub>. Found, %: C 60.28, H 7.99, N 16.64. NMR <sup>1</sup>H [200 MHz] δ: 0.81 (2H, m, C(3)HH, C(5)HH), 1.13 and 1.20 (12H, s and s, 2xCH<sub>3</sub>), 1.21 (3H, d, J = 6.5 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.86 (2H, d, J = 12.4 Hz, C(3)HH, C(5)HH), 2.54 (3H, s, Ph-SCH<sub>3</sub>), 2.88 (2H, m, CH<sub>2</sub>CHMe), 3.01 (1H, m, C(4)H), 3.47 (1H, m, CH<sub>2</sub>CHMe), 7.01 (2H, m, H<sub>arom</sub>), 7.34 (2H, m, H<sub>arom</sub>).

*2*-[5-(3-Methoxyphenylamino)-[1,2,4]thiadiazol-3-yl]-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**8**).

White powder. Yield 70%. Calcd, %: C 62.50, H 8.24, N 17.35 C<sub>21</sub>H<sub>33</sub>N<sub>5</sub>OS. Found, %: C 62.66, H 8.14, N 17.34. NMR <sup>1</sup>H [200 MHz] δ: 0.81 (2H, m, C(3)HH, C(5)HH), 1.09 and 1.17 (12H, c and c, 4xCH<sub>3</sub>), 1.20 (3H, d, J = 6.2 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.83 (2H, d, J = 12.2 Hz, C(3)HH, C(5)HH), 2.83 (2H, m, CH<sub>2</sub>CHMe), 2.97 (1H, m, C(4)H), 3.39 (1H, m, CH<sub>2</sub>CHMe), 3.81 (3H, c, Ph-OCH<sub>3</sub>), 6.73 (2H, m, H<sub>arom</sub>), 7.29 (2H, m, H<sub>arom</sub>).

*N*-[1-[5-(4-Fluorophenylamino)-[1,2,4]thiadiazol-3-yl]-propane-2-yl]-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**9**).

Light crystals. Yield 69%. Calcd, %: C 61.35, H 7.72, N 17.89 C<sub>20</sub>H<sub>30</sub>FN<sub>5</sub>S. Found, %: C 61.23, H 7.87, N 17.92. NMR <sup>1</sup>H [200 MHz] δ: 0.71 (2H, m, C(3)HH, C(5)HH), 1.02, 1.04 (12H, s, 4xCH<sub>3</sub>), 1.07 (3H, d, J = 6.6 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.72 (2H, dd, J = 3.0, 12.6 Hz, C(3)HH, C(5)HH), 2.72 (2H, dd, J = 2.3, 6.6 Hz, CH<sub>2</sub>CHMe), 2.91 (1H, m, C(4)H), 3.34 (1H, m, CH<sub>2</sub>CHMe), 7.04 (2H, t, J = 8.6 Hz, H<sub>arom</sub>), 7.58 (2H, dd, J = 4.7, 9.1 Hz, H<sub>arom</sub>).

*2*-[5-(5-Methylisoxazol-3-ylamino)-[1,2,4]thiadiazol-3-yl]-1-methylethyl-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (**10**).

White powder. Yield 56%. Calcd, %: C 57.11, H 7.99, N 22.20, C<sub>18</sub>H<sub>30</sub>N<sub>6</sub>OS. Found, %: C 56.91, H 8.10, N 22.44. NMR <sup>1</sup>H [200 MHz] δ: 0.74 (2H, m, C(3)HH, C(5)HH), 1.00 and 1.10 (12H, c and c, 4xCH<sub>3</sub>), 1.12 (3H, d, J = 6.2 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.76 (2H, d, J = 12.2 Hz, C(3)HH, C(5)HH), 2.33 (1H, s, C=CCH<sub>3</sub>), 2.83 (2H, m, CH<sub>2</sub>CHMe), 3.00 (1H, m,



C(4)H), 3.42 (1H, m, CH<sub>2</sub>CHMe), 4.12 (2H, br.s, 2NH), 5.87 (1H, s, C=CH).

**2-[5-Cyclopropylamino-[1,2,4]thiadiazol-3-yl]-1-methylethyl)-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (11).**

White powder. Yield 63%. Calcd, %: C 60.49, H 9.26, N 20.75 C<sub>17</sub>H<sub>31</sub>N<sub>5</sub>S. Found, %: C 60.66, H 9.11, N 20.74. NMR <sup>1</sup>H [200 MHz] δ: 0.76 (6H, m, C(3)HH, C(5)HH, CH<sub>2</sub>CH<sub>2</sub>), 1.07 and 1.13 (12H, c and c, 4xCH<sub>3</sub>), 1.13 (3H, d, J = 6.35 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.79 (2H, m, C(3)HH, C(5)HH), 2.61 (1H, m, NCH), 2.75 (2H, m, CH<sub>2</sub>CHMe), 2.91 (1H, m, C(4)H), 3.33 (1H, m, CH<sub>2</sub>CHMe), 6.87 (1H, br.s, NH).

**N-{1-[5-(4-Bromophenylamino-[1,2,4]thiadiazol-3-yl)-propane-2-yl]}-(2,2,6,6-tetramethylpiperidine-4-yl)-amine (12).**

Yellow powder. Yield 72%. Calcd, %: C 53.09, H 6.68, N 15.48 C<sub>20</sub>H<sub>30</sub>BrN<sub>5</sub>S. Found, %: C 53.28, H 6.56, N 15.64. NMR <sup>1</sup>H [200 MHz] δ: 0.73 (2H, m, C(3)HH, C(5)HH), 1.02 and 1.05 (12H, c and c, 4xCH<sub>3</sub>), 1.08 (3H, d, J = 6.5 Hz, CH<sub>2</sub>CHCH<sub>3</sub>), 1.72 (2H, dd, J = 3.1, 12.5 Hz, C(3)HH, C(5)HH), 2.75 (2H, dd, J = 2.3, 6.5 Hz, CH<sub>2</sub>CHMe), 2.91 (1H, m, C(4)H), 3.33 (1H, m, CH<sub>2</sub>CHMe), 7.40 and 7.52 (4H, d and d, J = 8.8 Hz, H<sub>arom</sub>).

## 2.2. Materials and Solvents

1-Octanol (1-Octanol, CH<sub>3</sub>(CH<sub>2</sub>)<sub>7</sub>OH, MW 130.2, lot 11K3688) ARG was from Sigma Chemical Co (USA). 1-Hexane (C<sub>6</sub>H<sub>14</sub>, MW 86.18, lot 07059903 C) ARG was from SDS (Peypin, France).

Bidistilled water (with electrical conductivity 2.1 μS cm<sup>-1</sup>) was used for preparation of buffer solutions.

Phosphate buffer pH 7.4 (I=0.15 mol/L) for distribution and solubility experiments was prepared by combining potassium dihydrogen phosphate solution (KH<sub>2</sub>PO<sub>4</sub>) (9.1 g in 1 L) and disodium hydrogen phosphate dodecahydrate solution (Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O) (23.6 g in 1 L) salts [26]. The pH values were measured by using pH meter FG2-Kit (Mettler Toledo, Switzerland) standardized with pH 1.68, 6.86 and 9.22 solutions.

Phosphate buffer saline for permeation experiments was prepared as described in [27] by mixing a 2.5% (weight/volume) sodium dihydrogen phosphate dehydrate solution with a 1.8% (weight/volume) disodium hydrogen phosphate dodecahydrate solution in a ratio of 1 to 4 in order to obtain a 74 mM phosphate solution.

The pH of the solution was measured by pH meter FG2-Kit (Mettler Toledo, Switzerland) and adjusted to physiologic conditions (pH 7.3–7.4) by the addition of sodium hydroxide. The osmolality of the buffer solution was measured by Semi-Micro Osmometer K-7400 (Herbert Knauer GmbH, Berlin, Germany) and adjusted to 280–290 mOsm by the addition of NaCl.

Permeapad™ barrier was prepared according to the description of di Cagno and Bauer-Brandl [27]. In brief, a thin layer of phosphatidylcholine (S-100) was applied on a support sheet (PützGmbH, Taunusstein, Germany). The final barrier was composed of support layer and a dried layer of lipid. The final barrier appears mechanically flexible and resistant, and can be cut to size by scissors or a punching tool.

## 2.3. Solubility experiments

All the experiments were carried out by the shake flask method [28] at 298 K (± 0.1 K). The essence of the above mentioned method includes determination of the compound concentration in the saturated solution. Glass ampoules containing the test substance and the solvent were placed into an air thermostat supplied by a stirring device. To ensure the equilibration condition, the dissolution profiles of the compounds were investigated. Thermodynamic equilibrium was determined from preliminary experiments including measuring the kinetic dependences of the concentration of the saturated solutions for the investigated substances. The time needed for reaching the plateau of drug concentration against time was considered a suitable equilibration time. The incubation time of 24 h was estimated to be high enough for the equilibrium to be reached and all the solubility experiments lasted 24 h before the solubility measurements were performed. An aliquot of the saturated solution was taken and centrifuged under temperature control (Biofuge stratus, Germany) for 5 minutes at 298 K temperature. The solid phase was removed through isothermal filtration by the filter MILLEX®HA 0.45 μm (Ireland). The saturated solution was diluted with the corresponding solvent to the required concentration. The molar solubility of drugs was determined by means of spectrophotometer Cary-50 (USA) in UV spectral region (λ = 200 – 400 nm) with an accuracy of 2–4 %. The experimental results are reported as an average value of at least three replicated experiments.

## 2.4. Permeation studies

### 2.4.1. Experiment design

Permeapad™ barrier adapted to Franz diffusion cell (SES GmbH-Analytical Systems, Germany) was applied to the determination of the permeability coefficients ( $P_{app}$ ) as recommended by di Cagno et al. [27]. In our experiment the volume of the drug solution in the lower (donor) chamber was 8 ml (approximately 5 mM concentration) and the volume of the buffer solution in the upper (acceptor) chamber was 1.5 ml. A surface area of the Permeapad™ barrier disposed between the lower and upper chambers was 1 cm<sup>2</sup>. Aliquots of 0.5 mL were periodically withdrawn from the upper chamber and after each withdrawal

## ARTICLE

## Journal Name

replaced by 0.5 ml of fresh buffer. The amount of the permeated substance was analyzed spectrophotometrically using Cary-50 (USA) spectrophotometer at the appropriate wavelength. Permeability coefficients through the support layer (without phospholipids) were determined similarly to those measured through the Permeapad™ barrier. The permeation experiments were performed at least in triplicate for each compound and the final  $P_{app}$ -value was determined from the average of the individual  $P_{app}$  calculated from each replicate. All experiments were carried out at 298 K ( $\pm 0.1$  K).

#### 2.4.2. Data analysis

The calculation procedure includes the following consecutive steps. Firstly, the cumulative amount of the permeated substance (Q), expressed in number of moles was calculated for each time interval and plotted against the time (t), expressed in seconds, taking into account the surface area of the barrier. The flux (J) - the slope of the linear regression was determined by the following equation:

$$J = \frac{dQ}{A \times dt} \quad (1)$$

Secondly, the obtained flux was applied to calculate the apparent permeability coefficient ( $P_{app}$ ) using the following equation:

$$P_{app} = \frac{J}{C_0} \quad (2)$$

where  $C_0$  is the substance concentration in the donor compartment.

At last, based on the permeability coefficients measured through the Permeapad™ barrier ( $P_{app}$ ) and support layer ( $P_{sup}$ ) the permeability coefficient through the lipid layer ( $P_{lip}$ ) was determined by the following equation:

$$\frac{1}{P_{app}} = \frac{1}{P_{lip}} + \frac{1}{P_{sup}} \quad (3)$$

where the inverse values of the permeability coefficients are the resistances (R) of the components (lipid and support) of the Permeapad™ barrier.

Standard deviation calculations and student's t-test were evaluated,  $P \leq 0.05$  was considered as significantly different. Thomson Tau test was applied in order to identify possible outlier values.

#### 2.5. Distribution coefficients determination

The shake flask method was used to determine the distribution coefficients ( $D^{oct/buf}$ ) in 1-octanol/buffer (pH 7.4) system [31-34]. The procedure was performed as follows. Buffer (pH 7.4) and 1-octanol were mixed vigorously for 24 h at 298 K ( $\pm 0.1$  K), to promote solvent saturation in both phases. The solvents were standing long enough to allow the phases to separate, and the compound was dissolved in the 1-octanol phase to obtain stock solution. The buffer saturated 1-octanol phase with the dissolved substance, and 1-octanol saturated buffer phase were placed in a glass vials and mixed during 24 hours at 298 K. The substance concentrations were determined spectrophotometrically using calibration curves. The distribution coefficient was calculated by the following equation:

$$D^{oct/buf} = C^{oct/buf} / C^{buf/oct} \quad (4)$$

where  $C^{oct/buf}$  and  $C^{buf/oct}$  are the molar concentrations of solute in the mutually saturated phases of 1-octanol and buffer. The accuracy of the distribution coefficient value was verified by checking the mass balance of the starting amount of compound  $i$  compared to the total amount of the compound partitioned between the two phases:

$$m^i = m^{oct/buf} + m^{buf/oct} \quad (5)$$

where  $m^i = C^i \cdot V^i$  is the starting mass (in moles) of the compound,  $m^{oct/buf} = C^{oct/buf} \cdot V^{oct/buf}$  is the mass of the substance dissolved in the water-saturated 1-octanol phase, and  $m^{buf/oct} = C^{buf/oct} \cdot V^{buf/oct}$  is the mass of the substance dissolved in the 1-octanol saturated water phase.

#### 2.6. Calcium-blocking property experiments

Synaptosomes of the rat brain cortex were isolated according to a standard procedure from the brain of newborn rats (9-10 days old). For accumulation of radioactive label,  $P_2$ -fraction of synaptosomes was suspended in the incubation buffer A: 132 mM NaCl, 5 mM KCl, 5 mM HEPES, 10 mM glucose, pH 7.4 (protein concentration – 1.5-2 mg/mL). Calcium concentration in the final volume was 1.25 mM. Glutamate concentration of 200  $\mu$ M was applied for stimulation of  $^{45}\text{Ca}^{2+}$  uptake. After 5 min incubation at 335 K, the process was terminated by the filtration on GF/B-filters (Whatman, UK), washed three times with cold buffer B (145 mM HEPES, 10 mM Tris, 5.4 mM Trilon B, pH 7.4). All the experiments were performed in triplicate. The samples were measured by means of fluid scintillator beta-analyzer SL-4000 (TriCarb 2800, Perkin Elmer).

In preliminary experiments, all the compounds were tested for the ability to inhibit glutamate stimulated Ca uptake at the concentration of 100  $\mu$ M. If the inhibition of Glu-Ca-uptake was 50% and more, then further studies

were carried out to determine the concentration dependence of inhibition. The amount of the uptake in synaptosomes  $^{45}\text{Ca}^{2+}$  was determined by the difference of the label content with and without uptake stimulation and expressed in % control (control - 100%). The corresponding value of  $K_{43/21}$  was calculated according to the following equation:

$$K_{43/21} = [(Ca_4 - Ca_3)/(Ca_2 - Ca_1)] \cdot 100\% \quad (6)$$

where  $Ca_1$  is the  $\text{Ca}^{2+}$  influx in the blank experiment (without glutamate and test compounds);  $Ca_2$  is the  $\text{Ca}^{2+}$  influx in the presence of glutamate only (Glu-Ca-uptake);  $Ca_3$  is the  $\text{Ca}^{2+}$  influx in the presence of test compound (without glutamate);  $Ca_4$  is the  $\text{Ca}^{2+}$  influx in the presence of both glutamate and test compound.

## 2.7. Calculation procedure

All the descriptors were calculated by program package HYBOT-PLUS (version of 2003) in Windows [35].

Equilibrium constants for the compounds synthesized were calculated by ACD/ChemSketch program [36]. Analysis of the equilibrium of charged state of the molecules was conducted by using program package presented in [37].

## 2.8. DSC experiments

The fusion temperatures of the compounds under investigation have been determined using a Perkin-Elmer Pyris 1 DSC differential scanning calorimeter (Perkin-Elmer Analytical Instruments, Norwalk, Connecticut, USA) with Pyris software for Windows NT. DSC experiments were conducted in an atmosphere of flowing ( $20 \text{ cm}^3 \cdot \text{min}^{-1}$ ) dry helium gas of high purity 0.99996 (mass fraction) using standard aluminum sample pans and a heating rate of  $2 \text{ K} \cdot \text{min}^{-1}$ . The accuracy of weight measurements was  $0.005 \text{ mg}$ . The DSC was calibrated with an indium sample from Perkin-Elmer (P/N 0319-0033). The value determined for the fusion enthalpy corresponded to  $28.48 \text{ J} \cdot \text{g}^{-1}$  (the reference value was  $28.45 \text{ J} \cdot \text{g}^{-1}$ ). The fusion temperature was  $429.5 \pm 0.1 \text{ K}$  (determined from ten measurements).

## 3. Results and discussion

### 3.1. Distribution experiments

The processes of distribution of compounds in biological media play an important role in drug delivery. These processes contribute significantly to membrane permeability determining the concentrations of drug compounds in the membrane/lipophilic compartments (biological medium phases). In addition, the distribution coefficient of the compounds in 1-octanol/buffer

( $D^{\text{oct/buf}}$ ) and 1-hexane/buffer ( $D^{\text{hex/buf}}$ ) systems are important characteristic's commonly used as a measure to describe many biological, chemical and physical processes. For example,  $D^{\text{oct/buf}}$  is often used to estimate the processes of intestinal tract membrane permeability, whereas  $D^{\text{hex/buf}}$  - the processes occurring in the BBB. Moreover, the parameter characterizing the difference between the 1-hexane phase to 1-octanol one ( $\log D^{\text{oct/hex}}$ ) is often used to distinguish the contributions of non-specific and specific interactions (donor-acceptor, hydrogen bonds) of the drug molecules with biological environments. Based on this, we have tried to describe the distribution of the selected class of compounds in the assigned model systems of immiscible solvents. The experimental data for  $D^{\text{oct/buf}}$  and  $D^{\text{hex/buf}}$  are presented in Table 2. It should be noted that for almost all the compounds (except **14**)  $D^{\text{hex/buf}}$ -values are less than 1. This indicates that these substances predominantly dissolve in the aqueous phase. On the contrary, in the 1-octanol/buffer system,  $D^{\text{oct/buf}}$ -values exceed 1 (except **10**), therefore, the substances prefer the 1-octanol phase.

We studied the trends between  $D^{\text{oct/buf}}$  and selected physicochemical descriptors. The main purpose of such search was connected with an idea to develop an algorithm for prediction of the experimental data on the basis of just molecular structure knowledge. The HYBOT [35] descriptors were selected as independent parameters, because they describe the interaction of drug molecules with biological media. It should be noted that the access to obtain the correlation equation was complemented by  $D^{\text{oct/buf}}$ -values for the related compounds which were prepared in the present work (group I) and those which have been published by us previously (group II) [23] (Table 1). However, the compounds of group I are in the ionized state in buffer (pH 7.4), whereas drugs of group II - in the form of neutral entities. The appropriate calculation of the equilibrium forms of the compounds by using the approach described in [37] and taking into account the calculated  $\text{pK}_a$  ( $\text{pK}_{a1} = 14.01 - 14.04$ ;  $\text{pK}_{a2} = 1.29 - 4.67$ ) [36] confirms the neutral forms of the compounds of group II. For this group of compounds the equality is realized:  $K_0^{\text{oct/buf}} = D^{\text{oct/buf}}$  and  $K_0^{\text{hex/buf}} = D^{\text{hex/buf}}$ .

In order to recalculate the distribution coefficients, the ionization state of their molecules was determined taking into account that the presently investigated thiadiazoles (group I) are bases and have several ionizable groups. The macroscopic dissociation constants were calculated according to [36].

Based on the calculated  $\text{pK}_a$ -values we determined the content of a number of the forms with different degrees of protonation in aqueous solutions depending on the pH of the media by using [37] taking into account the following ionization processes:



**Table 2** Experimental solubility<sup>a</sup> values of the compounds studied in buffer (  $S_2^{298}$ ,  $X_2^{298}$  ), distribution coefficients 1-octanol/buffer (pH 7.4) (  $D^{oct/buf}$  ) and 1-hexane/buffer (pH 7.4) (  $D^{hex/buf}$  ), partitioning coefficients for neutral molecules in 1-octanol/buffer (  $K_0^{oct/buf}$  ) and 1-hexane/buffer (  $K_0^{hex/buf}$  ), biological coefficient describing ability to inhibit Glu-Ca-uptake (  $K_{43/21}$  ), melting points (  $T_m$  ), molecular polarizability (  $\alpha$  ), acceptor molecular ability to create hydrogen bonds (  $\Sigma(C_a)$  ), acceptor molecular ability to create hydrogen bonds normalized to molecular polarizability (  $\Sigma(C_a)/\alpha$  ).

	$S_2^{298}$ / mol·l <sup>-1</sup>	$X_2^{298}$ / mol fract	$D^{oct/buf}$	$K_0^{oct/buf}$	$D^{hex/buf}$	$K_0^{hex/buf}$	$K_{43/21}$ / %	$T_m$ / K	$\alpha$	$\Sigma(C_a)$	$\Sigma(C_a)/\alpha$
1	7.45×10 <sup>-3</sup>	1.35×10 <sup>-4</sup>	12.60 ± 0.02	8.28×10 <sup>4</sup>	0.011 ± 0.003	72.3	305 ± 18	347.2 ± 0.5	47.113	7.68	0.163
2	5.57×10 <sup>-3</sup>	1.00×10 <sup>-4</sup>	11.01 ± 0.03	7.31×10 <sup>4</sup>	0.0031 ± 0.0002	20.6	214 ± 12	319.3 ± 0.5	47.02	7.75	0.165
3	9.24×10 <sup>-3</sup>	1.67×10 <sup>-4</sup>	5.0 ± 0.4	3.25×10 <sup>4</sup>	0.0064 ± 0.0005	42.0	189 ± 14	333.5 ± 0.5	45.094	7.76	0.172
4	4.86×10 <sup>-3</sup>	8.77×10 <sup>-5</sup>	6.9 ± 0.2	4.56×10 <sup>4</sup>	0.017 ± 0.004	112.8	177 ± 9	334.2 ± 0.5	47.02	7.83	0.166
5	3.72×10 <sup>-3</sup>	6.71×10 <sup>-5</sup>	1.83 ± 0.04	1.22×10 <sup>4</sup>	0.011 ± 0.005	73.0	172 ± 3	323.2 ± 0.5	45.185	7.69	0.17
6	8.37×10 <sup>-3</sup>	1.51×10 <sup>-4</sup>	6.6 ± 0.4	4.44×10 <sup>4</sup>	0.013 ± 0.001	89.3	128.0 ± 0.1	320.2 ± 0.5	45.092	7.73	0.171
7	1.20×10 <sup>-2</sup>	2.17×10 <sup>-4</sup>	4.4 ± 0.2	2.92×10 <sup>4</sup>	0.014 ± 0.002	92.9	117 ± 2	309.1 ± 0.5	47.792	8.16	0.171
8	2.13×10 <sup>-2</sup>	3.87×10 <sup>-4</sup>	3.5 ± 0.3	2.34×10 <sup>4</sup>	0.013 ± 0.003	86.3	112 ± 25	314.7 ± 0.5	45.366	8.33	0.184
9	1.82×10 <sup>-2</sup>	3.30×10 <sup>-4</sup>	7.6 ± 0.3	5.06×10 <sup>4</sup>	0.024 ± 0.003	158.0	112 ± 12	317.6 ± 0.5	43.166	7.66	0.177
10	1.41×10 <sup>-2</sup>	2.55×10 <sup>-4</sup>	0.375 ± 0.005	2.41×10 <sup>3</sup>	0.041 ± 0.002	258.6	108 ± 6	342.6 ± 0.5	41.113	8.48	0.206
11	3.10×10 <sup>-1</sup>	6.19×10 <sup>-3</sup>	1.21 ± 0.02	3.19×10 <sup>3</sup>	0.097 ± 0.004	655.5	86 ± 14	341.4 ± 0.5	38.328	7.32	0.191
12	3.49×10 <sup>-3</sup>	6.29×10 <sup>-5</sup>	2.041 ± 0.004	1.36×10 <sup>4</sup>	0.013 ± 0.001	85.0	82 ± 5	318.5 ± 0.5	45.883	7.69	0.168
13	nd <sup>b</sup>	nd <sup>b</sup>	610 ± 30 <sup>c</sup>	6.10×10 <sup>2</sup>	0.17 ± 0.01 <sup>c</sup>	0.17	51 ± 9 <sup>c</sup>	nd	29.307	5.21	0.178
14	4.83×10 <sup>-4d</sup>	8.70×10 <sup>-6d</sup>	1038 ± 10 <sup>c</sup>	1.038×10 <sup>3</sup>	1.68±0.07 <sup>c</sup>	1.68	87 ± 2 <sup>c</sup>	402.3 ± 0.2 <sup>e</sup>	29.214	5.28	0.181
15	nd	nd	226 ± 5 <sup>c</sup>	2.26×10 <sup>2</sup>	0.16 ± 0.04 <sup>c</sup>	0.16	75 ± 6 <sup>c</sup>	nd	27.288	5.29	0.194
16	4.45×10 <sup>-4d</sup>	8.00×10 <sup>-6d</sup>	1049 ± 50 <sup>c</sup>	1.049×10 <sup>3</sup>	0.56 ± 0.01 <sup>c</sup>	0.56	88 ± 16 <sup>c</sup>	374.5 ± 0.2 <sup>e</sup>	29.214	5.35	0.183
17	4.836×10 <sup>-4d</sup>	8.30×10 <sup>-6d</sup>	380 ± 5 <sup>c</sup>	3.80×10 <sup>2</sup>	0.46 ± 0.05 <sup>c</sup>	0.46	89 ± 13 <sup>c</sup>	408.4 ± 0.2 <sup>e</sup>	27.379	5.22	0.191
18	2.19×10 <sup>-3f</sup>	3.95×10 <sup>-5f</sup>	445 ± 28 <sup>c</sup>	4.45×10 <sup>2</sup>	0.31 ± 0.02 <sup>c</sup>	0.31	85 ± 4 <sup>c</sup>	390.1 ± 0.2 <sup>f</sup>	27.286	5.26	0.193
19	nd	nd	94 ± 13 <sup>c</sup>	9.4×10	0.48 ± 0.02 <sup>c</sup>	0.48	86.7 ± 0.5 <sup>c</sup>	nd	29.986	5.68	0.19
20	3.04×10 <sup>-3d</sup>	5.48×10 <sup>-5d</sup>	205 ± 5 <sup>c</sup>	2.05×10 <sup>2</sup>	0.58 ± 0.03 <sup>c</sup>	0.58	108 ± 11 <sup>c</sup>	363.4 ± 0.2 <sup>e</sup>	27.56	5.93	0.215
21	3.40×10 <sup>-3g</sup>	6.13×10 <sup>-4g</sup>	316 ± 3 <sup>c</sup>	3.16×10 <sup>2</sup>	0.32 ± 0.02 <sup>c</sup>	0.32	88 ± 16 <sup>c</sup>	376.3 ± 0.2 <sup>g</sup>	25.36	5.18	0.204
22	nd	nd	45 ± 1 <sup>c</sup>	4.5×10	0.18 ± 0.01 <sup>c</sup>	0.18	109 ± 20 <sup>c</sup>	nd	23.307	6.01	0.258

<sup>a</sup> relative standard uncertainty for solubility values  $u_r(S_2^{298}) = 0.03$ ;  $u_r(X_2^{298}) = 0.03$

<sup>b</sup> nd – not determined;

<sup>c</sup> Ref.(23);

<sup>d</sup> Ref.(20);

<sup>e</sup> Ref.(19);

<sup>f</sup> Ref.(17);

<sup>g</sup> Ref. (18);.



The  $pK_{a1} = 10.95$  corresponds to the protonation process of a nitrogen atom of NH-group of a pipyridine ring,  $pK_{a2} = 7.31 - 7.36$  – to the protonation of a bridge NH-group and  $pK_{a3} = 0.45 - 3.83$  – to the protonation process of a nitrogen atom of a thiadiazole ring.  $K_1$ ,  $K_2$  and  $K_3$  are equilibrium constants of the mentioned equations. The ionized/unionized particle distributions (%) as a function of pH for the compound **2** (as an example) is presented in Fig. 1SI (Supporting Information). It is found that all the compounds have similar dissociation constant values and, respectively, the protonation degree. As follows from the Fig. 1SI all the compounds are completely ionized at pH 7.4 and their molecules exist as approximately equal shares in the form of mono- and di-cations.

For ionizable compounds the distribution coefficient describing the overall lipophilicity of drugs is the ratio of the concentrations of all molecular forms of compounds existing in equilibrium in the aqueous phase and uncharged particles in 1-octanol. For the investigated thiadiazoles the equilibrium distribution can be represented by an equation by which the apparent distribution coefficient ( $D^{oct/buf}$ ) can be experimentally determined:

$$D^{oct/buf} = [L]_o / ([L]_w + [HL^+]_w + [HL^{2+}]_w + [HL^{3+}]_w) \quad (10)$$

where  $[L]_o$  – concentration of the neutral form of the compound in 1-octanol,  $[L]_w$ ,  $[HL^+]_w$ ,  $[HL^{2+}]_w$ , and  $[HL^{3+}]_w$  – the concentration of particles in an aqueous buffer solution. In view of the above equilibrium constants the equation can be represented as follows:

$$D^{oct/buf} = [L]_o / ([L]_w + [L][H^+]/K_1 + [L][H^+]^2/K_2 + [L][H^+]^3/K_3) \quad (11)$$

Further transformation leads to the equation, that relates the intrinsic distribution coefficients to the experimentally determined apparent distribution coefficients, hydrogen ion concentration ( $H^+$ ) and the dissociation constants,  $K_1$ ,  $K_2$  and  $K_3$ :

$$K_0^{oct/buf} = D^{oct/buf} \cdot (1 + [H^+]/K_1 + [H^+]^2/K_2 + [H^+]^3/K_3) \quad (12)$$

To make the conditions of the distribution processes comparable, we recalculated the values of  $D^{oct/buf}$  and  $D^{hex/buf}$  for the neutral forms of the compounds of group I by equation (12) (in Table 2, these coefficients are designated as  $K_0^{oct/buf}$  and  $K_0^{hex/buf}$ ).

The data in Table 2 show that the calculated intrinsic distribution coefficients (normalized to neutral species) of the studied thiadiazoles significantly exceed (more than 4 times) the experimental apparent ones (belong to the ionized forms of the molecules). The estimated ratios indicate greater lipophilicity of the neutral particles compared with the charged ones as expected.

The  $K_0^{oct/buf}$ -values have been analyzed concerning 32 descriptors. The suitable correlation was obtained for the independent variables which include polarizability ( $\alpha$ ) and acceptor molecular ability to create hydrogen bonds ( $\Sigma(C_a)$ ). The results of the correlation analysis can be presented by equation:

$$\log K_0^{oct/buf} = (0.16 \pm 0.39) + (0.16 \pm 0.02) \cdot \alpha - (0.37 \pm 0.14) \cdot \Sigma(C_a) \quad (13)$$

$$R = 0.9631; \sigma = 0.30; F = 121.6; n = 22$$

With an increase in the polarizability of the molecule the distribution coefficient grows. In other words, non-specific interactions are shown in more extent in 1-octanol phase as compared with aqueous one. While the molecular ability to create hydrogen bonds increases, the distribution coefficient in 1-octanol/buffer system decreases. The acceptor ability of the molecule to hydrogen bond formation should be revealed both in aqueous and 1-octanol phases. However, as follows from the derived equation (13), implementation of such bonds in aqueous phase is favorable. Such phenomenon can be due to a considerably smaller size of the water molecule in comparison with 1-octanol one, resulting in better conjugation between the solute and solvent molecules in the solution and higher probability of hydrogen bond formation. Appropriate analysis was performed for the distribution processes (calculation of the distribution coefficients for the uncharged species) of the selected compounds in 1-hexane/buffer system. Like in the previous case,  $\log K_0^{hex/buf}$ -value correlates in a best way with molecular polarizability and acceptor molecular ability to create hydrogen bonds ( $\Sigma(C_a)$ ):

$$\log K_0^{hex/buf} = - (5.2 \pm 0.6) + (0.037 \pm 0.032) \cdot \alpha - (0.69 \pm 0.23) \cdot \Sigma(C_a) \quad (14)$$

$$R = 0.9355; \sigma = 0.47; F = 66.6; n = 22$$

A comparison of equations (13) and (14) shows that  $K_0^{hex/buf}$  is approximately two times more sensitive to the variation of  $\Sigma(C_a)$ -descriptor than  $K_0^{oct/buf}$ . Most probably, this phenomenon can be explained by the specific interactions of the dissolved molecule with the solvent both in 1-octanol and buffer phases in 1-octanol/buffer system. Therefore, the contribution of the descriptor is partly equalized. As opposed to this, in 1-hexane/buffer system the dissolved molecules interact specifically only with one phase (buffer) which leads to a higher regression coefficient value. Thus, correlation equations (13) and (14) allow predicting the distribution coefficients for the studied class of compounds both in 1-octanol/water and 1-hexane/water systems. It is interesting to compare the distribution coefficients for the molecules of group I compounds (normalized to the neutral species) with those of group II which differ from each other by only one fragment (tetramethylpiperidine-fragment (group I) instead of hydroxyl-

## ARTICLE

## Journal Name

group (group II)) both in 1-octanol/water and 1-hexane/water systems. The results of the comparative analysis are given in Fig. 1. In 1-octanol/water system the difference of the distribution coefficients between group I and group II compounds is positive with an average  $\Delta \log K_0^{oct/buf}$ -value of  $2.58 \pm 0.20$  (excluding the pair of compounds 10/22). A similar tendency for 1-hexane/water system with an average  $\Delta \log K_0^{hex/buf}$ -value of  $2.47 \pm 0.23$  (excluding the pair of compounds 2/14) is observed. The derived values of  $\Delta \log K_0$  can be determined by: firstly, conformational differences of the molecules of groups I and II in water and 1-octanol and also water and 1-hexane phases; secondly, variation of the energy of hydrogen bond forming by a hydroxyl-group in the considered phases.

Analysis of  $D^{oct/hex}$ -values showed the existence of strong specific interactions of the drug molecules with the 1-octanol ( $1.5 < \log D^{oct/hex} < 3.6$ ). On this ground it may be assumed that the redistribution of molecules from blood plasma through the blood-brain barrier will be aggravated.




Fig. 1. Comparative analysis of distribution coefficients for the molecules of group I compounds (normalized to the neutral particles) with those of group II differ from each other by only one fragment (tetra-methylpiperidine-fragment (group I) instead of hydroxyl-group (group II)) both in 1-octanol/buffer and 1-hexane/buffer systems

### 3.2. Solubility in buffer and melting temperatures

The results of solubility experiments in buffer (pH 7.4) at 298 K and melting points of the compounds studied are presented in Table 2.

It is well known that the nature and position of the substituent may influence essentially the solubility of the compounds in structure analogues series. In this connection, the comparative solubility analysis for the synthesized thiadiazoles was performed. It was found that the solubility among the investigated compounds with tetramethylpiperidine fragment differs within two orders of magnitude. Compound (11) with a cyclopropyl-substituent has the maximal solubility, and compounds (12) and (5) with 4-bromophenyl- and 4-chlorophenyl- substituents, respectively, – the minimal ones. The replacement of a

chlorine-atom in para-position of the phenyl-ring by a methyl-group (compound 6) increases the solubility twice, and by methoxy-group (compound 8) or fluorine-atom (compound 9) – about 6 times. Changing a methylphenyl-substituent to methylisoxazole one slightly enhances the solubility (about 1.5 times). Interestingly, the introduction of the second chlorine atom in the meta- position of the phenyl ring of compound 5 (3,4-dichloro-derivative (1)) leads to double the solubility. At that, the solubility values of the structure isomers 3-chloro-4-methyl- (2) and 3-chloro-6-methyl- (4) are approximately equal. According to the solubility increase the investigated thiadiazoles are ranged in the following order: 12 (4-bromophenyl-)  $\leq$  5 (4-chlorophenyl-)  $<$  4 (3-chloro-6-methylphenyl-)  $\leq$  2 (3-chloro-4-methylphenyl-)  $\leq$  1 (3,4-dichlorophenyl-)  $\leq$  6 (4-methylphenyl-)  $\leq$  3 (3-chloro-4-fluorophenyl-)  $<$  7 (3-methylthiaphenyl-)  $\leq$  10 (methylisoxazol-)  $<$  9 (4-fluorophenyl-)  $\leq$  8 (4-methoxyphenyl-)  $<$  11 (cyclopropyl-). The solubility change regularities described above do not have a forecasting power. Therefore, it was important to obtain correlation equations between the solubility characteristics and any kinds of descriptors. HYBOT descriptors were used as physicochemical descriptors [25]. Crystal lattice thermodynamic characteristics are important parameters in solid substance dissolution. To take this factor into account, Yalkowsky et al. [38] introduced crystal melting temperature as a descriptor into the correlation equation (General Solubility Equation). Using this descriptor to analyze the solubility thermodynamics is justified by the fact that in many cases there is a correlation between the molecular crystal sublimation Gibbs energies and melting temperatures [39]. Besides, Yalkowsky et al. [38] described water solubility of the compounds using 1-octanol/water partitioning coefficients as descriptors (for imitation of hydration term). The descriptor series suggested by Yalkowsky was used to describe selected compounds solubility. As a result, the correlation equation looks as follows:

$$\log(S_2^{298}) = (0.4 \pm 1.5) - (0.19 \pm 0.14) \cdot \log(D^{oct/buf}) - (7.2 \pm 4.7) \cdot 10^{-2} \cdot T_m \quad (15)$$

$$R = 0.8071; \sigma = 0.31; F = 12.4; n = 15$$

Thus, the solubility of this class of compounds decreases with increasing the lipophilicity (due to reducing the hydration term) and the melting temperature growth (crystal lattice energy strengthening).

We also tried to find a correlation equation using HYBOT descriptors. Analysis of all the available descriptors found that the best correlations were obtained with polarizability ( $\alpha$ ) and acceptor molecular ability to create hydrogen bonds ( $\Sigma(C_a)$ ):

$$\Delta G_{sol}^{298}(Buffer)/kJ \cdot mol^{-1} = (38.3 \pm 2.6) + (0.23 \pm 0.13) \cdot \alpha + (-3.4 \pm 0.9) \cdot \Sigma(C_a) \quad (16)$$

$$R = 0.8493; \sigma = 1.54; F = 18.11; n = 17$$

Thus, the solubility-values of this class of compounds in a buffer solution can be estimated based on the knowledge of the structural formula only.

### 3.3. Permeability Properties

Recently, much attention has been paid on the development of better bioavailable drugs, which significantly reduce both dosage and side effects. Water solubility and permeability are considered to be the main parameters determining the bioavailability of drugs. Therefore, in our work we have tried to analyze these specified parameters. Permeapad™ barrier developed and described in [27] was chosen as a biomimetic artificial barrier to study the permeability. Advantages of Permeapad™ (compared to other artificial systems such as PAMPA, PVPA) are: a unique design, usability, functional stability in a wide range of pH values, compatibility with biomimetic media, adaptability to Franz diffusion cell and shelf life (more than 1 year). The apparent permeability coefficients through the Permeapad™ barrier ( $P_{app}$ ), support layer ( $P_{sup}$ ) and lipid layer ( $P_{lip}$ ) of the selected compounds, measured in a buffer solution (pH 7.4) at T=298 K are summarized in Table 3.

**Table 3** Apparent permeability coefficients measured through Permeapad™ barrier ( $P_{app}$ ), support layer ( $P_{sup}$ ) and extrapolated ( $P_{lip}$ , thus the contribution of the lipid layer to permeability resistance) of all compounds tested. Results are reported as average and standard deviation of replicate experiments (n = 3–5).

Compound	Apparent permeability coefficients		
	Permeapad™ barrier	Support layer	Lipid layer
	$P_{app} \times 10^6 / \text{cm} \cdot \text{s}^{-1}$	$P_{sup} \times 10^6 / \text{cm} \cdot \text{s}^{-1}$	$P_{lip} \times 10^6 / \text{cm} \cdot \text{s}^{-1}$
1	9.8 ± 0.8	15.7 ± 1.2	26.1 ± 4.0
2	13.5 ± 0.4	20.6 ± 0.1	39.5 ± 3.1
3	13.9 ± 2.3	25.4 ± 1.5	30.9 ± 7.3
4	14.4 ± 0.1	30.8 ± 3.0	26.9 ± 4.8
5	8.6 ± 0.9	21.9 ± 1.4	14.1 ± 0.2
6	12.2 ± 2.7	26.3 ± 1.9	22.8 ± 7.7
7	10.1 ± 0.5	22.8 ± 1.4	18.2 ± 1.9
8	9.4 ± 1.1	16.5 ± 1.3	21.7 ± 0.7
9	8.7 ± 1.1	16.3 ± 1.1	18.6 ± 2.0
10	3.4 ± 0.3	31.7 ± 4.4	3.7 ± 0.4
11	5.4 ± 0.4	27.6 ± 1.3	6.8 ± 0.6
12	6.5 ± 0.7	25.8 ± 2.2	9.5 ± 0.6

The results showed by [27] indicate that the permeability coefficients through the sheer support layer ( $P_{sup}$ ) were in all cases higher compared to the respective  $P_{app}$  measured for the entire Permeapad™ barrier. Our

data in Table 3 confirmed these conclusions. This fact is a clear indication of a significant retention effect induced by the lipids present in the artificial barrier and that the lipid layer is the dominant impediment to drug permeation. From this it can be concluded that for all compounds investigated the rate limiting step in permeability was represented by the lipid layer of the artificial barrier. Due to this fact it is proposed that for screening purposes, it is reasonable to go without measuring the  $P_{sup}$  - value.

The values of the permeation coefficients vary depending on the nature and number of substituents in the molecule. The maximum permeability coefficient ( $P_{lip}$ ) is observed for 3-chloro-4-methylphenyl- derivative (compound 2). Replacement of the methyl- group in this compound by a chlorine atom (dichlorophenyl-substituted compound (1)) leads to a small reduction of permeability. Thus, the permeability coefficient of 3-chloro-4-methylphenyl- substituted compound is about 1.5 times higher than that of its 3-chloro-6-methyl- isomer (4), indicating the effect of not only the nature but also the position of the substituent on the permeability of thiadiazoles. Significant reduction in permeability (minimum values among all compounds) occurs in substances with isoxazole (10) and cyclopropyl (11) substituents. In the order of permeability decreasing phenyl-substituted substances are ranged in the following order: 3-chloro-4-methyl- (2) > 3-chloro-4-fluoro- (3) > 3-chloro-6-methyl- (4) > 3,4-dichloro- (1) > 4-methyl- (6) > 3-methoxy- (8) > 4-fluoro- (9) > 3-methylthia- (7) > 4-chloro- (5) > 4-bromo- (12).

Due to the fact that the distribution and permeability processes for the selected compounds were studied in the same buffer solution (pH 7.4) (i.e. ionization degree of the particles and their ratio remained the same for the two designated processes), we tried  $D^{oct/buf}$  and  $D^{hex/buf}$  as descriptors for the permeability. In its turn,  $\log P_{app}$ ,  $\log P_{sup}$  and  $\log P_{lip}$  were selected as the analyzed functions. It should be noted that there are no any correlations between the selected functions and  $D^{hex/buf}$  (Fig. 2SI). In contrast, good correlations between  $\log P_{app}$  and  $\log D^{oct/buf}$ , as well as between  $\log P_{lip}$  and  $\log D^{oct/buf}$  (Fig. 2), exist, which can be described by the following equations:

$$\log P_{app} = - (5.25 \pm 0.05) + (0.37 \pm 0.06) \cdot \log(D^{oct/buf}) \quad (17)$$

$$R = 0.8737; \sigma = 9.46 \cdot 10^{-2}; n = 12$$

$$\log P_{lip} = - (5.13 \pm 0.05) + (0.63 \pm 0.08) \cdot \log(D^{oct/buf}) \quad (18)$$


$$R = 0.9351; \sigma = 0.11; n = 12$$

It is easy to see that  $\log P_{lip}$  is in 1.7 times more sensitive to  $\log(D^{oct/buf})$  changes in comparison with  $\log P_{app}$ . Thus, the supporting layer contributes to the sensitivity of this coefficient, but, on the other hand, the correlation parameters are not much deteriorated. Consequently,  $\log P_{app}$  can be used to analyze the lipid layer permeability.

## ARTICLE

## Journal Name

If we take into account that  $\log P_{lip}$  -value is burdened with two experimental errors (sequential execution of two experiments: permeability Permeapad<sup>TM</sup> barrier and Support layer), then  $P_{app}$  value is more accurate (and the total time of the experiment is halved).



**Fig.2.** Dependencies  $\log P_{app}$  vs  $\log D^{oct/buf}$  (a) and  $\log P_{lip}$  vs  $\log D^{oct/buf}$  (b)


As the next step it was interesting to analyze the relationships of HYBOT physicochemical descriptors with the obtained permeability characteristics. Testing all the descriptors revealed that only for  $\log P_{lip}$  and  $\log P_{app}$  (as in the previous cases for  $\log D^{oct/buf}$ ) the correlation with acceptor molecular ability to create hydrogen bonds normalized to molecular polarizability ( $\Sigma(C_a)/\alpha$ ) was observed. Results of the analysis for  $\log P_{lip}$  and  $\log P_{app}$  are presented in Fig. 3, whereas the relationship between  $\log P_{sup}$  and  $\Sigma(C_a)/\alpha$  is shown in Fig. 3SI. The received correlation can be described by the following equations:

$$\log P_{app} = -(2.95 \pm 0.48) - (11.9 \pm 2.7) \cdot \Sigma(C_a)/\alpha \quad (19)$$

$$R = 0.8089; \sigma = 0.11; n = 12$$

$$\log P_{lip} = -(1.47 \pm 0.80) - (18.9 \pm 4.5) \cdot \Sigma(C_a)/\alpha \quad (20)$$

$$R = 0.7961; \sigma = 0.19; n = 12$$



**Fig.3.** Dependencies  $\log P_{app}$  vs  $\Sigma(C_a)/\alpha$  (a) and  $\log P_{lip}$  vs  $\Sigma(C_a)/\alpha$  (b)

Thus, the permeability coefficients of this class of compounds may be predicted based solely on their structural formulas. Due to the small number of data points the paired correlation coefficients are not rather high. However, the improvement should be expected if the number of data points would increase. Just as in the case of the distribution coefficients the correlation equations for  $\log P_{app}$  and  $\log P_{lip}$  retain the same trends on the selected descriptor.

### 3.4. Integral Solubility - Permeability Properties

The available data on the solubility of the test compounds and the apparent permeability coefficients make it possible to evaluate the potential bioavailability of the substances, similarly to Sanphui et al. [40]. The approach consists of the estimation of diffusion fluxes through the lipid barrier ( $J_{lip}$ ) and Permeapad ( $J_{app}$ ) (area of 1 cm<sup>2</sup> and nominal thickness of 1 cm) on the assumption that at one end of the barrier the concentration is kept equal to thermodynamic solubility value, while at the other end - the substance is effectively removed and its concentration is practically equal to zero (sink conditions). In other words, the value of the product




of thermodynamic solubility ( $S_2^{298}$  in  $\text{mol}\cdot\text{l}^{-1}$ ), lipid permeability coefficient ( $P_{lip}$  in  $\text{cm}\cdot\text{s}^{-1}$ ) and Permeapad permeability coefficient ( $P_{app}$  in  $\text{cm}\cdot\text{s}^{-1}$ ) is analyzed. The analysis in logarithmic scale yields the following equations:

$$\log J_{lip} = \log(S_2^{298} \cdot P_{lip}) = \log S_2^{298} + \log P_{lip} \quad (21a)$$


$$\log J_{app} = \log(S_2^{298} \cdot P_{app}) = \log S_2^{298} + \log P_{app} \quad (21b)$$

Fig. 4 depicts the analyzed values. It is easy to see that according to the degree of decreasing the diffusion flux value (a measure of absorption) the compounds can be placed in the following order: for lipid layer – **11 > 8 > 9 > 3 > 2 > 7 > 1 > 6 > 4 > 5 ≈ 10 > 12**; for Permeapad – **11 > 8 > 9 > 3 > 7 > 6 > 2 ≈ 1 ≈ 4 > 10 > 5 > 12**.



**Fig. 4.** Experimental solubility data ( $S_2^{298}$ ), lipid layer permeability coefficients ( $P_{lip}$ ) and diffusion fluxes through lipid layer ( $J_{lip}$ ) for the compounds studied (a); Experimental solubility data ( $S_2^{298}$ ), Permeapad layer permeability coefficients ( $P_{app}$ ) and diffusion fluxes through Permeapad ( $J_{app}$ ) for the compounds studied (b)

With this range of changing the  $P_{lip}$  and  $P_{app}$ -values it turns out that the solubility values have the main contribution to the diffusion fluxes (Fig. 5).




**Fig. 5.** Dependences of  $\log J_{lip}$  vs  $\log S_2^{298}$  and  $\log J_{app}$  vs  $\log S_2^{298}$

### 3.5. Calcium-blocking Properties

The substances synthesized in the present study belong to essentially new group of compounds in a series of 1,2,4-thiadiazole derivatives which, being bioisosteric analogues of a model «ampakin» cyclothiazide, as proposed, can be the positive modulators of AMPA-receptors. First of all the ability of the synthesized substances to modulate glutamate-dependent Ca-uptake in synaptosomes of rat brain cortex was studied. The biological activity results ( $K_{43/21}$ -values) are presented in Table 2.

The obtained results clearly indicate that the variation of the structure of the substituent in the phenyl ring as well as the replacement of the phenyl ring by the cyclopropyl- and isoxazole-ones influence the ability of thiadiazole derivatives to inhibit calcium ion uptake. The biological activity ( $K_{43/21}$ -values) of the investigated substances is presented in Fig. 6 as a histogram to compare the compounds of groups I and II with the same substituents.



**Fig. 6.** Dependence  $\log(1/K_{43/21})$  vs.  $\log D^{\text{oct}/\text{buf}}$

The comparative analysis of the activity of thiadiazole derivatives newly synthesized and investigated by us before indicated that, on the whole, the presence of

## ARTICLE

## Journal Name

tetramethylpiperidine ring and additional NH-group instead of hydroxyl group facilitates a considerable growth of  $K_{43/21}$ -values (up to 6 times for the compounds with dichlorophenyl-substituent (compounds **1** and **13**)). However, it is interesting to note that if the replacement of the phenyl ring by isoxazole-one takes place, the activities of the respective structural analogues of groups I and II practically coincide. The analysis of the structure-activity relationship was conducted for thiadiazoles synthesized in the present study (Group I – compounds **1–12**). Compound **12** with a bromine atom as a substituent in para-position of the phenyl ring, revealing maximal inhibiting activity (minimal  $K_{43/21}$ ), was chosen as a reference. As seen from the histogram data, replacing a bromophenyl- substituent by cyclopropyl- and methyl-isoxazole ones leads to a slight consistent increase in  $K_{43/21}$  (compounds **11** and **10**). Further reduction of the inhibitory activity is observed for compounds with 4-fluorophenyl-, 3-methoxyphenyl- and 3-methylthiaphenyl- substituents, which have similar activity values (**9** (112%) = **8** (112%) ≤ **7** (117%)). Introduction of the methyl group (compound **6**) in a para- position of the phenyl ring contributes to a further increase of  $K_{43/21}$ . At that, the inhibitory activity of the compound with methylisoxazol- fragment (**10**) is higher than that of methylphenyl- substituted one (**6**).

The results presented in the histogram allowed us to evaluate the total biological potential of the compounds, not only as possible neuroprotectors, but also cognitive stimulants (nootrops). Thus, for the five compounds (**1**, **2**, **3**, **4**, and **5**) the values of  $K_{43/21} > 150$  suggest them for further testing as stimulants of cognitive functions using a recognition test. It is interesting to note that these compounds have chlorine atoms as substituents in the phenyl ring, at the same time, di-substituted substances with a chlorine atom in the meta-position reveal the maximum  $K_{43/21}$ -values and according to the reduction of the activation function of glutamate-dependent uptake of calcium ions the thiadiazoles can be arranged in the following order: **1** (3-chloro-4-chloro) > **2** (3-chloro-4-methyl) > **3** (3-chloro-4-fluoro-) > **4** (3-chloro-6-methyl) > **5** (4-chloro-).

### 3.6. Activity correlations

During scientific prognostication of physiological activity in QSAR [41] the influence of the following phenomena is usually assessed: electronic effects (influencing the ionization or polarity of the substance), steric characteristics of the structure (playing an important role for the evaluation of the binding strength of the investigated compound with the biological target), lipophilicity (ability to dissolve in fats, characterizing the properties of the substance to penetrate through the cell membranes). The distribution coefficient (and its logarithm) in 1-octanol/buffer system is well known to be the most

often used descriptor during the estimation of the quantitative structure-activity correlations.

In order to describe graphically the activity-lipophilicity dependence we applied an approach introduced in [42] for compounds of specific type of activity and close chemical structure according to equation:

$$\log(1/K_{ba}) = a_0 + a_1 \cdot \log D^{\text{oct/buf}} - a_2 \cdot (\log D^{\text{oct/buf}})^2 + a_3 \cdot \sigma + a_4 \cdot E_s \quad (22)$$

where  $K_{ba}$  – any experimental value characterizing the biological activity,  $D^{\text{oct/buf}}$  – distribution coefficient of the substance between 1-octanol and aqueous (buffer) phase,  $\sigma$  and  $E_s$  –parameters describe the electronic and steric impacts of the substituents, respectively,  $a_i$  – constants derived from processing the experimental data by the least square method.

Fig. 7 depicts the dependence of the biological activity of the investigated thiadiazoles on the lipophilicity of the molecules characterizing by the distribution coefficient in 1-octanol/buffer system.




Fig. 7. Dependence  $\log(1/K_{43/21})$  vs.  $\log D^{\text{oct/buf}}$

The correlation equation, in which the biological activity acts as a dependent variable and the distribution coefficient in 1-octanol/buffer system acts as an independent one, is as follows (compound **5** is excluded from the correlation):

$$\log(1/K_{43/21}) = -(1.91 \pm 0.05) + (0.04 \pm 0.10) \log D^{\text{oct/buf}} - (0.45 \pm 0.14) (\log D^{\text{oct/buf}})^2 \quad (23)$$

$$R = 0.7092; \sigma = 0.08; F = 13.19; n = 11$$

It should be noted that the obtained correlation is not high; this fact may be explained by the rather small number of the experimental points. Enhancement of the correlation can be expected at the extension of the number of the compounds investigated. However, the obtained regularities support the parabolic model to be an appropriate manner for describing the dependence of the specific activity of thiadiazoles synthesized in the present

study on the lipophilicity expressed by the distribution coefficient in 1-octanol/water system.

#### 4. Conclusions

Novel 1,2,4-thiadiazole derivatives as potent neuroprotectors were synthesized and identified. Their ability to inhibit the glutamate stimulated  $\text{Ca}^{2+}$  uptake was investigated *in vitro*. The structure of thiadiazole molecules was shown to influence the inhibition properties. Activity-lipophilicity correlation was estimated based on the parabolic model. The solubility of 1,2,4-thiadiazoles was measured in buffer solution pH 7.4 at 298 K. The distribution coefficients in 1-octanol/buffer (pH 7.4) and 1-hexane/buffer (pH 7.4) were determined. Apparent permeability coefficients were obtained in Franz diffusion cell using new Permeapad<sup>TM</sup> barrier. Search of the correlations between the distribution coefficients in 1-octanol/buffer and 1-hexane/buffer systems and physicochemical descriptors was performed. A suitable correlation was obtained for the independent variables which include polarizability ( $\alpha$ ) and acceptor molecular ability to create hydrogen bonds ( $\Sigma(C_a)$ ). The appropriate correlation equations were derived and predictive force of the models was assessed. Analysis of the relationships of physicochemical descriptors with the permeability characteristics was fulfilled. Testing all the descriptors revealed the correlation with  $(\Sigma(C_a)/\alpha)$ -parameter. The obtained correlation equations confirmed the possibility of permeability coefficients of this class of compounds to be predictable based only on their structural formulas. Potential bioavailability of the substances was determined by the solubility-permeability combination through the estimation of diffusion flux across the synthetic lipid membrane. The solubility values were revealed to have the main contribution to the diffusion fluxes of the investigated compounds. Results obtained in the present study will be useful when fast bioavailability screening of newly synthesized substances is required.

#### Acknowledgements

This work was supported by the Russian Science Foundation (№ 15-13-10017).

#### References


- 1 R. A. Jellinger, G. Ladurner, M. Windisch, *New Trends in the Diagnosis and Therapy of Alzheimer's Disease*. Wien-New-York: Springer-Verlag; 1994.
- 2 N. Guzior, A. Wieckowska, D. Panek, B. Malawska, *Curr. Med. Chem.*, 2015, **22** (3), 373.
- 3 A. Castro, T. Castano, A. Encinas, W. Porcal, C. Gil, *Bioorg & Med Chem.*, 2006, **14**, 1644.
- 4 A. Martinez, M. Alonso, A. Castro, C. Perez, F. J. Moren, *J. Med. Chem.*, 2002, **45**, 1292.

- 5 A. M. MacLeod, R. Baker, S. B. Freedman, S. Patel, K. J. Merchant, M. Roe, J. Saunders, *J. Med. Chem.*, 1990, **33**, 2052.
- 6 Y. Iizawa, K. Okonogi, R. Hayashi, T. Iwahi, T. Yamazaki, A. Imada, *Antimicrob. Agents Chemother.*, 1993, **37**, 100.
- 7 L. N. Petrova, S. O. Bachurin, *Bull. Expert. Biol. Med.*, 2006, **142**, 51.
- 8 J. N. Kew, J. A. Kemp, *Psychopharmacology*, 2005, **179** (1), 4.
- 9 L. Parnetti, U. Senin, P. Mecocci, *Drugs*, 1997, **53** (5), 752.
- 10 M. Husain, M. A. Mehta, *Trends in Cognitive Sciences*, 2011, **15** (1), 28.
- 11 E. Kerns, Li Di., *Druglike Properties: Concepts, Structure Design and Methods*. Academic Press; 2008.
- 12 A. Avdeef, *Absorption and Drug Development: Solubility, Permeability, and Charge State*. Hoboken: Wiley; 2003.
- 13 M. Masson, B. V. Sigurdadottir, K. Matthiasson, T. Loftsson, *Chem. and Pharm. Bull.*, 2005, **53**, 958.
- 14 A. Andrés, M. Rosés, C. Rafols, E. Bosch, S. Espinosa, V. Segarra, J. M. Huerta, *Eur. J. Pharm. Sci.*, 2015, **76**, 181.
- 15 J. E. Comer, High-throughput measurement of logD and pKa. In: van de Waterbeemd H, Lennernäs H, Artursson P, editors. *Drug Bioavailability: Estimation of Solubility, Permeability, Absorption, and Bioavailability*. Weinheim: Wiley-VCH; 2007.
- 16 P. Seiler, *Eur. J. Med. Chem.*, 1974, **9** (5), 473.
- 17 G. L. Perlovich, T. V. Volkova, A. N. Proshin, D. Yu. Sergeev, Bui T. Cong, L. N. Petrova, S. O. Bachurin, *J. Pharm. Sci.*, 2010, **99** (9), 3754.
- 18 G. L. Perlovich, A. N. Proshin, T. V. Volkova, Bui T. Cong, S. O. Bachurin, *Mol. Pharm.*, 2011, **8**, 1807.
- 19 A. O. Surov, Bui T. Cong, A. N. Proshin, P. Roussel, A. Idrissi, G. L. Perlovich, *J. Phys. Chem. B.*, 2013, **117**, 10414.
- 20 A. O. Surov, Bui T. Cong, T. V. Volkova, A. N. Proshin, G. L. Perlovich, *Phys. Chem. Chem. Phys.*, 2015, **17**, 20889.
- 21 A. O. Surov, Bui T. Cong, T. V. Volkova, A. N. Proshin, G. L. Perlovich, *J. Chem. Thermodynamics*, 2016, **96**, 57.
- 22 G. L. Perlovich, A. N. Proshin, T. V. Volkova, S. V. Kurkov, V. V. Grigoriev, L. N. Petrova, S. O. Bachurin, *J. Med. Chem.*, 2009, **52**, 1845.
- 23 G. L. Perlovich, A. N. Proshin, T. V. Volkova, L. N. Petrova, S. O. Bachurin, *Mol. Pharm.*, 2012, **9** (8), 2156.
- 24 S. V. Blokhina, M. V. Ol'khovich, A. V. Sharapova, A. N. Proshin, G. L. Perlovich, *J. Chem. & Eng. Data*, 2012, **57**, 1996.
- 25 A. J. Boulton, A. R. Katrizky, A. M. Hamid, *J. Chem. Soc.*, 1967, 2005.
- 26 Chemical equilibrium. Properties of solutions. In: Simanova CA, editors. *Manual chemist and technologist*. S.-Pt: Professional; 2004. p. 194.
- 27 M. di Cagno, H. A. Bibi, A. Bauer-Brandl, *Eur. J. Pharm. Sci.*, 2015, **73**, 29.
- 28 T. Higuchi, K. A. Connors, *Adv. in Anal. Chem. and Instr.*, 1965, **4**, 117.
- 29 M. Brandl, G. E. Flaten, A. Bauer-Brandl, *Passive diffusion across membrane*. Wiley Encyclopedia Chem Biol. 2009, **3**, 541-500.
- 30 E. B. Gharthey-Tagoe, J. S. Morgan, A. S. Neish, M. R. Prausnitz, *J. Contr. Release*, 2005, **103**, 177.
- 31 A. J. Leo, *J. Pharm. Sci.*, 1987, **76**, 166.
- 32 A. J. Leo, C. Hansch, D. Elkins, *Chem. Rev.*, 1972, **71**, 525.
- 33 A. Andrés, M. Rosés, C. Rafols, E. Bosch, S. Espinosa, V. Segarra, J. M. Huerta, *Eur. J. Pharm. Sci.*, 2015, **76**, 181.
- 34 OECD guideline for the testing of chemicals adopted by the Council on 27th July 1995 Partition Coefficient (n-octanol/water): Shake Flask Method. Available from: <http://www.oecd.org/chemicalsafety/risk-assessment/1948169.pdf>

## ARTICLE

## Journal Name

- 35 O. A. Raevsky, V. J. Grigor'ev, S. V. Trepalin. HYBOT (Hydrogen Bond Thermodynamics) program package, Registration by Russian State Patent Agency N 990090 of 26.02.99.
- 36 ACD/ChemSketch program. Available from: [http://eduinf.waw.pl/che/inne/prgchem/pages/chsk\\_eng.pdf](http://eduinf.waw.pl/che/inne/prgchem/pages/chsk_eng.pdf)
- 37 V. A. Borodin, E. V. Kozlovsky, V. P. Vasil'ev, *Russ. J. Inorg. Chem.*, 1982, **27**, 2169.
- 38 S. H. Yalkowsky, S. C. Valvani, *J. Pharm. Sci.*, 1980, **69**, 912.
- 39 G. L. Perlovich, A. M. Ryzhakov, V. V. Tkachev, L. Kr. Hansen, O. A. Raevsky, *Crystal Growth & Design*, 2013, **13** (9), 4002.
- 40 P. Sanphui, V. K. Devi, D. Clara, N. Malviya, G. S. Somnath, G. R. Desiraju, *Mol. Pharm.*, 2015, **12** (5), 1615.
- 41 C. Hansch, T. Fujita, *J. Am. Chem. Soc.*, 1964, **86**, 1616.
- 42 C. Hansch Quantitative structure-activity relationship in drug design. In: E.J. Ariens, editors. New York: Academic Press; 1971. p. 271-342.



Novel 1,2,4-thiadiazole derivatives as potent neuroprotectors were synthesized and their solubility, distribution coefficients, permeability and biological properties were studied.