

2-Arylidene Hydrazinecarbodithioates as Potent, Selective Inhibitors of #-Cystathionine-Lyase (CSE)

Abir Bhattacharjee, Antara Sinha, Kiira Ratia, Liang Yin, Loruham Delgado-Rivera, Pavel A. Petukhov, Gregory RJ Thatcher, and Duncan John Wardrop

ACS Med. Chem. Lett., **Just Accepted Manuscript** • DOI: 10.1021/acsmchemlett.7b00313 • Publication Date (Web): 21 Nov 2017

Downloaded from <http://pubs.acs.org> on November 24, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

2-Arylidene Hydrazinecarbodithioates as Potent, Selective Inhibitors of Cystathionine γ -Lyase (CSE)

Abir Bhattacharjee,^{†,‡} Antara Sinha,^{†,‡} Kiira Ratia,[#] Liang Yin,[#] Loruhamá Delgado-Rivera,[#] Pavel A Petukhov,[#] Gregory R. J. Thatcher,[#] and Duncan J. Wardrop*[†]

[†]Department of Chemistry, University of Illinois at Chicago, 845 W. Taylor St., Chicago, Illinois 60607, United States

[#]Department of Medicinal Chemistry & Pharmacognosy, University of Illinois College of Pharmacy, University of Illinois at Chicago, 833 S. Wood St., Chicago, IL 60612, United States

KEYWORDS. Cystathionine γ -lyase, cystathionine β -synthase, hydrogen sulfide, 2-pyridyl thiosemicarbazones.

ABSTRACT: Hydrogen sulfide is produced from L-cysteine by the action of both cystathionine γ -lyase (CSE) and cystathionine β -synthase (CBS) and increasingly has been found to play a profound regulatory role in a range of physiological processes. Mounting evidence suggests that upregulation of hydrogen sulfide biosynthesis occurs in several disease states, including rheumatoid arthritis, hypertension, ischemic injury, and sleep-disordered breathing. In addition to being critical tools in our understanding of hydrogen sulfide biology, inhibitors of CSE hold therapeutic potential for the treatment of diseases in which increased levels of this gasotransmitter plays a role. We describe the discovery and development of a novel series of potent CSE inhibitors that show increased activity over the benchmark inhibitor and, importantly, display high selectivity for CSE versus CBS.

In mammals, the pyridoxal-5'-phosphate (PLP)-dependent enzyme cystathionine γ -lyase (CSE) mediates the conversion of cystathionine to hydrogen sulfide (H_2S) through the intermediacy of L-cysteine (Cys).¹ Together with cystathionine β -synthase (CBS)² and, to a lesser extent, 3-mercaptopyruvate sulfurtransferase (3-MST),³ CSE is the principle source of endogenous H_2S which has been found to play an important regulatory role in a wide array of physiological processes such as inflammation,⁴ blood pressure homeostasis,⁵ neuromodulation,⁶ cytoprotection⁷ and aging.⁸

Upregulation of CSE and increased H_2S biosynthesis has been implicated in several disease states, including inflammatory joint disease, chronic obstructive pulmonary disease, Alzheimer's disease, and endotoxemia.⁹ Reducing H_2S levels through inhibition of the enzymes involved in its production has been found to hold promise as a therapeutic intervention.^{10,11} Notably, the benchmark CSE inhibitor, L-propargyl glycine (L-PAG, **1**) normalizes breathing and reduces hypoxia induced hypertension in rodent models of sleep-disordered breathing, suggesting that inhibition of H_2S biosynthesis within the carotid body may be a new approach to treat hypertension in patients with sleep apnea (Figure 1).¹² The development of inhibitors that display selectivity between CSE and CBS is also of importance because expression of these enzymes is more widespread and less tissue specific than once thought. The future delineation of the roles individually played by CSE and CBS will be dependent on the availability of selective inhibitors.




Figure 1. Chemical structure and IC_{50} values of selected inhibitors of cystathionine γ -lyase (CSE). Compounds **1-3** and **6** show some degree of selectivity for CSE over cystathionine β -synthase (CBS).

Since the discovery of compound **1** as a mechanism-based inhibitor ($IC_{50} = 40 \mu$ M)¹³ in 1976,¹⁴ few inhibitors of CSE have been identified and only a limited number of these display selectivity over CBS. Furthermore, many of these molecules have significant drawbacks, e.g. compound **1** also acts as an inhibitor of alanine monotrtransferase and aspartate aminotrtransferase at the concentrations employed,¹⁵ while β -cyanoalanine (BCA, **2**) is a neurotoxin.¹⁶ The inherent polarity of these amino acids is an additional drawback to their use since it leads to poor cell permeability.

The paucity of selective CSE inhibitors is a significant impediment to the study of H_2S pathways and has prompted considerable recent interest in this area. In 2016, the natural product L-

α -(2-aminoethoxyvinyl)glycine (AVG, **3**)¹⁷ was reported by Asimakopoulou and Panopoulos, to be a selective inhibitor of CSE ($IC_{50} = 1.1 \mu\text{M}$) versus CBS, although its selectivity against other PLP-dependent enzymes remains to be determined.¹³ In 2013, high throughput screening (HTS) efforts by Zu identified iminoquinolinone derivative NSC111041 (**4**) as an inhibitor of CSE ($IC_{50} = 6.3 \mu\text{M}$),¹⁸ while Pastore and Caliendo have recently reported the development of *N*-propargyl D-cysteine derivative **5** ($IC_{50} = 30 \mu\text{M}$).¹⁹ Interestingly D-penicillamine (**6**), which has previously been employed clinically to alleviate the symptoms of rheumatoid arthritis, has also been reported to selectively inhibit CSE, albeit weakly ($IC_{50} = 0.27 \text{mM}$).²⁰

As part of a program to develop therapeutic agents for sleep-disordered breathing, we recently initiated a search for small-molecule inhibitors of CSE that display selectivity over CBS. In this communication, we report the discovery of a series of α -pyridyl alkylthio(thiocarbonyl)hydrazones that not only potently inhibit CSE, but display high selectivity over CBS. Thiiosemicarbazones and their metal complexes display a range of biological activities, including antibacterial²¹ and anti-cancer properties,²²⁻²⁴ for which they are the subject of considerable interest.

To identify inhibitors, recombinant human CSE was screened against >100,000 compounds from several commercial collections, including ChemDiv, Chembridge, Maybridge, and Prestwick. The high-throughput primary assay monitored CSE activity by detecting production of L-cysteine from the cleavage of L-cystathionine, using the thiol-reactive fluorogenic probe, CPM (7-diethylamino-3-(4'-maleimidylphenyl)-4-methylcoumarin). Compounds were screened in duplicate at a concentration of 40 μM , and those displaying >30% enzyme inhibition (243 compounds) were validated in secondary assays. Remaining positive hits (22 compounds) were repurchased and then assessed for inhibitory activity against CSE- and CBS-catalyzed H₂S production, using the hydrogen sulfide selective probe, 7-azido-4-methylcoumarin (AzMC) (see Supporting Information).²⁵ From this screening effort, one compound, initially identified as structure **7**, which inhibited CSE at 40 μM but showed weak activity against CBS, was selected for further investigation (**Figure 2**). This hit was reconfirmed and found to display dose dependent inhibition of CSE, with an IC_{50} of 6 μM . We tested the reversibility of this compound by dialysis experiments (see Supporting Information) and found that compound **7** was a reversible inhibitor of CSE. Our independent synthesis of this compound (*vide infra*) indicates that its structure was incorrectly identified in the commercial library as a cyclic system, namely 2,3-dihydro-1,3,4-thiadiazole **7**, rather than Schiff base **9**.





Figure 2. Structural identity of high-throughput screening (HTS) hit. (a) Reported chemical structure of library hit compound **7**. (b) Revised chemical structure **9**, established through synthesis, and associated data of this cystathionine γ -lyase (CSE) inhibitor. 2-Pyridyl thiosemicarbazones can exist in syn (**8**) and anti (**9**) isomeric forms, which interconvert in protic solvents. The anti-isomers are thermodynamically favored. (c) CSE and CBS activity assays, measuring H₂S production in the presence and absence of compound **9** (40 μM) showing selectivity for CSE over CBS. Using this AMC assay, $IC_{50} \sim 1\text{-}2 \mu\text{M}$ for CSE.

Having identified **9** as a promising hit, we now sought to optimize the inhibitory activity of this compound, through chemical modification, and examine the structure-activity relationships (SAR). Thus, a library of thirty-two analogues (**17-55**) were synthesized following the general strategies outlined in **Scheme 1**. Employing the one-pot method of Busch,²⁶ reaction of carbon disulfide, hydrazine hydrate and alkyl halides **10** in the presence of base, generated the corresponding *S*-alkyl dithiocarbazates **11**, which underwent condensation with aldehydes and ketones **12** to generate hydrazones **15** as single diastereomers. Alternatively, treatment of hydrazones **13** with carbon disulfide generated 1,3,4-thiadiazolidine-2-thiones **14**,²⁷ which underwent ring-opening and exclusive *S*-alkylation to generate **15**. While alkylthio(thiocarbonyl)hydrazones can potentially exist in equilibrium with 2,3-dihydro-1,3,4-thiadiazoles **16**, no evidence for the presence of such *S,N*-acetals was found in the ¹³C NMR spectra of these products, which display characteristic carbodithioate and hydrazone signals at approximately 200 and 150 ppm, respectively. In further regard to the structure of **15**, although 2-pyridyl thiosemicarbazones can exist as syn and anti-isomers (**8** cf. **9**), detailed studies by Venkatachalam and co-workers have recently shown that in protic solvents, interconversion takes place and favors the anti-isomers.²⁸ In all cases, the ¹H and ¹³C NMR spectra of **15** displayed a set of signals corresponding to a single diastereomer.

Scheme 1. General Synthetic Scheme for the Preparation of 2-Arylidene Hydrazinecarbodithioates.^a



^aReagents and conditions: (a) R_3-X (**10**), CS_2 , $NH_2NH_2 \cdot H_2O$, KOH , $EtOH$, rt , 1-3 h; **12**, rt , 3 h (60-80%); (b) CS_2 , DMF , $reflux$, 6 h 80-90%; (c) R_3-X (**10**), KOH , $acetone$, H_2O , rt , 1 h (60-80%).

The results of our initial SAR study are summarized in Table 1.²⁹ Retaining the methyl group at R_3 , we first probed the im-

portance of the 2-pyridyl ring at R_1 for activity, finding that substitution with 3- and 4-pyridyl systems led to an approximately ten-fold decrease in potency in both cases. Substitution with larger electron-rich N -heterocycles (**29**, **30**) at R_1 proved to be unproductive, resulting in complete loss of activity. That the introduction of 2-pyridyl isosteres, including piperazine (**20**), thiazoles (**21**), and pyrazoles (**22**, **23**), at the C-2 position maintains activity further validates the structural requirement of a π -deficient 2-azaheterocyclic group at R_1 that may act as a hydrogen bond acceptor. While the presence of the 2-pyridyl group is clearly of importance, the influence of substituents on this ring appears not to be dramatic, as indicated in the series **24-28**; compound **28**, derived from 3-fluoropicolininaldehyde proved to be the most potent member of this group. Interestingly, analogs such as **19**, derived from ketones and consequently bearing non-hydrogen groups at R_2 display no significant inhibitory effects against CSE at concentrations up to 100 μM .

Table 1. Structure and Cystathionine γ -Lyase (CSE) Inhibitory Activity of Compound 8 & Compounds 17-30^a

Compd	R_1	R_2	IC_{50} (μM)	Compd	R_1	R_2	IC_{50} (μM)
8		H	6	24		H	5
17		H	50	25		H	6
18		H	40	26		H	10
19		Me	>100	27		H	12
20		H	15	28		H	5
21		H	7	29		H	>100
22		H	15	30		H	>100
23		H	6				


^a IC_{50} values were obtained as described in the Supporting Information. The values shown are the mean of a single experiment conducted in triplicate.

The results of our second SAR study are summarized in Table 2. Initially retaining the 2-pyridyl group at R_1 , the influence of the thioether substituent (R_3) was now explored. While replacement of the methyl group of initial hit **9** with simple alkyl, allyl and benzyl substituents ablated all CSE inhibitory activity (data not shown), introduction of an N -phenylacetamide group (**31**) proved to be more productive. Encouraged that this substituent resulted in equipotent activity with the parent system **9** and provided another point for introduction of structural diversity, we

now varied aryl substitution. Introduction of alkyl (**32**, **33**) and alkoxy groups (**34**) at the C-4 position of the acetanilide proved to be most productive, leading to a 3-fold increase in potency in the case of 4-methoxyphenyl acetamide **34**. The poor aqueous solubility of several derivatives (**35**, **36**, **47**, **51**) prevented analysis of SAR for these compounds. Prompted by the activity of 2-fluoropyridine derivative **28**, C-4 acetanilides derivatives **41-44** were prepared. Of this series, compound **43** proved to be the most potent CSE inhibitor found during our study with an IC_{50} of 1.2 μM . The beneficial influence of the C-4 methoxy

group is also apparent with chloropyrazole **54** and thiazole **55**, which have potency close to that found with 2-pyridyl derivative **34**. Interestingly, introduction of *N*-arylamide units at R_2 did increase the activity of 3-pyridyl derivatives **48** and **49** relative to their *S*-Me congener **17**.

In conclusion, through high throughput screening of commercial libraries and structure-based optimization efforts, we have identified a series of potent, small molecule inhibitors of cystathionine γ -lyase (CSE), one of the principle sources of endogenous H_2S .




- CSE $IC_{50} = 1.2 \mu M \pm 0.1$
- CBS $IC_{50} > 500 \mu M$
- Selectivity > 400:1


Figure 4. Profile of compound **43**, a potent, competitive inhibitor of cystathionine γ -lyase (CSE) which displays high selectivity versus cystathionine β -synthase (CBS).

By way of example, compound **43** (**Figure 4**) has potency for CSE inhibition greater than reported for L-PAG, the most widely used inhibitor of CSE, and displays at least 400-fold selectivity over inhibition of CBS. As such, it and other members of the series reported herein are potentially valuable pharmacological tools for the study of H_2S signaling.

Table 2. Structure and Cystathionine γ -Lyase (CSE) Inhibitory Activity of Compounds 31-55^a

Compd	R ₁	R ₂	IC ₅₀ (μ M)	Compd	R ₁	R ₂	IC ₅₀ (μ M)
31			6	44			6.0
32			2	45			10
33			3	46			15
34			2	47			>100 ^b
35			>100 ^b	48			25
36			>100 ^b	49			12
37			3.5	50			40
38			6	51			>100 ^b
39			1.9	52			3.0
40			2.5	53			6.0
41			2.0	54			2.7





^aIC₅₀ values were obtained as described in the Supporting Information. The values shown are the mean of a single experiment conducted in triplicate. ^bLack of activity may be due to low solubility.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: xxxxx/acsmchemlett.xxxxx.

Experimental details and characterization data for the reported compounds, NMR spectra, and biological data. (PDF)

AUTHOR INFORMATION

Corresponding Author

*Tel: 1 312-355-1035. E-mail: wardropd@uic.edu.

Author Contributions

All authors have given approval to the final version of the manuscript. ‡These authors contributed equally. A.B, A.S. and D.J.W. designed and synthesized compounds; K.R., L.Y., and L. D-R performed bioassays. P.A.P. carried out the docking studies. D.J.W. wrote the paper with input from coauthors.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This study was supported by the National Institutes of Health, National Heart, Lung, and Blood Institute (1UH2HL123610) and the UICentre for drug discovery.

ABBREVIATIONS

AVG, aminoethoxyvinylglycine; CBS, cystathionine- β -synthase; CSE, cystathionine γ -lyase; CPM, 7-diethylamino-3-(4'-maleimidylphenyl)-4-methylcoumarin; HTS, high throughput screening; H₂S, hydrogen sulfide; L-PAG, L-propargyl glycine; PLP, pyridoxal-5'-phosphate; SAR, structure-activity relationship.

REFERENCES

- (1) Rose, P.; Moore, P. K.; Zhu, Y. Z. H₂S Biosynthesis and Catabolism: New Insights from Molecular Studies. *Cell. Mol. Life Sci.* **2017**, *74*, 1391-1412.
- (2) Huang, C. W.; Moore, P. K. H₂S Synthesizing Enzymes: Biochemistry and Molecular Aspects. *Handb. Exp. Pharmacol.* **2015**, *230*, 3-25.
- (3) Bronowicka-Adamska P.; Wróbel M.; Magierowski M.; Magierowska K.; Kwiecień S.; Brzozowski T. Hydrogen Sulphide Production in Healthy and Ulcerated Gastric Mucosa of Rats. *Molecules* **2017**, *22*, 530.
- (4) Lo Faro, M. L.; Fox, B.; Whatmore, J. L.; Winyard, P. G.; Whiteman, M. Hydrogen Sulfide and Nitric Oxide Interactions in Inflammation. *Nitric Oxide* **2014**, *41*, 38-47.
- (5) Cacanyiova, S.; Berenyiova, A.; Kristek, F. The Role of Hydrogen Sulphide in Blood Pressure Regulation. *Physiol. Res.* **2016**, *65*, S273-S289.

- (6) Wang, Z.; Liu, D. X.; Wang, F. W.; Zhang, Q.; Du, Z. X.; Zhan, J. M.; Yuan, Q. H.; Ling, E. A.; Hao, A. J. L-Cysteine Promotes the Proliferation and Differentiation of Neural Stem Cells via the CBS/H₂S Pathway. *Neuroscience* **2013**, *237*, 106-117.

- (7) Zayachkivska, O.; Havryluk, O.; Hrycevyh, N.; Bula, N.; Grushka, O.; Wallace, J. L. Cytoprotective Effects of Hydrogen Sulfide in Novel Rat Models of Non-Erosive Esophagitis. *PLoS One* **2014**, *9*, e110688.

- (8) Miller, D. L.; Roth, M. B. Hydrogen Sulfide Increases Thermotolerance and Lifespan in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 20618-20622.

- (9) Whiteman, M.; Le Trionnaire, S.; Chopra, M.; Fox, B.; Whatmore, J. Emerging Role of Hydrogen Sulfide in Health and Disease: Critical Appraisal of Biomarkers and Pharmacological Tools. *Clin. Sci.* **2011**, *121*, 459-488.

- (10) Vandiver, M.; Snyder, S. H. Hydrogen Sulfide: A Gasotransmitter of Clinical Relevance. *J. Mol. Med.* **2012**, *90*, 255-263.

- (11) Szabó, C. Hydrogen Sulphide and its Therapeutic Potential. *Nat. Rev. Drug Discov.* **2007**, *6*, 917-935.

- (12) Yuan, G.; Peng, Y. J.; Khan, S. A.; Nanduri, J.; Singh, A.; Vasavda, C.; Semenza, G. L.; Kumar, G. K.; Snyder, S. H.; Prabhakar, N. R. H₂S Production by Reactive Oxygen Species in the Carotid Body Triggers Hypertension in a Rodent Model of Sleep Apnea. *Sci. Signaling* **2016**, *9*, ra80.

- (13) Asimakopoulou, A.; Panopoulos, P.; Chasapis, C. T.; Coletta, C.; Zhou, Z.; Cirino, G.; Giannis, A.; Szabo, C.; Spyroulias, G. A.; Papapetropoulos, A. Selectivity of Commonly used Pharmacological Inhibitors for Cystathionine γ Synthase (CBS) and Cystathionine β Lyase (CSEA). *Br. J. Pharmacol.* **2013**, *169*, 922-932.

- (14) Abeles, R. H.; Walsh, C. T. Acetylenic Enzyme Inactivators. Inactivation of γ -Cystathionase, in Vitro and in Vivo by Propargylglycine. *J. Am. Chem. Soc.* **1973**, *95*, 6124-6125.

- (15) Tanase, S.; Morino, Y. Irreversible Inactivation of Aspartate Aminotransferases during Transamination with L-Propargylglycine. *Biochem. Biophys. Res. Commun.* **1976**, *68*, 1301-1308.

- (16) Ressler, C.; Nigam, S. N.; Giza, Y. H. Toxic Principle in Vetch. Isolation and Identification of γ -L-Glutamyl-L- β -cyanoalanine from Common Vetch Seeds. Distribution in some Legumes. *J. Am. Chem. Soc.* **1969**, *91*, 2758-2765.

- (17) Pruess, D. L.; Scannell, J. P.; Kellett, M.; Ax, H. A.; Janecek, J.; Williams, T. H.; Stempel, A.; Berger, J. Antimetabolites Produced by Microorganisms. X. L-2-Amino-4-(2-aminoethoxy)-trans-3-butenoic Acid. *J. Antibiot.* **1974**, *27*, 229-233.

- (18) Zhou, Y.; Yu, J.; Lei, X.; Wu, J.; Niu, Q.; Zhang, Y.; Liu, H.; Christen, P.; Gehring, H.; Wu, F. High-Throughput Tandem-Microwell Assay Identifies Inhibitors of the Hydrogen Sulfide Signaling Pathway. *Chem. Commun.* **2013**, *49*, 11782-11784.

- (19) Corvino, A.; Severino, B.; Fiorino, F.; Frecentese, F.; Magli, E.; Perissutti, E.; Santagada, V.; Bucci, M.; Cirino, G.; Kelly, G.; Servillo, L.; Popowicz, G.; Pastore, A.; Caliendo, G. Fragment-Based De Novo Design of a Cystathionine γ -Lyase Selective Inhibitor Blocking Hydrogen Sulfide Production. *Sci. Rep.* **2016**, *6*, 34398.

- (20) Brancalone, V.; Esposito, I.; Gargiulo, A.; Vellecco, V.; Asimakopoulou, A.; Citi, V.; Calderone, V.; Gobbetti, T.; Perretti, M.; Papapetropoulos, A.; Bucci, M.; Cirino, G. D-Penicillamine Modulates Hydrogen Sulfide (H₂S) Pathway through Selective Inhibition of Cystathionine- γ -Lyase. *Br. J. Pharmacol.* **2016**, *173*, 1556-1565.

(21) Low, M. L.; Maigre, L.; Dorlet, P.; Guillot, R.; Pagès, J. M.; Crouse, K. A.; Policar, C.; Delsuc, N. Conjugation of a New Series of Dithiocarbamate Schiff Base Copper(II) Complexes with Vectors Selected to Enhance Antibacterial Activity. *Bioconjug. Chem.* **2014**, *25*, 2269-2284.

(22) Santini, C.; Pellei, M.; Gandin, V.; Porchia, M.; Tisato, F.; Marzano, C. Advances in Copper Complexes as Anticancer Agents. *Chem. Rev.* **2014**, *114*, 815-862.

(23) Yu, Y.; Kalinowski, D. S.; Kovacevic, Z.; Siafakas, A. R.; Jansson, P. J.; Stefani, C.; Lovejoy, D. B.; Sharpe, P. C.; Bernhardt, P. V.; Richardson, D. R. Thiosemicarbazones from the Old to New: Iron Chelators that are More than Just Ribonucleotide Reductase Inhibitors. *J. Med. Chem.* **2009**, *52*, 5271-5294.

(24) Stacy, A. E.; Palanimuthu, D.; Bernhardt, P. V.; Kalinowski, D. S.; Jansson, P. J.; Richardson, D. R. Structure-Activity Relationships of Di-2-pyridylketone, 2-Benzoylpyridine, and 2-Acetylpyridine Thiosemicarbazones for Overcoming Pgp-Mediated Drug Resistance. *J. Med. Chem.* **2016**, *59*, 8601-8620.

(25) Thorson, M. K.; Majtan, T.; Kraus, J. P.; Barrios, A. M. Identification of Cystathionine β -Synthase Inhibitors using a Hydrogen Sulfide-Selective Probe. *Angew. Chem. Int. Ed. Engl.* **2013**, *125*, 4739-4742.


(26) Shimada, K.; Otaki, A.; Yanakawa, M.; Mabuchi, S.; Yamakado, N.; Shimoguchi, T.; Inoue, K.; Kagawa, T.; Shoji, K.; Takikawa, Y. A Pummerer-Type Novel Ring Fission of 2-Methylsulfinyl-5,6-dihydro-4H-1,3,4-thiadiazine Derivatives: A Homologation of Aldehydes and Ketones. *Bull. Chem. Soc. Jpn.* **1996**, *69*, 1043-1054.

(27) Heugebaert, F. C.; Willems J. F. A Synthetic Approach to 1,3,4-Thiadiazolidine-2-thiones. *Tetrahedron* **1966**, *22*, 913-923.

(28) Venkatachalam, T. K.; Pierens, G. K.; Reutens, D. C. Synthesis, NMR Structural Characterization and Molecular Modeling of Substituted Thiosemicarbazones and Semicarbazones using DFT Calculations to Prove the Syn/Anti Isomer Formation. *Magn. Reson. Chem.* **2014**, *52*, 98-105.

(29) For a docking study of **43** and other members of this series with the active site of cystathionine γ -lyase, see Supporting Information.

Insert Table of Contents artwork here



Selective, competitive inhibitor of cystathionine γ -lyase (CSE)

CSE IC₅₀ = 1.2 μ M

CBS IC₅₀ > 500 μ M

CSE Selectivity >400:1