

Synthesis and SAR of a novel, potent and structurally simple LTD₄ antagonist of the quinoline class

Andreas von Sprecher^{*1}, Marc Gerspacher, Andreas Beck, Sabine Kimmel,
Hansruedi Wiestner, Gary P. Anderson, Ulrich Niederhauser, Natarajan
Subramanian and Michael A. Bray

Research Department, Novartis Pharma AG, CH-4002 Basel, Switzerland

Received 12 January 1998; accepted 16 March 1998

Abstract

The two geminal ethyl groups in the succinic acid moiety of CGP57698 (4-[3-(7-fluoro-2-quinolinyl-methoxy)phenyl-amino]-2,2-diethyl-4-oxo-butanoic acid) are responsible for the high *in vitro* and *in vivo* potency of this peptidoleukotriene antagonist of the quinoline type. The synthesis and structure activity relationships of CGP57698 and its analogs are described. © 1998 Elsevier Science Ltd. All rights reserved.


Keywords: Leukotrienes; Antagonists

Clinical trials with peptidoleukotriene (pLT) antagonists have supported the concept that pLT's (LTC₄, LTD₄ and LTE₄) are important mediators of human bronchial asthma [1]. These compounds have been tested against LTD₄-induced bronchoconstriction in either normal or asthmatic subjects. Antagonists producing greater than 20 fold shifts in the LTD₄ dose response curves can be expected to inhibit both the early and late phase antigen-induced bronchoconstriction as well as the subsequent antigen-induced airway hyperresponsiveness in sensitive subjects [1]. These compounds also act against exercise- and dry cold air-induced bronchoconstriction and bronchoconstriction induced by aspirin in patients sensitive to non-steroidal antiinflammatory drugs [1]. Such pLT antagonists are bronchodilators in moderate to severe asthmatics and this effect is additive with β -agonists [1]. Thus they improve lung function values (FEV₁ and PEF_R), reduce the need for rescue medication and improve daytime and night-time symptoms [1]. pLT antagonists are also effective in steroid treated patients (e.g. bronchodilation) [1]. pLT antagonists appear to be safe and no mechanism-based side effects have been reported so far. Where exactly pLT antagonists will fit into the treatment of asthma

¹ Fax +41 61 696 60 71; e-mail: andreas.von_sprecher@pharma.novartis.com

is subject of further clinical trials and has not yet been fully determined. Compounds sharing the above clinical profile come from different structural classes [2,3,4] and include ONO 1078 (pranlukast, OnonTM) (**1**) [5], ICI 204219 (zafirlukast, AccolateTM) (**2**) [6], MK-0476 (montelukast sodium, SingulairTM) (**3**) [7] and CGP45715A (iralukast sodium) (**4**) [8]. Iralukast (**4**) is a highly potent pLT antagonist which was in advanced clinical evaluation as an aerosol formulation. As a structural analog of LTD₄, Iralukast requires a demanding synthesis and has limited oral bioavailability. Our objectives for a follow-up compound of iralukast called for structural simplicity combined with high oral potency. pLT antagonists of the quinoline type were considered to be the right target, as the design of quinoline antagonists is relatively simple and such compounds have in general good oral activity [9,10]. Quinoline antagonists are characterized by a quinoline residue substituted in position 2 by a spacer consisting of two atoms and by a second aromatic unit attached to this spacer and carrying an acidic residue. Using this approach assembling of the appropriate structural elements led to the design and the synthesis of CGP57698 (**5**).

Figure 1
pLT antagonists from different structural classes




A number of analogs of CGP57698 (**5**) have been synthesized in order to determine the SAR in this series of quinoline pLT antagonists. Compounds **5** to **23** in Table 1 illustrate the influence of the substitution in the quinoline moiety on the biological activity. The 7-fluorine substitution in the quinoline residue of CGP57698 (**5**) is optimal in terms of the LTD₄ antagonist activity *in vitro* (Table 1) as well as *in vivo* (Table 2). Compounds **6** to **8** with fluorine in the positions 5,6 and 8 of the quinoline system and the di-fluoro compound **9** have somewhat reduced *in vitro* activities in the subnanomolar to nanomolar range. The 7-chloro

derivative **10** and the unsubstituted compound **12** showed high *in vitro* and *in vivo* potency, whereas the 7-bromo derivative **11** was already *in vitro* 20 times less active than CGP57698 (**5**). With an IC_{50} value of 3 nM *in vitro* the trifluoromethyl derivative **13** had surprisingly no oral activity *in vivo* up to a dose of 10 mg/kg. Compounds **14** to **23** with a variety of other quinoline substituents were considerably (45 times to more than 3 orders of magnitude) less potent than CGP57698 (**5**) *in vitro*. Polar groups like hydroxy or bulky groups like benzyl appear to be detrimental to the antagonist activity. The structure of the acidic residue is of great importance for the pLT antagonist potency of quinoline type antagonists as shown by compounds **24** to **31** in table 1. In CGP57698 (**5**) the acidic group is derived from succinic acid with two geminal ethyl groups in position 2, which are essential for the high potency of CGP57698 (**5**). The unsubstituted succinic acid derivative **24** was 3 orders of magnitude less potent *in vitro*. The activity is back in the nanomolar range with a selection of alkyl groups in position 2 of the succinic acid. An almost complete loss of activity occurred with a shift of the two geminal groups from position 2 to position 3 in the succinic acid residue (compound **31**). With compounds **32** to **34** in table 1 the influence of different two atom spacers on the antagonist activity has been explored. Inversion of the atom sequence in the spacer from -CH₂O- to -OCH₂- (compound **32**) practically abolished the *in vitro* activity. Replacement of oxygen by sulfur, while maintaining the normal atom sequence in the spacer (compound **33**) decreased the *in vitro* potency 20 times, whereas compound **34** with a double bond as spacer was equipotent to CGP57698 (**5**) *in vitro*.


Table 2 demonstrates the excellent potency of CGP57698 (**5**) in comparison with ICI 204219 *in vitro* as well as *in vivo*. The complete pharmacological profile of **5** will be published elsewhere [11]. Compounds **10** and **12** with subnanomolar activity *in vitro* in the guinea-pig ileum test had also *in vivo* comparable activity to CGP57698 (**5**) (table 2).

Table 1
SAR of CGP57698 (**5**) and its analogs **6–34**



compound	R ₁	R ₂	R ₃	R ₄	R ₅	X	IC ₅₀ (nM) ^a
CGP57698 (5)	7-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0.2
6	5-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0.3
7	6-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	2
8	8-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	6

Table 1 continued

compound	R ₁	R ₂	R ₃	R ₄	R ₅	X	IC ₅₀ (nM) ^a
9	6,7-di-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0.5
10	7-Cl	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0.8
11	7-Br	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	4
12	-H	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0.9
13	7-CF ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	3
14	7-CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	11
15	6-CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	24
16	6,7-di-CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	11
17	6-(CH ₂) ₃ -7	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	9
18	7-OCH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	80
19	7-OH	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0 @ 240
20	7-OCH ₂ Ph	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	0 @ 200
21	7-OCH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	80
22	7-OCF ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	9
23	7-COCH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	2500
24	7-F	-H	-H	-H	-H	-CH ₂ O-	200
25	7-F	-CH ₃	-CH ₃	-H	-H	-CH ₂ O-	1.4
26	7-F	-CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ O-	6.4
27	7-F	-n-propyl	-n-propyl	-H	-H	-CH ₂ O-	2
28	7-F	-n-butyl	-n-butyl	-H	-H	-CH ₂ O-	1
29	7-F	-(CH ₂) ₄ -		-H	-H	-CH ₂ O-	10
30	7-F	-(CH ₂) ₅ -		-H	-H	-CH ₂ O-	7.3
31	7-F	-H	-H	-CH ₂ CH ₃	-CH ₂ CH ₃	-CH ₂ O-	140
32	-H	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-OCH ₂ -	3900
33	7-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H	-CH ₂ S-	4
34	7-F	-CH ₂ CH ₃	-CH ₂ CH ₃	-H	-H		0.4

^a LTD₄ antagonism guinea-pig ileum [12] (n=3-5)

Table 2

In vitro and *in vivo* activity of CGP57698 (5), ICI 204219 (2) 10, 12 and 13

Compound	Inhib. of [³ H]-LTD ₄ binding to guinea-pig lung membranes IC ₅₀ (nM) [13] (n=4)	LTD ₄ antagonism guinea-pig ileum IC ₅₀ (nM) [12] (n=3-5)	LTD ₄ -induced bronchospasm guinea-pig ED ₅₀ (mg/kg) p.o. -2 h [14] (n=3-6)
CGP57698 (5)	6	0.2	0.07
ICI 204219 (2) ^a	44	5	0.55
10	64	0.8	0.11
12	9	0.9	0.42
13	not determined	3	inactive at 10 mg/kg p.o.


^a Values determined at Novartis Pharma AG, Basel

The structures shown in figure 2 complete the SAR picture of this series of quinoline pLT antagonists. The methyl ester **35** of CGP57698 (**5**) had weak LTD₄ antagonist activity with an IC₅₀ value of 110 nM *in vitro*. The corresponding tetrazole **36** showed similar *in vitro* activity as the parent compound. Interestingly the *N*-methyl derivative **37** was practically inactive. Inversion of the amide (compound **38**), replacement of the central phenyl nucleus by a naphthyl

residue or replacement of the quinoline system by a pyridyl ring (compounds **39** and **40**) had to be paid with considerable weaker activity *in vitro*.

Figure 2

More detailed SAR of quinoline pLT antagonists 35–40




In parentheses: IC₅₀ values for the inhibition of LTD₄-induced guinea-pig ileum contractions [12]

CGP57698 (**5**) and its analogs can be synthesized in only four steps (sequence a,b,c,d scheme 1). A modified Skraup reaction between 3-fluoroaniline, crotonaldehyde and the oxidant *p*-chloranil gave rise to 7-fluoroquinaldine (**41**) [15]. Side chain bromination under carefully controlled conditions followed by substitution with 3-aminophenol led to the aniline derivative **43**, which upon condensation with 2,2-diethylsuccinic anhydride (**44**) [16] yielded the desired product **5**. Alternatively side chain chlorination of the quinaldine **41** can be achieved *via* the *N*-oxide **42** (sequence e,f scheme 1). The four step sequence has been used to prepare CGP57698 (**5**) on a 10 kg scale.

Scheme 1

Synthesis of CGP57698 (**5**)



Reagents and conditions: a) *p*-Chloranil, 2-butanol/HCl, 60°, 3 h; b) 1. NaOH, H₂O, isopropylacetate; 2. NBS, AIBN, 78°, 5 h (X=Br); c) NaOCH₃, MeOH, 5–10° (X=Br); c') NaOCH₃, MeOH, DMF, 5–10°, 15 h (X=Cl); d) MTBE, 50°, 5 h; e) MCPBA, CH₂Cl₂, 5°, 16 h; f) Benzenesulfonylchloride, toluene, 50°, 8 h.

In summary CGP57698 (**5**) is a pLT antagonist of the quinoline type with equal potency against LTD₄ and LTE₄. The compound is in comparison to other pLT antagonists (e.g. ICI 204219, MK-0476) simpler to synthesize. CGP57698 (**5**) demonstrated good oral bioavailability (rats and marmosets) [11] and is biologically active in guinea-pig bronchoconstriction models *via* the oral route with high potency and long duration of action. Cardiovascular and central nervous system safety pharmacology as well as 3 month oral toxicity studies in rats and marmosets did not reveal prohibitive results for the further development of the compound [11]. The pharmacological profile indicates that CGP57698 (**5**) might be of value in the treatment of asthma and other pLT dependent diseases.

References

- [1] Ford-Hutchinson AW. Leukotriene antagonists and inhibitors: clinical applications. *Adv. Prostaglandin, Thromboxane, and Leukotriene Res.* 1995; 23: 69-74.
- [2] Brooks CDW, Summers JB. Modulators of leukotriene biosynthesis and receptor activation *J. Med. Chem.* 1996; 39: 2629-2654.
- [3] von Sprecher A, Beck A, Gerspacher M, Sallmann A, Anderson GP, Subramanian N, Niederhauser U, Bray MA. Strategies in the design of peptidoleukotriene antagonists. *J Lipid Mediators* 1993; 6 (1-3): 265-73.
- [4] von Sprecher A, Beck A, Gerspacher M, Bray MA. Peptidoleukotriene antagonists: state of the art. *Chimia* 1992; 46: 304-311.
- [5] Nakai H, Konno M, Kosuge S, Sakuyama S, Toda M, Arai Y, Obata T, Katsube N, Miyamoto T, Okegawa T, Kawasaki A. New potent antagonists of leukotriene C₄ and D₄. 1. Synthesis and structure-activity relationships. *J. Med. Chem.* 1988; 31: 84-2654.
- [6] Brown FJ. Discovery of Accolate™ (ICI 204,219), a peptide leukotriene antagonist for asthma. In: Merluzzi VJ, Adams J, editors. *The Search for Anti-Inflammatory Drugs*. Boston: Birkhäuser Boston, 1995: 161-189.
- [7] Labelle M, Belley M, Gareau Y, Gauthier JY, Guay D, Gordon R, Grossmann SG, Jones TR, Leblanc Y, McAuliffe M, McFarlane C, Masson P, Metters KM, Ouimet N, Patrick DH, Piechuta H, Rochette C, Sawyer N, Xiang YB, Pickett CB, Ford-Hutchinson AW, Zamboni R, Young RN. Discovery of MK-0476, a potent and orally active leukotriene D₄ receptor antagonist devoid of peroxisomal enzyme induction. *Bioorg. Med. Chem. Lett.* 1995; 5: 283-288.
- [8] von Sprecher A, Beck A, Sallmann A, Breitenstein W, Wiestner H, Kimmel S, Anderson GP, Subramanian N, Bray MA. Peptidoleukotriene antagonists: structural analogs of leukotriene D₄ with special emphasis on CGP45715A. *Drugs Fut.* 1991; 16: 827-843.
- [9] Galembo RA, Gavai A, Huang F-C. Recent developments in sulphidopeptide leukotriene receptor antagonists. *Curr. Opin. Ther. Patents.* 1992; 811-833.
- [10] Musser JH, Kreft AF. Substituted-[2-quinolinyl(bridged)aryl] compounds: modulators of eicosanoid biosynthesis and action. *Drugs Fut.* 1990; 15: 73-80.
- [11] von Sprecher A, Gerspacher M, Beck A, Anderson GP, Niederhauser U, Subramanian N, Ball HA, Gentsch C, Vassout A, Felner A, Bittiger H, Hauser K, Giese K, Kraetz J, Bray MA. CGP57698: a structurally simple, highly potent peptidoleukotriene (pLT) antagonist of the quinoline type. In: Sinzinger H, Vane JR, Samuelsson B, Paoletti R, Ramwell PW, Wong PY-K, editors. *Recent Advances in Prostaglandin, Thromboxane, and Leukotriene Research*. New York: Plenum Publishing Corporation, 1998: 169-172, in press.
- [12] Rang HP. *Br. J. Pharmacol. Chemother.* 1964; 22: 356.
- [13] Pong SS, De Haven RN. *Proc. Natl. Acad. Sci. U.S.A.* 1983; 80: 7415.
- [14] Amdur MO, Meade J. *Am. J. Physiol.* 1958; 192: 364.
- [15] Song Z, Mertzman M, Hughes DL. Improved synthesis of quinaldines by the Skraup reaction. *J. Heterocyclic Chem.* 1993; 30: 17-21.
- [16] Le Moal H, Foucaud A, Carrié R, Danion D, Fayat C. N° 152.- Recherches sur la structure et les propriétés physicochimiques des acides succiniques dissymétriquement substitués. II.- Préparation des esters-acides et diesters succiniques mono et α,α -disubstitués. *Bull. Soc. chim.* 1964; 828-830.