## A New 2*H*-Azirin-3-amine as a Synthon for $\alpha$ -Methyl Glutamate

by Florentine M. Hilty<sup>1</sup>), Kathrin A. Brun, and Heinz Heimgartner\*

Organisch-chemisches Institut der Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich

Dedicated to Professor Peter Stanetty on the occasion of his 60th birthday

The synthesis of a novel 2,2-disubstituted 2*H*-azirin-3-amine **10** as a building block for racemic Glu(2Me) is described. This synthon contains an ester group in the side chain. The reaction of **10** with thiobenzoic *S*-acid and the amino acid Z-Val-OH yielded the racemic monothiodiamide **17** and the dipeptide **18** as a mixture of diastereoisomers, respectively (*Scheme 2*). From **18**, each of the protecting groups was removed selectively (*Scheme 3*).

**1. Introduction.** – In the last few years, we have shown that 2*H*-azirin-3-amines ('3-amino-2*H*-azirines') are versatile synthons for 2,2-disubstituted glycines ( $\alpha$ , $\alpha$ -disubstituted  $\alpha$ -amino acids) in peptide synthesis. A useful method for the introduction of such amino acids into peptides is the so-called 'azirine/oxazolone method' [1], which proved to be a convenient preparative access to such peptides. This strategy has been extensively applied in the synthesis of linear oligopeptides [2–8], endothiopeptides [9–13], conformationally restricted cyclic peptides [14–17], and cyclic depsipeptides [17–26] containing 2,2-disubstituted glycines.

Recently, enantiomerically pure 2*H*-azirin-3-amines became available, such as the isovaline (Iva) synthons **1** and **2** [4][27], the Val(2Me), Leu(2Me), and the Ala(2cPent) synthons **3**, **4**, and **5** [27], the Phe(2Me) synthons **6** and **7** [27][28], as well as the synthons for Tyr(2Me) **8** and Dopa(2Me) **9** [29]. The latter two contain protected phenolic hydroxy groups, and are the first examples of enantiomerically pure building blocks with a functionalized side chain. All of these building blocks can be used for the synthesis of stereochemically pure peptides.

In the present paper, we describe the synthesis of a novel building block 10 for Glu(2Me), which contains an ester group as a new functional group in the side chain, and its applicability in the synthesis of model peptides. This racemic synthon is a first step towards the expansion of our library of enantiomerically pure 2*H*-azirin-3-amines.

**2. Results.** – 2.1. *Synthesis of the* 2H-*Azirine* **10**. The 2*H*-azirin-3-amine **10**, *i.e.*, a synthon for 2-methylglutamate (Glu(2Me)), was prepared in gram quantity according to *Scheme* 1.

The synthesis started from freshly distilled 3,4,5,6-tetrahydro-2*H*-pyran-2-one (**11**), which is commercially available. Methylation of **11** in  $\alpha$ -position to the C=O group by deprotonation with lithium disopropylamide (LDA), followed by treatment with MeI, yielded **12**. Instead of hexamethylphosphoric triamide (HMPA), 1,3-dimethylimida-

<sup>1)</sup> Part of the diploma thesis of F. M. H., Universität Zürich, 2003.

<sup>© 2004</sup> Verlag Helvetica Chimica Acta AG, Zürich



zolidin-2-one (DMI), which is of lower toxicological risk, was used as an additive [30]. Although *Li et al.* used lithiumhexamethyldisilazanide as a base for similar reactions [30], we preferred LDA in combination with DMI. Therefore, the yield (44%) was lower than reported (70%) [31].

Hydroxy amide **13** was synthesized directly from **12** by the reaction with *N*-methylaniline in the presence of AlCl<sub>3</sub> at room temperature. Due to its carcinogenic properties, the recommended solvent, 1,2-dichlorethane [32], was replaced with  $CH_2Cl_2$ . The yield in  $CH_2Cl_2$  (81%) is only slightly lower than in 1,2-dichlorethane (88%).

In the next step, the OH group of **13** was oxidized with ruthenium trichloride hydrate ( $RuCl_3 \cdot H_2O$ ) and sodium metaperiodate ( $NaIO_4$ ) to form the carboxylic acid **14**. This method, in which ruthenium tetroxide ( $RuO_4$ ) is the active species [33], has the advantage that  $RuCl_3 \cdot H_2O$  can be used in catalytic amounts, and the stochiometric oxidazing agent is  $NaIO_4$ . The task of  $NaIO_4$  is to re-oxidize the reduced forms of the Ru complex to  $RuO_4$ . As a solvent, a mixture of  $CCl_4$ , MeCN, and  $H_2O$  in the ratio 2:2:3 was used [34] in a first attempt. As  $CCl_4$  is toxic and ecologically undesirable, it was replaced with the same quantity of AcOEt [35]. As a result, the yield turned out to be lower (73%) than with  $CCl_4$  (84%).

Methylation of 14 with  $CH_2N_2$  gave the ester 15 in quantitative yield. The synthesis of 10 by the method of *Villalgordo* and *Heimgartner* [36][37] was unsuccessful, even though 15 is an *N*-alkyl-*N*-phenyl amide (*Scheme 1*). It is assumed that deprotonation



LDA = Lithium diisopropylamide; DMI = 1,3-dimethylimidazolidin-2-one; DPPCI = diphenylphosphorochloridate; DABCO = 1,4-diazabicyclo[2.2.2]octane

occurred in  $\alpha$ -position to the ester group instead of the  $\alpha$ -position to the amide group. According to another well-established method, the amide **15** was first converted to the corresponding thioamide **16** with *Lawesson* reagent in toluene at 130° in 93% yield. Finally, the synthesis of **10** was achieved by consecutive treatment of **16** with 2N COCl<sub>2</sub> solution in CH<sub>2</sub>Cl<sub>2</sub>, deprotonation with 1,4-diazabicyclo[2.2.2]octane (DABCO) in THF, and treatment with NaN<sub>3</sub> in THF/DMF, in 90% yield.

2.2. Reactions of **10** with PhCOSH and Z-L-Valine. To demonstrate that the new amino acid synthon **10** shows analogous chemical behavior as the already known 2*H*-azirin-3-amines (*cf.* [1]), it was reacted with PhCOSH [27-29][38][39] (*cf.* [11][12]) to give the monothiodiamide **17** in 99% yield (*Scheme 2*). The use of **10** as a synthon in peptide synthesis was shown by the reaction with Z-L-valine (*Scheme 2*), which led to the dipeptideamide **18** in 96% yield as a mixture of diastereoisomers. All attempts to separate the diastereoisomers failed.

2.3. Selective Cleavage of the Protecting Groups in Dipeptide 18. With the aim of proving the usefulness of the described coupling reaction, each of the protecting groups of the dipeptide 18 was removed selectively under standard or slightly modified conditions (*Scheme 3*). For example, the Z group was removed by hydrogenolysis to give the *N*-deprotected dipeptide 19 in quantitative yield.



Hydrolysis of the C-terminal amide group of **18** under standard conditions (3N HCl in THF/H<sub>2</sub>O 1:1) afforded a mixture of starting material (40%), and the dipeptides **20** (27%) and **21** (14%) with a deprotected carboxy group in the side chain and main chain, respectively. As the cleavage of the methyl ester in the side chain occurs so easily, it was assumed that the reaction follows the mechanism presented in *Scheme 4*: under

the reaction conditions, the tetrahydro-1,3-oxazepin-7-one **22** could be formed instead of the oxazolone **23**. Opening of the ring of **22** with  $H_2O$  to form **20** is expected to proceed smoothly.



Another possibility for the hydrolysis of dipeptide amides is the treatment with HCl gas in toluene, followed by hydrolysis with H<sub>2</sub>O. Thereby, the oxazolone **23** is formed as an intermediate (*Scheme 3*). After treatment of **18** with HCl gas for 13 min in toluene at 100°, oxazolone **23** was isolated in 71% yield after CC. In addition, 22% of starting material **18** was recovered, but no acid **21** was obtained. The oxazolone **23** was not hydrolized by stirring in H<sub>2</sub>O at room temperature overnight. Only after addition of one drop of 6N HCl and stirring at 50° for several hours, was the acid **21** obtained in 71% yield. The direct conversion of **18** to **21**, without purification of the intermediate oxazolone, gave the acid **21** in an overall yield of 70%, and 15% of the starting material **18** and 12% of the oxazolone **23** were isolated from the reaction mixture.

Finally, the selective hydrolysis of the methyl ester in **18** was achieved under standard conditions with LiOH in THF/MeOH/H<sub>2</sub>O 3:1:1 in quantitative yield (*Scheme 3*).

**3.** Conclusions. – The novel racemic 2,2-disubstituted 2H-azirin-3-amine **10** was prepared. This new synthon for  $\alpha$ -methylglutamate was successfully reacted with Z-protected value and thereby incorporated into a model dipeptide. Each protecting group could be removed selectively in good-to-excellent yield. Therefore, this synthon can easily be used in peptide synthesis as a building block for Glu(2Me). The synthesis of this racemic synthon is a first step towards the corresponding enantiomerically pure 2H-azirin-3-amine.

We thank the analytical departments of our institute for NMR and mass spectra, and for elemental analyses. Financial support of the Swiss National Science Foundation, F. Hoffmann-La Roche AG, Basel, the Stiftung für wissenschaftliche Forschung an der Universität Zürich, and the Prof. Dr. Hans E. Schmid-Stiftung is gratefully acknowledged.

## **Experimental Part**

1. General. See [27]. IR Spectra: Perkin-Elmer spectrometer. <sup>1</sup>H- (600 MHz) and <sup>13</sup>C-NMR (150.9 MHz) spectra: Bruker AMX-600 instrument.

2. Preparation of the  $\alpha$ -Methylglutamate Synthon **10**. 2.1. 3,4,5,6-Tetrahydro-3-methylpyran-2(2H)-one (**12**). A soln. of (i-Pr)<sub>2</sub>NH (15 ml, 106 mmol) in abs. THF (40 ml) was cooled to 0°; 1.6M BuLi in hexane (67 ml, 107 mmol) was added, and the mixture was stirred for 30 min, cooled to  $-65^{\circ}$ , and freshly distilled 3,4,5,6-tetrahydro-2H-pyran-2-one (**11**; 10.060 g, 100 mmol) was added at  $-65^{\circ}$  to  $-60^{\circ}$ . After stirring for 1 h, 1,3-

dimethylimidazolidin-2-one (DMI; 14 ml, 129 mmol) was added at  $-65^{\circ}$ , the mixture was stirred for 20 min, and MeI (7 ml, 112 mmol) was added at  $-65^{\circ}$ . After further stirring for 4 h at  $-65^{\circ}$ , the reaction was terminated by addition of a very small amount of H<sub>2</sub>O and AcOH. The org. layer was separated, and the aq. layer was extracted with AcOEt. The combined org. layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. CC (hexane/AcOEt 3 :2) yielded 4.098 g (44%) of **12**. Colorless oil. *R*<sub>f</sub> (hexane/AcOEt 3 :2) 0.43. IR (neat): 2939*m*, 1738*s*, 1462*m*, 1380*m*, 1243*m*, 1156*m*, 1117*m*, 1085*m*, 1029*m*, 1014*m*, 944*w*, 905*w*, 752*w*. <sup>1</sup>H-NMR: 4.40–4.25 (*m*, CH<sub>2</sub>O); 2.65–2.55 (*m*, MeCH); 2.2–2.05 (*m*, 1 H of MeCHCH<sub>2</sub>); 1.95–1.85 (*m*, CH<sub>2</sub>); 1.6–1.5 (*m*, 1 H of MeCHCH<sub>2</sub>); 1.24 (*d*, *J* = 6.9, Me). <sup>13</sup>C-NMR: 175.2 (*s*, CO); 68.3 (*t*, CH<sub>2</sub>O); 34.2 (*d*, CH); 26.7, 21.7 (2*t*, 2 CH<sub>2</sub>); 16.3 (*a*, Me).

2.2. 5-Hydroxy-2,N-dimethyl-N-phenylpentanamide (13). 2.2.1. Procedure 1. To a soln. of AlCl<sub>3</sub> (1.124 g, 8.43 mmol, 2 equiv.) in 1,2-dichlorethane (3 ml), N-methylaniline (1.75 ml, 16.1 mmol, 3.8 equiv.) was added at  $15-25^{\circ}$  (temp. control with ice bath). Thereby, the soln. turned black. Then, compound 12 (0.483 g, 4.23 mmol) in 3 ml of 1,2-dichlorethane was added at  $15-25^{\circ}$ , and the mixture was stirred for 5 h. To the grey-brown suspension, 5 ml of H<sub>2</sub>O was added, and the mixture was stirred for 30 min. The org. layer was separated, and the aq. layer was extracted with 1,2-dichlorethane and twice with CH<sub>2</sub>Cl<sub>2</sub>. The combined org. layers were dried (MgSO<sub>4</sub>) and evaporated. CC (hexane/AcOEt 1:1 to AcOEt) yielded 0.822 g (88%) of 13. Slightly brown solid.

2.2.2. *Procedure* 2. To a soln. of AlCl<sub>3</sub> (1.134 g, 8.5 mmol, 2 equiv.) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml), *N*-methylaniline (1.75 ml, 16.1 mmol, 3.8 equiv.) was added slowly at  $15-25^{\circ}$  (temp. control with ice bath). Thereby, the soln. turned black. Compound **12** (0.489 g, 4.28 mmol) was added at  $15-25^{\circ}$ , and the mixture was stirred for 5 h. To the grey-brown suspension, 5 ml of H<sub>2</sub>O was added, and the mixture was stirred for 30 min and passed through *Celite*. The layers were separated, and the aq. layer was extracted twice with CH<sub>2</sub>Cl<sub>2</sub>. The combined org. layers were dried (MgSO<sub>4</sub>) and evaporated. CC (hexane/AcOEt 1:1 to AcOEt) yielded 0.785 g (81%) of **13**. Slightly brown solid. M.p. 68.9–69.5°. *R*<sub>1</sub> (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 30:1) 0.29–0.16. IR (KBr): 3383s, 2971*m*, 2946*m*, 2923*m*, 2864*m*, 1637s, 1594s, 1495*m*, 1465*m*, 1432*m*, 1391*m*, 1370*w*, 1331*w*, 1274*m*, 1227*w*, 1176*w*, 1116*m*, 1067*m*, 1024*w*, 986*w*, 954*w*, 909*w*, 775*m*, 701s. <sup>1</sup>H-NMR: 7.45–7.3 (*m*, 3 H<sub>m</sub>, H<sub>p</sub>); 7.2–7.15 (*m*, 2 H<sub>o</sub>); 3.55–3.5 (*m*, CH<sub>2</sub>OH); 3.26 (*s*, MeN); 2.45–2.35 (*m*, CH); 1.8–1.7 (*m*, 1 H of CH<sub>2</sub>); 1.5–1.4 (*m*, CH<sub>2</sub>); 1.4–1.3 (*m*, 1 H of CH<sub>2</sub>); 1.04 (*d*, *J* = 6.7, Me). <sup>13</sup>C-NMR: 176.7 (*s*, CO); 144.0 (*s*, 1 arom. C); 129.7 (*d*, 2 C<sub>m</sub>); 127.7 (*d*, 1 C<sub>p</sub>); 127.3 (*d*, 2 C<sub>o</sub>); 62.4 (*t*, CH<sub>2</sub>OH); 37.3 (*q*, MeN); 36.2 (*d*, CH); 30.6, 30.3 (2*t*, 2 CH<sub>2</sub>); 18.3 (*q*, Me). CI-MS (NH<sub>3</sub>): 223 (15), 222 (100, [*M*+1]<sup>+</sup>), 220 (15), 204 (7, [*M*-OH]<sup>+</sup>). Anal. calc. for C<sub>13</sub>H<sub>19</sub>NO<sub>2</sub> (221.30): C 70.56, H 8.65, N 6.33; found: C 70.38, H 8.67, N 6.27.

2.3. 4-Methyl-5-[methyl(phenyl)amino]-5-oxopentanoic acid (14). Compound 13 (3.21 g, 14.5 mmol) and 12.71 g (59.4 mmol, 4.1 equiv.) of NaIO<sub>4</sub> were solved in a mixture of 24 ml of MeCN, 24 ml of AcOEt, and 36 ml of H<sub>2</sub>O. A small amount of RuCl<sub>3</sub> · H<sub>2</sub>O was added at r.t. After 4 h, the color of the suspension changed from light yellow to brown, which indicated the end of the conversion. H<sub>2</sub>O was added, and the aq. layer was extracted with AcOEt. The org. layers were combined, dried (Na<sub>5</sub>SO<sub>4</sub>), and evaporated. Recrystallization from AcOEt/ hexane 1:1 yielded 2.48 (73%) of 14. Colorless crystals. M.p. 116.1–116.7°.  $R_f$  (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 10:1) 0.34. IR (KBr): 3445w, 2961m, 1729s, 1611s, 1586s, 1497m, 1466m, 1454m, 1401m, 1333w, 1316w, 1274m, 1260m, 1193s, 1169m, 1115m, 1095w, 1073w, 1046w, 1027w, 1002w, 976w, 912w, 866w, 796w, 779m, 754w, 716w, 703m. <sup>1</sup>H-NMR: 7.45–7.3 (m, 2 H<sub>m</sub>, H<sub>p</sub>); 7.2–7.15 (m, 2 H<sub>o</sub>); 3.26 (s, MeN); 2.5–2.45 (m, MeCH); 2.35–2.2 (m, CH<sub>2</sub>); 2.0–1.9 (m, 1 H of CH<sub>2</sub>); 1.7–1.6 (m, 1 H of CH<sub>2</sub>); 1.05 (d, J = 6.8, Me). <sup>13</sup>C-NMR: 178.2 (s, COO); 175.9 (s, CON); 143.7 (s, 1 arom. C); 129.7 (d, 2 C<sub>m</sub>); 127.8 (d, 1 C<sub>p</sub>); 127.2 (d, 2 C<sub>o</sub>); 37.4 (q, MeN); 35.6 (d, CH); 31.5, 28.8 (2t, 2 CH<sub>2</sub>); 17.9 (q, Me). CI-MS (NH<sub>3</sub>): 237 (15), 236 (100,  $[M + 1]^+$ ). Anal. calc. for C<sub>13</sub>H<sub>17</sub>NO<sub>3</sub> (235.28): C 66.36, H 7.28, N 5.95; found: C 66.44, H 7.16, N 5.87.

2.4. *Methyl* 4-*Methyl*-5-[*methyl*(*phenyl*)*amino*]-5-oxopentanoate (**15**). To a soln. of **14** (2.506 g, 10.65 mmol) in abs. THF (25 ml), 40 ml of a *ca.* 4N soln. of  $CH_2N_2$  in  $Et_2O$  (prepared according to [40]) were added at 0°, the mixture was stirred until the yellow color disappeared. After 40 min, additional  $CH_2N_2$  soln. (10 ml) was added, and the mixture remained yellow. The ice bath was removed, and the mixture was stirred at r.t., until the yellow color disappeared again. After addition of another 4 ml of the  $CH_2N_2$  soln., the yellow color remained at r.t. for at least 90 min. Then, the excess of  $CH_2N_2$  was destroyed with AcOH, the solvent was evaporated, and the product was dried in high vacuum: 2.696 g (quant.) of **15**. Colorless oil. The product was used for the next step without further purification.  $R_t$  (hexane/AcOEt 2:1) 0.31. IR (neat): 2953s, 1738s, 1653s, 1596s, 1497s, 1436s, 1390s, 1327s, 1268s, 1170s, 1116s, 1074m, 1035m, 1002m, 987m, 918w, 897w, 842w, 800w, 776m, 752m, 703s. <sup>1</sup>H-NMR: 7.45 – 7.35 (*m*, 2 H<sub>m</sub>, H<sub>p</sub>); 7.2 – 7.15 (*m*, 2 H<sub>o</sub>); 3.59 (*s*, MeO); 3.26 (*s*, MeN); 2.5 – 2.4 (*m*, CH): 2.35 – 2.15 (*m*, CH<sub>2</sub>); 2.0 – 1.9 (*m*, 1 H of CH<sub>2</sub>); 1.7 – 1.6 (*m*, 1 H of CH<sub>2</sub>); 1.04 (*d*, *J* = 6.7, Me). <sup>13</sup>C-NMR: 175.9 (*s*, CON); 173.4 (*s*, COOMe); 143.8 (*s*, 1 arom. C); 129.7 (*d*, 2 C<sub>m</sub>); 127.7 (*d*, 1 C<sub>p</sub>); 127.2 (*d*, 2 C<sub>o</sub>); 51.2 (*q*, MeO); 37.3 (*q*, MeN); 35.6 (*d*, CH); 31.6, 29.1 (2*t*, 2 CH<sub>2</sub>); 17.9 (*q*, Me). CI-MS (NH<sub>3</sub>):

251 (15), 250 (100,  $[M + 1]^+$ ). Anal. calc. for C<sub>14</sub>H<sub>19</sub>NO<sub>3</sub> (249.31): C 67.45, H 7.68, N 5.62; found: C 67.22, H 7.93, N 5.54.

2.5. *Methyl* 4-*Methyl*-5-[*methyl*(*phenyl*)*amino*]-5-*thioxopentanoate* (**16**). To a soln. of **15** (2.599 g, 10.43 mmol) in toluene (10 ml), *Lawesson* reagent (2.57 g, 6.35 mmol, 1.2 equiv.) was added, and the mixture was stirred for 30 min at 130°. The excess of *Lawesson* reagent was precipitated with Et<sub>2</sub>O, the precipitate was filtered over *Celite*, and the filtrate was evaporated. CC (hexane/AcOEt 4 : 1) yielded 2.564 g (93%) of **16**. Pale brown oil.  $R_{\rm f}$  (hexane/AcOEt 4 : 1) 0.18. IR (neat): 2969*m*, 2930*m*, 2868*w*, 1737*s*, 1595*m*, 1493*s*, 1444*s*, 1385*s*, 1332*m*, 1269*m*, 1198*s*, 1112*m*, 1074*w*, 1037*m*, 1002*m*, 919*w*, 877*w*, 853*w*, 834*w*, 774*m*, 702*s*. <sup>1</sup>H-NMR: 7.5–7.35 (*m*, 2 H<sub>*m*</sub>, H<sub>*p*</sub>); 7.15–7.1 (*m*, 2 H<sub>*o*</sub>): 3.71, 3.58 (2*s*, MeO, MeN); 2.8–2.7 (*m*, CH); 2.3–2.1 (*m*, 3 H of 2 CH<sub>2</sub>); 1.8–1.55 (*m*, 1 H of 2 CH<sub>2</sub>); 1.32 (*d*, *J* = 6.6, Me). <sup>13</sup>C-NMR: 210.6 (*s*, CS); 173.3 (*s*, CO); 145.4 (*s*, 1 arom. C); 129.9, 128.4, 125.6 (3*d*, 5 arom. CH); 51.3 (*q*, MeO); 45.5 (*q*, MeN); 42.8 (*d*, CH); 32.7, 31.7 (2*t*, 2 CH<sub>2</sub>); 22.0 (*q*, Me). CI-MS (NH<sub>3</sub>): 268 (6), 267 (16), 266 (100, [*M*+1]<sup>+</sup>). Anal. calc. for C<sub>14</sub>H<sub>19</sub>NO<sub>2</sub>S (265.37): C 63.36, H 7.22, N 5.28, S 12.08; found: C 63.34, H 7.20, N 5.28, S 12.11.

2.6. Methyl 3-(3-Amino-2,N-dimethyl-N-phenyl-2H-azirin-2-yl)propanoate (10). To a soln. of 16 (2.558 g, 9.64 mmol) and 5 drops of abs. DMF in abs. CH<sub>2</sub>Cl<sub>2</sub> (12 ml) at 0°, 2N phosgene in toluene (6.5 ml, ca. 13 mmol, 1.3 equiv.) was added slowly, the ice bath was removed, the mixture stirred for 25 min, and the solvent evaporated. The residue was dissolved in abs. THF (12 ml), DABCO (1.096 g, 9.77 mmol) was added, and the soln. was stirred for 25 min at r.t. After filtration and addition of abs. DMF (12 ml), NaN<sub>3</sub> (1.256 g, 19.32 mmol, 2 equiv.) was added, the mixture was stirred for 3 d at r.t. and then filtered over Celite, and the filtrate was evaporated. CC (hexane/AcOEt 2:1) yielded 2.143 g (90%) of 10 as a yellow oil. In the <sup>1</sup>H- and <sup>13</sup>C-NMR spectra at 270 and 280 K, doubling of signals was observed, which almost disappeared at 300 K, and showed EXSY cross-peaks. Therefore, it is assumed that two conformers are detected at r.t. They were analyzed by HSQC and HMBC experiments at 270K. R<sub>f</sub> (hexane/AcOEt 2:1) 0.10. IR (neat): 2950w, 2920w, 1749s, 1656w, 1600s, 1502s, 1438m, 1375w, 1347w, 1319w, 1300m, 1286m, 1228m, 1198m, 1161m, 1112m, 1086w, 1071w, 1035w, 987w, 894w, 841w, 756m. <sup>1</sup>H-NMR (600 MHz, 270 K): 7.60 (d, J = 8.1, 2 H<sub>a</sub>, minor conformer); 7.45 – 7.4 (m, 2 H<sub>m</sub>); 7.2-7.1 (m, 1 H<sub>n</sub>); 7.05 (d, J = 7.9, 1 H<sub>n</sub>, major conformer); 3.54 (s, MeO, minor conformer); 3.53 (s, MeO, major conformer); 3.45 (s, MeN, major conformer); 3.43 (s, MeN, minor conformer); 2.35-2.3 (m, 1 H of CH<sub>2</sub>CO, minor conformer); 2.3-2.25 (m, 1 H of CH<sub>2</sub>CO, minor conformer, 1 H of CH<sub>2</sub>CO, major conformer, 1 H of CH<sub>2</sub>C(3), major conformer); 2.15-2.1 (m, CH<sub>2</sub>C(3), minor conformer, 1 H of CH<sub>2</sub>CO, major conformer); 2.05 - 2.0 (m, 1 H of CH<sub>2</sub>C(3), major conformer); 1.48 (s, Me, major conformer); 1.43 (s, Me, minor conformer). <sup>13</sup>C-NMR (151 MHz, 270 K): 173.9 (s, CO, minor conformer); 173.6 (s, CO, major conformer); 166.0 (s, C(3), major conformer); 164.9 (s, C(3), minor conformer); 142.1 (s, 1 arom. C, minor conformer); 142.1 (s, 1 arom. C, major conformer); 129.6 (d, 2 C<sub>m</sub>, major conformer); 129.2 (d, 2 C<sub>m</sub>, minor conformer); 123.2 (d, 1 C<sub>p</sub>, minor conformer); 123.1 (d, 1 C<sub>p</sub>, major conformer); 116.7 (d, 2 C<sub>o</sub>, minor conformer); 115.4 (d, 1 C<sub>o</sub>, major conformer); 51.6 (*q*, MeO); 45.9 (*s*, C(2), major conformer); 37.7 (*q*, MeN, minor conformer); 37.1 (*s*, C(2), minor conformer); 33.5 (q, MeN, major conformer); 32.1 (d, CH<sub>2</sub>C(2), major conformer); 31.2 (d, CH<sub>2</sub>C(2), minor conformer); 29.9 (d, CH<sub>2</sub>CO, major conformer); 29.8 (d, CH<sub>2</sub>CO, minor conformer); 24.5 (q, Me, major conformer); 23.5 (q, Me, minor conformer). CI-MS (NH<sub>3</sub>): 248 (16), 247 (100, [M+1]<sup>+</sup>). Anal. calc. for C14H18N2O2 (246.30): C 68.27, H 7.37, N 11.37; found: C 67.99, H 7.17, N 10.99.

3. Reactions of the α-Methylglutamate Synthon **10** with PhCOSH and Z-L-Valine. 3.1. With PhCOSH. Methyl 4-Methyl-5-[methyl(phenyl)amino]-4-[(phenylcarbonyl)amino]-5-thioxopentanoate (**17**). To PhCOSH (110 mg, 0.8 mmol), a soln. of **10** (182 mg, 0.74 mmol) in abs.  $CH_2Cl_2$  (5 ml) was added, and the mixture was stirred for 1 h at r.t. Prep. TLC (hexane/AcOEt 1:1) gave 281 mg (99%) of **17**. Pale yellow crystals. M.p. 133.9–134.9°.  $R_{\rm f}$  (hexane/AcOEt 2:1) 0.18. IR (KBr): 3444m, 3246m, 3065m, 3004m, 2956m, 2924m, 2853w, 1732s, 1641s, 1601m, 1578m, 1548s, 1490s, 1463s, 1442m, 1363s, 1330m, 1294s, 1254m, 1219m, 1196m, 1177m, 1093s, 1026w, 1006m, 980m, 934w, 898w, 857w, 803w, 773m, 706s. <sup>1</sup>H-NMR: 8.78 (br. s, NH); 7.8–7.75, 7.5–7.35 (2m, 10 arom. H); 3.76, 3.62 (2s, MeO, MeN); 2.95–2.9 (m, 1 H of 2 CH<sub>2</sub>); 2.4–2.35 (m, 2 H of 2 CH<sub>2</sub>); 2.3–2.2 (m, 1 H of 2 CH<sub>2</sub>); 1.68 (s, Me). <sup>13</sup>C-NMR: 206.8 (s, CS); 173.8 (s, COOMe); 164.6 (s, CON); ca. 147 (s, 1 arom. CN); 135.1 (s, 1 arom. C); 131.2, 129.5, 128.6, 128.3, 126.9, 126.5 (6d, 10 arom. CH); 65.0 (s, C(4)); 51.6 (q, MeO); the signal for MeN was not observed; 32.3, 29.3 (2t, 2 CH<sub>2</sub>); 2.60. (q, Me). ESI-MS (MeOH, Na1): 407 (100, [M + Na]<sup>+</sup>). Anal. calc. for  $C_{21}H_{24}N_2O_3$ S (384.49): C 65.60, H 6.29, N.729, S 8.34; found: C 65.57, H 6.38, N 6.62, S 8.13.

3.2. With Z-L-Valine. Methyl (RS)-4-[((S)-2-[[(Benzyloxy)carbonyl]amino]-3-methyl-1-oxobutyl)amino]-4-methyl-5-[methyl(phenyl)amino]-5-oxopentanoate (18). A soln. of 10 (1.03 g, 4.18 mmol) and Z-L-valine (1.11 g, 4.42 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) was stirred at r.t. for 24 h, and evaporated. CC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 50:1) yielded 1.99 g (96%) of 18. Colorless foam. M.p.  $103 - 104^{\circ}$ .  $R_{\rm f}$  (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 50:1) 0.16. IR (KBr): 3337m, 2963m, 1733s, 1672s, 1633s, 1594m, 1495s, 1454m, 1371m, 1233m, 1110m, 1026m, 774w, 703m. <sup>1</sup>H-NMR: 7.46 (br. s, NH of Glu(2Me), diastereoisomer B); 7.45–7.25 (*m*, 10 arom. H); 7.16 (br. *s*, NH of Glu(2Me), diastereoisomer A); 5.4–5.3 (*m*, NH of Val); 5.2–5.1 (*m*, PhCH<sub>2</sub>O); 4.0–3.9 (*m*, CH(2) of Val, B); 3.9–3.8 (*m*, CH(2) of Val, A); 3.65 (*s*, MeO, B); 3.64 (*s*, MeO, A); 3.27 (*s*, MeN, B); 3.26 (*s*, MeN, A); 2.7–2.45 (*m*, 1 H of 2 CH<sub>2</sub>); 2.45–2.15 (*m*, 2 H of 2 CH<sub>2</sub>); 2.15–2.0 (*m*, CH(3) of Val); 1.95–1.8 (*m*, 1 H of 2 CH<sub>2</sub>); 1.42 (*s*, Me(3) of Glu(2Me), B); 0.95–0.85 (*m*, 2Me(4) of Val). <sup>13</sup>C-NMR: 173.4, 171.9, 171.6, 169.2 (4*s*, COOMe, 2 CON); 156.1 (*s*, OCONH); 143.6, 143.4, 136.4 (3*s*, 2 arom. C); 129.5, 128.6, 128.43, 128.37, 128.0, 127.9 (6*d*, 10 arom. CH); 66.8 (*t*, PhCH<sub>2</sub>O); 61.8, 61.5 (2*s*, C(2) of Glu(2Me)); 60.5, 60.2 (2*d*, CH(2) of Val); 51.6, 51.5 (2*q*, MeO); 41.7, 41.6 (2*q*, MeN); 31.5, 30.8, 29.4, 29.2 (4*t*, 2 CH<sub>2</sub> of Glu(2Me)); 31.4, 31.3 (2*d*, CH(3) of Val); 23.2, 19.1, 19.0, 17.5, 17.2 (5*q*, Me(3) of Glu(2Me), 2 Me of Val). ESI-MS (MeOH, NaI): 520 (100,  $[M + Na]^+$ ). Anal. calc. for C<sub>27</sub>H<sub>38</sub>N<sub>3</sub>O<sub>6</sub> (497.58): C 65.17, H 7.09, N 8.44; found: C 65.06, H 7.15, N 8.51.

4. Deprotection of Dipeptide **18**. 4.1. Cleavage of the Z Group. Methyl (RS)-4-[[(S)-2-Amino-3-methyl-1oxobutyl]amino]-4-methyl-5-[methyl(phenyl)amino]-5-oxopentanoate (**19**). A soln. of dipeptide **18** (200 mg, 0.402 mmol) and a small amount of Pd/C (10% on activated charcoal) in MeOH (10 ml) was treated with H<sub>2</sub> for 2.5 h at r.t. The mixture was filtered over *Celite*, and the filtrate was evaporated. Prep. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 20:1) gave 147 mg (quant.) of **19**. Colorless, highly viscous substance.  $R_1$  (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 10:1) 0.25. IR (neat): 3322m, 2959s, 2874m, 1737s, 1638s, 1594m, 1495s, 1452m, 1368m, 1271m, 1222m, 1200s, 1174s, 1110m, 1082w, 1033w, 995w, 852w, 777w, 735w, 706m. <sup>1</sup>H-NMR: 8.15, 7.80 (2 br. s, NH of Glu(2Me)); 7.45 – 7.3 (m, 5 arom. H); 3.68, 3.65 (2s, MeO); 3.28, 3.26 (2s, MeN); 3.11, 3.01 (2d, J = 4.1 and 3.8, CH(2) of Val); 2.7 – 2.0 (m, 2 CH<sub>2</sub>, CH(3) of Val); 1.95 – 1.9 (m, NH<sub>2</sub> of Val); 1.48, 1.39 (2s, Me(3) of Glu(2Me)); 0.96, 0.93, 0.83, 0.79 (4d, J = 7.0, C), 6.9, and 6.9, 2 Me(4) of Val). <sup>13</sup>C-NMR: 173.5, 173.3, 172.4, 172.2, 171.6 (5s, 3 CO); 144.5, 144.1 (2s, arom. C); 129.4, 129.3, 128.1, 128.0, 127.8, 127.7 (6d, 5 arom. CH); 60.9 (s, C(2) of Glu(2Me)); 60.33, 60.25 (2d, CH(2) of Val); 51.6, 51.5 (2q, MeO); 41.6, 41.5 (2q, MeN); 32.9, 31.4 (2t, CH<sub>2</sub>); 30.8 (d, CH(3) of Val); 29.4, 29.1 (2t, CH<sub>2</sub>); 23.7, 23.3, 19.5, 19.4, 16.3 (5q, Me(3) of Glu(2Me)). ESI-MS (MeOH): 749 (23, [2 M + Na]<sup>+</sup>), 727 (40, [2 M + 1]<sup>+</sup>), 402 (17, [M + K]<sup>+</sup>), 386 (100, [M + Na]<sup>+</sup>), 364 (46, [M + 1]<sup>+</sup>), 257 (24, [M - N(Me)Ph]<sup>+</sup>).

4.2. Hydrolysis of the Amide Group. 4.2.1. Methyl 3-[(RS)-2-((S)-1-[[(Benzyloxy)carbonyl]amino]-2methylpropyl)-4-methyl-5-oxo-1,3-oxazol-4-yl]propanoate (23). A soln. of 18 (301 mg, 0.605 mmol) in toluene (60 ml) was heated to 105°. For 13 min, HCl (g) was bubbled through the mixture. During this procedure, the temp. fell to  $90-95^{\circ}$ . The remaining HCl (g) was removed by bubbling N<sub>2</sub> through the soln. for 20 min. The mixture was transferred into another flask with hexane, and the crystals of N-methylanilide chloride precipitated were filtered (55 mg, 0.39 mmol, 64%), and the resulting soln. was evaporated. CC (hexane/AcOEt 2:1) yielded 182 mg of 23, which still contained some AcOEt, and 67 mg of starting material 18 (22%). Calculation of the yield based on NMR integrals gave 168 mg 23 (71%) and 14 mg AcOEt. This material was hydrolized to the corresponding acid. For anal. purposes, 23 was dried in high vacuum. R<sub>f</sub> (hexane/AcOEt 2:1) 0.17. IR (neat): 3338m, 3066w, 3035w, 2966s, 2936m, 2877w, 1823s, 1732s, 1673s, 1526s, 1453m, 1375m, 1311s, 1233s, 1177s, 1146m, 1027m, 966m, 897s, 775w, 740w, 699m. <sup>1</sup>H-NMR: 7.35-7.3 (m, 5 arom. H); 5.35-5.25 (m, NH of Val); 5.15-5.1 (m, PhCH<sub>2</sub>O); 4.55-4.45 (m, CH(2) of Val); 3.65, 3.63 (2s, MeO); 2.4-2.05 (m, CH(3) of Val, 2 CH<sub>2</sub> of Glu(2Me); 1.40 (br. s, Me(3) of Glu(2Me)); 1.02, 0.98, 0.95 (3d, J = 6.8, 6.9, 6.9, 2 Me(4) of Val). <sup>13</sup>C-NMR: 179.5 (s, C(5)); 172.3 (s, COOMe); 163.0 (s, C(2)); ca. 156 (s, OCONH); 136.0 (s, 1 arom. C); 128.4, 128.1, 128.0 (3d, 5 arom. CH); 67.6 (s, C(4)); 67.1 (t, PhCH<sub>2</sub>O); 55.2, 54.7 (2d, CH(2) of Val); 51.6 (q, MeO); 32.3 (t, CH<sub>2</sub>); 30.7, 30.5 (2d, CH(3) of Val); 28.6 (t, CH<sub>2</sub>); 23.5, 23.3, 18.9, 18.8, 17.5, 17.1 (6q, Me(3) of Glu(2Me), 2 Me(4) of Val). ESI-MS (MeOH, NaI): 413 (100, [M + Na]<sup>+</sup>). Anal. calc. for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>6</sub> (390.44): C 61.53, H 6.71, N 7.17; found: C 61.78, H 6.80, N 7.04.

4.2.2. (RS)-2-[((S)-2-[[(Benzyloxy)carbonyl]amino]-3-methyl-1-oxobutyl)amino]-5-methoxy-2-methyl-5oxopentanoic Acid (21). 4.2.2.1. Procedure 1. From Oxazolone 23. A suspension of 158 mg of crude 23 (containing AcOEt, *i.e.*, 146 mg, 0.374 mmol of 23) in 2 ml H<sub>2</sub>O was stirred for 18 h at r.t., 2 ml of THF were added, and the suspension was stirred for 4 h at r.t. and for 1 h at 50°. As no hydrolysis was observed, one drop of 6N HCl was added. After stirring at 50° for 2 h, the hydrolysis was complete. Brine was added, and the soln. was extracted 3 × with AcOEt. The combined org. layers were dried (MgSO<sub>4</sub>) and evaporated: 161 mg (quant.) of crude 21. After Prep. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 10:1), 109 mg (71%) of pure 21 were obtained as a colorless foam.

4.2.2.2. Procedure 2. From Amide **18**. A soln. of **18** (152 mg, 0.305 mmol) in toluene (30 ml) was heated to 110°. For 20 min, HCl (g) was bubbled through the mixture. During this procedure, the temp. fell to  $100-95^{\circ}$ . The remaining HCl (g) was removed by bubbling N<sub>2</sub> through the soln. for 20 min. The mixture was transferred into another flask with hexane, and crystals of N-methylanilide chloride precipitated were filtered, and the resulting soln. was evaporated. This crude material was solved in 2 ml of THF and 2 ml of H<sub>2</sub>O, and 1 drop of 6N HCl was added. After stirring at 50° for 2.5 h, the hydrolysis was complete. Brine was added, and the soln. was extracted 3 × with AcOEt. The combined org. layers were dried (MgSO<sub>4</sub>), and evaporated. Prep. TLC

 $(CH_2Cl_2/MeOH 10:1)$  gave 87 mg (70%) of pure **21**, and a mixture of 23 mg (15%) of starting material **18** and 18 mg (12%) of oxazolone **23** (ratio determined by NMR).

4.2.2.3. Procedure 3. Under Standard Conditions from Amide **18**. A soln. of **18** (152 mg, 0.305 mmol) in 3N HCl (THF/H<sub>2</sub>O 1:1, 5 ml) was stirred for 1 h at r.t. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried (MgSO<sub>4</sub>), and evaporated. Prep. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 10:1) yielded 81 mg (40%) of starting material **18**, and a mixture of 53 mg (27%) of **20** and 23 mg (14%) of **21** (ratio determined by NMR).

Data of **21.** M.p.  $60-62^{\circ}$ . <sup>1</sup>H-NMR: *ca.* 8.9 (br., COOH); *ca.* 7.4 (br. *s*, NH of Glu(2Me)); 7.35–7.25 (*m*, 5 arom. H); 5.79 (*d*, *J* = 8.6, NH of Val); 5.11 (*s*, PhCH<sub>2</sub>O); 4.15–4.1 (*m*, CH(2) of Val); 3.64, 3.62 (2*s*, MeO); 2.55–2.0 (*m*, CH(3) of Val, 2 CH<sub>2</sub> of Glu(2Me)); 1.59 (*s*, Me(3) of Glu(2Me)); 1.0–0.9 (*m*, 2 Me(4) of Val). <sup>13</sup>C-NMR: *ca.* 176, *ca.* 174, 171.5 (3*s*, COOH, COOMe, CCONH); *ca.* 157 (*s*, OCONH); *ca.* 135 (*s*, 1 arom. C); 128.5, 128.2, 128.0 (3*d*, 5 arom. CH); 67.2 (*t*, PhCH<sub>2</sub>O); 60.6 (*d*, CH(2) of Val); 60.0 (*s*, C(2) of Glu(2Me)); 52.0 (*q*, MeO); 31.4 (*t*, 1 CH<sub>2</sub>); 31.0 (*d*, CH(3) of Val); 29.3 (*t*, 1 CH<sub>2</sub>); 22.9, 19.3, 19.1 (3*q*, Me(3) of Glu(2Me), 2 Me(4) of Val). ESI-MS (MeOH): 447 (8,  $[M+K]^+$ ), 432 (23), 431 (100,  $[M+Na]^+$ ). Anal. calc. for C<sub>20</sub>H<sub>28</sub>N<sub>2</sub>O<sub>7</sub>·0.2 H<sub>2</sub>O (412.05): C 58.30, H 6.95, N 6.80; found: C 58.26, H 6.95, N 6.69.

4.3. Hydrolysis of the Ester Group. 4.1.1. (RS)-4-[((S)-2-{[(Benzyloxy)carbonyl]amino]-3-methyl-1oxobutyl)amino]-4-methyl-5-[methyl(phenyl)amino]-5-oxopentanoic Acid (20). To a soln. of 20 (213 mg, 0.428 mmol) in 6 ml of a 3:1:1 mixture of THF, MeOH, and H<sub>2</sub>O, LiOH · H<sub>2</sub>O (54.5 mg, 1.30 mmol, 3 equiv.) was added. The mixture was stirred at r.t. for 2 h, and then neutralized with 6N HCl. The aq. layer was extracted with CH2Cl2, and the combined org. layers were washed with 1N HCl, dried (MgSO4), and evaporated: 213 mg (quant.) of 20. Colorless solid. M.p. 80-81°. R<sub>f</sub> (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 10:1) 0.31. IR (KBr): 3422s (br.), 3065m, 2965m, 2929m, 1717s, 1633s, 1593m, 1495s, 1455m, 1388m, 1261m, 1236m, 1110m, 1085m, 1028m, 801w, 774w, 738w, 701m. <sup>1</sup>H-NMR: 7.55 (br. s, NH of Glu(2Me)); 7.4-7.2 (m, 10 arom. H); 5.7-5.6 (m, NH of Val); 5.15-5.1 (m, PhCH<sub>2</sub>O); 3.95-3.85 (m, CH(2) of Val); 3.27, 3.24 (2s, MeN); 2.7-1.8 (m, CH(3) of Val, 2 CH<sub>2</sub> of Glu(2Me)); 1.43, 1.39 (2s, Me(3) of Glu(2Me)); 0.95 - 0.85 (m, 2 Me(4) of Val). <sup>13</sup>C-NMR: 176.4, ca. 172, ca. 171, 169.8 (4s, COOH, 2 CCON); 156.6, 156.5 (2s, OCONH); 143.4, 136.2 (2s, 2 arom. C); 129.6, 129.4, 128.4, 128.1, 128.0 (5d, 10 arom. CH); 67.0 (t, PhCH<sub>2</sub>O); 61.9, 61.2 (2s, C(2) of Glu(2Me)); 60.8, 60.2 (2d, CH(2) of Val); 41.7, 41.5 (2q, MeN); 31.4, 31.1 (2d, CH(3) of Val); 29.4, 28.9 (2t, 2 CH<sub>2</sub> of Glu(2Me)); 23.2, 23.1, 19.1, 17.7, 17.5 (5q,  $Me(3) of Glu(2Me), 2 Me(4) of Val). CI-MS (NH_3): 484 (12, [M+1]^+), 466 (9, [M-OH]^+), 395 (8), 394 (38$ 378 (10), 377 (45, [M - NMePh]<sup>+</sup>), 376 (8), 125 (8), 109 (8), 108 (100, PhCH<sub>2</sub>OH<sup>+-</sup>). Anal. calc. for C<sub>26</sub>H<sub>33</sub>N<sub>3</sub>O<sub>6</sub> · 0.5 H<sub>2</sub>O (501.58): C 63.40, H 6.96, N 8.53; found: C 63.33, H 6.79, N 8.29.

## REFERENCES

- [1] H. Heimgartner, Angew. Chem. 1991, 103, 271.
- [2] P. Wipf, H. Heimgartner, Helv. Chim. Acta 1990, 73, 13.
- [3] W. Altherr, H. Heimgartner, in 'Peptides 1990', Ed. E. Giralt, D. Andreu, ESCOM Science Publishers, Leiden, 1991, p. 107; W. Altherr, H. Heimgartner, in 'Peptides 1992', Ed. C. H. Schneider, A. N. Eberle, ESCOM Science Publishers, Leiden, 1993, p. 387; W. Altherr, Dissertation, Universität Zürich, 1994.
- [4] C. B. Bucher, H. Heimgartner, Helv. Chim. Acta 1996, 79, 1903.
- [5] R. Luykx, C. B. Bucher, A. Linden, H. Heimgartner, Helv. Chim. Acta 1996, 79, 527.
- [6] G. Suter, S. A. Stoykova, A. Linden, H. Heimgartner, Helv. Chim. Acta 2000, 83, 2961.
- [7] N. Pradeille, H. Heimgartner, J. Pept. Sci. 2003, 9, 827.
- [8] R. T. N. Luykx, A. Linden, H. Heimgartner, *Helv. Chim. Acta* 2003, 86, 4093; R. T. N. Luykx, Dissertation, Universität Zürich, 2000.
- [9] J. Lehmann, H. Heimgartner, Helv. Chim. Acta 1999, 82, 1899.
- [10] J. Lehmann, A. Linden, H. Heimgartner, Tetrahedron 1998, 54, 8721.
- [11] J. Lehmann, A. Linden, H. Heimgartner, Tetrahedron 1999, 55, 5359.
- [12] J. Lehmann, A. Linden, H. Heimgartner, Helv. Chim. Acta 1999, 82, 888.
- [13] R. A. Breitenmoser, H. Heimgartner, Helv. Chim. Acta 2001, 84, 786.
- [14] I. Dannecker-Dörig, H. Heimgartner, in 'Peptides 1990', Ed. E. Giralt, D. Andreu, ESCOM, Leiden, 1991, p. 460; I. Dannecker-Dörig, Dissertation, Universität Zürich, 1995.
- [15] F. S. Arnhold, Dissertation, Universität Zürich, 1997.
- [16] T. Jeremic, A. Linden, H. Heimgartner, *Chem. Biodiv.* 2004, in press; T. Jeremic, Dissertation, Universität Zürich, 2004.

- [17] H. Heimgartner, F. S. Arnhold, S. P. Fritschi, K. N. Koch, J. E. F. Magirius, A. Linden, J. Heterocycl. Chem. 1999, 36, 1539.
- [18] D. Obrecht, H. Heimgartner, Helv. Chim. Acta 1987, 70, 329.
- [19] D. Obrecht, H. Heimgartner, Helv. Chim. Acta 1990, 73, 221.
- [20] D. Obrecht, H. Heimgartner, Helv. Chim. Acta 1984, 67, 526.
- [21] J. E. F. Magirius, Dissertation, Universität Zürich, 1995.
- [22] J. M. Villalgordo, H. Heimgartner, Helv. Chim. Acta 1997, 80, 748.
- [23] K. N. Koch, A. Linden, H. Heimgartner, Tetrahedron 2001, 57, 2311.
- [24] K. N. Koch, A. Linden, H. Heimgartner, Helv. Chim. Acta 2000, 83, 233.
- [25] K. N. Koch, H. Heimgartner, Helv. Chim. Acta 2000, 83, 1881.
- [26] B. Iliev, A. Linden, H. Heimgartner, Helv. Chim. Acta 2003, 86, 3215.
- [27] K. A. Brun, A. Linden, H. Heimgartner, Helv. Chim. Acta 2001, 84, 1756.
- [28] C. B. Bucher, A. Linden, H. Heimgartner, Helv. Chim. Acta 1995, 78, 935.
- [29] K. A. Brun, A. Linden, H. Heimgartner, Helv. Chim. Acta 2002, 85, 3422. [30] B. Li, R. A. Buzon, M. J. Castaldi, Org. Process Res. Devel. 2001, 5, 609.
- [31] M. Oikawa, H. Oikawa, A. Ichihara, Tetrahedron 1995, 51, 6237. [32] P. Lesimple, D. C. H. Bigg, Synthesis 1991, 4, 306.
- [33] W. P. Griffith, S. V. Ley, G. P. Whitcombe, A. D. White, J. Chem. Soc., Chem. Commun. 1987, 1625.
- [34] P. H. J. Carlsen, T. Katsuki, V. S. Martin, K. B. Sharpless, J. Org. Chem. 1981, 46, 3936.
- [35] M. Prashad, Y. Lu, H.-Y. Kim, B. Hu, O. Repic, T. J. Blacklock, Synth. Commun. 1999, 29, 2937.
- [36] J. M. Villalgordo, H. Heimgartner, Tetrahedron 1993, 49, 7215.
- [37] J. M. Villalgordo, H. Heimgartner, Helv. Chim. Acta 1993, 76, 2830.
- [38] C. Jenny, H. Heimgartner, Helv. Chim. Acta 1986, 69, 374.
- [39] C. Strässler, A. Linden, H. Heimgartner, Helv. Chim. Acta 1997, 80, 1528.
- [40] H. G. O. Becker, 'Organikum', 20. bearb. und erw. Aufl., Barth, Heidelberg; Leipzig, 1996, p. 599.

Received June 25, 2004