

Article

Benzoxazolone Carboxamides as Potent Acid Ceramidase Inhibitors: Synthesis and Structure-Activity Relationship (SAR) Studies

Anders Bach, Daniela Pizzirani, Natalia Realini, Valentina Vozella, Debora Russo, Ilaria Penna, Laurin Melzig, Rita Scarpelli, and Daniele Piomelli

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.5b01188 • Publication Date (Web): 11 Nov 2015

Downloaded from <http://pubs.acs.org> on November 12, 2015

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

1
2
3
4
5
6
7 Benzoxazolone Carboxamides as Potent Acid
8
9
10
11 Ceramidase Inhibitors: Synthesis and Structure-
12
13
14
15 Activity Relationship (SAR) Studies
16
17
18
19

20 Anders Bach^{†,‡,§}, Daniela Pizzirani,^{†,§} Natalia Realini,[†] Valentina Vozella,[†] Debora Russo,[†] Ilaria
21
22 Penna,[†] Laurin Melzig,^{†,‡} Rita Scarpelli,[†] Daniele Piomelli^{†,§,*}
23
24
25

26
27 [†]Drug Discovery and Development, Fondazione Istituto Italiano di Tecnologia, Via Morego 30,
28
29 I-16163 Genova, Italy
30
31

32
33 [§]Departments of Anatomy and Neurobiology, Biological Chemistry and Pharmacology,
34
35 University of California, Irvine, California 92697-4625, United States
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Ceramides are lipid-derived intracellular messengers involved in the control of senescence, inflammation, and apoptosis. The cysteine amidase, acid ceramidase (AC), hydrolyses these substances into sphingosine and fatty acid and, by doing so, regulates their signalling activity. AC inhibitors may be useful in the treatment of pathological conditions, such as cancer, in which ceramide levels are abnormally reduced. Here, we present a systematic SAR investigation of the benzoxazolone carboxamides, a recently described class of AC inhibitors that display high potency and systemic activity in mice. We examined a diverse series of substitutions on both benzoxazolone ring and carboxamide side chain. Several modifications enhanced potency and stability, and one key compound with a balanced activity-stability profile (**14**) was found to inhibit AC activity in mouse lungs and cerebral cortex after systemic administration. The results expand our arsenal of AC inhibitors thereby facilitating the use of these compounds as pharmacological tools and their potential development as drug leads.

INTRODUCTION

Acid ceramidase (AC) is a cysteine amidase involved in the metabolism of ceramides, bioactive lipid molecules that belong to the class of sphingolipids.¹ Sphingolipids act as cellular messengers throughout the body, and are involved in the survival, growth and differentiation of cells,² as well as in pathophysiological process such as inflammation and neuropathic pain.³ AC plays a pivotal role in balancing sphingolipid-mediated signalling.⁴ The enzyme is mainly, albeit not uniquely located in lysosomes,⁵ where it cleaves the amide bond of ceramides producing the aliphatic amino alcohol sphingosine along with fatty acid (typically 14-26 carbon atoms in length) (Figure 1). Sphingosine is a metabolic precursor of sphingosine-1-phosphate, which enhances cell survival and proliferation,⁶ while ceramides exert pro-senescent and pro-apoptotic effects in both normal and tumor cells.^{4, 6c, 7} Interestingly, ceramide concentrations increase in cells that are stressed with chemotherapeutic agents, DNA damage and ionizing radiation,^{7f} either through stimulation of *de novo* ceramide biosynthesis, hydrolysis of sphingomyelin or the salvage pathway;^{1a} moreover, the cytotoxic effects of certain anticancer drugs partly depend on *de novo* ceramide biosynthesis.⁸ AC is over-expressed in several cancer types⁹ and in prostate cancer this up-regulation renders cells more resistant to chemotherapy and radiotherapy, while genetic or pharmacological AC inhibition restores sensitivity to therapy *in vivo*.¹⁰ Thus, modulating the ceramide/sphingosine-1-phosphate axis by inhibiting AC could provide a new strategy for future therapeutics against cancer and, possibly, other pathologies in which ceramide metabolism is dysregulated. To test this hypothesis, several AC inhibitors have been developed over the past decades.¹¹

We have recently disclosed a small set of benzoxazolone carboxamides as the first example of systemically active inhibitors of intracellular AC activity.¹² We demonstrated that these

1
2
3 molecules covalently inhibit AC through *S*-acylation of its catalytic nucleophile, Cys-143, with
4 the benzoxazolone ring acting as leaving group. Preliminary structure-activity relationship
5 (SAR) studies showed that a secondary carboxamide moiety is mandatory for activity, as in the
6 initial hit compound **1** (h-AC IC₅₀ = 64 nM) (Figure 2), and that introducing a bromine atom at
7 position 6 on the heterocyclic scaffold leads to a 2-fold increase in inhibitory potency (**2**, IC₅₀ =
8 31 nM), diminishing at the same time chemical and metabolic stability in aqueous buffer and
9 mouse plasma. Also, we observed that replacing bromine with a *p*-fluorophenyl group, as in 6-
10 (4-fluorophenyl)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (**3**, ARN14974)¹²
11 allows for balancing potency (IC₅₀ = 79 nM) with enhanced chemical and metabolic stability
12 (Figure 2). We found that compound **3** inhibits AC in intact cells and *in vivo*, causing a
13 substantial reduction in AC activity in multiple organs and resulting in expected changes of
14 ceramide and sphingosine levels.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 In the present study we expanded our initial SAR study around compound **1** with the objective
33 of demonstrating whether the benzoxazolone carboxamide scaffold is generally capable of
34 providing potent AC inhibitors. We synthesized a series of benzoxazolone carboxamides to
35 explore the roles of both position and stereoelectronic properties of substituents on the
36 heterocyclic scaffold, as well as the effect of side chain modifications on AC inhibition (Figure
37 2). Finally, we tested select derivatives for their stability in aqueous buffer and demonstrated *in*
38 *vivo* AC inhibitory activity of the optimised compound, 6-chloro-2-oxo-*N*-(4-phenylbutyl)-1,3-
39 benzoxazole-3-carboxamide (**14**).
40
41
42
43
44
45
46
47
48
49
50

51 CHEMISTRY

52
53
54
55 A series of benzoxazolone carboxamide derivatives (**4-36**) bearing different substituents on
56 the bicyclic ring system were synthesized, as outlined in Schemes 1-5. In analogy to hit
57
58
59
60

compound **1**, we initially coupled substituted benzoxazolones with the commercial 4-phenylbutyl isocyanate, in the presence of 4-dimethylaminopyridine (DMAP), under basic conditions (Method A).¹³ First, we prepared a set of derivatives (**4-19**) bearing groups with different stereoelectronic properties on the heterocyclic ring (Scheme 1). Coupling reactions proceeded smoothly to give the final carboxamides **4-19** in good yield, with the exception of compounds **4** and **7**, for which no or low conversion was observed, seemingly due to the substituent at position 4 leading to chemical instability. In most cases, the starting benzoxazolone scaffolds were commercially available (**37b**, **37e-f**, **37h-k**); alternatively the properly substituted heterocycles (**37a**, **37c-d**, **37g**, **37l-p**) were obtained from the corresponding 2-aminophenols by intramolecular cyclization reaction in the presence of 1,1'-carbonyldiimidazole (CDI) (Scheme 1).¹⁴

Secondly, we introduced aromatic substituents on the benzoxazolone scaffold by Suzuki-Miyaura coupling reaction between the appropriate bromo heterocycles (**37a-c**, **38**) and commercially available boronic acids (Scheme 2). Couplings at position 5 and 6 of the benzoxazolone system were accomplished in satisfactory yields, while the reactions to get 4- and 7-phenyl benzoxazolones **39a** and **39d** led to a complex mixture, and the intermediates were used as crudes in the following step. *N*-acylation of **39a-g** with 4-phenylbutyl isocyanate yielded the targeted compounds **20-26**.

Next, we synthesized a small set of benzoxazolone carboxamide derivatives (**27-30**) with alkyl/alkenyl-cyclohexyl/phenyl groups at position 6. The appropriate vinyl boronic acid was reacted with 6-bromo-benzoxazolone **38** under Suzuki-Miyaura reaction conditions to afford the corresponding alkenylic derivatives (**40a-b**) in good yields. Hydrogenation of the double bond

yielded compounds **41a-b** quantitatively. Coupling of **40a-b** and **41a-b** to 4-phenylbutyl isocyanate gave the targeted derivatives **27-28** and **29-30**, respectively, in good yield (Scheme 3).

To further expand the series of 6-substituted derivatives, compounds **31-33** bearing electron-donating and bulky alkoxy groups were prepared as shown in Scheme 4. The benzoxazolone **42** was obtained by orthogonal protection of the commercial 6-hydroxy derivative **43** (Scheme 4).¹⁵ *O*-silyl group cleavage by treatment with tetrabutylammonium fluoride (TBAF) gave the *N*-protected 6-hydroxy derivative **44**, which was readily alkylated with commercial alcohols under Mitsunobu reaction conditions, using triphenylphosphine and di-*tert*-butylazodicarboxylate (DBAD). DBAD was degraded under mild conditions in trifluoroacetic acid (TFA),¹⁶ affording *N*-protected-6-alkoxy benzoxazolones **45a-c** in good yield (Scheme 4). Cleavage of the methoxyethoxymethyl (MEM) group with refluxing TFA¹⁵ led only to partial deprotection (**46a-c**); however, subsequent treatment in dimethylformamide (DMF) at 153 °C provided efficient and clean reaction conditions for obtaining benzoxazolones **47a-c**. Coupling with 4-phenylbutyl isocyanate delivered the desired final compounds **31-33** (Scheme 4).

Scheme 5 shows the syntheses of benzoxazolone carboxamides **34-36** bearing a benzoyl group at either position 5 or 6. Acylation of the benzoxazolone ring could be efficiently achieved by means of Friedel-Craft reaction. To get compound **34**, the commercial *N*-acetyl 2-aminophenol was treated with a mild electrophile, such as the complex AlCl₃·DMF in the presence of benzoyl chloride to regioselectively acylate position 4 (Scheme 5A). Subsequent *N*-acetyl group cleavage and ring closure with CDI afforded the targeted 5-benzoyl-3*H*-1,3-benzoxazol-2-one **49**, which was then coupled to 4-phenylbutyl isocyanate. To obtain compounds **35-36**, the unsubstituted benzoxazolone **50** was first *N*-acylated with benzoyl chloride or 4-chlorobenzoyl chloride to give **51a** and **51b**, respectively (Scheme 5B). Heating in

the presence of AlCl_3 promoted the migration of acyl group from the nitrogen to the carbon atom at position 6 (**52a-b**), according to a “Fries-like” rearrangement mechanism.¹⁷ Final *N*-acylation of nitrogen *N*3 with 4-phenylbutyl isocyanate led to the targeted compounds.

Finally, a series of unsubstituted benzoxazolone derivatives with different side chains at the carboxamide moiety was accessed as depicted in Scheme 6. Compounds **53-66** were obtained by activating benzoxazolone **50** with triphosgene, either in pyridine (Method B) (**54, 62-63, 66**) or in DCM in the presence of Et_3N (Method C) (**53, 55-61, 64-65**), and *in situ* quenching with the proper amine (**67a-n**) to incorporate the desired side chains (Scheme 6). Non-commercial amines (**67a-j**) were synthesized through a two-step sequence starting from alcohols **68a-i** and bromide **69** under Mitsunobu and Gabriel reaction conditions, respectively (Scheme 7). Hydrazine-mediated cleavage of phthalimide derivatives **70a-j** led to the desired amines as hydrochloride salts. Non-commercial alcohols **68a-e** were readily obtained by reduction of the corresponding ester (**71a**) or acids (**71b-e**).

RESULTS AND DISCUSSION

We recently disclosed benzoxazolone carboxamides as the first class of potent and systemically active inhibitors of intracellular AC.¹² We demonstrated that these compounds covalently inhibit AC, and identified derivative **3** as the first chemical probe that may be used to study the impact of AC inhibition *in vivo* (Figure 2). A preliminary SAR study around hit compound **1** revealed that a secondary carboxamide moiety is mandatory for activity, and substitution at position 6 of the benzoxazolone ring allows for modulation of the inhibitory potency against AC. These results prompted us to expand the class of benzoxazolone carboxamides by systemically investigating the effects of different substituents on the heterocyclic scaffold as well as of modifications of the side chain. The new compounds were

1
2
3 tested for their ability to inhibit the hydrolysis of *N*-[(1*S*,2*R*)-2-hydroxy-1-(hydroxymethyl)-4-(2-
4 oxochromen-7-yl)oxy-butyl]dodecanamide by recombinant human AC (h-AC) in a fluorescence-
5 based assay.¹⁸
6
7

8
9
10 To probe for stereoelectronic effects on the benzoxazolone system, we initially
11 monosubstituted **1** with bromine or methyl groups at positions 4, 5, 6 or 7 (Table 1). As
12 previously observed with the 6-bromo derivative **2** ($IC_{50} = 31$ nM), introduction of bromine at
13 either position 5 or 7 led to a 2- to 4-fold increase in potency relative to **1** (**5**, $IC_{50} = 22$ nM and **6**,
14 $IC_{50} = 18$ nM). Bromine introduction at position 4 yielded the unstable derivative **4**, and the
15 same result was obtained when a methyl group was introduced at the same position (**7**). As for
16 the small set of methyl-substituted derivatives, with the exception of compound **10** bearing a
17 methyl at position 7 and having an IC_{50} of 23 nM, the introduction of this group led to a decrease
18 in potency when the methyl was placed in position 5 (**8**, $IC_{50} = 117$ nM) or to an equipotent
19 compound, compared to **1**, when the methyl was placed in position 6 (**9**, $IC_{50} = 66$ nM). Next, we
20 expanded the set of substituted analogs at positions 5 and 6 investigating compounds **11-19**,
21 which were 2- to 20-fold more potent than **1** (Table 1). The 5-fluoro derivative **11** ($IC_{50} = 12$
22 nM) turned out to be ~3-fold more potent than the corresponding 6-substituted analog **12** ($IC_{50} =$
23 31 nM). The same trend in potency associated to halide introduction at either position 5 or 6 was
24 observed for the chloro-substituted compounds **13** and **14** (and for the bromine-substituted **5** and
25 **2**). Replacement of halides with strong electron-withdrawing groups such as trifluoromethyl and
26 nitro groups diminished the difference in potency between 5 and 6-substituted derivatives (**15-**
27 **18**). Of particular interest were the results obtained with 5-nitro benzoxazolone carboxamide **17**
28 and the corresponding 6-nitro analog **18**, which represent the first single-digit nanomolar
29 compounds in this class, both inhibiting AC with an IC_{50} value of 3 nM. Finally, we explored
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

whether incorporating simultaneously two favorable substitutions would provide additive effects. We observed that the disubstituted compound **19** was potent ($IC_{50} = 18$ nM), but not significantly more so than the related monosubstituted analogs **13** and **14** (Table 1).

As a continuation of the SAR study around the benzoxazolone system, we prepared a small series of derivatives bearing a phenyl ring (Table 2). First, we systematically explored the introduction of this moiety at positions 4, 5, 6 and 7. Interestingly, stability issues did not prevent isolation and testing of the 4-phenyl derivative **20**, which turned out to be 75 folds more potent than **1** ($IC_{50} = 0.8$ nM). Derivatives **21-23** bearing a phenyl ring at positions 5, 6, and 7, respectively, were approximately as potent as **1**. Next, we focused on positions 5 and 6, and prepared a small series of aryl-substituted derivatives (**24-36**). Overall, decorating the phenyl ring with fluorine or methoxy group in the *para* position (**3**, **24** and **25-26**) did not improve potency over the unsubstituted benzoxazolone derivative **1**. However, replacement of the phenyl with aryl ketones (**34-36**) led to 2- to 5-fold increases in potency relative to **1**, with compound **36** ($IC_{50} = 14$ nM) being the most potent within this small series. Finally, introduction of an alkylic or alkenylic spacer between the benzoxazolone system and the phenyl ring as well as the replacement of the phenyl with a cyclohexyl led to a loss in potency, with compounds **27-30** (Scheme 3) showing no inhibitory activity against AC at the concentrations tested (50 and 500 nM). Similarly, compounds **31-33** (Scheme 4) bearing various alkoxy groups at position 6 were inactive.

We next examined the role of the side chain by preparing a series of unsubstituted benzoxazolone carboxamides. First, we evaluated the length of the carbon linker between the benzoxazolone and the phenyl ring of **1**. Compounds **53-56**, in which the spacer was progressively increased from two to six methylene units, were tested for their inhibitory activity

(Table 3). The short linker with two methylene units led to a 2-fold decrease in potency (**53**, IC_{50} = 121 nM) compared to **1**, while compounds **54-56** – which contain a 3-, 5- or 6-carbon atom linker, respectively – turned out to be as potent as **1**. Replacement of one methylene unit of the four-carbon atom chain of **1** with a heteroatom (**62-64**) was detrimental for activity, with the exception of the thioether derivative **63**, which was equipotent to **1**. Next, to evaluate the role of the phenyl ring, we replaced it with a naphthyl (**57**) or thiophene (**65**) group, observing an increase in potency with the latter derivative (**65**, IC_{50} = 37 nM). Introduction of substituents with different electronic properties in the *para* position of the phenyl ring was generally well tolerated, as shown by compounds **59-61**, with the exception of derivative **58** (IC_{50} = 312 nM), which turned out to be 5-fold less potent than **1**. Interestingly, the most potent compound of this series showed the same side chain length of **1**, but with the phenyl ring situated one carbon atom from the *N*3-carboxamide group (**66**, IC_{50} = 19 nM).

Previously, we reported that introducing bromine in position 6 (**2**) destabilizes compounds and leads to lower half-life in PBS and blood plasma compared to **1**, but that having a *p*-fluorophenyl in position 6 (**3**) enhanced chemical and metabolic stability dramatically. Conversely, stability under standard AC assay conditions (pH 4.5) was not affected by these substitutions.¹² In order to gain a more comprehensive understanding of the structure-stability relationship of benzoxazolone carboxamides, compounds **14**, **19-26** and **35-36** were tested for stability in aqueous buffer at physiological pH (Table 4). Representative compounds (**14**, **17-18**, **21**, **35-36**) were also evaluated for their stability at pH 4.5 (Table S1, Supporting Information). At acidic pH, all compounds were stable, as previously observed with derivatives **1-3**. The analysis at physiological pH showed that, in contrast to bromine, chlorine in position 6 (**14**) or positions 5 and 6 (**19**) did not induce instability relative to the unsubstituted compound **1**.

Aromatic substitutions in positions 5-7 enhanced stability in all cases (**3**, **21-26**), with *p*-fluorophenyl in position 6 (**3**) being the most efficient substituent and *p*-methoxyphenyl in the same position (**26**) the least efficient one. Noticeably, incorporating a phenyl group in position 4 resulted in a very unstable compound (**20**), which tallies well with the chemical instability observed with other compounds (**4**, **7**) with substitutions in the same position. Finally, compound **35** with the benzoyl group in position 6 demonstrated enhanced stability, and interestingly its close analog **36** was even more stable with a half-life 6-fold higher than **1** (Table 4). Overall, these results demonstrate that stability properties are not simply explained by the electronegativity of the substituents on the benzoxazolone ring, and that increased activity is not necessarily linked with decreased stability. Indeed, we see that it is possible to increase potency without decreasing stability, as exemplified by compounds **14** and **19**, and that it is even possible to enhance both potency and stability relative to **1** as seen for compounds **35** and **36**. However, the most potent compound in this series, compound **20** ($IC_{50} = 0.8$ nM, Table 2) also showed the lowest half-life, providing a case in which activity and stability are seemingly related.

To test whether the explored structural modifications of the benzoxazolone carboxamide scaffold could also provide novel systemically active compounds, we selected derivative **14** for further pharmacological studies *in vivo*, based on its balanced profile in terms of potency, stability, solubility and drug-likeness. Indeed, compound **14** shows an improved potency compared to **3**, and an adequate solubility in the vehicle used for systemic administration. In this regard, other slightly more potent and stable derivatives of this series, such as compounds **35-36**, were found to be poorly soluble and thus difficult to handle in *in vivo* studies. As additional criteria of selection, we considered the drug-likeness of **14** in comparison with other considerably more potent compounds, such as the single-digit nanomolar inhibitors **17** and **18**.

The nitro-substituted benzoxazolone ring makes these derivatives important for SAR purposes but not promising as candidates for further development, due to the potential *in vivo* toxicity of aromatic nitro groups.

Compound **14** was injected in mice at the dose of 10 mg kg⁻¹ (i.p.) and AC activity was evaluated at different time points in lysosomal fractions prepared from lung tissue and cerebral cortex. We selected the lungs because of the high basal AC activity in this tissue, and the cerebral cortex to evaluate the ability of **14** to reach the brain. As shown in Figure 3, the compound significantly inhibited AC activity in both compartments. The effect was long lasting, as AC inhibition was still observed 24 hours after administration, although a slight recovery in activity was seen at this time point (Figure 3A-C). In the lungs, AC inhibition was accompanied by an increase in the levels of ceramide and dihydroceramide (namely d18:1/16:0 and d18:0/16:0) for up to 6 hours; sphingosine and sphingosine 1-phosphate were also decreased at 3 hours (Figure 3B). No statistically significant changes in sphingolipid levels were observed in the cerebral cortex (Figure 3D), presumably owing to the relative low level of AC inhibition achieved in that brain region. The results indicate that **14** engages its intended target in two relevant tissues, with results that are comparable to those previously obtained with compound **3** under the same experimental conditions.¹²

CONCLUSIONS

The present study describes a systematic SAR investigation of a recently disclosed class of AC inhibitors that features the benzoxazolone carboxamide scaffold.¹² We show that this scaffold is able to provide stable systemically active AC inhibitors, and may thus represent a good starting point for further discovery. We present various synthetic approaches to obtain diversified analogs in high yields and with great feasibility. We show that electron-withdrawing

groups in positions 5-7 of the benzoxazolone scaffold (**2**, **5-6**, **10-19**) lead to enhanced inhibitory activity toward AC. This is consistent with the mechanism of action of these agents, as electron-withdrawing groups are likely to make the carboxamide carbonyl more prone to nucleophilic attack by AC's active site cysteine. Introducing a phenyl group at position 4 gave a highly potent but very unstable compound (**20**), which correlates with the chemical instability seen with other molecules substituted in the same position (**4** and **7**). Additionally, we investigated a range of aromatic substitutions on positions 5-7 of the benzoxazolone scaffold (**21-26**), which all provide stable compounds but do not improve activity towards AC. Also, we demonstrated that long and bulky aliphatic groups are not favorable for inhibitory activity, as seen with compounds **27-30** and **31-33**. Instead, we found that substitutions with benzoyl (**34-35**) or 4-chlorobenzoyl (**36**) groups combine good potency and high stability: the most favorable of these derivatives, **36**, shows a 5-6-fold improvement in AC inhibition relative to **1** and **3** and a 6-fold improvement in stability relative to **1**. Finally, we addressed the role of the carboxamide side chain and demonstrated that the inhibitory potency of **1** could be improved 2-3 folds by incorporating the phenyl group within the linker (**66**) or replacing it with thiophene (**65**). Based on potency, stability and adequate solubility, we chose to investigate compound **14** for *in vivo* activity. We show that following systemic administration in mice, this compound inhibits AC in peripheral tissues as well as in the brain, leading to the expected variations in the sphingolipid profile. This provides essential evidence that AC inhibition can be successfully obtained in live animals using potent and reasonably stable benzoxazolone carboxamides. Compound **14** might be used as a tool to unravel the biological role of AC and define the potential value of this enzyme as a therapeutic target.

EXPERIMENTAL SECTION

Chemistry. General. All commercial available reagents and solvents were used as purchased from vendors without further purification. Dry solvents (pyridine, DCM) were purchased from Sigma-Aldrich. Automated column chromatography purifications were done using a Teledyne ISCO apparatus (CombiFlash® Rf) with pre-packed silica gel columns of different sizes (from 4 g up to 40 g). Mixtures of increasing polarity of cyclohexane and ethyl acetate (EtOAc) were used as eluents (unless otherwise stated). Microwave heating was performed using Explorer®-48 positions instrument (CEM). Hydrogenation reactions were performed using H-Cube™ continuous hydrogenation equipment (SS-reaction line version), employing disposable catalyst cartridges (CatCart™) preloaded with the required heterogeneous catalyst. NMR experiments were run on a Bruker Avance III 400 system (400.13 MHz for ¹H, and 100.62 MHz for ¹³C), equipped with a BBI probe and Z-gradients. Spectra were acquired at 300 K, using deuterated dimethylsulfoxide (DMSO-*d*₆) or deuterated chloroform (CDCl₃) as solvents. Chemical shifts for ¹H and ¹³C spectra were recorded in parts per million using the residual non-deuterated solvent as the internal standard (for CDCl₃: 7.26 ppm, ¹H and 77.16 ppm, ¹³C; for DMSO-*d*₆: 2.50 ppm, ¹H; 39.52 ppm, ¹³C). UPLC-MS analyses were run on a Waters ACQUITY UPLC-MS system consisting of a SQD (Single Quadrupole Detector) Mass Spectrometer equipped with an Electrospray Ionization interface and a Photodiode Array Detector. PDA range was 210-400 nm. Analyses were performed on an ACQUITY UPLC HSS T3 C₁₈ column (50 × 2.1 mm ID, particle size 1.8 μm) with a VanGuard HSS T3 C₁₈ pre-column (5 × 2.1 mm ID, particle size 1.8 μm). Mobile phase was either 10 mM NH₄OAc in H₂O at pH 5 adjusted with AcOH (A) and 10 mM NH₄OAc in MeCN-H₂O (95:5) at pH 5 (B). Electrospray ionization in positive and negative mode was applied. All final compounds showed ≥ 95% purity by NMR (¹H, ¹³C, ¹H-¹H COSY,

¹H-¹³C HSQC) and UPLC-MS (UV). DMSO stock solutions of final compounds (10 mM) used for biological tests were evaluated prior to tests (NMR, LC-MS), and concentration was assessed by quantitative ¹H-NMR.

General Procedure for the Synthesis of Final Benzoxazolone *N*-Carboxamides 4-36 via Method A (Scheme 1-5, Table 1-2). The properly substituted 3*H*-1,3-benzoxazol-2-one (1.0 equiv) was dissolved in dry pyridine (6 mL per mmol benzoxazol-2-one). DMAP (1.1 equiv) was added and the reaction mixture stirred under nitrogen atmosphere at rt for 30 min. 4-Phenylbutyl isocyanate (1.1 equiv) was added and the resulting mixture was stirred for 15 h. The solvent was removed under reduced pressure, and the compound was purified by silica gel column chromatography using the Teledyne ISCO apparatus (cyclohexane/EtOAc).

5-Bromo-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (5). The reaction was carried out following method A, using **37b** (100 mg, 0.467 mmol) and 4-phenylbutyl isocyanate (90 mg, 0.088 mL, 0.514 mmol). White solid (88 mg, 48%). ¹H NMR (400 MHz, CDCl₃) δ 1.64 – 1.77 (m, 4H), 2.67 (t, *J* = 7.2 Hz, 2H), 3.44 (td, *J* = 6.8, 5.6 Hz, 2H), 7.11 (d, *J* = 8.6 Hz, 1H), 7.16 – 7.21 (m, 3H), 7.26 – 7.31 (m, 2H), 7.37 (dd, *J* = 8.6, 2.1 Hz, 1H), 7.97 (t, *J* = 4.6 Hz, 1H), 8.27 (d, *J* = 2.0 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 28.7, 29.1, 35.6, 40.4, 111.3, 117.9, 119.0, 126.0, 127.6, 128.5 (4C), 129.2, 140.8, 142.0, 149.5, 152.9. MS (ESI) *m/z*: 212 and 214 [M - CONH(CH₂)₄Ph][–].

7-Bromo-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (6). The reaction was carried out following method A, using **37c** (140 mg, 0.654 mmol) and 4-phenylbutyl isocyanate (126 mg, 0.123 mL, 0.720 mmol). White solid (143 mg, 56%). ¹H NMR (400 MHz, CDCl₃) δ 1.61 – 1.78 (m, 4H), 2.67 (t, *J* = 7.1 Hz, 2H), 3.44 (q, *J* = 6.4 Hz, 2H), 7.13 – 7.21 (m, 4H), 7.26 – 7.31 (m, 2H), 7.38 (dd, *J* = 8.3, 1.1 Hz, 1H), 7.99 (t, *J* = 6.0 Hz, 1H), 8.02 (dd, *J* = 8.2, 1.0 Hz,

1H). ¹³C NMR (101 MHz, CDCl₃) δ 28.7, 29.1, 35.6, 40.4, 102.4, 114.7, 126.0, 126.2, 128.0, 128.5 (4C), 129.0, 140.0, 142.0, 149.6, 152.4. MS (ESI) *m/z*: non-ionizable compound under routine conditions used.

4-Methyl-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (7). The reaction was carried out following method A, using **37d** (94 mg, 0.630 mmol) and 4-phenylbutyl isocyanate (121 mg, 0.119 mL, 0.693 mmol). Clear colorless oil (49 mg, 24%). ¹H NMR (400 MHz, CDCl₃) δ 1.64 – 1.79 (m, 4H), 2.53 (s, 3H), 2.68 (t, *J* = 7.2 Hz, 2H), 3.45 (q, *J* = 6.6 Hz, 2H), 7.04 – 7.09 (m, 2H), 7.11 – 7.21 (m, 4H), 7.26 – 7.31 (m, 2H), 7.60 (t, *J* = 4.5 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 21.5, 28.6, 29.1, 35.6, 40.9, 107.6, 124.8, 126.0, 126.4, 126.6, 128.4, 128.5 (2C), 128.5 (2C), 142.1, 143.0, 149.3, 154.0. MS (ESI) *m/z*: 325 [M + H]⁺, 342 [M + NH₄]⁺.

5-Methyl-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (8). The reaction was carried out following method A, using **37e** (50 mg, 0.334 mmol) and 4-phenylbutyl isocyanate (64 mg, 0.063 mL, 0.367 mmol). White solid (99 mg, 91 %). ¹H NMR (400 MHz, CDCl₃) δ 1.64 – 1.77 (m, 4H), 2.41 (s, 3H), 2.67 (t, *J* = 7.2 Hz, 2H), 3.44 (td, *J* = 6.7, 5.5 Hz, 2H), 7.00 – 7.04 (m, 1H), 7.08 – 7.11 (m, 1H), 7.16 – 7.21 (m, 3H), 7.26 – 7.30 (m, 2H), 7.90 (br s, 1H), 8.08 (t, *J* = 4.9 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 21.7, 28.7, 29.2, 35.6, 40.2, 109.5, 116.1, 125.0, 126.0, 128.0, 128.5 (2C), 128.5 (2C), 135.2, 139.9, 142.0, 150.1, 153.6. MS (ESI) *m/z*: 325 [M + H]⁺, 342 [M + NH₄]⁺.

6-Methyl-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (9). The reaction was carried out following method A, using **37f** (50 mg, 0.334 mmol) and 4-phenylbutyl isocyanate (64 mg, 0.063 mL, 0.367 mmol). White solid (73 mg, 68%). ¹H NMR (400 MHz, CDCl₃) δ 1.63 – 1.78 (m, 4H), 2.41 (s, 3H), 2.67 (t, *J* = 7.2 Hz, 2H), 3.44 (td, *J* = 6.7, 5.5 Hz, 2H), 7.03 – 7.08 (m, 2H), 7.15 – 7.21 (m, 3H), 7.26 – 7.30 (m, 2H), 7.89 – 7.92 (m, 1H), 8.04 (t, *J* = 5.4 Hz, 1H).

¹³C NMR (101 MHz, CDCl₃) δ 21.6, 28.7, 29.2, 35.6, 40.2, 110.5, 115.2, 125.6, 125.7, 126.0, 128.5 (2C), 128.5 (2C), 135.0, 142.0, 142.1, 150.0, 153.5. MS (ESI) *m/z*: 325 [M + H]⁺, 342 [M + NH₄]⁺.

7-Methyl-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (10). The reaction was carried out following method A, using **37g** (83 mg, 0.556 mmol) and 4-phenylbutyl isocyanate (107 mg, 0.105 mL, 0.612 mmol). White solid (128 mg, 71%). ¹H NMR (400 MHz, CDCl₃) δ 1.63 – 1.78 (m, 4H), 2.39 (s, 3H), 2.67 (t, *J* = 7.1 Hz, 2H), 3.44 (q, *J* = 6.6 Hz, 2H), 7.04 (d, *J* = 7.8 Hz, 1H), 7.12 – 7.20 (m, 4H), 7.26 – 7.31 (m, 2H), 7.88 (dd, *J* = 8.0, 1.2 Hz, 1H), 8.09 (t, *J* = 5.5 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 14.5, 28.7, 29.2, 35.6, 40.2, 113.11, 120.5, 124.9, 126.0, 126.1, 127.8, 128.5 (2C), 128.5 (2C), 140.5, 142.0, 150.1, 153.5. MS (ESI) *m/z*: 325 [M + H]⁺, 342 [M + NH₄]⁺.

5-Fluoro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (11). The reaction was carried out following method A, using **37h** (61 mg, 0.398 mmol) and 4-phenylbutyl isocyanate (76.7 mg, 0.075 mL, 0.438 mmol). White solid (111 mg, 85%). ¹H NMR (400 MHz, CDCl₃) δ 1.64-1.78 (m, 4H), 2.67 (t, *J* = 7.1 Hz, 2H), 3.44 (q, *J* = 6.6 Hz, 2H), 6.94 (td, *J* = 9.0, 2.7 Hz, 1H), 7.15 – 7.21 (m, 4H), 7.26 – 7.31 (m, 2H), 7.85 (dd, *J* = 8.5, 2.7 Hz, 1H), 8.01 (t, *J* = 6.3 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 28.6, 29.1, 35.5, 40.3, 104.3 (d, *J* = 31.3 Hz), 110.5 (d, *J* = 9.5 Hz), 111.2 (d, *J* = 25.1 Hz), 126.0, 128.5 (4C), 128.7 (d, *J* = 14.0 Hz), 137.8 (d, *J* = 2.2 Hz), 142.0, 149.5, 153.4, 159.8 (d, *J* = 242.4 Hz). MS (ESI) *m/z*: 152 [M - CONH(CH₂)₄Ph]⁻.

6-Fluoro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (12). The reaction was carried out following method A, using **37i** (51 mg, 0.334 mmol) and 4-phenylbutyl isocyanate (64 mg, 0.063 mL, 0.367 mmol). White solid (87 mg, 79%). ¹H NMR (400 MHz, CDCl₃) δ 1.63-1.78 (m, 4H), 2.67 (t, *J* = 7.2 Hz, 2H), 3.44 (td, *J* = 6.7, 5.6 Hz, 2H), 6.96 – 7.04 (m, 2H), 7.16 –

7.21 (m, 3H), 7.25 – 7.31 (m, 2H), 7.96 (t, $J = 5.7$ Hz, 1H), 8.00 – 8.05 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.1, 35.6, 40.3, 99.1 (d, $J = 28.8$ Hz), 111.8 (d, $J = 23.4$ Hz), 116.3 (d, $J = 9.0$ Hz), 124.3 (d, $J = 2.6$ Hz), 126.0, 128.5 (4C), 141.8, 142.0 (d, $J = 2.3$ Hz), 149.6, 153.2, 159.7 (d, $J = 245.2$ Hz). MS (ESI) m/z : 329 $[\text{M} + \text{H}]^+$, 346 $[\text{M} + \text{NH}_4]^+$.

5-Chloro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (13). The reaction was carried out following method A, using **37j** (68 mg, 0.401 mmol) and 4-phenylbutyl isocyanate (76.7 mg, 0.075 mL, 0.438 mmol). Clear colorless oil (127 mg, 92%). ^1H NMR (400 MHz, CDCl_3) δ 1.64 – 1.79 (m, 4H), 2.67 (t, $J = 7.1$ Hz, 2H), 3.44 (td, $J = 6.7, 5.5$ Hz, 2H), 7.13 – 7.24 (m, 5H), 7.26 – 7.31 (m, 2H), 7.98 (t, $J = 5.2$ Hz, 1H), 8.12 (d, $J = 2.1$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.1, 35.6, 40.4, 110.8, 116.3, 124.7, 126.0, 128.5 (4C), 128.9, 130.8, 140.3, 142.0, 149.5, 153.0. MS (ESI) m/z : 345 and 347 $[\text{M} + \text{H}]^+$.

6-Chloro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (14). The reaction was carried out following method A, using **37k** (56 mg, 0.334 mmol) and 4-phenylbutyl isocyanate (64 mg, 0.063 mL, 0.367 mmol). White solid (77 mg, 67%). ^1H NMR (400 MHz, CDCl_3) δ 1.64 – 1.78 (m, 4H), 2.67 (t, $J = 7.2$ Hz, 2H), 3.44 (td, $J = 6.7, 5.6$ Hz, 2H), 7.16 – 7.20 (m, 3H), 7.24 – 7.31 (m, 4H), 7.93 – 8.02 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.1, 35.6, 40.3, 110.9, 116.4, 125.4, 126.1, 126.8, 128.5 (4C), 130.2, 142.0, 142.1, 149.6, 152.9. MS (ESI) m/z : 168 and 170 $[\text{M} - \text{CONH}(\text{CH}_2)_4\text{Ph}]^-$.

2-Oxo-*N*-(4-phenylbutyl)-5-(trifluoromethyl)-1,3-benzoxazole-3-carboxamide (15). The reaction was carried out following method A, using **37l** (128 mg, 0.630 mmol) and 4-phenylbutyl isocyanate (121 mg, 0.119 mL, 0.693 mmol). White solid (166 mg, 70%). ^1H NMR (400 MHz, CDCl_3) δ 1.65 (m, 4H), 2.68 (t, $J = 7.1$ Hz, 2H), 3.46 (q, $J = 6.5$ Hz, 2H), 7.16 – 7.21 (m, 3H), 7.26 – 7.31 (m, 2H), 7.34 (d, $J = 8.5$ Hz, 1H), 7.53 – 7.57 (m, 1H), 7.96 (t, $J = 5.7$ Hz, 1H), 8.40

(d, $J = 1.9$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.1, 35.5, 40.4, 110.2, 113.5 (q, $J = 4.0$ Hz), 122.2 (q, $J = 3.9$ Hz), 123.8 (q, $J = 272.0$ Hz), 126.0, 127.8 (q, $J = 33.3$ Hz), 128.5, 128.5 (4C), 142.0, 143.8, 149.4, 152.8. MS (ESI) m/z : 379 $[\text{M} + \text{H}]^+$, 396 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-*N*-(4-phenylbutyl)-6-(trifluoromethyl)-1,3-benzoxazole-3-carboxamide (16). The reaction was carried out following method A, using **37m** (81 mg, 0.399 mmol) and 4-phenylbutyl isocyanate (76.7 mg, 0.075 mL, 0.438 mmol). White solid (91 mg, 60%). ^1H NMR (400 MHz, CDCl_3) δ 1.64 – 1.79 (m, 4H), 2.68 (t, $J = 7.1$ Hz, 2H), 3.46 (q, $J = 6.6$ Hz, 2H), 7.15 – 7.21 (m, 3H), 7.26 – 7.31 (m, 2H), 7.51 (d, $J = 1.7$ Hz, 1H), 7.55 – 7.59 (m, 1H), 7.98 (t, $J = 5.9$ Hz, 1H), 8.20 (d, $J = 8.4$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.6, 29.1, 35.6, 40.4, 107.6 (q, $J = 4.0$ Hz), 115.9, 122.6 (q, $J = 3.9$ Hz), 123.7 (q, $J = 272.1$ Hz), 126.0, 127.2 (q, $J = 33.3$ Hz), 128.5 (4C), 130.9, 141.6, 141.9, 149.3, 152.8. MS (ESI) m/z : 202 $[\text{M} - \text{CONH}(\text{CH}_2)_4\text{Ph}]^-$.

5-Nitro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (17). The reaction was carried out following method A, using **37n** (124 mg, 0.688 mmol) and 4-phenylbutyl isocyanate (133 mg, 0.130 mL, 0.757 mmol). White solid (150 mg, 61%). ^1H NMR (400 MHz, CDCl_3) δ 1.65 – 1.79 (m, 4H), 2.68 (t, $J = 7.1$ Hz, 2H), 3.47 (td, $J = 6.7, 5.6$ Hz, 2H), 7.16 – 7.21 (m, 3H), 7.26 – 7.31 (m, 2H), 7.37 (d, $J = 8.9$ Hz, 1H), 7.88 (t, $J = 5.3$ Hz, 1H), 8.24 (dd, $J = 8.9, 2.4$ Hz, 1H), 8.98 (d, $J = 2.4$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.6, 29.1, 35.5, 40.5, 110.2, 112.0, 121.1, 126.1, 128.5 (2C), 128.5 (2C), 128.6, 141.9, 145.3, 145.6, 148.9, 152.6. MS (ESI) m/z : 354 $[\text{M} + \text{H}]^+$.

6-Nitro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (18). The reaction was carried out following method A, using **37o** (72 mg, 0.400 mmol) and 4-phenylbutyl isocyanate (76.7 mg, 0.075 mL, 0.438 mmol). White solid (86 mg, 60%). ^1H NMR (400 MHz, CDCl_3) δ

1
2
3 1.64 – 1.79 (m, 4H), 2.68 (t, $J = 7.0$ Hz, 2H), 3.47 (q, $J = 6.4$ Hz, 2H), 7.15 – 7.21 (m, 3H), 7.26
4 – 7.31 (m, 2H), 7.94 (t, $J = 5.8$ Hz, 1H), 8.15 (t, $J = 1.2$ Hz, 1H), 8.23 – 8.28 (m, 2H). ^{13}C NMR
5 (101 MHz, CDCl_3) δ 28.6, 29.0, 35.5, 40.5, 106.3, 115.6, 121.5, 126.1, 128.5 (2C), 128.6 (2C),
6 133.3, 141.3, 141.9, 144.7, 149.0, 152.7. MS (ESI) m/z : non-ionizable compound under routine
7 conditions used.

8
9
10
11
12
13
14
15 **5,6-Dichloro-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (19).** The reaction
16 was carried out following method A, using **37p** (121 mg, 0.593 mmol) and 4-phenylbutyl
17 isocyanate (114 mg, 0.112 mL, 0.652 mmol). White solid (203 mg, 90%). ^1H NMR (400 MHz,
18 CDCl_3) δ 1.63 – 1.78 (m, 4H), 2.67 (t, $J = 7.1$ Hz, 2H), 3.44 (td, $J = 6.7, 5.5$ Hz, 2H), 7.16 – 7.21
19 (m, 3H), 7.26 – 7.31 (m, 2H), 7.37 (s, 1H), 7.84 (t, $J = 5.8$ Hz, 1H), 8.22 (s, 1H). ^{13}C NMR (101
20 MHz, CDCl_3) δ 28.6, 29.1, 35.6, 40.4, 112.0, 117.3, 126.1, 127.5, 128.5 (4C), 128.6, 129.3,
21 140.4, 141.9, 149.2, 152.6. MS (ESI) m/z : non-ionizable compound under routine conditions
22 used.

23
24
25
26
27
28
29
30
31
32
33
34 **2-Oxo-4-phenyl-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (20).** The reaction was
35 carried out following method A, using crude **39a** (140 mg) and 4-phenylbutyl isocyanate (127
36 mg, 0.124 mL, 0.724 mmol). In total, four portions of 4-phenylbutyl isocyanate were added with
37 about 3 h in between. Waxy white solid (18 mg, 6% over steps from **37a**). ^1H NMR (400 MHz,
38 CDCl_3) δ 1.47 – 1.58 (m, 2H), 1.59 – 1.68 (m, 2H), 2.61 (t, $J = 7.5$ Hz, 2H), 3.20 (td, $J = 6.8, 5.7$
39 Hz, 2H), 7.09 (t, $J = 5.1$ Hz, 1H), 7.15 – 7.25 (m, 4H), 7.26 – 7.38 (m, 9H). ^{13}C NMR (101 MHz,
40 CDCl_3) δ 28.5, 29.1, 35.6, 40.8, 108.8, 124.9, 125.0, 126.0, 126.4, 127.3, 127.3 (2C), 127.6,
41 128.4 (2C), 128.5 (2C), 128.5 (3C), 139.4, 142.1, 143.6, 147.9. MS (ESI) m/z : 387 $[\text{M} + \text{H}]^+$, 404
42 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-5-phenyl-N-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (21). The reaction was carried out following method A, using **39b** (14.9 mg, 0.0705 mmol) and 4-phenylbutyl isocyanate (33 mg, 0.032 mL, 0.187 mmol). Clear colourless sticky oil (17 mg, 63%). ^1H NMR (400 MHz, CDCl_3) δ 1.65 – 1.80 (m, 4H), 2.68 (t, J = 7.1 Hz, 2H), 3.46 (q, J = 6.4 Hz, 2H), 7.15 – 7.21 (m, 3H), 7.25 – 7.31 (m, 3H), 7.33 – 7.38 (m, 1H), 7.41 – 7.47 (m, 3H), 7.57 – 7.61 (m, 2H), 8.08 (t, J = 5.3 Hz, 1H), 8.34 (d, J = 1.9 Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.2, 35.6, 40.3, 110.1, 114.2, 123.6, 126.0, 127.5 (2C), 127.7, 128.5 (2C), 128.5 (2C), 128.6, 129.0 (2C), 139.0, 140.4, 141.3, 142.0, 150.0, 153.5. MS (ESI) m/z : 210 $[\text{M} - \text{CONH}(\text{CH}_2)_4\text{Ph}]^-$.

2-Oxo-6-phenyl-N-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (22). The reaction was carried out following method A, using **39c** (29 mg, 0.138 mmol) and 4-phenylbutyl isocyanate (29 mg, 0.028 mL, 0.165 mmol). White solid (33 mg, 62%). ^1H NMR (400 MHz, CDCl_3) δ 1.65 – 1.79 (m, 4H), 2.68 (t, J = 7.2 Hz, 2H), 3.47 (td, J = 6.7, 5.6 Hz, 2H), 7.16 – 7.21 (m, 3H), 7.25 – 7.31 (m, 2H), 7.34 – 7.40 (m, 1H), 7.43 – 7.50 (m, 4H), 7.55 – 7.59 (m, 2H), 8.06 (t, J = 5.3 Hz, 1H), 8.11 (d, J = 8.4 Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.2, 35.6, 40.3, 108.7, 115.8, 124.1, 126.0, 127.3 (2C), 127.3, 127.9, 128.5 (2C), 128.6 (2C), 129.1 (2C), 138.6, 140.05, 142.0, 142.4, 149.9, 153.4. MS (ESI) m/z : 387 $[\text{M} + \text{H}]^+$, 404 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-7-phenyl-N-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (23). The reaction was carried out following method A, using crude **39d** (399 mg) and 4-phenylbutyl isocyanate (145 mg, 0.141 mL, 0.825 mmol). The reaction was stirred for 68 h. Yellow solid (52 mg, 18% over 2 steps from **37c**). ^1H NMR (400 MHz, CDCl_3) δ 1.65 – 1.80 (m, 4H), 2.68 (t, J = 7.1 Hz, 2H), 3.47 (q, J = 6.6 Hz, 2H), 7.16 – 7.21 (m, 3H), 7.25 – 7.32 (m, 2H), 7.33 – 7.36 (m, 1H), 7.38 – 7.44 (m, 2H), 7.46 – 7.52 (m, 2H), 7.69 – 7.73 (m, 2H), 8.07 (dd, J = 7.9, 1.3 Hz, 1H), 8.11 (t, J = 5.6 Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.2, 35.6, 40.3, 114.6, 124.4, 124.5, 125.4,

126.0, 128.5 (4C), 128.5 (3C), 128.7, 129.0 (2C), 134.4, 139.0, 142.0, 150.0, 153.3. MS (ESI) m/z : 387 $[M + H]^+$, 404 $[M + NH_4]^+$.

5-(4-Fluorophenyl)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (24). The reaction was carried out following method A, using **39e** (26 mg, 0.113 mmol) and 4-phenylbutyl isocyanate (33 mg, 0.032 mL, 0.187 mmol). White solid (26 mg, 58%). 1H NMR (400 MHz, $CDCl_3$) δ 1.64 – 1.78 (m, 4H), 2.68 (t, J = 7.1 Hz, 2H), 3.46 (q, J = 6.5 Hz, 2H), 7.09 – 7.16 (m, 2H), 7.17 – 7.21 (m, 3H), 7.26 – 7.31 (m, 3H), 7.39 (dd, J = 8.4, 1.9 Hz, 1H), 7.51 – 7.57 (m, 2H), 8.07 (t, J = 5.5 Hz, 1H), 8.28 (d, J = 1.9 Hz, 1H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 28.7, 29.2, 35.6, 40.3, 110.2, 114.4, 115.9 (d, J = 21.5 Hz, 2C), 123.5, 126.0, 128.5 (2C), 128.5 (2C), 128.7, 129.1 (d, J = 7.7 Hz, 2C), 136.5, 136.6, 141.3, 142.0, 150.0, 153.4, 162.8 (d, J = 246.9 Hz). MS (ESI) m/z : 228 $[M - CONH(CH_2)_4Ph]^-$.

5-(4-Methoxyphenyl)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (25). The reaction was carried out following method A, using **39f** (41 mg, 0.170 mmol) and 4-phenylbutyl isocyanate (33 mg, 0.032 mL, 0.187 mmol). White solid (33 mg, 47%). 1H NMR (400 MHz, $CDCl_3$) δ 1.65 – 1.78 (m, 4H), 2.68 (t, J = 7.1 Hz, 2H), 3.46 (q, J = 6.6 Hz, 2H), 3.86 (s, 3H), 6.94 – 6.99 (m, 2H), 7.15 – 7.21 (m, 3H), 7.24 – 7.30 (m, 3H), 7.40 (dd, J = 8.4, 1.9 Hz, 1H), 7.50 – 7.54 (m, 2H), 8.09 (t, J = 5.3 Hz, 1H), 8.29 (d, J = 1.8 Hz, 1H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 28.7, 29.2, 35.6, 40.3, 55.5, 110.0, 114.1, 114.4 (2C), 123.1, 126.0, 128.5 (2C), 128.5 (2C), 128.6 (2C), 128.6, 132.9, 138.6, 140.9, 142.0, 150.1, 153.5, 159.5. MS (ESI) m/z : 417 $[M + H]^+$, 416 $[M + NH_4]^+$.

6-(4-Methoxyphenyl)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (26). The reaction was carried out following method A, using **39g** (36 mg, 0.149 mmol) and 4-phenylbutyl isocyanate (29 mg, 0.028 mL, 0.165 mmol). White solid (52 mg, 84%). 1H NMR (400 MHz,

CDCl₃) δ 1.65 – 1.78 (m, 4H), 2.68 (t, J = 7.1 Hz, 2H), 3.46 (td, J = 6.8, 5.6 Hz, 2H), 3.86 (s, 3H), 6.96 – 7.01 (m, 2H), 7.15 – 7.21 (m, 3H), 7.26 – 7.31 (m, 2H), 7.41 (d, J = 1.5 Hz, 1H), 7.44 (dd, J = 8.4, 1.7 Hz, 1H), 7.48 – 7.52 (m, 2H), 8.03 – 8.09 (m, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 28.7, 29.2, 35.6, 40.3, 55.5, 108.2, 114.6 (2C), 115.7, 123.6, 126.0, 126.8, 128.3 (2C), 128.5 (2C), 128.6 (2C), 132.6, 138.2, 142.0, 142.5, 149.9, 153.4, 159.7. MS (ESI) m/z : non-ionizable compound under routine conditions used.

2-Oxo-*N*-(4-phenylbutyl)-6-[(*E*)-styryl]-1,3-benzoxazole-3-carboxamide (27). The reaction was carried out following method A, using **40a** (80 mg, 0.337 mmol) and 4-phenylbutyl isocyanate (65 mg, 0.064 mL, 0.371 mmol). White solid (72 mg, 52%). ¹H NMR (400 MHz, CDCl₃) δ 1.64 – 1.79 (m, 4H), 2.68 (t, J = 7.1 Hz, 2H), 3.45 (q, J = 6.6 Hz, 2H), 7.09 (br s, 2H), 7.16 – 7.21 (m, 3H), 7.24 – 7.31 (m, 3H), 7.35 – 7.42 (m, 4H), 7.49 – 7.53 (m, 2H), 8.02 (d, J = 8.2 Hz, 1H), 8.02 – 8.06 (m, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 28.7, 29.2, 35.6, 40.3, 107.2, 115.7, 124.0, 126.0, 126.7 (2C), 127.3, 127.5, 128.1, 128.5 (2C), 128.6 (2C), 128.9 (2C), 129.6, 134.8, 136.9, 142.0, 142.5, 149.8, 153.4. MS (ESI) m/z : non-ionizable compound under routine conditions used.

6-[(*E*)-2-Cyclohexylvinyl]-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (28). The reaction was carried out following method A, using **40b** (58 mg, 0.238 mmol) and 4-phenylbutyl isocyanate (46 mg, 0.045 mL, 0.262 mmol). White solid (81 mg, 82%). ¹H NMR (400 MHz, CDCl₃) δ 1.13 – 1.39 (m, 5H), 1.63 – 1.86 (m, 9H), 2.08 – 2.18 (m, 1H), 2.67 (t, J = 7.1 Hz, 2H), 3.44 (q, J = 6.4 Hz, 2H), 6.16 (dd, J = 15.9, 6.8 Hz, 1H), 6.33 (dd, J = 15.9, 1.2 Hz, 1H), 7.15 – 7.27 (m, 5H), 7.26 – 7.30 (m, 2H), 7.93 (d, J = 8.3 Hz, 1H), 8.04 (t, J = 5.6 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 26.1 (2C), 26.3, 28.7, 29.2, 33.0 (2C), 35.6, 40.26, 41.3, 106.8,

115.4, 123.3, 126.0, 126.3, 126.6, 128.5 (2C), 128.5 (2C), 135.6, 137.9, 142.0, 142.3, 149.9, 153.5. MS (ESI) m/z : 419 $[M + H]^+$, 436 $[M + NH_4]^+$.

2-Oxo-6-phenethyl-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (29). The reaction was carried out following method A, using **41a** (48 mg, 0.201 mmol) and 4-phenylbutyl isocyanate (46 mg, 0.045 mL, 0.262 mmol). White solid (65 mg, 78%). 1H NMR (400 MHz, $CDCl_3$) δ 1.65 – 1.80 (m, 4H), 2.70 (t, J = 7.2 Hz, 2H), 2.91 – 3.03 (m, 4H), 3.46 (q, J = 6.4 Hz, 2H), 7.03 (d, J = 1.6 Hz, 1H), 7.07 (dd, J = 8.2, 1.6 Hz, 1H), 7.14 – 7.17 (m, 2H), 7.18 – 7.24 (m, 4H), 7.27 – 7.33 (m, 4H), 7.95 (d, J = 8.2 Hz, 1H), 8.07 (t, J = 5.7 Hz, 1H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 28.7, 29.2, 35.6, 37.9, 38.0, 40.2, 110.0, 115.4, 125.2, 126.0, 126.1, 126.3, 128.5 (2C), 128.6 (4C), 128.6 (2C), 138.1, 141.1, 141.9, 142.1, 150.0, 153.5. MS (ESI) m/z : non-ionizable compound under routine conditions used.

6-(2-Cyclohexylethyl)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (30). The reaction was carried out following method A, using **41b** (42 mg, 0.173 mmol) and 4-phenylbutyl isocyanate (40 mg, 0.033 mL, 0.194 mmol). White solid (55 mg, 75%). 1H NMR (400 MHz, $CDCl_3$) δ 0.87 – 0.98 (m, 2H), 1.10 – 1.29 (m, 4H), 1.45 – 1.56 (m, 2H), 1.62 – 1.78 (m, 9H), 2.62 – 2.69 (m, 4H), 3.44 (q, J = 6.6 Hz, 2H), 7.05 (s, 1H), 7.07 (d, J = 1.2 Hz, 1H), 7.15 – 7.20 (m, 3H), 7.26 – 7.31 (m, 2H), 7.92 (d, J = 8.2 Hz, 1H), 8.05 (t, J = 5.8 Hz, 1H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 26.4 (2C), 26.8, 28.7, 29.2, 33.3, 33.4 (2C), 35.6, 37.3, 39.5, 40.2, 109.8, 115.3, 125.0, 125.8, 126.0, 128.5 (2C), 128.5 (2C), 140.3, 142.0, 142.1, 150.1, 153.5. MS (ESI) m/z : non-ionizable compound under routine conditions used.

6-Butoxy-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (31). The reaction was carried out following method A, using **47a** (49 mg, 0.238 mmol) and 4-phenylbutyl isocyanate (46 mg, 0.045 mL, 0.262 mmol). White solid (67 mg, 74%). 1H NMR (400 MHz, $CDCl_3$) δ 0.98

(t, $J = 7.4$ Hz, 3H), 1.44 – 1.55 (m, 2H), 1.63 – 1.81 (m, 6H), 2.67 (t, $J = 7.2$ Hz, 2H), 3.44 (q, $J = 6.4$ Hz, 2H), 3.95 (t, $J = 6.5$ Hz, 2H), 6.78 (d, $J = 2.3$ Hz, 1H), 6.79 – 6.82 (m, 1H), 7.14 – 7.21 (m, 3H), 7.26 – 7.31 (m, 2H), 7.91 (d, $J = 8.7$ Hz, 1H), 8.00 (t, $J = 5.2$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 13.14.0, 19.3, 28.7, 29.2, 31.3, 35.6, 40.2, 68.7, 97.6, 111.1, 116.0, 121.4, 126.0, 128.5 (2C), 128.5 (2C), 142.0, 142.6, 150.0, 153.5, 156.9. MS (ESI) m/z : 383 $[\text{M} + \text{H}]^+$, 400 $[\text{M} + \text{NH}_4]^+$.

6-(2-Cyclohexylethoxy)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (32).

The reaction was carried out following method A, using **47b** (50 mg, 0.190 mmol) and 4-phenylbutyl isocyanate (37 mg, 0.036 mL, 0.209 mmol). White solid (45 mg, 55%). ^1H NMR (400 MHz, CDCl_3) δ 0.91 – 1.04 (m, 2H), 1.10 – 1.33 (m, 3H), 1.43 – 1.54 (m, 1H), 1.62 – 1.80 (m, 11H), 2.67 (t, $J = 7.2$ Hz, 2H), 3.43 (q, $J = 6.4$ Hz, 2H), 3.98 (t, $J = 6.7$ Hz, 2H), 6.77 (d, $J = 2.3$ Hz, 1H), 6.79 – 6.82 (m, 1H), 7.15 – 7.20 (m, 3H), 7.26 – 7.30 (m, 2H), 7.91 (d, $J = 8.7$ Hz, 1H), 8.00 (t, $J = 5.7$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 26.4 (2C), 26.7, 28.7, 29.2, 33.4 (2C), 34.7, 35.6, 36.6, 40.2, 67.0, 97.6, 111.1, 116.0, 121.4, 126.0, 128.5 (2C), 128.5 (2C), 142.1, 142.6, 150.01, 153.5, 156.9. MS (ESI) m/z : 260 $[\text{M} - \text{CONH}(\text{CH}_2)_4\text{Ph}]^-$.

2-Oxo-6-phenethyloxy-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (33). The

reaction was carried out following method A, using **47c** (65 mg, 0.254 mmol) and 4-phenylbutyl isocyanate (49 mg, 0.048 mL, 0.279 mmol). White solid (71 mg, 65%). ^1H NMR (400 MHz, CDCl_3) δ 1.61 – 1.77 (m, 4H), 2.67 (t, $J = 7.2$ Hz, 2H), 3.10 (t, $J = 7.0$ Hz, 2H), 3.43 (q, $J = 6.4$ Hz, 2H), 4.17 (t, $J = 7.0$ Hz, 2H), 6.78 (d, $J = 2.2$ Hz, 1H), 6.79 – 6.82 (m, 1H), 7.15 – 7.21 (m, 4H), 7.22 – 7.36 (m, 6H), 7.91 (d, $J = 8.5$ Hz, 1H), 7.99 (t, $J = 5.6$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.2, 35.6, 35.8, 40.2, 69.7, 97.8, 111.2, 116.0, 121.7, 126.0, 126.8, 128.5 (2C),

128.5 (2C), 128.7 (2C), 129.1 (2C), 130.4, 138.0, 142.1, 142.6, 150.0, 156.6. MS (ESI) m/z : 431 [M + H]⁺.

5-Benzoyl-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (34). The reaction was carried out following method A, using **49** (100 mg, 0.418 mmol) and 4-phenylbutyl isocyanate (81 mg, 0.079 mL, 0.460 mmol). White solid (127 mg, 73%). ¹H NMR (400 MHz, CDCl₃) δ 1.62 – 1.79 (m, 4H), 2.67 (t, J = 7.2 Hz, 2H), 3.44 (td, J = 6.7 Hz, 2H), 7.14 – 7.23 (m, 3H), 7.24 – 7.32 (m, 2H), 7.35 (d, J = 8.4 Hz, 1H), 7.50 (d, J = 7.8 Hz, 2H), 7.56 – 7.65 (m, 1H), 7.79 (dd, J = 1.5, 8.3 Hz, 3H), 7.98 (t, J = 5.4 Hz, 1H), 8.51 (d, J = 1.7 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 28.7, 29.4, 35.6, 40.4, 110.0, 117.7, 126.0, 127.4, 128.0, 128.5 (4C), 128.6 (2C), 130.1 (2C), 132.8, 135.1, 137.4, 142.0, 144.5, 149.5, 153.1, 195.2. MS (ESI) m/z : 415 [M + H]⁺, 432 [M + NH₄]⁺.

6-Benzoyl-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (35). The reaction was carried out following method A, using **52a** (100 mg, 0.418 mmol) and 4-phenylbutyl isocyanate (81 mg, 0.079 mL, 0.461 mmol). White solid (142 mg, 82%). ¹H NMR (400 MHz, CDCl₃) δ 1.61 – 1.79 (m, 4H), 2.68 (t, J = 6.6 Hz, 2H), 3.47 (q, J = 6.3 Hz, 2H), 7.15 – 7.21 (m, 3H), 7.26 – 7.32 (m, 2H), 7.46 – 7.54 (m, 2H), 7.58 – 7.65 (m, 1H), 7.69 – 7.80 (m, 4H), 8.03 (t, J = 5.3 Hz, 1H), 8.16 (d, J = 8.6 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 28.7, 29.1, 35.6, 40.4, 111.6, 115.2, 126.1, 128.1, 128.5 (4C), 128.6 (2C), 130.0 (2C), 131.6, 132.8, 134.4, 137.4, 141.8, 142.0, 149.5, 153.2, 194.9. MS (ESI) m/z : 238 [M - CONH(CH₂)₄Ph]⁻.

6-(4-Chlorobenzoyl)-2-oxo-*N*-(4-phenylbutyl)-1,3-benzoxazole-3-carboxamide (36). The reaction was carried out following method A, using **52b** (169 mg, 0.616 mmol) and 4-phenylbutyl isocyanate (119 mg, 0.116 mL, 0.678 mmol). White solid (66 mg, 26%). ¹H NMR (400 MHz, CDCl₃) δ 1.65 – 1.79 (m, 4H), 2.68 (t, J = 7.0 Hz, 2H), 3.47 (q, J = 6.3 Hz, 2H), 7.16

– 7.22 (m, 3H), 7.26 – 7.31 (m, 2H), 7.47 – 7.51 (m, 2H), 7.69 – 7.75 (m, 4H), 8.02 (t, $J = 5.4$ Hz, 1H), 8.17 (d, $J = 8.2$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.7, 29.1, 35.6, 40.4, 111.5, 115.2, 126.1, 128.0, 128.5 (4C), 129.0 (2C), 131.4 (2C), 131.7, 133.9, 135.6, 139.4, 141.8, 141.9, 149.4, 153.1 193.7. MS (ESI) m/z : 272 and 274 $[\text{M} - \text{CONH}(\text{CH}_2)_4\text{Ph}]^-$.

General Procedure for the Synthesis of Benzoxazolone *N*-Carboxamides **54, **62-63**, and **66** via Method B (Scheme 6, Table 3).** Triphosgene (1.0 equiv) was placed in an oven-dried flask, cooled to 0 °C, and dissolved in dry pyridine (5 mL per mmol triphosgene). 3*H*-1,3-benzoxazol-2-one (**50**, 1.0 equiv) dissolved in dry pyridine (1.9 mL per mmol **50**) was added, and the reaction mixture was stirred under nitrogen for 30 min at rt. The solution was cooled to 0 °C and the proper amine (1.5 equiv), dissolved in dry pyridine (3.5 mL per mmol amine), was added. The reaction was allowed to warm to rt and stirred for 3 h. The solvent was removed under reduced pressure, and the compound was purified by column chromatography using the Teledyne ISCO apparatus (cyclohexane/EtOAc).

2-Oxo-*N*-(3-phenylpropyl)-1,3-benzoxazole-3-carboxamide (54**).** The reaction was carried out following method B, using **50** (70 mg, 0.520 mmol) and 3-phenylpropan-1-amine **67l** (106 mg, 0.111 mL, 0.784 mmol). White solid (65 mg, 42%). ^1H NMR (400 MHz, CDCl_3) δ 1.99 (p, $J = 7.4$ Hz, 2H), 2.73 (t, $J = 7.6$ Hz, 2H), 3.46 (q, $J = 6.6$ Hz, 2H), 7.16 – 7.32 (m, 8H), 8.07 (d, $J = 7.5$ Hz, 1H), 8.07 – 8.14 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 31.0, 33.2, 39.9, 110.0, 115.7, 124.7, 125.1, 126.2, 128.1, 128.5 (2C), 128.6 (2C), 141.1, 141.9, 150.0, 153.3. MS (ESI) m/z : 297 $[\text{M} + \text{H}]^+$, 314 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-*N*-[2-(benzyloxy)ethyl]-1,3-benzoxazole-3-carboxamide (62**).** The reaction was carried out following method B, using **50** (70 mg, 0.520 mmol) and 2-(benzyloxy)ethan-1-amine hydrochloride **67m** (584 mg, 3.11 mmol, 6.0 equiv). The reaction was stirred for 22 h. White

solid (89 mg, 55%). ^1H NMR (400 MHz, CDCl_3) δ 3.62 – 3.69 (m, 4H), 4.58 (s, 2H), 7.22 – 7.29 (m, 4H), 7.31 – 7.39 (m, 4H), 8.03 – 8.08 (m, 1H), 8.36 (br s, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 40.4, 68.3, 73.4, 110.0, 115.7, 124.6, 125.1, 127.9 (3C), 128.1, 128.6 (2C), 137.9, 141.9, 150.1, 153.1. MS (ESI) m/z : 313 $[\text{M} + \text{H}]^+$, 330 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-*N*-[2-(benzylthio)ethyl]-1,3-benzoxazole-3-carboxamide (63). The reaction was carried out following method B, using **50** (70 mg, 0.520 mmol) and 2-(benzylthio)ethan-1-amine hydrochloride **67n** (159 mg, 0.780 mmol). White powder (58 mg, 34%). ^1H NMR (400 MHz, CDCl_3) δ 2.69 (t, $J = 6.6$ Hz, 2H), 3.55 (q, $J = 6.5$ Hz, 2H), 3.77 (s, 2H), 7.17–7.37 (m, 8H), 8.03–8.07 (m, 1H), 8.29 (t, $J = 6.2$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 31.0, 36.3, 39.4, 110.1, 115.7, 124.7, 125.1, 127.3, 128.1, 128.7 (2C), 129.0 (2C), 138.1, 141.9, 149.9, 153.1. MS (ESI) m/z : 329 $[\text{M} + \text{H}]^+$, 346 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-*N*-[(4-propylphenyl)methyl]-1,3-benzoxazole-3-carboxamide (66). The reaction was carried out following method B, using **50** (60 mg, 0.444 mmol) and amine **67e** (125 mg, 0.673 mmol). White powder (40 mg, 29%). ^1H NMR (400 MHz, CDCl_3) δ 0.94 (t, $J = 7.3$ Hz, 3H), 1.58 – 1.71 (m, 2H), 2.58 (dd, $J = 8.4, 6.8$ Hz, 2H), 4.58 (d, $J = 5.7$ Hz, 2H), 7.17 (d, $J = 8.1$ Hz, 2H), 7.21 – 7.32 (m, 5H), 8.07 – 8.14 (m, 1H), 8.35 – 8.46 (m, 1H). ^{13}C NMR (101 MHz, DMSO) δ 13.9, 24.7, 37.8, 44.1, 110.0, 115.8, 124.7, 125.1, 128.1, 129.0, 134.6, 141.9, 142.5, 150.0, 153.2. MS (ESI) m/z : 311 $[\text{M} + \text{H}]^+$, 328 $[\text{M} + \text{NH}_4]^+$.

General Procedure for the Synthesis of Benzoxazolone *N*-Carboxamides 53, 55–61 and 64–65 via Method C (Scheme 6, Table 3). In an oven-dried flask, triphosgene (1.0 equiv) was dissolved in dry DCM (5 mL per mmol triphosgene). A solution of 3*H*-1,3-benzoxazol-2-one (**50**, 1.0 equiv) and Et_3N (4.0 equiv) in dry DCM (5.6 mL per mmol **50**) was added at 0 °C, and the reaction was stirred for 1 h at rt under nitrogen. Then a solution of the appropriate amine (1.5

equiv) and Et₃N (1.5 equiv) in dry DCM (6.5 mL per mmol amine) was added at 0 °C, and the reaction was stirred at rt for 3 h. The reaction mixture was diluted with DCM (20 mL per mmol **50**) and quenched with sat aq NH₄Cl (30 mL per mmol **50**). The two phases were separated and the aq layer was extracted with DCM (3 × 20 mL per mmol **50**). The combined organic phases were dried over Na₂SO₄, evaporated on celite or silica, and the compound was purified by silica gel column chromatography using the Teledyne ISCO apparatus (cyclohexane/EtOAc).

2-Oxo-N-(phenethyl)-1,3-benzoxazole-3-carboxamide (53). The reaction was carried out following method C, using **50** (60 mg, 0.444 mmol) and 2-phenylethan-1-amine **67k** (80 mg, 0.084 mL, 0.660 mmol). White solid (79 mg, 64%). ¹H NMR (400 MHz, CDCl₃) δ 2.96 (t, *J* = 7.2 Hz, 2H), 3.69 (td, *J* = 5.8, 7.2 Hz, 2H), 7.19 – 7.29 (m, 6H), 7.34 (dd, *J* = 5.8, 8.9 Hz, 2H), 8.05 – 8.09 (m, 1H), 8.09 – 8.17 (m, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 35.9, 41.8, 110.0, 115.7, 124.7, 125.1, 126.9, 128.1, 128.9, 128.9, 138.4, 141.9, 149.9, 153.2. MS (ESI) *m/z*: 283 [M + H]⁺, 300 [M + NH₄]⁺. MS (ESI) *m/z*: 281[M - H]⁻.

2-Oxo-N-(5-phenylpentyl)-1,3-benzoxazole-3-carboxamide (55). The reaction was carried out following method C, using **50** (68 mg, 0.503 mmol) and amine **67j** (100 mg, 0.501 mmol). White solid (83 mg, 51%). ¹H NMR (400 MHz, CDCl₃) δ 1.42 – 1.51 (m, 2H), 1.65 – 1.76 (m, 4H), 2.66 (t, *J* = 7.6 Hz, 2H), 3.45 (td, *J* = 7.1, 5.9 Hz, 2H), 7.16 – 7.22 (m, 3H), 7.24 – 7.32 (m, 5H), 8.06 – 8.13 (m, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 26.4, 29.3, 31.0, 35.7, 40.2, 109.9, 115.6, 124.5, 125.0, 125.7, 128.0, 128.3, 128.4, 141.7, 142.3, 149.8, 153.2. MS (ESI) *m/z*: 325 [M + H]⁺.

2-Oxo-N-(6-phenylhexyl)-1,3-benzoxazole-3-carboxamide (56). The reaction was carried out following method C, using **50** (40 mg, 0.300 mmol) and amine **67f** (95 mg, 0.440 mmol). White solid (32 mg, 31%). ¹H NMR (400 MHz, CDCl₃) δ 1.32 – 1.52 (m, 4H), 1.59 – 1.71 (m,

4H), 2.51 – 2.74 (m, 2H), 3.31 – 3.51 (m, 2H), 7.13 – 7.20 (m, 3H), 7.21 – 7.31 (m, 5H), 8.02 – 8.11 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 26.8, 29.0, 29.5, 31.4, 36.0, 40.4, 110.0, 115.7, 124.6, 125.1, 125.8, 128.2, 128.4, 128.5, 141.9, 142.7, 149.9, 153.3. MS (ESI) m/z : 339 $[\text{M} + \text{H}]^+$, 356 $[\text{M} + \text{NH}_4]^+$.

***N*-[4-(2-Naphthyl)butyl]-2-oxo-1,3-benzoxazole-3-carboxamide (57).** The reaction was carried out following method C, using **50** (26 mg, 0.190 mmol) and amine **67a** (68 mg, 0.300 mmol). White solid (36 mg, 52 %). ^1H NMR (400 MHz, CDCl_3) δ 1.66 – 1.79 (m, 2H), 1.78 – 1.91 (m, 2H), 2.84 (t, $J = 7.4$ Hz, 2H), 3.47 (td, $J = 7.0, 5.7$ Hz, 2H), 7.21 – 7.30 (m, 3H), 7.34 (dd, $J = 8.4, 1.7$ Hz, 1H), 7.38 – 7.49 (m, 2H), 7.60 – 7.66 (m, 1H), 7.74 – 7.84 (m, 3H), 8.01 – 8.12 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.5, 29.2, 35.7, 40.2, 110.1, 115.7, 124.6, 125.1, 125.3, 126.0, 126.6, 127.3, 127.5, 127.7, 128.1, 128.1, 132.2, 133.7, 139.5, 141.9, 150.0, 153.3. MS (ESI) m/z : non-ionizable compound under routine conditions used.

***N*-[4-(4-Nitrophenyl)butyl]-2-oxo-1,3-benzoxazole-3-carboxamide (58).** The reaction was carried out following method C, using **50** (70 mg, 0.518 mmol) and amine **67g** (180 mg, 0.780 mmol). White solid (29 mg, 16%). ^1H NMR (400 MHz, CDCl_3) δ 1.64 – 1.82 (m, 4H), 2.78 (t, $J = 7.4$ Hz, 2H), 3.47 (q, $J = 6.7$ Hz, 2H), 7.21 – 7.32 (m, 3H), 7.32 – 7.36 (m, 2H), 8.03 – 8.07 (m, 1H), 8.10 (t, $J = 5.0$ Hz, 1H), 8.15 – 8.17 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.2, 29.2, 35.5, 40.0, 110.1, 115.7, 123.9 (2C), 124.8, 125.2, 128.1, 129.3 (2C), 141.9, 146.6, 149.8, 150.0, 153.3. MS (ESI) m/z : non-ionizable compound under routine conditions used.

***N*-[4-(4-Fluorophenyl)butyl]-2-oxo-1,3-benzoxazole-3-carboxamide (59).** The reaction was carried out following method C, using **50** (70 mg, 0.518 mmol) and amine **67b** (160 mg, 0.786 mmol). White solid (80 mg, 47%). ^1H NMR (400 MHz, CDCl_3) δ 1.64 – 1.80 (m, 4H), 2.67 (t, $J = 7.0$ Hz, 2H), 3.39 – 3.57 (m, 2H), 6.92 – 7.05 (m, 2H), 7.11 – 7.21 (m, 2H), 7.23 – 7.27 (m,

2H), 7.27 – 7.32 (m, 1H), 7.83 – 8.38 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.8, 29.1, 34.8, 40.2, 110.0, 115.20 (d, $J = 21.4$ Hz, 2C), 115.7, 124.7, 125.1, 128.1, 129.8 (d, $J = 7.9$ Hz, 2C), 137.6, 141.9, 150.0, 153.3, 161.4 (d, $J = 242.8$ Hz, 1C). MS (ESI) m/z : 329 $[\text{M} + \text{H}]^+$, 346 $[\text{M} + \text{NH}_4]^+$.

***N*-[4-(4-Methoxyphenyl)butyl]-2-oxo-1,3-benzoxazole-3-carboxamide (60).** The reaction was carried out following method C, using **50** (70 mg, 0.52 mmol) and amine **67h** (168 mg, 0.78 mmol). White solid (42 mg, 24%). ^1H NMR (400 MHz, CDCl_3) δ 1.61 – 1.75 (m, 4H), 2.61 (t, $J = 6.9$ Hz, 2H), 3.44 (q, $J = 6.5$ Hz, 2H), 3.78 (s, 3H), 6.79 – 6.85 (m, 2H), 7.07 – 7.12 (m, 2H), 7.21 – 7.33 (m, 3H), 8.03 – 8.09 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 28.9, 29.1, 34.7, 40.3, 55.4, 110.0, 113.9 (2C), 115.7, 124.6, 125.1, 128.2, 129.4 (2C), 134.1, 141.9, 150.0, 153.3, 158.0. MS (ESI) m/z : 341 $[\text{M} + \text{H}]^+$, 358 $[\text{M} + \text{NH}_4]^+$.

2-Oxo-*N*-[4-(*p*-tolyl)butyl]-1,3-benzoxazole-3-carboxamide (61). The reaction was carried out following method C, using **50** (50 mg, 0.370 mmol) and amine **67c** (110 mg, 0.551 mmol). White solid (47 mg, 39%). ^1H NMR (400 MHz, CDCl_3) δ 1.63 – 1.83 (m, 4H), 2.31 (s, 3H), 2.63 (t, $J = 7.1$ Hz, 2H), 3.44 (td, $J = 6.7, 5.5$ Hz, 2H), 7.04 – 7.15 (m, 4H), 7.22 – 7.29 (m, 3H), 8.01 – 8.13 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 21.1, 28.8, 29.2, 35.1, 40.3, 110.0, 115.8, 124.6, 125.1, 128.2, 128.4, 129.2, 135.5, 138.9, 141.9, 150.0, 153.3. MS (ESI) m/z : non-ionizable compound under routine conditions used.

2-Oxo-*N*-[2-(benzyl(methyl)amino)ethyl]-1,3-benzoxazole-3-carboxamide (64). The reaction was carried out following method C, using **50** (70 mg, 0.518 mmol) and amine **67i** (216 mg, 0.777 mmol). Clear colorless oil (70 mg, 41%). ^1H NMR (400 MHz, CDCl_3) δ 2.28 (s, 3H), 2.64 (t, $J = 6.0$ Hz, 2H), 3.54 (q, $J = 5.7$ Hz, 2H), 3.58 (s, 2H), 7.18 – 7.27 (m, 4H), 7.27 – 7.33 (m, 2H), 7.36 – 7.40 (m, 2H), 8.03 – 8.07 (m, 1H), 8.45 (t, $J = 4.9$ Hz, 1H). ^{13}C NMR (101 MHz,

CDCl₃) δ 37.9, 41.9, 55.3, 62.6, 110.0, 115.7, 124.5, 125.0, 127.3, 128.2, 128.5 (3C), 129.1 (2C), 141.9, 150.0, 153.0. MS (ESI) m/z : 326 [M + H]⁺.

2-Oxo-N-[4-(2-thienyl)butyl]-1,3-benzoxazole-3-carboxamide (65). The reaction was carried out following method C, using **50** (70 mg, 0.518 mmol) and amine **67d** (150 mg, 0.782 mmol). White solid (83 mg, 51%). ¹H NMR (400 MHz, CDCl₃) δ 1.67 – 1.84 (m, 4H), 2.89 (t, J = 7.2 Hz, 2H), 3.46 (q, J = 6.7 Hz, 2H), 6.80 (d, J = 3.3 Hz, 1H), 6.91 (dd, J = 5.1, 3.4 Hz, 1H), 7.11 (dd, J = 5.1, 1.2 Hz, 1H), 7.22 – 7.29 (m, 3H), 8.04 – 8.10 (m, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 29.0, 29.1, 29.6, 40.1, 110.0, 115.7, 123.2, 124.5, 124.7, 125.1, 126.9, 128.1, 141.9, 144.8, 150.0, 153.3. MS (ESI) m/z : 317 [M + H]⁺, 334 [M + NH₄]⁺.

Pharmacology. Preparation of enzyme-enriched lysate. Cells were suspended in 20 mM Tris HCl (pH 7.5) with 0.32 M sucrose, sonicated and centrifuged at 800g for 30 min at 4 °C. Supernatants were then centrifuged at 12.000g for 30 min at 4 °C. Pellets were re-suspended in PBS buffer (pH 7.4) and subjected to three freeze-thaw cycles at - 80 °C. The suspension was finally centrifuged at 105.000g for 1 h at 4 °C and protein concentration was measured in the supernatant with bicinchoninic acid based protein assay.

Fluorescence-Based *In Vitro* Assay. Compounds were assessed for their ability to inhibit hydrolysis of the fluorogenic substrate, *N*-[(1S,2R)-2-hydroxy-1-(hydroxymethyl)-4-(2-oxochromen-7-yl)oxy-butyl]dodecanamide, which is converted to umbelliferone by AC in the fluorescence-based *in vitro* assay, as previously described.^{12, 18} Lysosomal lysate, enriched with AC, was prepared from Hek293 cells stably expressing human AC and preincubated with test compounds and a positive control (diluted 20× from DMSO stock solutions at different concentrations) for 10 min. Then, the fluorogenic substrate (diluted 40× from EtOH stock solution, final concentration 5 μ M) was added and the mixture was incubated for 3 h at 37 °C,

1
2
3 stopped with MeOH and treated with NaIO₄ (fresh solution in 100 mM glycine/NaOH buffer pH
4 10.6) followed by 2 h incubation at 37 °C in the dark. Fluorescence intensities were measured
5
6 (excitation/emission: 355/460 nm) and plotted as a function of compound concentrations. IC₅₀
7
8 values were calculated by non-linear regression analysis using GraphPad Prism 5 (GraphPad
9
10 Software Inc., CA, USA) applying a standard slope curve fitting.
11
12
13
14

15 **PBS Stability Assay.**¹² Compounds were incubated at 10 μM in phosphate-buffered saline
16 (pH 7.4; 1% DMSO) at 37 °C under shaking. Compounds were sampled at various time points
17 and analyzed on a Xevo triple-quad UPLC system using a BEH C18 reversed phase column and
18 a linear gradient of MeCN in water. Stability was evaluated from the corresponding MRM
19 (multiple reaction monitoring) peak areas plotted versus time. The corresponding decay profile
20 was fitted with Prism to derive the corresponding half-life values.
21
22
23
24
25
26
27
28

29 **Animals and Treatments.** Male C57BL/6 mice (20–35 g, Charles River) were group-housed
30 at rt on a 12 h light/dark cycle. Water and standard chow pellets were freely available. Drugs
31 were dissolved in 15% polyethylene glycol, 15% Tween-80 and 70% saline (injection volume,
32 10mL/kg; i.p.). Animals were sacrificed 30 min, 3 h, 6 h, 12 h, 16 h and 24 h after drug
33 administration; tissues were collected, frozen in liquid nitrogen and stored at - 80 °C. All
34 procedures were performed in accordance with the Italian regulations on the protection of
35 animals used for experimental and other scientific purposes (D.M. 116192), and European
36 Economic Community regulations (O.J. of E.C. L 358/1 12/18/1986).
37
38
39
40
41
42
43
44
45
46
47

48 **AC Activity in Lungs and Cerebral Cortex.** AC activity in lung and cerebral cortex
49 lysosomal extracts was measured by LC-MS as previously described.¹⁹ Briefly, tissues (10-20
50 mg) were suspended in 20 mM Tris HCl (pH 7.5) with 0.32 M sucrose, sonicated and
51 centrifuged at 800g for 15 min at 4 °C. Supernatants were then centrifuged at 12.000g for 30 min
52
53
54
55
56
57
58
59
60

at 4 °C. Pellets were re-suspended in PBS buffer (pH 7.4) and subjected to two freeze-thaw cycles at - 80 °C. The suspension was finally centrifuged at 105.000g for 1 h at 4 °C and protein concentration was measured in the supernatant with bicinchoninic acid based protein assay. Lysosomal preparations from tissues were diluted in assay buffer (100 mM sodium phosphate, 0.1% Nonidet P-40, 150 mM NaCl, 3 mM DTT, 100 mM sodium citrate, pH 4.5). Reactions were started by the addition of 50 µM C12-ceramide (Nu-Chek Prep, Elysian, MN) and carried on for 1 h at 37 °C. Reactions were stopped by addition of a mixture of chloroform/MeOH (2:1) containing 1 nmol 11-lauroleic acid (NuChek Prep). The organic phases were collected, dried under nitrogen and analyzed by UPLC-MS (Acquity, Waters) in the negative-ion mode monitoring the reaction product (lauric acid, m/z : 199) using 11-lauroleic acid as internal standard. Lipids were eluted on an Acquity UPLC BEH C18 column (50mm length, 2.1 mm i.d., 1.7 µm pore size, Waters) column at 0.5 mL-min⁻¹ for 1.5 min with a gradient of MeCN and water, both containing 0.25% acetic acid and 5 mM ammonium acetate (70% to 100% MeCN in 0.5 min, 100% MeCN for 0.5 min, 70% MeCN for 0.4 min). The column temperature was 40 °C. Electrospray ionization (ESI) was in the negative mode, capillary voltage was 1 kV and cone voltage was 50 V. N₂ was used as drying gas at a flow rate of 500 L/h and at a temperature of 400 °C. The $[M - H]^-$ ion was monitored in the selected-ion monitoring mode (m/z values: lauric acid 199, 11-lauroleic acid 197.35). Calibration curves were generated with authentic lauric acid (Nu Check Prep).

Sphingolipid levels measurement. Sphingolipids (namely, dihydroceramide d18:0/16:0, ceramide 18:1/16:0, sphingosine and sphingosine 1-phosphate) were quantified by LC-MS analysis according to the method previously described.²⁰

ASSOCIATED CONTENT

Supporting Information

Detailed experimental procedures of intermediates (**37a**, **c-d**, **g**, **l-p**; **39a-g**; **40a-b**; **41a-b**; **42**; **44**; **45a-c**; **46a-c**; **47a-c**; **48-49**; **51a-b**; **52a-b**; **67a-j**; **70a-j**).

AUTHOR INFORMATION

Corresponding Author

*E-mail: piomelli@uci.edu. Phone: +1(949)824-9179.

Present Addresses

[‡]Department of Drug Design and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, Universitetsparken 2, DK-2100 Copenhagen, Denmark.

[†]Bachem AG, Hauptstrasse 144, 4416 Bubendorf, Switzerland.

Author Contributions

‡These authors contributed equally. The manuscript was written through contributions of all authors and all authors have given approval to the final version of the manuscript.

Funding Sources

A.B. was generously supported by a Carlsberg Foundation fellowship.

Notes

The authors declare the following competing financial interests. Two patent applications protecting the class of compounds disclosed in this paper were filed by the following authors: Piomelli, D.; Pizzirani, D.; Bach, A.; Realini, N.; Melzig, L.; Scarpelli, R.

ACKNOWLEDGMENT

The authors thank Luca Goldoni for NMR technical support, Andrea Armirotti and Giuliana Ottonello for analytical stability data, Abdul Basit for sphingolipid levels measurement method and Silvia Venzano for handling compounds.

ABBREVIATIONS

AC, acid ceramidase; AlCl_3 , aluminum chloride; DMAP, 4-dimethylaminopyridine; DMSO, dimethyl sulfoxide; h-AC, human acid ceramidase.

REFERENCES

- (1) (a) Hannun, Y. A.; Obeid, L. M. Principles of bioactive lipid signalling: lessons from sphingolipids. *Nat. Rev.Mol. Cell Biol.* **2008**, *9*, 139-150; (b) Bartke, N.; Hannun, Y. A. Bioactive sphingolipids: metabolism and function. *J. Lipid Res.* **2009**, *50 Suppl*, S91-96.
- (2) (a) Ogretmen, B.; Hannun, Y. A. Biologically active sphingolipids in cancer pathogenesis and treatment. *Nat. Rev.Cancer* **2004**, *4*, 604-616; (b) Gangoiti, P.; Camacho, L.; Arana, L.; Ouro, A.; Granado, M. H.; Brizuela, L.; Casas, J.; Fabrias, G.; Abad, J. L.; Delgado, A.; Gomez-Munoz, A. Control of metabolism and signaling of simple bioactive sphingolipids: Implications in disease. *Prog. Lipid Res.* **2010**, *49*, 316-334; (c) Dimanche-Boitrel, M. T.; Rebillard, A., Sphingolipids and response to chemotherapy. *Handb. Exp. Pharmacol.* **2013**, 73-91.
- (3) (a) Salvemini, D.; Doyle, T.; Kress, M.; Nicol, G., Therapeutic targeting of the ceramide-to-sphingosine 1-phosphate pathway in pain. *Trends Pharmacol. Sci.* **2013**, *34*, 110-118; (b) Patti, G. J.; Yanes, O.; Shriver, L. P.; Courade, J. P.; Tautenhahn, R.; Manchester, M.; Siuzdak, G. Metabolomics implicates altered sphingolipids in chronic pain of neuropathic origin. *Nat. Chem. Biol.* **2012**, *8*, 232-234.
- (4) Mao, C.; Obeid, L. M. Ceramidases: regulators of cellular responses mediated by ceramide, sphingosine, and sphingosine-1-phosphate. *Biochim. Biophys. Acta* **2008**, *1781*, 424-434.
- (5) (a) Hu, Z. Z.; Valencia, J. C.; Huang, H.; Chi, A.; Shabanowitz, J.; Hearing, V. J.; Appella, E.; Wu, C. Comparative bioinformatics analyses and profiling of lysosome-related organelle proteomes. *Int. J. Mass Spectrom.* **2007**, *259*, 147-160; (b) Bernardo, K.; Hurwitz, R.; Zenk, T.; Desnick, R. J.; Ferlinz, K.; Schuchman, E. H.; Sandhoff, K. Purification, characterization, and biosynthesis of human acid ceramidase. *J. Biol. Chem.* **1995**, *270*, 11098-11102.

- (6) (a) Spiegel, S.; Milstien, S. Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nat. Rev. Mol. Cell Biol.* **2003**, *4*, 397-407; (b) Takabe, K.; Spiegel, S. Export of sphingosine-1-phosphate and cancer progression. *J. Lipid Res.* **2014**, *55*, 1839-1846; (c) Huang, W. C.; Chen, C. L.; Lin, Y. S.; Lin, C. F. Apoptotic sphingolipid ceramide in cancer therapy. *J. Lipids* **2011**, *2011*, 565316; (d) Bielawska, A.; Linardic, C. M.; Hannun, Y. A. Ceramide-mediated biology. Determination of structural and stereospecific requirements through the use of *N*-acyl-phenylaminoalcohol analogs. *J. Biol. Chem.* **1992**, *267*, 18493-18497.
- (7) (a) Maceyka, M.; Spiegel, S. Sphingolipid metabolites in inflammatory disease. *Nature* **2014**, *510*, 58-67; (b) Pettus, B. J.; Chalfant, C. E.; Hannun, Y. A. Ceramide in apoptosis: an overview and current perspectives. *Bioch. Biophys. Acta* **2002**, *1585*, 114-125; (c) Morales, A.; Lee, H.; Goni, F. M.; Kolesnick, R.; Fernandez-Checa, J. C. Sphingolipids and cell death. *Apoptosis* **2007**, *12*, 923-939; (d) Nussbaumer, P. Medicinal chemistry aspects of drug targets in sphingolipid metabolism. *ChemMedChem* **2008**, *3*, 543-551; (e) Adan-Gokbulut, A.; Kartal-Yandim, M.; Iskender, G.; Baran, Y. Novel agents targeting bioactive sphingolipids for the treatment of cancer. *Curr. Med. Chem.* **2013**, *20*, 108-122; (f) Wymann, M. P.; Schneider, R. Lipid signalling in disease. *Nat. Rev. Mol. Cell Biol.* **2008**, *9*, 162-176; (g) Morad, S. A.; Cabot, M. C. Ceramide-orchestrated signalling in cancer cells. *Nat. Rev. Cancer* **2013**, *13*, 51-65.
- (8) (a) Bose, R.; Verheij, M.; Haimovitz-Friedman, A.; Scotto, K.; Fuks, Z.; Kolesnick, R. Ceramide synthase mediates daunorubicin-induced apoptosis: an alternative mechanism for generating death signals. *Cell* **1995**, *82*, 405-414; (b) Chauvier, D.; Morjani, H.; Manfait, M., Ceramide involvement in homocamptothecin- and camptothecin-induced cytotoxicity and apoptosis in colon HT29 cells. *Int. J. Oncol.* **2002**, *20*, 855-863; (c) Biswal, S. S.; Datta, K.; Acquah-Mensah, G. K.; Kehrer, J. P. Changes in ceramide and sphingomyelin following

fludarabine treatment of human chronic B-cell leukemia cells. *Toxicology* **2000**, *154*, 45-53; (d) Chalfant, C. E.; Rathman, K.; Pinkerman, R. L.; Wood, R. E.; Obeid, L. M.; Ogretmen, B.; Hannun, Y. A. De novo ceramide regulates the alternative splicing of caspase 9 and Bcl-x in A549 lung adenocarcinoma cells. Dependence on protein phosphatase-1. *J. Biol. Chem.* **2002**, *277*, 12587-12595.

(9) (a) Seelan, R. S.; Qian, C.; Yokomizo, A.; Bostwick, D. G.; Smith, D. I.; Liu, W. Human acid ceramidase is overexpressed but not mutated in prostate cancer. *Genes Chromosomes Cancer* **2000**, *29*, 137-146; (b) Norris, J. S.; Bielawska, A.; Day, T.; El-Zawahri, A.; ElOjeimy, S.; Hannun, Y.; Holman, D.; Hyer, M.; Landon, C.; Lowe, S.; Dong, J. Y.; McKillop, J.; Norris, K.; Obeid, L.; Rubinchik, S.; Tavassoli, M.; Tomlinson, S.; Voelkel-Johnson, C.; Liu, X. Combined therapeutic use of AdGFP FasL and small molecule inhibitors of ceramide metabolism in prostate and head and neck cancers: a status report. *Cancer Gene Ther.* **2006**, *13*, 1045-1051; (c) ElOjeimy, S.; Liu, X.; McKillop, J. C.; El-Zawahry, A. M.; Holman, D. H.; Cheng, J. Y.; Meacham, W. D.; Mahdy, A. E.; Saad, A. F.; Turner, L. S.; Cheng, J.; T, A. D.; Dong, J. Y.; Bielawska, A.; Hannun, Y. A.; Norris, J. S. Role of acid ceramidase in resistance to FasL: therapeutic approaches based on acid ceramidase inhibitors and FasL gene therapy. *Mol. Ther.* **2007**, *15*, 1259-1263; (d) Musumarra, G.; Barresi, V.; Condorelli, D. F.; Scire, S. A bioinformatic approach to the identification of candidate genes for the development of new cancer diagnostics. *Biol. Chem.* **2003**, *384*, 321-327.

(10) (a) Mahdy, A. E.; Cheng, J. C.; Li, J.; ElOjeimy, S.; Meacham, W. D.; Turner, L. S.; Bai, A.; Gault, C. R.; McPherson, A. S.; Garcia, N.; Beckham, T. H.; Saad, A.; Bielawska, A.; Bielawski, J.; Hannun, Y. A.; Keane, T. E.; Taha, M. I.; Hammouda, H. M.; Norris, J. S.; Liu, X. Acid ceramidase upregulation in prostate cancer cells confers resistance to radiation: AC inhibition, a

potential radiosensitizer. *Mol. Ther.* **2009**, *17*, 430-438; (b) Camacho, L.; Meca-Cortes, O.; Abad, J. L.; Garcia, S.; Rubio, N.; Diaz, A.; Celia-Terrassa, T.; Cingolani, F.; Bermudo, R.; Fernandez, P. L.; Blanco, J.; Delgado, A.; Casas, J.; Fabrias, G.; Thomson, T. M. Acid ceramidase as a therapeutic target in metastatic prostate cancer. *J. Lipid Res.* **2013**, *54*, 1207-1220.

(11) Saied, E. M.; Arenz, C., Inhibitors of ceramidases. *Chem. Phys. Lipids* **2015**, <http://dx.doi.org/10.1016/j.chemphyslip.2015.07.009>.

(12) Pizzirani, D.; Bach, A.; Realini, N.; Armirotti, A.; Mengatto, L.; Bauer, I.; Girotto, S.; Pagliuca, C.; De Vivo, M.; Summa, M.; Ribeiro, A.; Piomelli, D. Benzoxazolone carboxamides: potent and systemically active inhibitors of intracellular acid ceramidase. *Angew. Chem. Int. Ed. Engl.* **2015**, *54*, 485-489.

(13) Pizzirani, D.; Pagliuca, C.; Realini, N.; Branduardi, D.; Bottegoni, G.; Mor, M.; Bertozzi, F.; Scarpelli, R.; Piomelli, D.; Bandiera, T. Discovery of a new class of highly potent inhibitors of acid ceramidase: synthesis and structure-activity relationship (SAR). *J. Med. Chem.* **2013**, *56*, 3518-3530.

(14) (a) Nachman, R. J. Convenient Preparation of 2-Benzoxazolinones with 1,1-Carbonyldiimidazole. *J. Heterocycl. Chem.* **1982**, *19*, 1545-1547; (b) Moon, J. K.; Kim, J. H.; Shibamoto, T. Photodegradation pathways and mechanisms of the herbicide metamifop in a water/acetonitrile solution. *J. Agric. Food Chem.* **2010**, *58*, 12357-12365.

(15) Carato, P.; Yous, S.; Sellier, D.; Poupaert, J. H.; Lebegue, N.; Berthelot, P. Efficient and selective deprotection method for *N*-protected 2(3*H*)-benzoxazolones and 2(3*H*)-benzothiazolones. *Tetrahedron* **2004**, *60*, 10321-10324.

- (16) Pelletier, J. C.; Kincaid, S., Mitsunobu reaction modifications allowing product isolation without chromatography: application to a small parallel library. *Tetrahedron Lett.* **2000**, *41*, 797-800.
- (17) Ucar, H.; Van Derpoorten, K.; Depovere, P.; Lesieur, D.; Isa, M.; Masereel, B.; Delarge, J.; Poupaert, J. H. "Fries like" rearrangement: a novel and efficient method for the synthesis of 6-acyl-2(3*H*)-benzoxazolones and 6-acyl-2(3*H*)-benzothiazolones. *Tetrahedron* **1998**, *54*, 1763-1772.
- (18) Bedia, C.; Casas, J.; Garcia, V.; Levade, T.; Fabrias, G. Synthesis of a novel ceramide analogue and its use in a high-throughput fluorogenic assay for ceramidases. *ChemBioChem* **2007**, *8*, 642-648.
- (19) Realini, N.; Solorzano, C.; Pagliuca, C.; Pizzirani, D.; Armirotti, A.; Luciani, R.; Costi, M. P.; Bandiera, T.; Piomelli, D. Discovery of highly potent acid ceramidase inhibitors with in vitro tumor chemosensitizing activity. *Sci. Rep.* **2013**, *3*, 1035.
- (20) Basit, A.; Piomelli, D.; Armirotti, A. Rapid evaluation of 25 key sphingolipids and phosphosphingolipids in human plasma by LC-MS/MS. *Anal. Bioanal. Chem.* **2015**, *407*, 5189-5198.

FIGURES, SCHEMES AND TABLES




Figure 1. The biochemical pathway of ceramide to sphingosine 1-phosphate mediated by the enzymes acid ceramidase (AC) and sphingosine kinase (SK).




Figure 2. Benzoxazolone carboxamides **1-3** and illustration of the design strategy for structure-activity relationship (SAR) studies.






Figure 3. Effects of compound 14 (10 mg kg⁻¹, i.p.) on AC activity and sphingolipid levels in mice lungs (A, B) and cerebral cortex (C, D) after 30 min, 3 h, 6 h, 12 h, 16 h and 24 h. Activity is expressed as percentage of vehicle (basal AC activity in lungs: 98 ± 7.1 pmol/min/mg; and cerebral cortex: 74 ± 8.6 pmol/min/mg). Values are reported as the mean ± S.E.M (n=6). Repeated experiments gave similar results. ***p<0.001 vs vehicle, One-Way ANOVA followed by Dunnet's test.

Scheme 1. Synthesis of differently substituted-benzoxazolone carboxamides **4-19**





37a: R = 4-Br (88%)	4: R = 4-Br (n.o.) ^a
37b: R = 5-Br	5: R = 5-Br (48%)
37c: R = 7-Br (89%)	6: R = 7-Br (56%)
37d: R = 4-CH ₃ (85%)	7: R = 4-CH ₃ (24%)
37e: R = 5-CH ₃	8: R = 5-CH ₃ (91%)
37f: R = 6-CH ₃	9: R = 6-CH ₃ (68%)
37g: R = 7-CH ₃ (74%)	10: R = 7-CH ₃ (71%)
37h: R = 5-F	11: R = 5-F (85%)
37i: R = 6-F	12: R = 6-F (79%)
37j: R = 5-Cl	13: R = 5-Cl (92%)
37k: R = 6-Cl	14: R = 6-Cl (67%)
37l: R = 5-CF ₃ (85%)	15: R = 5-CF ₃ (70%)
37m: R = 6-CF ₃ (90%)	16: R = 6-CF ₃ (60%)
37n: R = 5-NO ₂ (92%)	17: R = 5-NO ₂ (61%)
37o: R = 6-NO ₂ (87%)	18: R = 6-NO ₂ (60%)
37p: R = 5,6-Cl (80%)	19: R = 5,6-Cl (90%)

Method A: 4-phenylbutylisocyanate, DMAP, pyridine.
^an.o.: not obtained.


Scheme 2. Synthesis of phenyl-substituted benzoxazolone carboxamides 20-26


Scheme 3. Synthesis of 6-substituted benzoxazolone carboxamides **27-30**



Scheme 4. Synthesis of 6-alkoxy benzoxazolone carboxamides 31-33


Scheme 5. Synthesis of benzoyl-benzoxazolone carboxamides **34-36**




Scheme 6. Synthesis of benzoxazolone carboxamide derivatives **53-66**

Method B: 1) triphosgene, pyridine; 2) R-NH₂ (**67e**, **67l-n**), pyridine. Method C: 1) triphosgene, Et₃N, DCM; 2) R-NH₂ (**67e-d**, **67f-k**), Et₃N, DCM.

Scheme 7. Synthesis of amines **67a-j**^a



^aYields are not reported when compounds were used as crude in the following step. Alcohols **68f-i** and amines **67k-n** were commercially available.


Table 1. Inhibitory potencies (IC_{50}) of compounds on human AC activity^a

Compounds	R	IC_{50} (nM) \pm S.E.M.
1	H	64 ± 7^b
2	6-Br	31 ± 9^b
4	4-Br	n.d.
5	5-Br	22 ± 11
6	7-Br	18 ± 11
7	4-Me	n.d.
8	5-Me	117 ± 26
9	6-Me	66 ± 10
10	7-Me	23 ± 5
11	5-F	12 ± 4
12	6-F	31 ± 5
13	5-Cl	15 ± 7
14	6-Cl	33 ± 4
15	5-CF ₃	18 ± 2
16	6-CF ₃	20 ± 9
17	5-NO ₂	3 ± 2
18	6-NO ₂	3 ± 1
19	5,6-Cl	18 ± 8

^a IC_{50} values are reported as mean values of two or more determinations.

^b IC_{50} values from ref.¹² n.d: not determined due to low chemical stability.


Table 2. Inhibitory potencies (IC₅₀) of compounds on human AC activity^a



Compounds	R	IC ₅₀ (nM) ± S.E.M.
3	6-(<i>p</i> -F-Ph)	79 ± 31 ^b
20	4-Ph	0.8 ± 0.1
21	5-Ph	42 ± 21
22	6-Ph	84 ± 35
23	7-Ph	91 ± 10
24	5-(<i>p</i> -F-Ph)	56 ± 14
25	5-(<i>p</i> -OCH ₃ -Ph)	67 ± 58
26	6-(<i>p</i> -OCH ₃ -Ph)	67 ± 5
34	5-(COPh)	24 ± 6
35	6-(COPh)	29 ± 1
36	6-CO(<i>p</i> -Cl-Ph)	14 ± 4

^aIC₅₀ values are reported as mean values of two or more determinations.


^bIC₅₀ values from ref.¹²

Table 3. Inhibitory potencies (IC_{50}) of compounds on human AC activity^a

Compounds	R	IC_{50} (nM) \pm S.E.M.
1	$(CH_2)_4Ph$	64 ± 7^b
53	$(CH_2)_2Ph$	121 ± 19
54	$(CH_2)_3Ph$	55 ± 6
55	$(CH_2)_5Ph$	54 ± 12
56	$(CH_2)_6Ph$	51 ± 16
57		71 ± 13
58		312 ± 90
59		75 ± 1
60		63 ± 22
61		25 ± 14
62		649 ± 270
63		65 ± 26
64		No inhibition ^c
65		37 ± 21
66		19 ± 2

^a IC_{50} values are reported as mean values of two or more determinations.^b IC_{50} values from ref.¹² ^cTested at 5 μ M.

Table 4. Chemical stability of compounds **1-3**, **14**, **19-26** and **35-36** in PBS (pH 7.4)^a




Compounds	R	Buffer stability <i>t</i> _{1/2} (min)
1	H	45 ^b
2	6-Br	24 ^b
3	6-(<i>p</i> -F-Ph)	>300 ^b
14	6-Cl	58
19	5,6-Cl	55
20	4-Ph	7
21	5-Ph	296
22	6-Ph	>300
23	7-Ph	>300
24	5-(<i>p</i> -F-Ph)	>300
25	5-(<i>p</i> -OCH ₃ -Ph)	176
26	6-(<i>p</i> -OCH ₃ -Ph)	132
35	6-(COPh)	98
36	6-CO(<i>p</i> -Cl-Ph)	274

^aDetermined at room temperature by LC-MS analysis.

^b*t*_{1/2} values from ref.¹²

Table of Contents graphic



FIGURES, SCHEMES AND TABLES




Figure 1. The biochemical pathway of ceramide to sphingosine 1-phosphate mediated by the enzymes acid ceramidase (AC) and sphingosine kinase (SK).




Figure 2. Benzoxazolone carboxamides **1-3** and illustration of the design strategy for structure-activity relationship (SAR) studies.






Figure 3. Effects of compound **14** (10 mg kg⁻¹, i.p.) on AC activity and sphingolipid levels in mice lungs (A, B) and cerebral cortex (C, D) after 30 min, 3 h, 6 h, 12 h, 16 h and 24 h. Activity is expressed as percentage of vehicle (basal AC activity in lungs: 98 ± 7.1 pmol/min/mg; and cerebral cortex: 74 ± 8.6 pmol/min/mg). Values are reported as the mean ± S.E.M (n=6). Repeated experiments gave similar results. ***p<0.001 vs vehicle, One-Way ANOVA followed by Dunnet's test.


Scheme 1. Synthesis of differently substituted-benzoxazolone carboxamides **4-19**

Method A: 4-phenylbutylisocyanate, DMAP, pyridine.


^an.o.: not obtained.


Scheme 2. Synthesis of phenyl-substituted benzoxazolone carboxamides **20-26**




Scheme 3. Synthesis of 6-substituted benzoxazolone carboxamides 27-30

Scheme 4. Synthesis of 6-alkoxy benzoxazolone carboxamides **31-33**




Scheme 5. Synthesis of benzoyl-benzoxazolone carboxamides 34-36

Scheme 6. Synthesis of benzoxazolone carboxamide derivatives **53-66**




Method B: 1) triphosgene, pyridine; 2) R-NH₂ (**67e**, **67l-n**), pyridine. Method C: 1) triphosgene, Et₃N, DCM; 2) R-NH₂ (**67e-d**, **67f-k**), Et₃N, DCM.

Scheme 7. Synthesis of amines **67a-j**^a

^aYields are not reported when compounds were used as crude in the following step. Alcohols **68f-i** and amines **67k-n** were commercially available.


Table 1. Inhibitory potencies (IC₅₀) of compounds on human AC activity^a



Compounds	R	IC ₅₀ (nM) ± S.E.M.
1	H	64 ± 7 ^b
2	6-Br	31 ± 9 ^b
4	4-Br	n.d.
5	5-Br	22 ± 11
6	7-Br	18 ± 11
7	4-Me	n.d.
8	5-Me	117 ± 26
9	6-Me	66 ± 10
10	7-Me	23 ± 5
11	5-F	12 ± 4
12	6-F	31 ± 5
13	5-Cl	15 ± 7
14	6-Cl	33 ± 4
15	5-CF ₃	18 ± 2
16	6-CF ₃	20 ± 9
17	5-NO ₂	3 ± 2
18	6-NO ₂	3 ± 1
19	5,6-Cl	18 ± 8

^aIC₅₀ values are reported as mean values of two or more determinations.

^bIC₅₀ values from ref.¹² n.d: not determined due to low chemical stability.


Table 2. Inhibitory potencies (IC_{50}) of compounds on human AC activity^a

Compounds	R	IC_{50} (nM) \pm S.E.M.
3	6-(<i>p</i> -F-Ph)	79 ± 31^b
20	4-Ph	0.8 ± 0.1
21	5-Ph	42 ± 21
22	6-Ph	84 ± 35
23	7-Ph	91 ± 10
24	5-(<i>p</i> -F-Ph)	56 ± 14
25	5-(<i>p</i> -OCH ₃ -Ph)	67 ± 58
26	6-(<i>p</i> -OCH ₃ -Ph)	67 ± 5
34	5-(COPh)	24 ± 6
35	6-(COPh)	29 ± 1
36	6-CO(<i>p</i> -Cl-Ph)	14 ± 4

^a IC_{50} values are reported as mean values of two or more determinations.

^b IC_{50} values from ref.¹²

Table 3. Inhibitory potencies (IC₅₀) of compounds on human AC activity^a




Compounds	R	IC ₅₀ (nM) ± S.E.M.
1	(CH ₂) ₄ Ph	64 ± 7 ^b
53	(CH ₂) ₂ Ph	121 ± 19
54	(CH ₂) ₃ Ph	55 ± 6
55	(CH ₂) ₅ Ph	54 ± 12
56	(CH ₂) ₆ Ph	51 ± 16
57		71 ± 13
58		312 ± 90
59		75 ± 1
60		63 ± 22
61		25 ± 14
62		649 ± 270
63		65 ± 26
64		No inhibition ^c
65		37 ± 21
66		19 ± 2

^aIC₅₀ values are reported as mean values of two or more determinations.

^bIC₅₀ values from ref.¹² ^cTested at 5 μM.

Table 4. Chemical stability of compounds **1-3**, **14**, **19-26** and **35-36** in PBS (pH 7.4)^a



Compounds	R	Buffer stability $t_{1/2}$ (min)
1	H	45 ^b
2	6-Br	24 ^b
3	6-(<i>p</i> -F-Ph)	>300 ^b
14	6-Cl	58
19	5,6-Cl	55
20	4-Ph	7
21	5-Ph	296
22	6-Ph	>300
23	7-Ph	>300
24	5-(<i>p</i> -F-Ph)	>300
25	5-(<i>p</i> -OCH ₃ -Ph)	176
26	6-(<i>p</i> -OCH ₃ -Ph)	132
35	6-(COPh)	98
36	6-CO(<i>p</i> -Cl-Ph)	274

^aDetermined at room temperature by LC-MS analysis.

^b $t_{1/2}$ values from ref.¹²