

# Click Syntheses of 1,2,3-Triazolylbiferrocenyl Dendrimers and the Selective Roles of the Inner and Outer Ferrocenyl Groups in the Redox Recognition of $\text{ATP}^{2-}$ and $\text{Pd}^{2+}$ \*\*

Rodrigue Djeda, Amalia Rapakousiou, Liyuan Liang, Nicola Guidolin, Jaime Ruiz, and Didier Astruc\*

Ferrocenyl (Fc) dendrimers<sup>[1]</sup> and polymers<sup>[2]</sup> have attracted much attention owing to their multielectron redox properties and functions as biosensors,<sup>[3]</sup> virus-like receptors,<sup>[4a]</sup> models of molecular batteries,<sup>[4b]</sup> and colorimetric sensors.<sup>[2]</sup> Fc-terminated dendrimers belong to the large family of redox-active metallodendrimers that may eventually mimic related nanometer-sized biological processes and provide useful energy-relevant devices.<sup>[5]</sup>

In typical examples, Nishihara and co-workers have extensively studied the efficient electrodeposition of acylbiFc nanodevices in gold nanoparticles,<sup>[6]</sup> and recently, the group of Reinhoudt and Ravoo elegantly demonstrated that small acylbiFc dendrimers form inclusion complexes with self-assembled monolayered  $\beta$ -cyclodextrin as “molecular printboards”.<sup>[7]</sup> Biferrocene itself can be incorporated without covalent bonding as a biferrocenium charge-transfer complex with the arylimino groups of Yamamoto’s arylazidomethine dendrimers.<sup>[8]</sup> From these studies, it has become clear that the redox properties of the biFc derivatives, which include three easily accessible oxidation states, are much richer than those of the Fc group.

We have envisaged taking advantage of the possibility of stabilizing class II mixed-valence biferrocenium cations and exploiting specific properties of the two Fc groups therein.<sup>[9,10]</sup> Herein we report 1) the  $\text{Cu}^{\text{I}}$ -catalyzed azide alkyne “click” cycloaddition ( $\text{CuAAC}$ )<sup>[11]</sup> of the known alkyne ethynylbiferrocene **1** with five generations of azido-terminated dendrimers containing 3<sup>n</sup> terminal tethers, from  $n = 2$  ( $G_0$ , 9 termini) to  $n = 6$  ( $G_4$ , 729 termini), leading to the formation of large 1,2,3-triazolylFc-terminated dendrimers, 2) the isolation and full

characterization of the first mixed-valence dendrimer, 3) the redox recognition, with positive dendritic effects, of both the  $\text{ATP}^{2-}$  anion using the outer Fc groups of the dendrimer and  $\text{Pd}^{\text{II}}$  using the inner Fc groups of the dendrimers, and 4) the role of the mixed-valence metallodendrimers.

The new complex **1** was synthesized in 50 % overall yield from acetylbiFc<sup>[12]</sup> in a reaction that parallels that known for the synthesis of ethynylferrocene<sup>[13]</sup> (see the Supporting Information). The  $\text{CuAAC}$  reaction (Scheme 1) was carried out using  $\text{CuSO}_4$ /sodium ascorbate as the  $\text{Cu}^{\text{I}}$  source between the terminal alkyne **1** and five generations  $G_0$  to  $G_4$  of arene-centered dendrimers dend- $\text{N}_3$  constructed according to 1 $\rightarrow$ 3 connectivity<sup>[14]</sup> and containing in theory respectively 9 ( $G_0$ ), 27 ( $G_1$ ), 81 ( $G_2$ ), 243 ( $G_3$ ), and 729 ( $G_4$ ) azido termini.<sup>[15,16]</sup>

The five triazolylbiFc dendrimers of generations  $G_0$ – $G_4$  are yellow solids that are soluble in dichloromethane and THF but insoluble in hydrocarbons, diethyl ether, and acetonitrile. They are air-stable and thermally stable, except that  $G_3$ –243 slowly becomes insoluble in all solvents, which we tentatively attribute to supramolecular polymerization owing to interpenetration of the terminal tethers among dendrimers. This irreversible catenation is selective to the interbranch spacing for this precise dendritic generation, and remarkably this phenomenon is not observed when the interbranch spacing is larger (in  $G_2$ –81) or smaller (in  $G_4$ –729).

Besides standard  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy, infrared spectroscopy ( $G_0$ – $G_4$ ), and MALDI TOF mass spectrometry ( $G_0$ –9:  $m/z$  calcd 5086.98; found 5087.11,  $M^+$ , 100 %), these metallodendrimers were also characterized by elemental analysis, size exclusion chromatography (SEC, Figure 1), and dynamic light scattering (DLS); the latter two techniques show the mass and size progression (Table 1).

The DLS measurements show that the size increase is reduced between  $G_3$ –243 and  $G_4$ –729 as compared to large size increases between the generations up to  $G_3$ , which must be due to steric congestion at the dendrimer surface in  $G_4$ –729 forcing extensive backfolding of the rather large triazolylbiFc termini inside the dendrimer.<sup>[17]</sup>


Cyclic voltammetry (CV) of the five metallodendrimers was recorded on Pt in  $\text{CH}_2\text{Cl}_2$  using 0.1 M  $[\text{nBu}_4\text{N}][\text{PF}_6]$  as the supporting electrolyte and decamethylferrocene ( $\text{FcH}^*$ ) as the internal reference. Each dendrimer shows two reversible waves at 0.43 and 0.75 V vs.  $\text{FcH}^{*+0}$  (Figure 2a), with the adsorption of the dendrimers on the electrode surface dramatically growing with increasing dendrimer generation. For instance,  $\Delta(E_{\text{pc}} - E_{\text{pa}})$ , which has a value of 60 mV for  $G_0$ –9

[\*] R. Djeda, A. Rapakousiou, L. Liang, Dr. J. Ruiz, Prof. D. Astruc  
Institut des Sciences Moléculaires, UMR CNRS N°5255  
Université Bordeaux 1, 33405 Talence Cedex (France)  
Fax: (+33) 5-4000-2995  
E-mail: d.astruc@ism.u-bordeaux1.fr  
Homepage: <http://astruc.didier.free.fr>


N. Guidolin  
Laboratoire de Chimie des Polymères Organiques  
UMR CNRS N° 5629  
Université Bordeaux 1, 33607 Pessac Cedex (France)

[\*\*] Helpful assistance and discussion of Mössbauer data with Prof. Azzedine Bousseksou (LCC, Toulouse), EPR data with Mattieu Duttine (ICMCB, Pessac), and syntheses with Dr. A. K. Diallo (ISM), and financial support from the Université Bordeaux I, the CNRS, and the ANR are gratefully acknowledged.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201004756>.



**Scheme 1.** The CuAAC reaction and structures of dendrimers  $G_0\text{-}9$ ,  $G_1\text{-}27$ , and  $G_2\text{-}81$ .



**Figure 1.** Size exclusion chromatograms of the triazolybiferrocenyl dendrimers ( $t_R$  = retention time). The polydispersity indices (PDI) are 1.01 ( $G_0\text{-}9$ ), 1.02 ( $G_1\text{-}27$ ), 1.15 ( $G_2\text{-}81$ ), and 1.08 ( $G_3\text{-}263$ ).  $G_4\text{-}729$  appeared to be too large for SEC analysis, but its chloromethylsilyl-terminated  $G_4$  precursor provided a satisfactory SEC trace with PDI = 1.03 (see the Supporting Information).



**Figure 2.** Cyclic voltammograms of  $G_2\text{-}81$ : a) in  $\text{CH}_2\text{Cl}_2$ ,  $[\text{nBu}_4\text{N}][\text{PF}_6]$  0.1 M; see Table 2 for data and conditions; b) progressive adsorption upon scanning around the biFc area; c) spitting of the first CV wave upon addition of ATP (the second wave is not represented, because its scanning destroys the first one); d) addition of  $\text{Pd}(\text{OAc})_2$  provoking the splitting of the second wave.

**Table 1:** Hydrodynamic diameters of the metallodendrimers (obtained by DLS), and calculated diffusion coefficient, volume, and density.

Product <sup>[a]</sup>	MM <sup>[b]</sup>	$2 \times R_h$ <sup>[c]</sup>	$D$ <sup>[d]</sup> [ $\text{m}^2 \text{s}^{-1}$ ]	Volume <sup>[e]</sup> [ $\text{m}^3$ ]	$d$ <sup>[f]</sup> [ $\text{kg m}^{-3}$ ]
$G_1\text{-}27$	16 927	8.6	$10.84 \times 10^{-14}$	$3.33 \times 10^{-25}$	84
$G_2\text{-}81$	52 522	14.7	$6.34 \times 10^{-14}$	$16.63 \times 10^{-25}$	52
$G_3\text{-}243$	159 306	25.2	$3.70 \times 10^{-14}$	$83.79 \times 10^{-25}$	32
$G_4\text{-}729$	479 928	27.2	$3.43 \times 10^{-14}$	$105.36 \times 10^{-25}$	76

[a] The hydrodynamic diameters of  $G_0\text{-}9$  could not be obtained by DLS because it is below the lower limit of the technique. [b] MM: molecular mass ( $\text{g mol}^{-1}$ ). [c]  $2 \times R_h$ : hydrodynamic diameter (nm) measured in THF at 25 °C. [d]  $D$ : diffusion coefficient. [e] considering the globular shape of the dendrimer as a perfect sphere ( $V = (4/3)\pi R_h^3$ ). [f]  $d$ : density.

at 25 °C, progressively drops to 0 for  $G_3\text{-}243$  and  $G_4\text{-}729$  (Table 2), signifying complete adsorption at the first scan. This result is an advantage for the facile formation of robust metallodendrimer-modified electrodes (Figure 2b) upon scanning around the biFc potential zone. The 1,2,3-triazolyl group is an electron-withdrawing substituent for the inner Fc

**Table 2:** Cyclic voltammetry data for the triazolybiferrocenyl dendrimers.<sup>[a]</sup>


Product	$E_{1/2}$ [V] <sup>[b]</sup>	$\Delta$ <sup>[b]</sup>	$I_{pc}/I_{pa}$ <sup>[b]</sup>	$E_{1/2}$ (V) <sup>[c]</sup>	$\Delta$ <sup>[c]</sup>	$I_{pc}/I_{pa}$ <sup>[c]</sup>
G <sub>0</sub> -9	0.430	0.060	1.25	0.740	0.060	3
G <sub>1</sub> -27	0.430	0.045	1.6	0.745	0.030	3.2
G <sub>2</sub> -81	0.430	0.005	1.8	0.745	0.010	1.3
G <sub>3</sub> -243	0.430	0.000	1.9	0.750	0.000	1.5
G <sub>4</sub> -729	0.430	0.000	2	0.740	0.000	1.8

[a] Solvent: CH<sub>2</sub>Cl<sub>2</sub>; electrolyte [nBu<sub>4</sub>N][PF<sub>6</sub>] 0.1 M; scan rate: 0.200 V s<sup>-1</sup>; 25 °C; working and counter electrodes: Pt; quasi-reference electrode: Ag; all half-wave potentials  $E_{1/2}$  are given versus the internal reference system [Fe( $\eta^5$ -C<sub>5</sub>Me<sub>3</sub>)<sub>2</sub>]<sup>+/0</sup>. [b] ( $E_{pc}$ – $E_{pa}$ ) for the first wave. [c] ( $E_{pc}$ – $E_{pa}$ ) for the second wave. See also Ref. [19b].

group, whereas the outer Fc group only bears the electron-releasing inner Fc group. Thus, if the mixed valence is localized (see below), the first oxidation wave at 0.43 V vs. FcH<sup>+</sup> can be assigned to the outer Fc groups that are easier to oxidize, whereas the second one at 0.74 V vs. FcH<sup>+</sup> can be assigned to the inner Fc groups. This assignment is confirmed by the remarkable selectivity of redox molecular recognition studies with the inner and the outer Fc groups.

Indeed, addition of the adenosyl triphosphate salt [nBu<sub>4</sub>N]<sub>2</sub>[ATP] to the electrochemical cell containing G<sub>2</sub>-81 provokes a splitting of the outer Fc CV wave at 0.43 V owing, to some extent, to ion-pairing interaction<sup>[18]</sup> between [ATP]<sup>2-</sup> and [outer Fc]<sup>+</sup>, reflected by the new part of this CV wave (Figure 2c). Geiger and co-workers have shown that ion pairing can also have an important influence on the separation between the two CV waves of binuclear or multinuclear redox systems. For instance, these authors demonstrated that exchanging the ion-pairing BF<sub>4</sub> or PF<sub>6</sub> anion in the supporting electrolyte for the non-ion-pairing anions BAR<sub>4</sub> (Ar = C<sub>6</sub>F<sub>5</sub> or *meta*-C<sub>6</sub>H<sub>3</sub>(CF<sub>3</sub>)<sub>2</sub>) provoked an increase of the splitting between the CV waves of biferrocene, oligoferrocenes, and other polynuclear complexes.<sup>[19a–c]</sup> We have compared the cyclic voltammograms of biferrocene<sup>[9,19d]</sup> and benzyltriazo-lybiferrocene<sup>[20]</sup> with those of the triazolybiferrocenyl dendrimers using both electrolytes [nBu<sub>4</sub>N][PF<sub>6</sub>] and [nBu<sub>4</sub>N][B(*meta*-C<sub>6</sub>H<sub>3</sub>(CF<sub>3</sub>)<sub>2</sub>)<sub>4</sub>], and we also observed a large increase of the  $E_2$ – $E_1$  value (480 mV) using the later. However, this enhancement is subjected to a negative dendritic effect,<sup>[19b]</sup> as opposed to the dramatic positive dendritic effect found in the recognition of the ATP anion described above, owing to synergistic additional supramolecular interactions that are enhanced by dendritic effects.

On the other hand, addition of Pd(OAc)<sub>2</sub> provokes the splitting of the CV wave at 0.74 V owing to coordination of the triazolyl ligand<sup>[20]</sup> attached to the inner Fc group, while the CV wave of the outer Fc group is left unchanged (Figure 2d). This splitting by 80 mV is due to the dendritic effect, because it is not observed at all with benzyltriazo-lybiferrocene.<sup>[20]</sup> If the Pd<sup>II</sup> cations are introduced as [Pd(MeCN)<sub>4</sub>][PF<sub>6</sub>]<sub>2</sub>, an analogous phenomenon is still observed, but a purple precipitate immediately forms, owing to the oxidized dendrimers that stabilize the Pd nanoparticles (PdNPs). The oxidized dendrimer and the PdNPs absorb in the UV/Vis spectrum around 600 nm (Figure 3), and these PdNPs stabi-




**Figure 3.** A) UV/Vis spectra of G<sub>1</sub>-27 and [G<sub>1</sub>-27][PF<sub>6</sub>]<sub>27</sub>. B) EPR spectra of [G<sub>1</sub>-27][PF<sub>6</sub>]<sub>27</sub>: a) experimental; b) simulated;  $g_{||} = 3.413$ ,  $g_{\perp} = 1.885$ ,  $T = 4$  K; B = magnetic field.

lized by the G<sub>0</sub>–G<sub>2</sub> biferrocenium dendrimers absorb at 530–550 nm (see the Supporting Information). The PdNPs formed in this way efficiently catalyze the Miyaura–Suzuki coupling of PhB(OH)<sub>2</sub> with PhI at 20 °C for 24 h at a concentration of 0.1 % Pd catalyst. These PdNPs are stable for a few hours. Although PdNPs did not form upon addition of Pd(OAc)<sub>2</sub> to the electrochemical cell, they formed upon further addition of methanol that reduced Pd<sup>II</sup> to G<sub>1</sub>-27-protected PdNPs. At 0.1 % Pd vs. substrates, yields of 33 % to 42 % of C–C coupling product were obtained in 24 h at 25 °C for PdNPs stabilized by G<sub>0</sub>–G<sub>2</sub> dendrimers, and this yield reached 53 % of C–C coupled product upon diluting the G<sub>1</sub>-PdNPs ten times to a concentration of 0.01 % Pd. Both the generation near-independency and the yield increase upon dilution are consistent with a leaching mechanism that generates catalytically active Pd atoms in solution,<sup>[21]</sup> which has already been observed with dendrimer-supported PdNPs.<sup>[21b,c]</sup>


In contrast to triazolybiferrocenyl dendrimers, triazolybiferrocenyl-terminated dendrimers did not reduce [Pd(MeCN)<sub>4</sub>][PF<sub>6</sub>]<sub>2</sub>, thus showing the better efficiency of the outer Fc groups to reduce [Pd(MeCN)<sub>4</sub>][PF<sub>6</sub>]<sub>2</sub> compared to triazoly-substituted Fc groups either in triazolybiferrocenyl-terminated dendrimers or in triazolybiferrocenyl-terminated dendrimers. This finding also shows that mixed-valence triazolybiferrocenyl-terminated dendrimers stabilize catalytically active PdNPs.

To determine the electronic structure of these mixed-valence triazolybiferrocenyl-terminated dendrimers, we oxidized G<sub>1</sub>-27 with [FcH][PF<sub>6</sub>], which has a redox potential only slightly more positive (by 70 mV) than the redox potential of the outer Fc groups of the metallodendrimers. We therefore reasoned that precipitation of the mixed-valence polycationic metallodendrimer should completely drive the redox equilibrium toward its formation. Indeed, the addition of 27 equivalents [FcH][PF<sub>6</sub>] to G<sub>1</sub>-27 in CH<sub>2</sub>Cl<sub>2</sub> and subsequent addition of diethyl ether provided soluble ferrocene and the dark-blue dendritic complex [G<sub>1</sub>-27][PF<sub>6</sub>]<sub>27</sub> (Scheme 2), which was isolated as a powder subsequent to reprecipitation by addition of diethyl ether to a MeCN solution. [G<sub>1</sub>-27][PF<sub>6</sub>]<sub>27</sub> was soluble in MeCN but almost insoluble in CH<sub>2</sub>Cl<sub>2</sub>, and it was characterized by satisfactory elemental analysis and by UV/Vis and EPR spectroscopy (Figure 3), which showed the



Scheme 2.

classic spectrum of  $d^5$  ferrocenium, and by Mössbauer spectroscopy under zero field at 77 K (Figure 4) and 293 K (see the Supporting Information). Both spectra show a localized mixed-valence state, as for parent biferrocenium itself<sup>[9]</sup> at this frequency ( $10^{-8}$  s).



**Figure 4.** Mössbauer spectra of  $[G_1-27][PF_6]_{27}$  at zero field and 80 K. Outer quadrupole doublet (inner Fc,  $Fe^{II}$ ): isomer shift (IS) = 0.506(3), quadrupole splitting (QS) = 2.224(6). Central doublet (outer ferrocenium group): IS = 0.51(1); QS = 0.491(17). See the spectrum at 293 K in the Supporting Information, also showing localized mixed valence.

The complex  $[G_1-27][PF_6]_{27}$  may be considered as belonging to class II of mixed-valence complexes according to the Robin–Day classification,<sup>[22]</sup> because the first coordination spheres of the two iron centers are identical with cyclopentadienyl rings, even if the ring substituents are different, provoking a dissymmetry of the two valence potential wells.

In conclusion, the first “click” synthesis, isolation, and characterization of five generations of triazolybiFc dendrimers has been carried out. Their electronic structure and role as the first mixed-valence dendrimers have also been determined. The mixed-valence species can be considered to belong to class II from the Mössbauer data, despite a splitting of the CV waves that reaches 480 mV using the supporting electrolyte  $[nBu_4N][B\{meta-C_6H_3(CF_3)_2\}_4]$ . Redox recognition is also remarkable, as it selectively involves the first CV wave for  $ATP^{2-}$  (outer Fc) and the second one for  $Pd^{II}$  in  $Pd(OAc)_2$  and  $[Pd(MeCN)_4][PF_6]_2$  (inner Fc) with dendritic effects, and both the neutral and mixed-valence dendrimers stabilize catalytically active Pd nanoparticles.

## Experimental Section

“Click” synthesis of the 1,2,3-triazolybiFc dendrimers  $G_n-3^{n+2}$ , ( $n = 0-4$ ): Ethynylbiferrocene **1** (1.5 equiv per branch,) and the azido-terminated dendrimer were dissolved in oxygen-free THF (50 mL), and oxygen-free water (50 mL) was added. A solution of 1 M  $CuSO_4$

(1 equiv per branch) was added at 0°C, with subsequent dropwise addition of a freshly prepared solution of 1 M sodium ascorbate (2 equiv per branch). The brown solution at 0°C changed to orange at room temperature. The reaction mixture was allowed to stir for 24 h under  $N_2$  at room temperature. Then  $CH_2Cl_2$  (100 mL) was added followed by the addition of an aqueous solution of ammonia. The mixture was allowed to stir for 10 min in order to remove all the copper salts trapped inside the dendrimer. The organic phase was washed twice with water, dried over sodium sulfate, and filtered, and the solvent was removed under vacuum. The product was then washed with methanol to remove the excess alkyne and precipitated from  $CH_2Cl_2$  with methanol and then with pentane, giving a yellow powder.  $G_0-9$ : 0.134 g, yield 80%. Elemental analysis calcd (%) for  $C_{261}H_{291}Fe_{18}N_{27}Si_9$ : C 61.56, H 5.72; found: C 60.58, H 7.91. The  $G_1-G_4$  dendrimers were obtained analogously (see the Supporting Information).

Synthesis of the mixed-valence dendrimer  $[G_1-27][PF_6]_{27}$ : A solution of ferrocenium hexafluorophosphate (0.043 g, 0.128) in  $CH_2Cl_2$  (50 mL) was added to a solution of  $G_1-27$  (0.080 g, 0.005) in  $CH_2Cl_2$  (5 mL) at room temperature, formation of a dark-blue precipitate was observed, and the mixture was allowed to stir under  $N_2$  for 1 h at room temperature. Complete precipitation of the product was obtained by dropwise addition of diethyl ether (50 mL) to the reaction mixture. After filtration under  $N_2$  on Celite, the solid was dried under vacuum to give  $[G_1-27][PF_6]_{27}$  as a dark-blue powder (0.062 g, 66.3% yield). Elemental analysis calcd (%) for  $C_{882}H_{1029}Fe_{54}N_{81}O_9Si_{36}P_{27}F_{162}$ : C 50.70, H 5.01; found: C 49.74, H 5.05. From the filtrate, 0.023 g (96%) ferrocene was recovered.

Received: July 31, 2010

Published online: September 20, 2010

**Keywords:** dendrimers · metallocenes · mixed-valent compounds · nanoparticles · redox chemistry

- [1] For reviews on Fc dendrimers and their electrochemical properties, see: C. M. Casado, I. Cuadrado, M. Morán, B. Alonso, B. Garcia, B. Gonzales, J. Losada, *Coord. Chem. Rev.* **1999**, 185–186, 53–79; S. H. Hwang, C. D. Shreiner, C. N. Moorefield, G. R. Newkome, *New J. Chem.* **2007**, 31, 1027–1038; D. Astruc, C. Ornelas, J. Ruiz, *Acc. Chem. Res.* **2008**, 41, 841–856.
- [2] P. Nguyen, P. Gomez-Elipse, I. Manners, *Chem. Rev.* **1999**, 99, 1515–1548; J. C. Eloi, L. Chabanne, G. R. Whittell, I. Manners, *Mater. Today* **2008**, 11, 28–36. For the electrochemistry of multi-Fc polymers, see: R. Rulkens, A.-J. Lough, I. Manners, S. R. Lovelace, C. Grant, W. E. Geiger, *J. Am. Chem. Soc.* **1996**, 118, 12683–12695.
- [3] J. Losada, M. P. G. Armada, M. Zamora, B. Alonso, I. Cuadrado, C. M. Casado, *Bioelectrochemistry* **2006**, 69, 65; J. Losada, M. Zamora, P. G. Armada, I. Cuadrado, B. Alonso, C. M. Casado, *Anal. Bioanal. Chem.* **2006**, 385, 1209; F. J. Martinez, B. Gonzalez, B. Alonso, J. Losada, M. P. Garcia-Armada, C. M. Casado, *J. Inorg. Organomet. Polym. Mater.* **2008**, 18, 51.
- [4] M.-C. Daniel, J. Ruiz, D. Astruc, *J. Am. Chem. Soc.* **2003**, 125, 1150–1151; C. Ornelas, J. Ruiz, C. Belin, D. Astruc, *J. Am. Chem. Soc.* **2009**, 131, 590–601.
- [5] For reviews on redox-active metallodendrimers, see: G. R. Newkome, E. He, C. N. Moorefield, *Chem. Rev.* **1999**, 99, 1689–1746; M. Venturi, S. Serroni, A. Juris, S. Campagna, V. Balzani in *Dendrimers, Topics Curr. Chem.*, Vol. 197 (Ed.: F. Vögtle), Springer, Heidelberg, **1998**, pp. 193–228; C. M. Casado, M. Morán, B. Alonso, J. Losada, *Coord. Chem. Rev.* **1999**, 193–195, 395–445; D. Astruc, E. Boisselier, C. Ornelas, *Chem. Rev.* **2010**, 110, 1857–1959.
- [6] T. Horikoshi, M. Itoh, M. Kurihara, K. Kubo, H. Nishihara, *J. Electroanal. Chem.* **1999**, 473, 113–116; H. Nishihara, *Bull.*



- Soc. Chem. Jpn.* **2001**, *74*, 19–29; M. Yamada, H. Nishihara, *Chem. Commun.* **2002**, 2578–2579; M. Yamada, H. Nishihara, *Eur. Phys. J.* **2003**, *24*, 257–260; M. Yamada, H. Nishihara, *Langmuir* **2003**, *19*, 8050–8056; M. Yamada, H. Nishihara, *ChemPhysChem* **2004**, *5*, 555–559; M. Yamada, T. Tadera, K. Kubo, H. Nishihara, *J. Phys. Chem. B* **2003**, *107*, 3703–3711; M. Muraa, H. Nishihara, *J. Inorg. Organomet. Polym.* **2005**, *15*, 147–156.
- [7] C. A. Nijhuis, K. A. Dolatowska, B. J. Ravoo, J. Huskens, D. N. Reinhoudt, *Chem. Eur. J.* **2007**, *13*, 69–80.
- [8] Y. Ochi, M. Suzuki, T. Imaoka, M. Murata, H. Nishihara, Y. Einaga, K. Yamamoto, *J. Am. Chem. Soc.* **2010**, *132*, 5061–5069.
- [9] D. O. Cowan, F. Kaufman, *J. Am. Chem. Soc.* **1970**, *92*, 219–220; D. O. Cowan, F. Kaufman, *J. Am. Chem. Soc.* **1971**, *93*, 3889–3893; C. Levanda, D. O. Cowan, K. Bechgaard, *J. Am. Chem. Soc.* **1975**, *97*, 1980–1981; M. J. Powers, T. J. Meyer, *J. Am. Chem. Soc.* **1978**, *100*, 4393–4398.
- [10] M. Yamada, *Bull. Soc. Chim. Jpn.* **2009**, *82*, 152–170.
- [11] R. F. Kovar, M. D. Rausch, *J. Org. Chem.* **1972**, *35*, 351–366.
- [12] G. Doisneau, G. Balavoine, T. Fillebeen-Khan, *J. Organomet. Chem.* **1992**, *425*, 113–117.
- [13] V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem.* **2002**, *114*, 2708; *Angew. Chem. Int. Ed.* **2002**, *41*, 2596; C. W. Tornøe, C. Christensen, M. Meldal, *J. Org. Chem.* **2002**, *67*, 3057–3064; M. Meldal, C. W. Tornøe, *Chem. Rev.* **2008**, *108*, 2952–3015. For a review of “click” dendrimers, see: G. Franc, A. Kakkar, *Chem. Commun.* **2008**, 5267–5276.
- [14] G. R. Newkome, Z. Yao, G. R. Baker, V. K. Gupta, *J. Org. Chem.* **1985**, *50*, 2003–2004; G. R. Newkome, *Pure Appl. Chem.* **1998**, *70*, 2337–2343; V. V. Narayanan, G. R. Newkome, *Top. Curr. Chem.* **1998**, *197*, 19–33.
- [15] a) Generation 0 of azido dendrimers resulted from  $\pi$ -C<sub>5</sub>H<sub>5</sub>Fe-induced nona-allylation of mesitylene including removal of the organoiron group using visible-light photolysis,<sup>[15b]</sup> and subsequent hydrosilylation with HSi(CH<sub>2</sub>Cl)Me<sub>2</sub> and azidation using NaN<sub>3</sub>,<sup>[16]</sup> b) F. Moulines, L. Djakovitch, R. Boese, B. Gloaguen, W. Thiel, J.-L. Fillaut, M.-H. Delville, D. Astruc, *Angew. Chem.* **1993**, *105*, 1132–1134; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1075–1077; D. Catheline, D. Astruc, *J. Organomet. Chem.* **1983**, *248*, c9–c12; V. Sartor, L. Djakovitch, J.-L. Fillaut, F. Moulines, F. Neveu, V. Marvaud, J. Guittard, J.-C. Blais, D. Astruc, *J. Am. Chem. Soc.* **1999**, *121*, 2929–2930.
- [16] a) Subsequent generations were constructed with a known iteration consisting of a Williamson reaction of the iodomethyl-silyl-terminated dendrimers with the phenol triallyl dendron  $p$ -HO-C<sub>6</sub>H<sub>4</sub>C(CH<sub>2</sub>CH=CH<sub>2</sub>)<sub>3</sub>, followed by the same hydrosilylation and azidation reactions.<sup>[16b]</sup> b) J. Ruiz, G. Lafuente, S. Marcen, C. Ornelas, S. Lazare, J.-C. Blais, E. Cloutet, D. Astruc, *J. Am. Chem. Soc.* **2003**, *125*, 7250–7257; C. Ornelas, J. Ruiz, E. Cloutet, S. Alves, D. Astruc, *Angew. Chem.* **2007**, *119*, 890–895; *Angew. Chem. Int. Ed.* **2007**, *46*, 872–877; D. Astruc, J. Ruiz, *Tetrahedron* **2010**, *66*, 1769–1785.
- [17] P. G. de Gennes, H. Hervet, *J. Phys. Lett.* **1983**, *44*, 351–360; K. Naidoo, S. J. Hugues, J. R. Moss, *Macromolecules* **1999**, *32*, 331; D. Pötschke, M. Ballauf, P. Lindner, M. Fischer, F. Vögtle, *Macromolecules* **1999**, *32*, 4079–.
- [18] a) For a review of the distinct factors involving redox recognition of oxoanions by polyFc hosts, see: P. D. Beer, *Chem. Commun.* **1996**, 689–696; P. D. Beer, *Acc. Chem. Res.* **1998**, *31*, 71–80; P. D. Beer, P. A. Gale, *Angew. Chem.* **2001**, *113*, 502–532; *Angew. Chem. Int. Ed.* **2001**, *40*, 486–516; M. C. Daniel, J. Ruiz, J.-C. Blais, N. Daro, D. Astruc, *Chem. Eur. J.* **2003**, *9*, 4371–4379; D. Astruc, M.-C. Daniel, J. Ruiz, *Chem. Commun.* **2004**, 2637–2649; b) for ATP sensing by Fc sensors, see: O. Reynes, J. C. Moutet, J. Pecaut, G. Royal, E. Saint-Aman, *New J. Chem.* **2002**, *26*, 9–12; c) for ion-pairing interactions, see: A. Loupy, B. Tchoubar, D. Astruc, *Chem. Rev.* **1992**, *92*, 1141–1165; for dendritic ion-pairing effects, see: C. Ornelas, J. Ruiz, D. Astruc, *Organometallics* **2009**, *28*, 4431–4437.
- [19] a) N. Camine, U.-T. Mueller-Westerhoff, W. E. Geiger, *J. Organomet. Chem.* **2001**, 637–639, 823–826; F. Barrière, N. Camine, W. E. Geiger, *J. Am. Chem. Soc.* **2002**, *124*, 7262–7263; b) F. Barrière, W. E. Geiger, *J. Am. Chem. Soc.* **2006**, *128*, 3980–3989; c) F. Barrière, W. E. Geiger, *Acc. Chem. Res.* **2010**, *43*, 1030–1039. d) The  $E_2$ – $E_1$  value is found to be only slightly larger for these monomeric biferrocenes (by 35 mV) than for the dendrimers with [nBu<sub>4</sub>N][PF<sub>6</sub>], but more significantly larger for the monomers than for the dendrimers (by 120 mV) with [nBu<sub>4</sub>N][B{meta-C<sub>6</sub>H<sub>3</sub>(CF<sub>3</sub>)<sub>2</sub>}]<sub>4</sub>.
- [20] S. Badèche, J.-C. Daran, J. Ruiz, D. Astruc, *Inorg. Chem.* **2008**, *47*, 4903–4908.
- [21] a) J. G. de Vries, *Dalton Trans.* **2006**, 421–429; b) A. K. Diallo, C. Ornelas, L. Salmon, J. Ruiz, D. Astruc, *Angew. Chem.* **2007**, *119*, 8798–8802; *Angew. Chem. Int. Ed.* **2007**, *46*, 8644–8648; c) C. Ornelas, J. Ruiz, L. Salmon, D. Astruc, *Adv. Synth. Catal.* **2008**, *350*, 837–845.
- [22] M. B. Robin, B. Melvin, P. Day, *Adv. Inorg. Chem. Radiochem.* **1968**, *10*, 247–422.