UNEXPECTED SYNTHESIS OF 4-R-PHENYLALLYLIDENE MELDRUM'S ACID DERIVATIVES

Braulio Insuasty,¹* Harlen Torres,^{1,2} Rodrigo Abonia,¹ Jairo Quiroga,¹* John Low,³ Adolfo Sanchez,⁴ Justo Cobo⁴ and Manuel Nogueras⁴

¹Grupo de Investigación de Compuestos Heterocíclicos, Department of Chemistry, Universidad del Valle, A. A. 25360, Cali, Colombia; ²Universidad Nacional de Colombia Sede Palmira, Chapinero via Candelaria, Palmira- Colombia; ³Department of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland; ⁴Department of Inorganic and Organic Chemistry, Universidad de Jaén, 23071 Jaén, Spain.

Abstract: Six novel cinnamylidene Meldrum's acid derivatives 3 have been obtained from the reaction of benzaldehydes 1 and Meldrum's acid 2, by heating under reflux in 2-ethoxyethanol and catalytic amounts of triethylamine. Analysis of the NMR-bidimensional experiments and X-ray diffraction of some derivatives 3 allowed us to determine unambiguously its structures.

Introduction

From a synthetic viewpoint, Meldrum's acid and its derivatives are used most often in the preparation of arylidene-condensation products (i.e. compounds that have a strongly polarized double bond).¹ At the simplest level, Meldrum's acid can act as a methylene synthon from hydrolysis, loss of acetone molecule and exhaustive decarboxylation.²

The use of arylidene derivatives of Meldrum's acid in the synthesis of heterocyclic compounds has received little attention, although despite their relatively easy preparation and the mild reaction conditions required in their reactions made them quite valuable starting materials for synthetic purposes.³ For example, it has been reported that the condensation of arylidene Meldrum's acid derivatives with heterocyclic monoamines and α , β -unsaturated carbonyl or β -dicarbonyl compounds affords coumarin⁴ or pyridine⁵ derivatives respectively.

Results and Discussion

As an attempt to prepare benzylidene derivatives of Meldrum's acid and use them in the reaction with heterocyclic amines, a mixture of equimolar amounts of benzaldehyde 1a and Meldrum's acid 2 was heated at reflux in 5 mL of 2-ethoxyethanol and catalytic amounts of triethylamine. A new compound precipitated from the hot solution was fully characterized through spectroscopic and analytical methods as the 3-phenylpropylidene Meldrum's acid derivative 3a in which apparently two molecules of 2 are involved in its formation (Scheme 1). The reaction was then optimized using a molar ratio 1:2 of 1a and 2 increasing substantially the reaction yield. These

results prompted us to explore other aldehydes (1b-f), showing a similar behavior and yielding their corresponding products **3b-f** in acceptable to good yields.



Scheme 1

All compounds **3a-f** were extensively characterized by their ¹H, ¹³C nmr and DEPT spectra (including 2D-techniques like COSY and HMBC), mass spectra, elemental analysis and X-ray diffraction.

All signals in the ¹H nmr spectra are consistent with the proposed structure for compounds 3, where the most relevant feature is the presence of the proton signals of the fragment =CH(1')-CH(2')=CH(3')-Ar at δ 8.15-8.18, 8.25-8.42 and 7.35-7.43 ppm respectively, each one showing a doublet with J between 15.3 – 16.1 Hz. While the aromatic protons appear at δ = 7.08-8.31.

The ¹³C nmr spectra shows a characteristic pattern for the fragment =CH(1')-CH(2')=CH(3')-Ar at δ 155.2-157.7, 121.7-127.3 and 149.9-155.0 ppm respectively. The other aliphatic and aromatic carbon atoms were also assigned.

All mass spectra are characterized by showing their molecular ions. This finding could reflects the relatively stability of these structures which is the base of their possible applications in organic synthesis.

Finally, we confirmed the proposed structure for compounds 3 by direct synthesis of the compound 3a from the reaction of cinnamaldehyde and Meldrum's acid in the presence of Cdl₂ (Scheme 2). This methodology has previously been successfully used by us for the synthesis of arylidene Meldrum's acid derivatives.^{5b-d}





Furthermore, the crystalline structures for two of such compounds **3b** and **3d** have been recently reported by us,⁶ confirming the proposed structures. Crystalline structures of two arylidenederivatives of Meldrum's acid were described in the work.⁷

In summary, we have synthesized and characterized six new 3-arylallylidene derivatives of Meldrum's acid in acceptable to good yields, starting with benzaldehydes under mild conditions. This finding open additional possibilities of potential applications of this kind of compounds, for example as $\square\square$ -unsaturated carboxylic systems in cyclocondensation reactions with heterocyclic monoamines.

EXPERIMENTAL

Melting points were determined in a Buchi Melting Point Apparatus and are uncorrected. The ¹H- and ¹³C nmr spectra were run on a Bruker DPX 300 spectrometer operating at 300 MHz and 75 MHz respectively, using CDCl₃ as solvent and tetramethylsilane as internal standard. The mass spectra were scanned on a Hewlett Packard HP Engine-5989 spectrometer (equipped with a direct inlet probe) and operating at 70 eV. The elemental analysis has been obtained using a LECO CHNS-900 equipment.

General procedure for the preparartion of the 5-[3-(4-R-phenyl)allylidene]-2,2-dimethyl-[1,3]dioxane-4,6-diones 3a-f.

A solution of benzaldehyde **1a-f** (2.0 mmoles) and Meldrum's acid **2** (4.0 mmoles) in 2ethoxyethanol (5 ml) and catalytic amounts of triethylamine was refluxed during 5-30 minutes (tlc control), after cooling the resulting precipitate was filtered, washed with ethanol, dryed and purified by column chromatography on silica-gel and chloroform as eluent.

5-[3-Phenylallylidene]-2,2-dimethyl-[1,3]dioxane-4,6-dione 3a

This compound was obtained according to general procedure as yellow crystals, mp 101 °C, yield 68 %. This compound also was obtained by the direct heating of mixture of cinnamaldehyde (2.0 mmoles) and Meldrum's acid 2 (2.0 mmoles) in presence of Cdl₂ (mp 101 °C, yield 55 %) ¹H nmr (DMSO-d₆): $\delta = 1.77$ (s, 6H, (CH₃)₂C), 7.43-7.68 (m, 5H, C₆H₅), 7.43 (d, 1H, H-3'), 8.32 (dd, J_{H1'-H2'} = 12.0 Hz, J_{H2'-H3'} = 14.9 Hz, 1H, H-2'), 8.18 (d, 1H, H-1'). ¹³C nmr (DMSO-d₆): $\delta = 27.0$ (CH₃),

104.5 (C-2), 111.6 (C-5), 123.9 (C-2'), 128.8 (C_o), 129.3 (C_m), 131.0 (C_p), 134.8 (C_i), 154.1 (C-3'), 156.9 (C-1'), 160.2 (C=O), 162.1 (C=O). The mass spectrum shows the following peaks: ms: (70 eV) m/z (%) = 258 (M⁺, 22), 201 (24), 200 (95), 172 (40), 155 (23), 144 (53), 132 (34), 131 (53), 128 (100), 127 (44), 116 (24), 115 (27), 102 (40), 77 (26), 51 (23), 43 (95), 39 (15). Anal. Calcd. for $C_{15}H_{14}O_4$: C, 69.76; H, 5.46. Found: C, 69.81; H, 5.56.

5-[3-(4-Methylphenyl)allylidene]-2,2-dimethyl-[1,3]dioxane-4,6-dione 3b

This compound was obtained according to general procedure as orange crystals, mp 128 °C, yield 65 %. ¹H nmr (DMSO-d₆): $\delta = 1.76$ (s, 6H, (CH₃)₂C), 2.41 (s, 3H, CH₃), 7.57 (d, 2H, H_o), 7.24 (dd, 2H, H_m), 7.40 (d, 1H, H-3'), 8.28 (dd; $J_{\text{H1'-H2'}} = 12.0$ Hz, $J_{\text{H2'-H3'}} = 14.8$ Hz, 1H, H-2'), 8.17 (d, 1H, H-1'). ¹³C nmr (DMSO-d₆): $\delta = 21.1$ (CH₃), 27.0 (4-CH₃C₆H₄), 104.4 (C-2), 110.8 (C-5), 123.0 (C-2'), 129.0 (C_o), 129.9 (C_m), 142.2 (C_p), 132.2 (C₁), 154.5 (C-3'), 157.2 (C-1'), 160.3 (C=O), 162.2 (C=O). The mass spectrum shows the following peaks: ms: (70 eV) m/z (%) = 272 (M⁺, 32), 215 (23), 214 (88), 1186 (27), 158 (46), 155 (46), 142 (67), 141 (100), 131 (62), 129 (20), 127 (19), 116 (15), 115 (81), 65 (11), 63 (14), 43 (82), 39 (16).

Anal. Calcd. for C₁₆H₁₆O₄: C, 70.58; H, 5.92. Found: C, 70.49; H, 5.87.

5-[3-(4-Methoxyphenyl)allylidene]-2,2-dimethyl-[1,3]dioxane-4,6-dione 3c

This compound was obtained according to general procedure as orange crystals, mp 147 °C, yield 52 %. ¹H nmr (DMSO-d₆): $\delta = 1.76$ (s, 6H, (CH₃)₂C), 3.87 (s, 3H, CH₃O), 7.64 (d, 2H, H_o), 6.95 (dd, 2H, H_m), 7.39 (d, 1H, H-3'), 8.25 (dd, $J_{\text{H1'-H2'}} = 11.9$ Hz, $J_{\text{H2'-H3'}} = 14.9$ Hz, 1H, H-2'), 8.16 (d, 1H, H-1'). ¹³C NMR (DMSO-d₆): $\delta = 26.9$ (CH₃), 55.5 (OCH₃), 104.3 (C-2), 109.5 (C-5), 121.7 (C-2'), 131.2 (C_o), 114.9 (C_m), 156.5 (C_p), 127.2 (C_i), 155.5 (C-3'), 157.7 (C-1'), 160.4 (C=O), 162.4 (C=O). The mass spectrum shows the following peaks: ms: (70 eV) m/z (%) = 288 (M⁺, 34), 231 (14), 230 (51), 185 (15), 174 (17), 162 (19), 159 (13), 158 (76), 143 (18), 131 (14), 128 (20), 116 (11), 115 (100), 89 (17)m 63 (15), 43 (58), 39 (10).

Anal. Calcd. for C₁₆H₁₆O₅: C, 66.66; H, 5.59. Found: C, 66.65; H, 5.51.

5-[3-(4-Chlorophenyl)allylidene]-2,2-dimethyl-[1,3]dioxane-4,6-dione 3d

This compound was obtained according to general procedure as orange crystals, mp 182 °C, yield 70 %. ¹H nmr (DMSO-d₆): $\delta = 1.77$ (s, 6H, (CH₃)₂C), 7.40 (d, 2H, H_o), 7.60 (dd, 2H, H_m), 7.36 (d, 1H, H-3'), 8.29 (dd, $J_{H1'-H2'} = 12.0$ Hz, $J_{H2'-H3'} = 15.1$ Hz, 1H, H-2'), 8.15 (d, 1H, H-1'). ¹³C NMR (DMSO-d₆): $\delta = 27.0$ (CH₃), 104.9 (C-2), 112.1 (C-5), 124.4 (C-2'), 129.4 (C_o), 130.4 (C_m), 136.0 (C_p), 133.7 (C_i), 152.2 (C-3'), 156.4 (C-1'), 160.1 (C=O), 162.0 (C=O). The mass spectrum shows the following peaks: ms: (70 eV) m/z (%) = 294/292 (M⁺, 10/26), 236 (31), 235 (22), 206 (30), 189 (18), 178 (35), 166 (17), 164 (26), 162 (69), 161 (15), 155 (16), 131 (67), 127 (78), 126 (37), 115 (19), 101 (17), 77 (25), 75 (24), 51 (13), 43 (100), 39 (10).

Anal. Calcd. for C₁₅H₁₃ClO₄: C, 61.55; H, 4.48. Found: C, 61.62; H, 4.44.

5-[3-(4-Bromophenyl)allylidene]-2,2-dimethyl-[1,3]dioxane-4,6-dione 3e

This compound was obtained according to general procedure as orange crystals, mp 182 °C, yield 74 %. ¹H nmr (DMSO-d₆): $\delta = 1.77$ (s, 6H, (CH₃)₂C), 7.52 (d, 2H, H_o), 7.58 (dd, 2H, H_m), 7.35 (d, 1H, H-3'), 8.30 (dd, $J_{H1'-H2'} = 12.0$ Hz, $J_{H2'-H3'} = 15.2$ Hz, 1H, H-2'), 8.15 (d, 1H, H-1'). ¹³C NMR (DMSO-d₆): $\delta = 27.0$ (CH₃), 104.6 (C-2), 112.2 (C-5), 124.5 (C-2'), 130.6 (C_o), 132.3 (C_m), 125.0 (C_p), 134.0 (C_i), 152.3 (C-3'), 156.4 (C-1'), 160.1 (C=O), 162.0 (C=O). The mass spectrum shows the following peaks: ms: (70 eV) m/z (%) = 338/336 (M⁺, 12/13), 336 (13), 281 (11), 280 (44), 279 (12), 278 (39), 252 (15), 250 (14), 224 (14), 222 (13), 208 (16), 206 (16), 155 (16), 131 (52), 127 (81), 126 (35), 115 (19), 101 (20), 77 (27), 75 (20), 51 (16), 50 (14), 43 (100), 39 (10). Anal. Calcd. for C₁₅H₁₃BrO₄: C, 53.43; H, 3.89. Found: C, 53.50; H, 3.86.

5-[3-(4-Nitrophenyl)allylidene]-2,2-dimethyl-[1,3]dioxane-4,6-dione 3f

This compound was obtained according to general procedure as orange crystals, mp 196 °C, yield 77 %. ¹H nmr (DMSO-d₆): $\delta = 1.79$ (s, 6H, (CH₃)₂C), 7.82 (d, 2H, H_o), 8.29 (dd, 2H, H_m), 7.43 (d, 1H, H-3'), 8.42 (dd, $J_{\text{H1'-H2'}} = 11.9$ Hz, $J_{\text{H2'-H3'}} = 15.5$ Hz, 1H, H-2'), 8.17 (d, 1H, H-1'). ¹³C NMR (DMSO-d₆): $\delta = 27.1$ (CH₃), 104.9 (C-2), 114.1 (C-5), 127.3 (C-2'), 124.3 (C_o), 129.6 (C_m), 140.9 (C_p), 148.3 (C_i), 149.9 (C-3'), 155.2 (C-1'), 160.0 (C=O), 161.8 (C=O). The mass spectrum shows the following peaks: ms: (70 eV) m/z (%) = 303 (M⁺, 10), 246 (28), 245 (100), 217 (32), 189 (43), 161 (14), 160 (58), 154 (25), 131 (15), 130 (18), 127 (47), 126 (35), 115 (29), 101 (71), 77 (32), 75 (20), 63 (14), 59 (14), 51 (18), 43 (92), 39 (10).

Anal. Calcd. for C₁₅H₁₃NO₆: C, 59.41; H, 4.32; N, 4.62. Found: C, 59.35; H, 4.23; N, 4.54.

Acknowledgements

Authors thank The Colombian Institute for Science and Research (COLCIENCIAS), UNIVERSIDAD DEL VALLE and "Ministerio de Educación, Cultura y Deportes (Programa de Cooperación con Iberoamérica, AECI)" of Spain for financial support.

References

- 1. (a) J. Prousek, Collect. Czech. Chem. Commun. 1991, 56, 1361; (b) P. Schuster, O. E. Polansky and F. Wessely, Monatsh. Chem. 1964, 95, 53.
- 2. (a) F. X. Smith and G. G. Evans, J. Heterocyclic Chem. 1976, 13, 1025; (b) J. R. Wilt, G. A. Reynolds and J. A. Van Allan, Tetrahedron 1973, 29, 795.
- (a) For a revision on the synthetic use of Meldrum's acid see: Chen, B.-C. *Heterocycles* 1991, 32, 529; (b) Tietze, L.; Beifuss, U. In Comprehensive Organic Synthesis; Trost, B. M.; Fleming, I.; Heathcock, C. H., Eds: Pergamon Press: Oxford, 1991; vol. 2, pp. 341.
- 4. Margaretha, P. Angew. Chem. 1972, 84, 308; (c) Bigi, F, Carloni, S, Ferrari, L, Maggi, R, Mazzacani A, Sartori, G. Tetrahedron Letters, 2001, 32, 5203.

- (a) A. Morales, E. Ochoa, M. Suárez, Y. Verdecia, L. González, N. Martín, M. Quinteiro, C. Seoane and J. L. Soto, J. Heterocyclic Chem. 1996, 33, 103; (b) J. Quiroga, A. Hormaza, B. Insuasty, M. Nogueras, A. Sánchez, N. Hanold and H. Meier, J. Heterocyclic Chem. 1997, 34, 521; (c) J. Quiroga, A. Hormaza, B. Insuasty and M. Márquez, J. Heterocyclic Chem. 1998, 35, 409; (d) J. Quiroga, B. Insuasty, A. Hormaza, P. Cabildo, R. M. Claramunt and J. Elguero, Heterocyclic Commun. 1999, 5 (2), 115; (e) R. Rodríguez, M. Suárez, E. Ochoa, N. Martín, M. Quinteiro, C. Seoane and J. L. Soto, J. Heterocyclic Chem. 1996, 33, 45; (f) R. Rodríguez, M. Suárez, E. Ochoa, B. Pita, R. Espinosa, N. Martín, M. Quinteiro, C. Seoane and J. L. Soto, J. Heterocyclic Chem. 1997, 34, 957.
- J. N. Low, J. Cobo, M. Nogueras, A. Sanchez, B. Insuasty and H. Torres, Acta Cryst. 2002, C58, 39.
- 7. Novoa de Armas, H.; Blaton, N. M.; Peeters, O. M.; De Ranter, C. J.; Suarez, M.; Ochoa, E.; Verdecia, Y.; Salfran, E. J. Chem. Cryst., 2000, 30, 189.

Received on Jun 20, 2004