# AGRICULTURAL AND FOOD CHEMISTRY

ARTICLE

pubs.acs.org/JAFC

# Antioxidative Caffeoylquinic Acids and Flavonoids from *Hemerocallis fulva* Flowers

Yun-Lian Lin,\* Chung-Kuang Lu, Yeh-Jeng Huang, and Hong-Jhang Chen

National Research Institute of Chinese Medicine, Taipei 112, Taiwan

**ABSTRACT:** Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ )-induced reactive oxygen species (ROS) production in HepG2 was used to screen hepatocyte protective compounds from the flowers of *Hemerocallis fulva*. Three new polyphenols, *n*-butyl 4-*trans*-O-caffeoylquinate (1), kaempferol 3-O-{ $\alpha$ -L-rhamnopyranosyl(1→6)[ $\alpha$ -L-rhamnopyranosyl(1→2)]}- $\beta$ -D-galactopyranoside (2), and chrysoeriol 7-O-[ $\beta$ -D-glucuronopyranosyl(1→2)(2-O-*trans*-feruloyl)- $\beta$ -D-glucuronopyranoside (3), together with four caffeoylquinic acid derivatives (4–7), eight known flavones (8–15), one naphthalene glycoside, stelladerol (16), one tryptophan derivative (17), adenosine (18), and guanosine (19) were isolated from the bioactive fractions of the aqueous ethanol extract of *H. fulva* flowers. The structures of isolated compounds were characterized by means of spectroscopic data. Compounds 1–3 were described as first isolated natural products. Among the above-mentioned compounds, the caffeoylquinic acid drivatives are the major components with potent free radical scavenging activity in HepG2 cells and are for the first time isolated from *H. fulva* flowers. A convenient ultraperformance liquid chromatography (UPLC) method was also developed to simultaneously separate and identify caffeoyl-quinic acids and flavonoids promptly.

KEYWORDS: Hemerocallis fulva, flowers, caffeoylquinic acids, flavonoids, TNF-α, free radical scavenger, UPLC

## INTRODUCTION

Excess production of reactive oxygen species (ROS) and/or defective cellular antioxidant systems have been implicated in many pathological conditions including chronic liver injury and fibrogenesis.<sup>1–3</sup> Many inflammatory cytokines were expressed through the production of ROS during chronic liver damage.<sup>3</sup> Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), a pro-inflammatory cytokine, plays a central role in hepatocyte cell death during liver injury.<sup>4</sup> Multiple studies have shown that TNF- $\alpha$  signaling is associated with enhanced generation of ROS, which significantly contributes to TNF- $\alpha$ -induced cell death.<sup>5,6</sup> Phenols such as lignans and flavonoids derived from fruits and vegetables have been regarded as hepatoprotective agents.<sup>7–9</sup>

*Hemerocallis fulva* L. (daylily) belongs to the Liliaceae family.<sup>10</sup> Both fresh and dried flowers of the plant have been commonly used as a vegetable in eastern Asia. Different kinds of compounds including lactams,<sup>11,12</sup> carotenoids,<sup>13</sup> steroidal saponins,<sup>14</sup> anthraquinones,<sup>15,16</sup> and flavones<sup>17</sup> have been isolated from the aerial parts or roots or flowers of *Hemerocallis* spp. As a part of our interest in liver protective constituents in edible flowers, further investigation on the chemical and biological studies of *H. fulva* flowers was conducted. This paper reports on the isolation and structural elucidation of three new compounds from *H. fulva* flowers and the free radical scavenging activity of the isolates in HepG2 cells. Meanwhile, a convenient ultraperformace liquid chromatography (UPLC) method was developed to simultaneously separate and identify caffeoylquinic acids and flavonoids.

#### MATERIALS AND METHODS

**General Experimental Procedures.** Infrared (IR) spectra were recorded on a Nicolet Avatar 320 FT-IR spectrophotometer (Thermo Electron, Akron, OH). Optical rotation was measured on a Jasco P-2000

polarimeter (Hachioji, Tokyo, Japan). Ultraviolet (UV) spectra were measured on a Hitachi U-3310 spectrophotometer (Hitachi, Tokyo, Japan). Nuclear magnetic resonance (NMR) spectra were run in CD<sub>3</sub>OD or DMSO-*d*<sub>6</sub> on a Varian Unity INOVA-500 or VNMRS 600 (Varian, Palo Alto, CA). ESIMS and HRESIMS mass spectra were recorded on a Finnigan MAT LCQ and Finnigan Mat 95S, respectively. UPLC was performed on a Waters AcQuity Ultra Performance LC (Milford, MA). Silica gel (230-400 mesh) and a semipreparative Si column (LiChrosorb Si-60) (Merck, Darmstadt, Germany) and Sephadex LH-20 (Amersham Biosciences, Uppsala, Sweden) were used for column chromatography. Solvents (analytical grade) were purchased from Merck. Dulbecco's modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were from Gibco BRL (Grand Isand, NY). Dimethyl sulfoxide (DMSO) and analytical grade solvents were from Merck. Trypan blue, 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT), and TNF- $\alpha$  were purchased from Sigma (St. Louis, MO). CM-H<sub>2</sub>DCF-DA was from Invitrogen (Carlsbad, CA). All other chemicals were of analytical grade and purchased from commercial suppliers.

**Determination of Total Phenolic Content.** The total phenolic content was determined according to the Folin–Ciocalteu method.<sup>18</sup> Briefly, 200  $\mu$ L of each sample (2000  $\mu$ g/mL) was separately mixed with 2 mL of distilled water, 1 mL of Folin–Ciocalteu reagent, and 5 mL of 20% sodium carbonate. After incubation in the dark for 20 min at room temperature, the absorbance was measured at 735 nm with a microplate reader (Molecular Devices, Sunnyvale, CA). The amount of Folin–Ciocalteu reagent was substituted by the same amount of distilled water in the blank. Gallic acid (0, 200, 400, 600, 800, and 1000  $\mu$ g/mL) was used to make a calibration curve. The total phenolic content was expressed as milligram gallic acid equivalents (GAE) per gram of extract.

Received:	March 22, 2011
Revised:	May 31, 2011
Accepted:	July 15, 2011
Published:	July 15, 2011

**Determination of Total Flavonoid Content.** The AlCl<sub>3</sub> method was used to determine the total flavonoid content.<sup>18</sup> Briefly, 0.5 mL of each sample (2000  $\mu$ g/mL) was separately mixed with 0.1 mL of 10% aluminum chloride, 0.1 mL of 1 M potassium acetate, and 2.8 mL of distilled water. After incubation at room temperature for 30 min, the absorbance was measured at 415 nm with a microplate reader (Molecular Devices). The amount of 10% aluminum chloride was substituted by the same amount of distilled water in the blank. Catechin (0, 100, 200, 400, 600, and 800  $\mu$ g/mL) was used to make a calibration curve. The total flavonoid content was expressed as milligram catechin equivalents (CE) per gram of extract.

**Evaluation of Reactive Oxygen Species in HepG2 Cells.** ROS in HepG2 cells were measured by the conversion of nonfluorescent 2',7'-dichlorofluorescein diacetate (DCF-DA) into 2',7' dichlorofluorescein (DCF). PBS-rinsed cells were treated with TNF- $\alpha$  (200 ng/mL) with or without extract or isolated compound for 2 h and then incubated with 8  $\mu$ M CM-H<sub>2</sub>DCFDA for 30 min at 37 °C. After PBS washing, fluorometric analysis was performed at the indicated time points with the excitation and emission wavelengths at 485 and 530 nm, respectively, in a FLEX Station (Molecular Devices Corp.).<sup>19</sup>

**Cell Viability Assay.** HepG2 cells were cultured in DMEM medium (containing 10% FBS, pH 7.2–7.4). During 24 h of incubation, cells in serum-free medium were exposed to TNF- $\alpha$  in the absence or presence of extract or isolated pure compounds at the indicated concentrations for 6 h by MTT assay. The cell viability of various test groups was determined by the following equation: (absorbance of the test group/absorbance of the control)  $\times$  100%. All test samples mentioned above were dissolved in DMSO. The final concentration of DMSO was <0.1% and the group of 0.1% DMSO was used as the control group.<sup>20</sup>

Isolation and Purification Process. The air-dried flowers of H. fulva (10 kg) were purchased from Taitung, Taiwan, in 2009. The dried flowers (10 kg) were soaked with aqueous ethanol (80% EtOH) at 60 °C overnight (3  $\times$  80 L). The combined extracts were evaporated under reduced pressure in a rotary vacuum evaporator. The dry extract (HF, 952 g) was sucessively partitioned with ethyl acetate (EtOAc) and nbutanol (BuOH) to give EtOAc- (HF-EtOAc, 353 g), BuOH- (HF-BuOH, 136 g), and water-soluble fractions (HF-H<sub>2</sub>O, 450 g). The EtOAc-soluble fraction was then subjected to silica gel (230-400 mesh) column chromatography with successive elution by a Hex/EtOA/ MeOH gradient solvent system to obtain six fractions: A (57 g, Hex), B (112 g, 25% EtOAc/Hex), C (79 g, 50% EtOAc/Hex), D (27 g, 75% EtOAc/Hex), E (24 g, EtOAc), and F (48 g, MeOH). Fractions D and E were separated on a Sephadex LH-20 column (MeOH) repeatedly to give 1 (38 mg), 4 (0.95 g), 8 (32 mg), and 9 (35 mg). The H<sub>2</sub>O-soluble fraction was chromatogramed over a Diaion HP-20 column and eluted with a gradient of MeOH in H2O. Fraction 75% MeOH/H2O eluate was further purified on a Sephadex LH-20 column (80% MeOH/H<sub>2</sub>O) to yield 18 (252 mg) and 19 (123 mg). The MeOH eluates of the H<sub>2</sub>Osoluble fraction (18 g) and HF-EtOAc-F (52 g) were combined with the BuOH-soluble fraction and further separated on Sephadex LH-20 (MeOH) and RP-18 (preparative and semipreparative) columns repeatedly to give 2 (127 mg), 3 (36 mg), 5 (1.79 g), 6 (64 mg), 7 (85 mg), 10 (78 mg), 11 (141 mg), 12 (108 mg), 13 (43 mg), 14 (34 mg), 15 (18 mg), 16 (25 mg), and 17 (34 mg).

*n-Butyl* 4-*trans-O-caffeoylquinate* (**1**): colorless amorphous powder;  $[\alpha]^{25}_{D} = -8.08^{\circ}$  (MeOH, *c* 0.15); IR (KBr)  $\nu_{max}$  (cm<sup>-1</sup>), 3406, 1718, 1686, 1627, 1508, 1441, 1069, 1049, 934; UV (MeOH)  $\lambda_{max}$  nm (log  $\varepsilon$ ), 220 (3.82), 233 (3.78), 243 (3.78), 295 (3.89), 329 (3.98); <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD),  $\delta$  0.93 (3H, *t*, *J* = 7.2 Hz, H-4'), 1.41 (2H, hepta, *J* = 7.2 Hz, H-3'), 1.64 (2H, hepta, *J* = 7.2 Hz, H-2'), 2.01 (1H, dd, *J* = 13.2, 10.2 Hz, H-6a), 2.06 (1H, m, H-2a), 2.16-2.20 (2H, m, H-2b, -6b), 4.13 (2H, hepta, *J* = 7.2 Hz, H-1'), 4.28 (1H, ddd, *J* = 10.2, 9.6, 4.8 Hz,  $w_{1/2} = 20.0$  Hz, H-3), 4.31 (1H, m,  $J_{w1/2} = 7.6$  Hz, H-5), 4.83 (1H, dd, *J* = 9.6, 2.4 Hz, H-4), 6.35 and 7.62 (1H each, d, *J* = 16.0 Hz, H- $\alpha$ , H- $\beta$ ), 6.78 (1H, d, *J* = 7.8 Hz, H-S"), 6.94 (1H, dd, *J* = 7.8, 2.0 Hz, H-6"), 7.06 (1H, d, *J* = 2.0 Hz, H-2"); <sup>13</sup>C NMR (125 MHz, CD<sub>3</sub>OD),  $\delta$  14.0 (q, C-4'), 20.0 (t, C-3'), 31.6 (t, C-2'), 38.3 (t, C-6), 42.3 (t, C-2), 65.6 (d, C-3), 66.4 (t, C-1'), 69.2 (d, C-5), 76.5 (s, C-1), 78.8 (d, C-4), 115.2 (d, C- $\alpha$ ), 115.2 (d, C-2"), 116.5 (d, C-5"), 123.0 (d, C-6"), 127.8 (s, C-1"), 146.7 (s, C-3"), 147.1 (d, C- $\beta$ ), 149.4 (d, C-4"), 169.0 (s, C-9"), 175.3 (s, C-7) Key HMBC correlations: H-1', H-2, and H-6/C-7; H-4, H- $\alpha$ , and H- $\beta$ /C-9". ESIMS, *m*/*z* 409 [M – H]<sup>-</sup>. HRESIMS, 409.1536 [M + Na]<sup>+</sup>(calcd 409.1499 for C<sub>20</sub>H<sub>25</sub>O<sub>9</sub>).

Kaempferol 3-O-{ $\alpha$ -L-rhamnopyranosyl(1 $\rightarrow$ 6)-[ $\alpha$ -L-rhamnopyranosyl- $(1\rightarrow 2)$ ]- $\beta$ -D-galactopyranoside (**2**): yellow amorphous; IR (KBr)  $\nu_{max}$  $(\text{cm}^{-1})$ , 3410, 1667, 1608, 1505, 1069; UV (MeOH)  $\lambda_{\text{max}}$  nm  $(\log \varepsilon)$ , 265 (3.96), 348 (3.89); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  0.79 and 0.94 (3H each, d, J = 6.0 Hz, rhamnosyl H-6<sup>''''</sup> and 6<sup>'''</sup>), 4.30 and 5.04 (1H each, br s, anomeric H-1<sup>'''</sup> and -1<sup>'''</sup>), 5.47 (1H, d, J = 7.5 Hz, anomeric H-1"), 6.18 and 6.38 (1H each, d, J = 2.0 Hz, H-6 and -8), 6.86 and 7.93 (2H each, d, J = 8.5 Hz, H-(3'(5')) and -2'(6')), 10.10, 10.80, and 12.63 (1H each, br s, OH-4', OH-7, OH-5); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>), δ 17.3 (q, C-6<sup>''''</sup>), 17.7 (q, C-6<sup>'''</sup>), 66.9 (t, galactopyranosyl C-6"), 68.2 (d, rhamnopyranosyl, C-5""), 68.3 (d, rhamnopyranosyl, C-5""), 70.3 (d, rhamnopyranosyl C-2"(2"")), 70.5 (d, rhamnopyranosyl C-3""), 70.6 (d, rhamnopyranosyl, C-3""), 70.6 (d, galactopyranosyl C-4"), 71.8 (d, rhamnopyranosy C-4"" (4"")), 75.7 (d, galactopyranosyl C-3"), 77.1 (d, galactopyranosyl C-2"), 77.2 (d, galactopyranosyl C-5"), 93.5 (d, C-8), 98.6 (d, C-6), 98.7 (d, galactopyranosyl C-1"), 100.6 (d, rhamnosyl, C-1""), 100.8 (d, rhamnosyl, C-1""), 104 (s, C-10), 115.0 (d, C-3'(5')), 120.9 (s, C-1'), 130.7 (d, C-2'(6')), 132.6 (s, C-3), 156.4 (s, C-9), 156.9 (s, C-2), 159.8 (C-4'), 161.2 (s, C-5), 164.0 (s, C-7), 177.2 (s, C-4). Key HMBC correlations: H-6, H-8, 7-OH (δ 10.80)/C-7; 5-OH (δ 12.63)/C-5, C-6, C-10; 4'-OH (δ 10.10), H-2'(6'), H-3'(5')/C-4'; H-1"/C-3, C-2", C-3"; H-1"'/C-2", C-2<sup>'''</sup>, C-3<sup>'''</sup>, C-5<sup>'''</sup>; H-1<sup>''''</sup>/C-6<sup>''</sup>, C-2<sup>''''</sup>, C-3<sup>''''</sup>, C-5<sup>''''</sup>; H-6<sup>'''</sup>/C-4<sup>'''</sup>, C-5<sup>'''</sup>; H-6<sup>''''</sup>/C-4<sup>''''</sup>, C-5<sup>''''</sup>. ESIMS: m/z 739 [M – H]<sup>-</sup>.

Chrysoeriol 7-O-[(2-O-trans-feruloylglucuronopyranosyl)  $(1 \rightarrow 2)$ ]- $O-\beta$ -D-qlucuronopyranoside (3): yellow amorphous powder; IR (KBr)  $v_{\text{max}}$  (cm<sup>-1</sup>), 3358, 1716, 1657, 1605, 1593, 1514, 1262, 1187, 1159, 1088, 1056; UV (MeOH)  $\lambda_{max}$  nm (log  $\varepsilon$ ), 248 (3.95), 270 (3.88), 334 (4.00); (+NaOAc)  $\lambda_{max}$  nm (log  $\varepsilon$ ), 249 (3.95), 270 (3.88), 335 (4.00); (+AlCl<sub>3</sub>)  $\lambda_{\text{max}}$  nm (log  $\varepsilon$ ), 277 (3.95), 296 (3.88), 338 (3.97), 387 (3.85); (+AlCl<sub>3</sub>+HCl)  $\lambda_{max}$  nm (log  $\varepsilon$ ), 278 (3.95), 296 (3.88), 337 (3.98), 387 (3.85); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  3.54 (1H, m, H-2"), 4.63 (1H, t, J = 8.5 Hz, H-2""), 4.90 (1H, d, J = 8.5 Hz, anomeric H-1<sup>'''</sup>), 5.37 (1H, d, J = 7.5 Hz, anomeric H-1<sup>''</sup>), 6.39 and 7.49 (1H each, d, J = 15.5 Hz, feruloyl H- $\alpha$  and H- $\beta$ ), 6.41 (1H, d, J = 2.0 Hz, H-6), 6.76 (1H, d, J = 8.0 Hz, feruloyl H-5), 6.78 (1H, d, J = 2.0 Hz, H-8), 6.93 (1H, d, J = 8.5 Hz, H-5'), 6.94 (1H, s, H-3), 7.06 (1H, dd, J = 8.0, 2.0 Hz, feruloyl H-6), 7.24 (1H, d, J = 2.0 Hz, feruloyl H-2), 7.56 (1H, dd, J = 8.5, 2.0 Hz, H-6'), 7.57 (1H, d, J = 2.0 Hz, H-2'), 9.55 (1H, br s, feruloyl 4-OH), 9.99 (1H, br s, 4'-OH), 12.96 (1H, br s, 5-OH); <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>), δ 55.7 (q, OCH<sub>3</sub>), 56.0 (q, OCH<sub>3</sub>), 71.4 (d, glucuronopyranosyl-C-4"), 71.9 (d, glucuronopyranosyl-C-4"'), 73.4 (d, glucuronopyranosyl-C-2"), 73.9 (d, glucuronopyranosyl-C-3"), 74.9 (d, glucuronopyranosyl-C-5"), 75.0 (d, glucuronopyranosyl-C-3""), 75.6 (d, glucuronopyranosyl-C-5"), 79.9 (d, glucuronopyranosyl-C-2""), 94.9 (d, C-8), 97.2 (glucuronopyranosyl-C-1"), 99.2 (d, C-6), 100.7 (d, glucuronopyranosyl-C-1""), 103.5 (s, C-3), 105.5 (s, C-10), 110.4 (d, C-2'). 111.2 (d, feruloyl-C-2), 115.0 (d, feruloyl-C-α), 115.5 (d, feruloyl-C-5), 115.8 (d, C-5'), 120.5 (d, C-6'), 121.5 (s, feruloyl C-1), 122.9 (d, feruloyl C-6), 125.8 (s, C-1'), 144.6 (d, feruloyl C- $\beta$ ), 147.9 (s, C-3'), 148.1 (s, feruloyl C-3'), 149.2 (s, feruloyl C-4), 150.9 (s, C-4'), 156.9 (s, C-9), 161.2 (s, C-5), 162.2 (s, C-7), 164.2 (s, C-2), 165.8 (s, feruloyl C=O), 169.9 (s, glucuronopyranosyl C-6"), 170.1 (s, glucuronopyranosyl C-6""), 182.1 (s, C-4). Key HMBC correlations:

 Table 1. Phenol and Flavonoid Contents of the Aqueous

 Ethanol Extract and the Subtractions of H. fulva

extract of <i>H. fulva</i> flowers	total phenols (mg gallic acid equiv/g extracts)	total flavonoids (mg catechin equiv/g extracts)
HF	$30.9\pm0.7$	$15.1\pm0.9$
HF-EA	$25.3\pm2.7$	$10.4\pm2.3$
HF-BuOH	$34.6\pm2.7$	$19.5\pm1.3$
HF-H <sub>2</sub> O	$28.8\pm0.7$	$14.6\pm0.5$

H-3/C-2, C-4; 5-OH/C-5, C-6 and C-10; H-8/C-6, C-7, C-9 and C-10; H-2<sup>'''</sup>/feruloyl C=O and C-1<sup>'''</sup>; H-2<sup>''</sup>/C-1<sup>''</sup>, C-1<sup>'''</sup>, C-3<sup>''</sup>, C-4<sup>''</sup>; H-1<sup>''</sup>/C-7; H-1<sup>'''</sup>/C-2<sup>'''</sup>. Key NOE correlations: OCH<sub>3</sub> ( $\delta_{\rm H}$  3.73)/ feruloyl H-2; OCH<sub>3</sub> ( $\delta_{\rm H}$  3.79)/H-2<sup>'</sup>. ESIMS: *m/z* 827 [M – H]<sup>-</sup>, 651 [M – H – 175]<sup>-</sup>, 527 [M – H – 300]<sup>-</sup>, 299. HRESIMS: 827.1831[M – H]<sup>+</sup>(calcd 827.1670 for C<sub>38</sub>H<sub>35</sub>O<sub>21</sub>).

*Hydrolysis* of **2** and **3**: Compound **2** (10 mg) was dissolved in 5% HCl/ aqueous ethanol (2 mL) at 80 °C for 30 min. After neutralization with NaHCO<sub>3</sub>, the mixture was evaporated. The residue was resuspended in H<sub>2</sub>O and then filtered to yield kaempferol (3 mg). The sugar components in the filtrate were identified by TLC (on Si gel, developed with ethyl acetate/butanone/formic acid/H<sub>2</sub>O = 5:3:1:1) as D-galactose ( $R_f$  0.20) and L- rhamnose ( $R_f$  0.55) in comparison with authentic samples. Compound **3** (5 mg) was conducted by basic hydrolysis (5% NaOH in 50% aqueous ethanol at 80 °C) for 1 h to give compound **15** and ferulic acid.

UPLC Analysis of Isolated Caffeoylquinic Acids and Flavonoids. UPLC analyses were carried out on a Waters AcQuity Ultra Performance LC, which was equipped with a UPLC column (ACQUITY BEH C18, 1.7  $\mu$ m, 2.1 × 100 mm) and a guard column (VanGuard BEH C18, 1.7  $\mu$ m, 2.1 × 5 mm) at a constant flow rate of 0.4 mL/min. Mobile phases A (acetonitrile) and B (0.1% formic acid) were run on a programmed protocol: 0–3 min, 2–18% A; 3–5 min, 18–18% A; 5–7 min, 18–40% A; 7–9 min, 40–70% A. The DAD was set at a wavelength between 200 and 400 nm. The chromatographs were plotted at an absorbance of 254 nm. An aliquot of sample (1  $\mu$ L) was directly introduced into the column through the sample manager.

**Statistical Analysis.** Data are expressed as the mean  $\pm$  SEM. Oneway analysis of variance (ANOVA) was used. The statistical significance between control and compound-treated group was evaluated by Student's *t* test. A *p* < 0.05 was considered to be statistically significant.

### RESULTS AND DISCUSSION

Phenol and Flavonoid Compositions. Phenolic compounds are ubiquitously distributed in the plant kingdom. Phenolic compounds such as caffeoylquinic acid derivatives and flavonoids are the most abundant polyphenols in human diet and have been shown to possess different kinds of bioactivities.<sup>21-24</sup> Levels of phenolic contents and flavonoids were shown to exhibit antioxidant capacity and redox properties, such as reducing agents, hydrogen donors, and reactive oxidative species scavengers.<sup>25</sup> As Table 1 shows, the total phenolic contents of HF, HF-EA, HF-BuOH, and H<sub>2</sub>O were  $30.9 \pm 0.7$ ,  $25.3 \pm 2.7$ ,  $34.6 \pm 2.7$ , and  $28.8 \pm 0.7$  mg gallic acid equiv (GAE)/g of extract, respectively. The flavonoid contents of three subfractions were 10.4  $\pm$  2.3, 19.5  $\pm$  1.3, and 14.6  $\pm$  0.5 mg catechin equiv (CE)/g of extracts), respectively. Only the HF-BuOH subfraction has a flavnoid content higher than that in crude extract  $(15.1 \pm 0.9 \text{ mg})$ CE/g of extracts). Total flavonoid/total phenolic contents in the crude extract and three subfractions were 0.49 (HF), 0.41 (HF-

EtOAc), 0.56 (HF-BuOH) and 0.51 (HF- $H_2O$ ), respectively. The results suggest that both flavonoids and phenols play major roles in ROS scavenging in the active fractions.

**Compound Isolation and Structural Identification.** Nineteen compounds including three new compounds (1-3), together with four known caffeoylquinic acid derivatives, chlorogenic acid methyl ester (4),<sup>26,27</sup> chlorogenic acid (5),<sup>26</sup> 4-caffeoylquinic acid (6),<sup>28</sup> and 5-caffeoylquinic acid (7),<sup>28</sup> eight known flavones, including kaempferol (8),<sup>26</sup> quercetin (9),<sup>26</sup> astragalin (10),<sup>26</sup> isoquercitrin (11),<sup>26</sup> kaempferol 3-O-rutinoside (12),<sup>29</sup> rutin (13),<sup>29,30</sup> quercetin 3-O-{ $\alpha$ -L-rhamnopyranosyl- $(1\rightarrow 6)[\alpha$ -L-rhamnopyranosyl $(1\rightarrow 2)]$ - $\beta$ -D-galactopyranoside (14),<sup>31</sup> chrysoeriol 7-O- $\beta$ -D-glucuronopyranosyl $(1\rightarrow 2)$ -O- $\beta$ -Dglucuronopyranoside (15),<sup>28</sup> a naphthalene glycoside, stelladerol (16),<sup>16</sup> a tryptophan derivative, lycoperodine-1 (17),<sup>32</sup> adenosine (18),<sup>26</sup> and guanosine (19) were isolated from high phenolic subfractions of the aqueous ethanol extract of *H. fulva* flowers (Figure 1). The structures of known compounds were elucidated by spectral methods and their spectroscopic data compared with those of an authentic sample or with published data. Compound 4 is probably an artifact during the isolation.

Compound 1 was obtained as a colorless amorphous powder,  $[\alpha]^{25}_{D} = -8.08^{\circ}$  (MeOH, c 0.15), and gave a molecular ion peak at  $m/z 409 [M - H]^{-}$  in agreement with a molecular formula of C<sub>20</sub>H<sub>25</sub>O<sub>9</sub> by HRESIMS and DEPT NMR. The IR spectrum showed absorption bands for hydroxyl (3406 cm<sup>-1</sup>), ester (1718 cm<sup>-1</sup>), conjugated carbonyl (1686 cm<sup>-1</sup>), and an aro-matic group (1627 and 1508 cm<sup>-1</sup>). <sup>1</sup>H and <sup>13</sup> C NMR indicated signals for a butyroyl [ $\delta_{\rm H}$  0.93 (3H, t, *J* = 7.2 Hz), 1.41 and 1.64 and 4.13 (2H each, hepta, J = 7.2 Hz);  $\delta_{\rm C}$  14.0 (q), 20.0 (t), 31.6 (t), and 66.4 (t)], two methylenes [ $\delta_{\rm H}$  2.01 (1H, dd, J = 13.2, 10.2 Hz), 2.06 (1H, m), and 2.16–2.20 (2H, m);  $\delta_{\rm C}$  38.3 (t, C-6) and 42.3 (t, C-2)], three oxymethines [ $\delta_{\rm H}$  4.28 (1H, ddd, J = 10.2, 9.6, 4.8 Hz), 4.31 (1H, m), 4.83 (1H, dd, J = 9.6, 2.4 Hz);  $\delta_{\rm C}$ 65.6 (d, C-3), 69.2 (d, C-5), and 78.8 (d, C-4)], one transcaffeoyl [ $\delta_{\rm H}$  6.35 and 7.62 (1H each, d, J = 16.0 Hz,), 6.78 (1H, d, J = 7.8 Hz), 6.94 (1H, dd, J = 7.8, 2.0 Hz), and 7.06 (1H, d, J = 2.0Hz);  $\delta_{\rm C}$  115.2 (d), 115.2 (d), 116.5 (d), 123.0 (d), 127.8 (d), 146.7 (s), 147.1 (d), 149.4 (s), and 169.0 (s)], a quaternary oxygenated carbon [ $\delta_{\rm C}$  76.5 (s)], and a carbonyl [ $\delta_{\rm C}$  175.3 (s)]. From the data mentioned above, compound 1 was suggested as the *n*-butyl ester of 4-caffeoylquinic acid. The coupling constants of H<sub>3</sub> and H<sub>4</sub> (J = 9.6 Hz) with a large half-width value ( $J_{w1/2}$  = 20.0 Hz), and a small one between  $H_4$  and  $H_5$  (J = 2.4 Hz) in the quinate moiety indicated they had trans-diaxial protons and axial-equatorial gauche configuration, respectively.<sup>33</sup> The acylation of hydroxyl groups produced the downfield shifts of H-1'  $[\delta_{\rm H} 4.13 \ (2\text{H}, \text{hepta}, J = 7.2 \text{ Hz})]$  and H-4  $[\delta_{\rm H} 4.83 \ (1\text{H}, \text{dd}, J =$ 9.6, 2.4 Hz)], which displayed the presence of acyloxy groups at both positions. This assignment was further confirmed by HMBC correlations from H-1', H-2, and H-6 to C-7 and from H-4, H- $\alpha$ , and H- $\beta$  to C-9".

Compound 2 showed the *quasi* molecular ion peak at m/z 739  $[M - H]^-$ . <sup>1</sup>H and <sup>13</sup>C NMR signals were assigned by 1D-TOCSY, <sup>1</sup>H-<sup>1</sup>H-COSY, HMQC, and HMBC NMR experiments and suggested kaempferol, a galactopyranosyl, and two rhamnopranosyl units were included in the skeleton. The upfield shift (about 1.8 ppm) of C-3 ( $\delta_{\rm C}$  132.6) and a lower field shift (about 9.7 ppm) of C-2 ( $\delta_{\rm C}$  156.9) in the kaempferol moiety (compared to kaempferol (8)) indicated *O*-glycosidation at C-3.<sup>21</sup> The coupling constants of the anomeric proton signals at  $\delta_{\rm H}$  5.47 (1H, d, *J* = 7.5 Hz), 4.30, and 5.04 (1H each, br s), and



Figure 1. Structures of the isolated compounds.

two methyl group at  $\delta_{\rm H}$  0.79 and 0.94 (3H each, d, *J* = 6.0 Hz) revealed **2** with one β-D-galactosyl and two α-L-rhamnosyl units. Acidic hydrolysis of **2** yielded kaempferol, D-galactose, and Lrhamnose. The downfield shift of C-2 ( $\delta_{\rm C}$  77.1) and C-6 ( $\delta_{\rm C}$ 66.9) of the galactopyranosyl indicated the interglycosidic linkage at both positions. The assignment was further supported by *J*<sub>2</sub> and *J*<sub>3</sub> correlations between H-1" ( $\delta_{\rm H}$  5.47) and C-3 ( $\delta_{\rm C}$ 132.6), between H-1"'' ( $\delta_{\rm H}$  5.04) and C-2" ( $\delta_{\rm C}$  77.1), and between H-1"''' ( $\delta_{\rm H}$  4.30) and C-6"( $\delta_{\rm C}$  66.9). The <sup>1</sup>H and <sup>13</sup>C NMR spectral data of **2** were similar to those of 14 except for signals assignable to the aglycone moiety in place of kaempferol in **2**. The structure was therefore assigned as kaempferol 3-*O*-{α-L-rhamnopyranosyl(1→6)-[α-L-rhamnopyranosyl(1→2)]}-β-Dgalactopyranoside.

The molecular formula of compound 3 was determined to be  $C_{38}H_{36}O_{21}$  on the basis of <sup>13</sup>C and DEPT spectra and the HRESIMS molecular ion at m/z 827.1831  $[M - H]^+$ . The UV characteristic absorption bands at 232, 270, and 333 nm indicated compound 2 to be a flavone derivative.<sup>34</sup> The presence of a signal at  $\delta_{\rm H}$  12.96 and a bathochromic shift of 30 nm of band II (248 nm shifts to 278 nm) in the presence of AlCl<sub>3</sub>/HCl indicated that C-5 was hydroxylated.<sup>34</sup> Meanwhile, the absence of a 7-hydroxyl group was elucidated by no bathochromic shift on adding shift reagent NaOAc.<sup>34</sup> The <sup>1</sup>H, <sup>13</sup>C, and DEPT spectra revealed the presence of chrysoeriol, two glucuronopyranosyl, and trans-feruloyl moieties. Alkali hydrolysis gave compound 15 and ferulic acid. Longrange correlations between H-2<sup>*III</sup></sup> [\delta\_{\rm H} 4.63 (t, <i>J* = 8.5 Hz)] and carbonyl of feruloyl ( $\delta_{\rm C}$  165.8), between H-1<sup>*III*</sup> [ $\delta_{\rm H}$  4.90 (d, *J* =</sup> 8.5 Hz)] and C-2" ( $\delta_{\rm C}$  79.9), and between H-1" [ $\delta_{\rm H}$  5.37 (d, J = 7.5 Hz) and C-7 ( $\delta_{\rm C}$ 162.2) elucidated their connectivities. The structure was further confirmed by the NOE correlations between methoxyl at  $\delta_{\rm H}$  3.73 and feruloyl H-2 at  $\delta_{\rm H}$  7.24 and between methoxyl at  $\delta_{\rm H}$  3.79 and H-2' at  $\delta_{\rm H}$  7.57 (d, J = 2.0 Hz).



**Figure 2.** Effects of the ethanol extract of *H. fulva* on TNF-α-induced HepG2 cell ROS production. HepG2 cells were seeded at 8 ×10<sup>4</sup> cells/ well in 96 wells overnight. PBS-rinsed cells were treated with TNF-α (10 ng/mL) with or without HF extract for 2 h and then incubated with 8  $\mu$ M of CM-H<sub>2</sub>DCFDA for 30 min at 37 °C. The fluorometric analysis was measured with the excitation and emission wavelengths at 485 and 530 nm. Each column represents the mean ± SEM of three experiments. Resveratrol was used as a positive control. (#) *P* < 0.05, compared with control; (\*) *P* < 0.05, compared with TNF-α-induced group.

Effects of Subfractions and Isolated Polyphenols on ROS scavenging in HepG2 Cells. Flavonoids, phenethyl glycosides,





(A) PBS-rinsed cells were treated with TNF- $\alpha$  (10 ng/mL) with or without test compound for 2 h AND then incubated with 8  $\mu$ M CM-H<sub>2</sub>DCFDA for 30 min at 37 °C. The fluorometric analysis was measured with the excitation and emission wavelengths at 485 and 530 nm. (B) cell viability in TNF- $\alpha$ -induced HepG2 cells was tested by MTT assay. Each column represents the mean  $\pm$  SEM of three experiments. Resveratrol was used as a positive control. (#) P < 0.05, compared with control group; (\*) P < 0.05, compared with TNF- $\alpha$ -induced group.

naphthalene glycosides, and carotenoids have been isolated and claimed as the antioxidative principles of daylily (*Hemerocallis* spp.) flowers<sup>13,16</sup> or leaves.<sup>17</sup> In this study, caffeoylquinate derivatives including *n*-butyl 4-*trans*-O-caffeoylquinate (1, 38 mg), methyl chloriginate (4, 0.95 g), chlorogenic acid (5, 1.79 g), 4-caffeoylquinic acid (6, 64 mg), and 5-caffeoylquinic acid (7, 85 mg) were isolated as the major constituents in *H. fulva* flowers. Chlorogenic acid and the related caffeoylquinic acids from natural resources have been focused on various bioactivities including antioxidative effect.<sup>22–24</sup> As Figure 2 shows, the

crude extract of the flowers of *H. fulva* and three subfractions significantly inhibited TNF- $\alpha$ -induced ROS production in HepG2 cells at the concentration of 50  $\mu$ g/mL and had no direct cytotoxicity (data not shown). The HF-BuOH subfraction preserved the most potent activity and diminished ROS to a level similar to that of the control group. The isolated caffeoylquinic acids and flavonoids (1–17) were also tested for ROS scavenging activity (Figure 3A). Among those compounds, caffeoylquinic acid derivatives 1 and 4–7, flavonoids 3 and 10–14, and anthraquinone 16 significantly inhibited



**Figure 4.** UPLC chromatogram of (A) isolated caffeoylquinic acids and flavonoids and (B) crude aqueous ethanol extract. (1) *n*-Butyl 4-caffeoylquinate; (2) kaempferol 3-O-{α-L-rhamnopyranosyl(1→6)-[α-Lrhamnopyranosyl(1→2)]}-β-D-galactopyranoside; (3) chrysoeriol 7-O-[(2-O-trans-feruloylglucuronopyranosyl) (1→2)]-O-β-D-glucuronopyranoside; (4) methyl chlorogenate; (5) chlorogenic acid; (6) 4-caffeoylquinic acid; (7) 5-caffeoylquinic acid; (8) kaempferol; (9) quercetin; (10) astragalin; (11) isoquercitrin; (12) kaempferol 3-O-rutinoside; (13) rutin; (14) quercetin 3-O-{α-L-rhamnopyranosyl(1→6)[α-Lrhamnopyranosyl(1→2)]}-β-D-galactopyranoside; (15) chrysoeriol 7-Oβ-D-glucuronopyranosyl(1→2)-O-β-D-glucuronopyranoside; (16) stelladerol; (17) lycoperodine-1.

ROS production in HepG2 cells at 50  $\mu$ M without direct toxicity under the working concentration (Figure 3B). From our results, both caffeoylquinic acid derivatives and flavonoids are responsible for the ROS scavenging activity in *H. fulva*.

Identification of Caffeoylquinic Acids and Flavonoids by UPLC. The isolated caffeoylquinic acids and flavonoids were analyzed by a UPLC system. A complete UV chromatogram (254 nm) obtained with the standard solution is shown in Figure 4. Peaks were identified by comparison with the retention time and DAD spectra of the authentic samples (Figure 4). The results indicated that caffeoylquinic acids and flavonoids gave good resolutions except rutin (13) and chrysoeriol 7-O- $\beta$ -D-glucuronopyranosyl(1 $\rightarrow$ 2)-O- $\beta$ -D-glucuronopyranoside (15) overlapping. Our results suggest that UPLC analysis can be considered as a convenient analytical method for caffeoylquinic acids and flavonoids simultaneously and promptly.

This is the first report that *H. fulva* flowers contain high levels of caffeoylquinic acids and possess potent ROS scavenging activity. Our results suggest that caffeoylquinic acids, especially chlorogenic acid in *H. fulva* flowers, at least partially contributed to their ROS scavenging effect. Thus, *H. fulva* flowers may be a source of natural ROS scavenger and beneficial to the health of consumers.

### AUTHOR INFORMATION

#### Corresponding Author

\*Phone: 886-2-2820-1999, ext. 6531. Fax: 886-2-2826-4266. E-mail: yllin@nricm.edu.tw.

### ABBREVIATIONS USED

HF, aqueous ethanolic extract of the flowers of *H. fulva*; EtOAc, ethyl acetate; BuOH, *n*-butanol; HF-EtOAc, ethyl acetate-soluble subfraction of HF; HF-BuOH, *n*-butanol-soluble subfraction of HF; HF-H<sub>2</sub>O, H<sub>2</sub>O-soluble subfraction of HF; Hex, hexane; MeOH, methanol; GAE, gallic acid equivalents; CE, catechin equivalents; IR, infrared spectra; UV, ultraviolet spectra; NMR, nuclear magnetic resonance; ESIMS, electrospray ionization mass spectra; UPLC, ultraperformance liquid chromatography; DMSO, dimethyl sulfoxide; DMEM, Dulbecco's modified Eagle's medium; FBS, fetal bovine serum; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; MTT, 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2*H*-tetrazolium bromide; ROS, reactive oxygen species; DCF-DA, 2',7'-dichlorofluorescein diacetate; ANOVA, one-way analysis of variance.

#### REFERENCES

(1) Droge, W. Free radical in the physiological control of cell function. *Physiol. Rev.* **2002**, *82*, 47–95.

(2) Friedman, S. L. Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. *J. Biol. Chem.* 2000, 275, 2247–2250.

(3) Baroni, G. S.; D'Ambrosio, L.; Ferretti, G.; Casini, A.; Sario, A. D.; Salzano, R.; Ridolfi, F.; Saccomanno, S.; Jezequel, A. M.; Benedetti, A. Fibrogenic effect of oxidative stress on rat hepatic stellate cells. *Hepatology* **1998**, *27*, 720–726.

(4) Wullaert, A.; Heyninck, K.; Beyaert, R. Mechanisms of crosstalk between TNF-induced NF-κB and JNK activation in hepatocytes. *Biochem. Pharmacol.* 2006, *72*, 1090–1101.

(5) Sakon, S.; Xue, X.; Takekawa, M.; Sasazuki, T.; Okazaki, T.; Kojima, Y.; Piao, J. H.; Yagita, H.; Okumura, K.; Doi, T.; Nakano, H. NF- $\kappa$ B inhibits TNF- $\alpha$ -induced accumulation of ROS that mediate prolonged MAPK activation and necrotic cell death. *EMBO J.* **2003**, 22, 3898–3909.

(6) Kamata, H.; Honda, S.; Maeda, S.; Chang, L.; Hirata, H.; Karin, M. Reactive oxygen species promote TNF $\alpha$ -induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* **2005**, *120*, 649–661.

(7) Gebhardt, R. Oxidative stress, plant-derived antioxidants and liver fibrosis. *Planta Med.* **2002**, *68*, 289–296.

(8) Prasad, S.; Phromnoi, K.; Yadav, V. R.; Chaturvedi, M. M.; Aggarwal, B. B. Targeting inflammatory pathways by flavonoids for prevention and treatment of cancer. *Planta Med.* **2010**, *76*, 1044–1063.

(9) Lin, Y. L.; Wu, C. H.; Huang, Y. J.; Luo, M. H.; Wang, C. N.; Shiao, M. D.; Huang, Y. T. In vitro protective effects of salvianolic acid B on primary hepatocytes and hepatic stellate cells. *J. Ethnopharmacol.* **2006**, *105*, 215–222.

(10) Yang, T. I. *A List of Plants in Taiwan;* Natural Publishing: Taipei, Taiwan, 1982; p 353.

(11) Inoue, T.; Iwagoe, K.; Konishi, T.; Kiyosawa, S.; Fujiwara, Y. Novel 2,5-dihydrofuryl-7-lactam derivatives from *Hemerocallis fulva* L. var. *kwanzo* Regel. *Chem. Pharm. Bull.* **1990**, *38*, 3187–3189.

(12) Inoue, T.; Konishi, T.; Kiyosawa, S.; Fujiwara, Y. 2,5-Dihydrofuryl-γ-lactam derivatives from from *Hemerocallis fulva* L. var. *kwanso* Regel. II. *Chem. Pharm. Bull.* **1994**, *42*, 154–155.

(13) Tai, C. Y.; Chen, B. H. Analysis and stability of carotenoids in the flowers of daylily (*Hemerocallis disticha*) as affected by various treatments. *J. Agric. Food Chem.* **2000**, *48*, 5962–5968.

(14) Konishi, T.; Fujiwara, Y.; Konoshima, T.; Kiyosawa, S.; Nishi,
 M.; Miyahara, K. Steroidal saponins from *Hemerocallis fulva* var. *Kwanso. Chem. Pharm. Bull.* 2001, 49, 318–320.

(15) Cichewicz, R. H.; Kim, K. C.; McKerrow, J. H.; Nair, M. G. Kwanzoquinones A–G and other constituents of *Hemerocallis fulva* 'Kwanso' roots and their activity against the human pathogenic trematode *Schistosoma mansoni. Tetrahedron* **2002**, *58*, 8597–8606.

(16) Cichewicz, R. H.; Nair, M. G. Isolation and characterization of stelladerol, a new antioxidant naphthalene glycoside, and other antioxidant glycosides from edible daylily (*Hemerocallis*) flowers. *J. Agric. Food Chem.* **2002**, *50*, 87–91.

(17) Zhang, Y.; Cichewicz, R. H.; Nair, M. G. Lipid peroxidation inhibitory compounds from daylily (*Hemerocallis fulva*) leaves. *Life Sci.* **2004**, *75*, 753–763.

(18) Moyo, M.; Ndhlala, A. R.; Finnie, J. F.; Van Staden, J. Phenolic composition, antioxidant and acetylcholinesterase inhibitory activities of *Sclerocarya birrea* and *Harpephyllum caffrum* (Anacardiaceae) extracts. *Food Chem.* **2010**, *123*, 67–76.

(19) Tsai, M. K.; Lin, Y. L.; Huang, Y. T. Effects of salvianolic acids, A and B on oxidative stress and hepatic fibrosis in rats. *Toxicol. Appl. Pharmacol.* **2010**, *242*, 155–164.

(20) Lin, Y. L.; Wu, C. H.; Luo, M. H.; Huang, Y. J.; Wang, C. N.; Shiao, M. S.; Huang, Y. T. Effect of salvianolic acid B on oxidative damage in primary hepatocytes and hepatic stellate cells. *J. Ethnopharmacol.* **2006**, *105*, 215–222.

(21) Matsuda, H.; Morikawa, T.; Ando, S.; Toguchida, I.; Yoshikawa, M. Structural requirements of flavonoids for nitric oxide production inhibitory activity and mechanism of action. *Bioorg. Med. Chem.* **2003**, *11*, 1995–2000.

(22) David dos Santos, M.; Almeida, M. C.; Lopes, N. P.; Petto de Souza, G. E. Evaluation of the anti-inflammation, analgesic and anti-pyretic activities of the natural polyphenol chlorogenic acid. *Biol. Pharm. Chem.* **2006**, *29*, 2236–2240.

(23) Bouayed, J.; Rammal, H.; Dicko, A.; Younos, C.; Soulimani, R. Chlorogenic acid, a polyphenol from *Prunus domestica* (Mirabelle), with coupled anxiolytic and antioxidant effects. *J. Neurol. Sci.* 2007, 262, 77–84.

(24) Lin, L. C.; Kuo, Y. C.; Chou, C. J. Immunomodulatory principles of *Dichrocephala bicolor*. J. Nat. Prod. **1999**, 62, 405–408.

(25) Robards, K.; Prenzler, P.; Tucker, G.; Swatsitang, P.; Glover, W. Phenolic compounds and their role in oxidative processes in fruits. *Food Chem.* **1999**, *66*, 401–436.

(26) Lin, Y. L.; Wang, W. Y.; Kuo, Y. H.; Chen, C. F. Nonsteroidal constituents from *Solanum incanum*. J. Chin. Soc. **2000**, 47, 247–251.

(27) Lin, L. C.; Kuo, Y. C.; Chou, C. J. Immunomodulatory principles of *Dichrocephala bicolor*. J. Nat. Prod. **1999**, 62, 405–408.

(28) Lin, Y. L.; Wang, W. Y. Chemical constituents of *Biophytum* sensitivum. Chin. Pharm. J. **2003**, 55, 71–75.

(29) Lin, Y. L.; Lee, H. P.; Huang, R. L.; Ou, J. C.; Kuo, Y. H. Studies on the constituents of *Hyptis rhomboides*. *Chin. Pharm. J.* **1993**, 45, 61–68.

(30) Lin, Y. L.; Kuo, Y. H.; Shiao, M. S.; Chen, C. C.; Ou, J. C. Flavonoid glycosides from *Terminalia catappa*. J. Chin. Chem. Soc. **2000**, 47, 253–256.

(31) Yasukawa, K.; Sekine, H.; Takido, M. Two flavonol glycosides from *Lysimachia fortunei*. *Phytochemistry* **1989**, *28*, 2215–2216.

(32) Yahara, S.; Uda, N.; Yoshio, E.; Yae, E. Steroidal alkaloid glycosides from tomato (*Lycopersicon esculentum*). J. Nat. Prod. 2004, 67, 500–502.

(33) Corse, J.; Lundin, R. E.; Sondheimer, E.; Waiss, A. C. Conformation analyses of D-(-)-quinic acid and some of its derivatives by nuclear magnetic resonance. *Phytochemistry* **1966**, *5*, 767–776.

(34) Mabry, T. J.; Markham, K. R.; Thomas, M. B. *The Systematic Identification of Flavonoids*; Springer-Verlag: New York, 1970; pp 41–61.