

SYNTHESIS OF 2,6-DISUBSTITUTED PYRIDINES BY LITHIATION OF PYRAZOLO[1,5-*a*]PYRIDINES

Akira MIYASHITA,*^a Yasushi SATO,^a Sayuri WATANABE,^a Ken-ichi TANJI,^b and Takeo HIGASHINO^a

School of Pharmaceutical Sciences^a and School of Food & Nutritional Sciences,^b
University of Shizuoka, 52-1 Yada, Shizuoka 422, Japan

Lithiation of 3-(4,4-dimethyl-2-oxazolin-2-yl)pyrazolo[1,5-*a*]pyridine (**1a**) with 2 molar equivalents of *n*-BuLi followed by reaction with benzaldehyde yielded α -(4,4-dimethyl-2-oxazolidinylidene)-6-(α -hydroxybenzyl)-2-pyridineacetonitrile (**2a**). Upon similar treatment of other electrophiles, the corresponding 2,6-disubstituted pyridines **2** were produced. The formation of the pyridines proceeded through lithiation, reaction with an electrophile, and ring-cleavage of the pyrazole ring.

KEYWORDS pyrazolo[1,5-*a*]pyridine; lithiation; 2,6-disubstituted pyridine; electrophile; ring cleavage

We have reported that the lithiation of pyrazolo[1,5-*a*]pyridine with *n*-BuLi proceeded selectively at the 7-position, and a subsequent reaction with electrophiles gave 7-substituted derivatives.¹⁾ However, it is difficult to produce 2-substituted pyrazolo[1,5-*a*]pyridines by lithiation. We expected that the introduction of a functional group at the 2-position on a pyrazolo[1,5-*a*]pyridine could be achieved by use of the *ortho*-directing effect of a substituent at the 3-position. Several substituents, such as 2-oxazolin-2-yl, *N,N*-dimethylaminomethyl, and carbamoyl groups, efficiently promote *ortho*-lithiation.²⁾ In order to introduce a functional group at the 2-position, we synthesized 3-(4,4-dimethyl-2-oxazolin-2-yl)pyrazolo[1,5-*a*]pyridine (**1a**).³⁾

Treatment of **1a** with 2 molar equivalents of *n*-BuLi at -78°C for 0.5 h followed by the reaction with benzaldehyde unexpectedly yielded α -(4,4-dimethyl-2-oxazolidinylidene)-6-(α -hydroxybenzyl)-2-pyridineacetonitrile (**2a**).⁴⁾ The structure of **2a** was supported by the following result: the characteristic absorptions at 2186 cm^{-1} (CN), 3178 cm^{-1} , and 3442 cm^{-1} (OH or NH) in the IR spectrum and characteristic signals at 9.62 ppm, 5.77 ppm, and 2.70 ppm due to NH, and CHOH in the ^1H -NMR spectrum were observed. Elemental analysis and MS measurement also supported the proposed structure.⁵⁾




Chart 1

In order to examine the scope and limitations of this method for synthesis of 2,6-disubstituted pyridines, we used various electrophiles

such as propionaldehyde, benzophenone, methyl propyl ketone, benzonitrile, and *N*-butyrylpiperidine, and the corresponding 2,6-disubstituted pyridines **2** were obtained in moderate yields. Moreover, treatment of 3-(*N,N*-diethylcarbamoyl)pyrazolo[1,5-*a*]pyridine (**1b**) with 2 molar equivalents of *n*-BuLi followed by reaction with benzaldehyde gave ring-opening product **3**. A similar result was obtained in the reaction of **4**, whose structure includes a pyrazolo[1,5-*a*]pyridine moiety.

Lithiation of **1a** with equimolar *n*-BuLi at -78°C followed by reaction with benzaldehyde produced only the 7-substituted pyrazolo[1,5-*a*]pyridine **7a**. Subsequent lithiation of 7-substituted pyrazolo[1,5-*a*]pyridine **7a** led to the 2,6-disubstituted pyridine **2a**. Furthermore, lithiation of **1a** with 2 molar equivalents of *n*-BuLi at -78°C for 0.5 h gave ring-opening product **6a**; however, lithiation with equimolar *n*-BuLi under the same conditions recovered the starting **1a**. Similar results were obtained in the lithiation at -50°C and -10°C. We considered that the pyridine **2a** is produced by the sequential reaction involving lithiation, reaction with benzaldehyde, and cleavage of the pyrazole ring, as shown in Chart 2. But the details are not clear.





Chart 2

Lithiation of 3-(*N,N*-dimethylaminomethyl)pyrazolo[1,5-*a*]pyridine with 2 molar equivalents of *n*-BuLi followed by reaction with benzaldehyde produced only the 7-substituted pyrazolo[1,5-*a*]pyridine in 72% yield. This result indicates that ring-opening of the pyrazolo[1,5-*a*]pyridines requires an electron-withdrawing substituent at the 3-position. Sufficient acidity of the C²-hydrogen achieved the ring-opening through the formation of 2-lithiopyrazolo[1,5-*a*]pyridine by chelation and the electron-withdrawing effect of the substituent.

In addition, we examined the ring-cleavage of several pyrazolo[1,5-*a*]pyridines with or without an electron-withdrawing group at the 3-position using a base. When 3-cyano-, 3-ethoxycarbonyl, and 3-benzoylpyrazolo[1,5-*a*]pyridines were treated with NaH in DMF at 140°C for 2 h, the corresponding ring-cleavage products, α-cyano- (**6c**), α-ethoxycarbonyl- (**6d**), α-benzoyl-2-pyridineacetonitrile (**6e**), were produced in moderate yields. On the other hand, similar treatment of unsubstituted pyrazolo[1,5-*a*]pyridine with either NaH and *n*-BuLi failed to cleave the ring. The ring-cleavage of the pyrazole ring by base action has already been reported.⁶⁾




	R	base	yield (%)
6c	CN	NaH / DMF	39
6d	COOEt	NaH / DMF	35
6e	COPh	NaH / DMF	49
6f	H	NaH / DMF	—
6f	H	<i>n</i> -BuLi / THF	—

Chart 3

We found a new method for synthesis of 2,6-disubstituted pyridines from pyrazolo[1,5-*a*]pyridines. In general, the direct introduction of carbon-functional groups at the 2- or 6-position on pyridine ring by lithiation is difficult;⁷ nevertheless, pyrazolo[1,5-*a*]pyridines are easily lithiated at the 7-position, which corresponds to the 2- or 6-position on a pyridine ring.

REFERENCES AND NOTES

- 1) T. Sasahara, K. Tanji, T. Saito, T. Higashino, Abstract of Papers, *The 110th Annual Meeting of Pharmaceutical Society of Japan, Sapporo*, Aug. 1990, 2, 131.
- 2) a) B. J. Wakefield, *"Organolithium Methods"* p. 32, Academic Press (London), 1988; b) A. I. Meyers, E. D. Mihelich *J. Org. Chem.*, **40**, 3158 (1975).
- 3) Starting compound 1a was synthesized through two steps, as shown below.



- 4) A typical procedure: To a solution of 1a (300 mg, 1.4 mmol) in 10 ml of THF was slowly added *n*-BuLi (1.8 ml, 1.6 mol / l in hexane) at -78°C, and the solution was further stirred for 30 min. A solution of benzaldehyde (456 mg, 4.2 mmol) in 1 ml of THF was added to the solution, and the resulting mixture was stirred for 30 min at -78°C. After the reaction mixture was stirred at room temperature for 1 h, the mixture was poured into H₂O. The mixture was extracted with CHCl₃, and the extract was dried over Na₂SO₄ and concentrated. Purification of the residue by column chromatography on SiO₂ gave 2a.
- 5) 2a: *Anal.* Calcd for C₁₉H₁₉N₃O₂: C, 71.01; H, 5.96; N, 13.07. Found: C, 71.00; H, 5.87; N, 13.05. IR (KBr) cm⁻¹: 2186 (CN), 3178 (OH), 3442 (NH). ¹H-NMR (CDCl₃) δ: 9.62 (1H, bs, NH), 7.60 (1H, t, *J* = 7.8 Hz, pyridine), 7.39-7.30 (5H, m, Ph), 7.21 (1H, d, *J* = 7.8 Hz, pyridine), 7.16 (1H, d, *J* = 7.8 Hz, pyridine), 5.77 (1H, s, CH), 4.14 (2H, s, CH₂), 2.70 (1H, bs, OH), 1.29 (3H, s, Me), 1.21 (3H, s, Me). ¹³C-NMR (CDCl₃) δ: 167.6 (s), 159.5 (s), 154.9 (s), 143.5 (s), 137.3 (d), 128.8 (d), 128.0 (d), 127.0 (d), 119.6 (s), 117.2 (d), 114.2 (d), 79.3 (t, CH₂), 76.7 (d, -CH-OH), 59.2 (s), 26.77 (q), 26.72 (q). The spectral data supported the oxazolidinylidene structure 2a, but the stereochemistry has not yet been determined.
- 6) a) B. K. Martsokha, A. M. Simonov, *Chem. Heterocycl. Compd. (Engl. Transl.)*, **1**, 531 (1965); b) D. J. Gale, J. F. K. Wilshire, *Aust. J. Chem.*, **26**, 2683 (1973); c) T. Higashino, Y. Matsushita, M. Takemoto, E. Hayashi, *Chem. Pharm. Bull.*, **31**, 3951 (1983).
- 7) a) Y. Kondo, N. Murata, T. Sakamoto, *Heterocycles*, **37**, 1467 (1994); b) G. Queguiner, F. Marsais, V. Snieckus, J. Epsztajin, *Adv. Heterocycl. Chem.*, **52**, 187 (1991).

(Received September 30, 1994; accepted November 18, 1994)