

Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn19>

Synthesis of Nucleoside 3'-Thiophosphates in One Step Procedure

Maciej B. Szczepanik^a, Laurent Désaubry^a & Roger A. Johnson^a

^a Department of Physiology and Biophysics, Health Sciences Center, State University of New York at Stony Brook, Stony Brook, NY, 11794, USA

Published online: 04 Oct 2006.

To cite this article: Maciej B. Szczepanik, Laurent Désaubry & Roger A. Johnson (1999) Synthesis of Nucleoside 3'-Thiophosphates in One Step Procedure, Nucleosides and Nucleotides, 18:4-5, 951-953, DOI: [10.1080/15257779908041610](https://doi.org/10.1080/15257779908041610)

To link to this article: <http://dx.doi.org/10.1080/15257779908041610>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS OF NUCLEOSIDE 3'-THIOPHOSPHATES IN ONE STEP PROCEDURE

Maciej B. Szczepanik, Laurent Désaubry, Roger A. Johnson

Department of Physiology and Biophysics, Health Sciences Center,
State University of New York at Stony Brook, Stony Brook, NY 11794, USA

ABSTRACT:


A mild and efficient one-step method of thiophosphorylation was devised for acid-sensitive nucleosides. The procedure is based on thiophosphorylation of nucleoside magnesium alkoxide by 2-chloro-2-thio-1,3,2-dioxaphospholane. The utility and efficiency of this method combined with deprotection of the resulting cyclic triester were demonstrated by its application to the synthesis of both adenosine 3'- and 5'-thiophosphates. The procedure does not require protection of the exocyclic amino group and can be successfully used for the thiophosphorylation of nucleosides that are unusually sensitive to depurination.

Continuing our studies on the structure activity relationship of inhibitors of adenylyl cyclase, we synthesized 3'-thiophosphate analogs of 2'-d-3'AMP and 2',5'-dd-3'AMP, two modestly potent inhibitors of this enzyme.^{1,2}

A new, efficient method for introduction of a thiophosphate group in one step was devised. The procedure is based on thiophosphorylation of nucleoside magnesium alkoxide by 2-chloro-2-thio-1,3,2-dioxaphospholane. The utility and efficiency of this method combined with deprotection of the resulting cyclic triester has been demonstrated by its application to the synthesis of both adenosine 3'- and 5'-thiophosphates. The procedure does not require protection of the exocyclic amino group and can be successfully used for the preparation of labile nucleotides such as 2',5'-dd-3'AMPS, unavailable via the phosphoramidite approach.

* Correspondence should be sent to: RAJ, email: rjohnson@ccmail.sunysb.edu, telephone: 516-444-3040; facsimile: 516-444-3432.

Scheme



Sequential treatment of *N*-unprotected 2',5'-dd-Ado³ **1a** with *tert*-butylmagnesium chloride and 2-chloro-2-thio-1,3,2-dioxaphospholane⁴ **4** yielded the thiophosphate **2a** (Scheme). The cyclic triester **2a** could be easily deprotected by treatment with NaCN in DMSO under vacuum followed by ethanolic NaOH solution to give 2',5'-dd-3'AMPS **3a** in 61% overall yield.⁵ Analogous treatment of 5'-TBDMS-2'-deoxyadenosine **1b** and 2',3'-O-isopropylideneadenosine gave 2'-d-3'AMPS and 5'AMPS in overall yields 31% and 47%, respectively. The structures of all products were confirmed by ¹H-NMR, ³¹P-NMR, MS, and UV spectroscopic analyses.⁶

As expected, 2'-d-3'AMPS and 2',5'-dd-3'AMPS were found to inhibit adenylyl cyclase, with potencies comparable to those of the respective 3'-phosphates.⁷ Both 2'-d-3'AMPS and 2',5'-dd-3'AMPS were stable to rat liver hydrolytic enzymes, which agrees with the generally higher stability of thiophosphates as opposed to phosphates. But, somewhat surprisingly, the 3'-thiophosphates were not effective competitive inhibitors of the hydrolysis of 2'-d-3'AMP.

The availability of these thiophosphates can lead to the preparation of tools for biochemical and pharmacological studies. This includes the synthesis of stable, chiral

derivatives useful in studies of enzyme mechanism or conjugation to reactive functionalities to create unique probes such as affinity labels.

Acknowledgment: We wish to thank Dr J. Marecek and T. Fischer for their valuable help. This work was supported by NIH research grant DK 38828.

REFERENCES AND NOTES

1. Désaubry, L.; Shoshani, I.; Johnson R.A., *J. Biol. Chem.* **1996**, 271, 14028-14034.
2. Johnson, R.A.; Yeung, S-M.H.; Stübner, D.; Bushfield, M.; Shoshani, I., *Mol. Pharmacol.* **1989**, 35, 681-688.
3. Désaubry, L.; Shoshani, I.; Johnson, R.A., *Nucleosides & Nucleotides* **1995**, 14, 1453-1460.
4. Yamasaki, T., Sato, T., *Science Repts. Research Insts. Tohoku Univ.* **1954**, Ser. A, 6, 384, or *Chem. Abstr.* **1956**, 50, 314.
5. Thuong, N.T., Chabrier, P., *Bull. Soc.Chim. Fr.* **1975**, 9-10, 2083-2088.
6. Szczepanik M.B., Desaubry, L., Johnson, R.A., accepted for publication in *Tetrahedron Letters*.
7. IC₅₀ values are as follows: 2'-d-3'AMP: 1.2 μM, 2'-d-3'AMPS: 3.1 μM, 2',5'-dd-3'AMP: 0.46 μM, 2',5'-dd-3'AMPS: 0.60 μM.