

Hitting a moving target: simulation and crystallography study of ATAD2 bromodomain blockers

Aymeric Dolbois, Laurent Batiste, Lars Wiedmer, Jing Dong, Manuela Brütsch, Danzhi Huang, Nicholas M. Deearain, Dimitrios Spiliotopoulos, Iván Cheng-Sánchez, Eleen Laul, Cristina Nevado, Pawel Sledz, and Amedeo Caflisch

ACS Med. Chem. Lett., **Just Accepted Manuscript** • Publication Date (Web): 30 Jun 2020

Downloaded from pubs.acs.org on June 30, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Hitting a moving target: simulation and crystallography study of ATAD2 bromodomain blockers

Aymeric Dolbois,^{1†} Laurent Batiste,^{2†} Lars Wiedmer,² Jing Dong,^{2†} Manuela Brütsch,¹

Danzhi Huang,² Nicholas M. Deerain,² Dimitrios Spiliotopoulos,² Iván Cheng-Sánchez,¹

Eleen Laul,¹ Cristina Nevado,^{1} Paweł Śledź,^{2*} Amedeo Caflisch^{2*}*

Department of Chemistry¹ and Department of Biochemistry,² University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

ABSTRACT: Small molecule ligand binding to ATAD2 bromodomain is investigated here through the synergistic combination of molecular dynamics and protein crystallography. A previously unexplored conformation of the binding pocket upon rearrangement of the gatekeeper residue Ile1074 has been found. Further, our investigations reveal how minor

1
2
3 structural differences in the ligands result in binding with different plasticity of the ZA loop
4
5
6
7 for this difficult-to-drug bromodomain.
8
9

10
11
12 KEYWORDS: ATAD2 bromodomain, conformational selection, fuzzy binding, protein X-
13
14
15 ray crystallography, molecular dynamics
16
17
18
19
20
21
22

23 Contemporary drug discovery often relies on the availability of structural data to guide
24
25
26 the design of ligands and hit optimization.¹ This task becomes challenging when the
27
28
29 protein target is flexible and its conformational changes need to be considered.^{2,3} Proper
30
31
32 understanding of protein flexibility and conformational selection in small molecule binding
33
34
35 events can be leveraged to the benefit of lead discovery campaigns, in particular with the
36
37
38 help of atomistic simulation approaches.⁴⁻¹⁴ Reader domains involved in epigenetic
39
40
41 regulation (regulation of gene expression without alterations to the DNA sequence), are
42
43
44 highly flexible and adaptive receptor targets.¹⁵⁻¹⁷ Given their potential implication in the
45
46
47 control of genetic information transfer within the cell, the quest for potent and selective
48
49
50
51
52
53
54
55
56
57
58
59
60

ligand binders that interact with these proteins has been the focus of intense efforts both in academia and industry over the past decade.


A particularly difficult-to-drug target in this series is the ATAD2 (ATPase family AAA domain-containing 2) bromodomain, a highly flexible and challenging protein-protein interaction domain implicated in multiple cancers.¹⁸⁻²³ Despite its high therapeutic potential, only two probes have been recently reported.²⁴⁻²⁷ ATAD2 is considered poorly druggable mainly due to a shallower, more polar and flexible KAc binding pocket as compared to other bromodomains. Here, we report a detailed structural investigation on the binding interactions to the acetyl-lysine (KAc)'s reader ATAD2 through a combination of molecular dynamics (MD) simulations and protein crystallography for several de novo designed ligands. Our study sheds light on the multiple factors underlying the plasticity of this bromodomain, including the local disorder of the hydrophobic side chains involved in binding as well as the remarkable flexibility and adaptability of the ZA loop.

Our investigation commenced with acetylthiazole hit **1** (Table 1), which was identified by protein structure-based pharmacophore search in the available commercial chemical space.²⁸ While several sub-classes of 2-amino thiazole (2-AT) are considered Pan Assay

1
2
3
4 INterference CompoundS (PAINS), additional experimental and computational evidence
5
6
7 summarized herein below made us confident in our choice of this fragment for further
8
9
10 optimization.^{29,30} Our key discovery hypothesis was the interaction between the
11
12
13 acetylthiazole and the conserved asparagine residue (Asn1064), as well as between the
14
15
16
17 primary ammonium group of the alanine moiety and Asp1071 (Figure 1). The stability of
18
19
20 these interactions was confirmed by MD simulations (for a movie of the MD simulation,
21
22
23 see SI), and the binding of compound **1** was validated both experimentally via TR-FRET
24
25
26
27 (336 μ M, see SI) and crystallographically (Fig. 1A). As predicted, **1** occupies the KAc
28
29
30 binding pocket of ATAD2 surrounded by the hydrophobic residues Val1008, Val1013,
31
32
33 Val1018, Tyr1021, Tyr1063 and Ile1074. The amino group of the alanine moiety of **1**
34
35
36
37 indeed reaches out to Asp1071, making a relatively weak salt bridge (3.6 Å).
38
39
40
41

42 Based on the structure of compound **1** structure, a series of derivatives was designed
43
44
45 and subsequently prepared to explore the binding landscape of this highly flexible and
46
47
48 thus challenging target. The compounds synthesized within this campaign have been
49
50
51 summarized in Table 1 (See experimental section. 2D structures, PDB codes for protein-
52
53
54
55 ligand complexes as well as IC₅₀ or K_d values measured by TR-FRET or ITC can also be
56
57
58
59
60

found in Table 1 and in the SI). Within this compound series, we first set out to rigidify the structure and improve the salt bridge observed for compound **1** by presenting the secondary amino group to the Asp1071. To this end a compound bearing a piperazine was synthesized (**2**). The X-Ray structure of this ligand in complex with ATAD2 showed the desired interaction of the piperazine group with Asp1071, with its distal nitrogen atom forming a higher quality interaction than the original fragment (2.6 Å – Fig. 1B). In contrast, the proximal nitrogen assumed a close position to the amino group of **1** and did not engage with the aspartate residue. A guanidinium derivative of **1** was also prepared (**3**), which formed a network of hydrogen bonds between Tyr1063 and Asp1071 (Fig. 1C). This led to rearrangement of the side chains of the gatekeeper residue Ile1074 as well as that of Val1008, translating into slight remodeling of the binding pocket and tilting of the thiazole ring to maintain hydrophobic interactions with these residues. Similar rearrangement of the binding site was seen in the case of a close derivative **4** with an additional methyl group introduced to improve hydrophobic interactions to Tyr1063 (Fig. 1D). While some interactions seemed to have improved with this approach, no significant change in binding affinity could be observed for these compounds.

Table 1. Binders of ATAD2 bromodomain described in this study.

Compound n°	R ₁	R ₂	IC ₅₀ ¹ (μM)	PDB code	Compound n°	R ₁	R ₂	IC ₅₀ ¹ (μM)	PDB code
1	H		336	5F36	10			> 150	6HI7
2	H		> 150	6EPU	11			> 150	6HI8
3	H		140	6EPX	12			> 150	6EPT
4	H		> 150	6HI3	13			> 150	6HIA
5			> 150	6EPV	14			80	6HIB
6			> 150	6EPJ	15			> 150	6HIC
7			> 150	6HI4	16			> 150	6HID
8			> 150	6HI5	17	H		15 25 ²	6HIE
9			95	6HI6	18			> 150	/

¹ Measured in a Time-Resolved (TR)-FRET assay. ² Measured by ITC.




Figure 1. A-D – Crystal structures of the ATAD2 bromodomain in the complex with compounds 1-4. 2Fo-Fc electron density maps are shown in blue. In all figures, conserved water molecules and water molecules involved in bridging polar interactions (dashed lines) between protein and ligand are shown (red spheres).

To better understand the conformational rearrangements within the pocket and their potential impact on binding, we reanalyzed large datasets of all-atom simulations that we previously collected for ATAD2 bromodomain, both of its apo form as well as of the one

1
2
3 bound to KAc-containing peptide.³¹ The rearranged conformations of residues Val1008
4
5
6
7 and Ile1074 (Fig. 2A) were populated in the runs with both apo and holo ATAD2 and
8
9
10 appeared independent of the movements of the ZA loop, hinting at a conformational
11
12
13 selection mechanism in ligand binding. Moreover, while both observed conformations of
14
15
16
17 Val1008 turned out to appear with similar frequency, there was a clear preference for the
18
19
20
21 conformation of the gatekeeper Ile1074 that opens a remodelled cavity (Fig. 2B). While
22
23
24 this preference was limited in terms of estimated energy difference (~1-2 kcal/mol), we
25
26
27
28 envisaged the possibility of attaining better ligands by binding to the more populated
29
30
31 conformation of the gatekeeper Ile1074.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. A. Superposition of structures of compounds **1** and **3** (shown in black). B. Populations of different conformers of the residues Val1008 and Ile1074 according to the MD simulations. Individual data points represent the conformational states in crystal structures deposited in the PDB as of 01.03.2019. C. Co-crystal structure of compound **5** bound to ATAD2 bromodomain.

We decided to cyclize the original fragment **1** and ‘grow’ it deeper into the pocket to gain stabilization of the acetyl group, whose high mobility has been observed in the MD simulations. (see Supporting Information for details). Furthermore, most bromodomains,

1
2
3
4 like ATAD2, contain up to four water molecules in their KAc binding pocket. As shown by
5
6
7 others³² different acetyl mimic can perturb this water network and induce selectivity over
8
9
10 other bromodomains. Therefore, we hypothesized that improved selectivity could be
11
12
13 attained for a reshaped pocket by the cyclic acetyl mimic, as suggested by docking.
14
15
16
17 Compound **5** was synthesized confirming this hypothesis according to the X-Ray structure
18
19
20 shown in Fig. 2C. Still, no appreciable affinity improvement could be quantified for this
21
22
23 compound and thus alternative modifications of the acetylthiazole core were
24
25
26
27 subsequently explored.
28
29
30

31
32 Based on the initial exploration of the ligand blueprints and interactions within the
33
34
35 binding site, the molecules were grown in the direction of the ZA loop. Recently, our
36
37
38 groups have reported a novel molecular diversification software named AutoCouple.³³
39
40
41
42 AutoCouple generates combinatorial virtual library of binder candidates that are novel and
43
44
45 synthetically accessible in one formal step by coupling a validated fragment with
46
47
48 commercially available building blocks. The library, fairly exhaustive in terms of easily
49
50
51 accessible chemical derivative space, is then evaluated for possible binding to the target
52
53
54
55
56 by the means of tethered docking, and candidates most likely to bind are shortlisted.
57
58
59
60

Guided by these predictions, we prepared a series of derivatives of compound **1** via Pd-catalyzed Suzuki cross-coupling reactions (compounds **6-10**) to pick up additional interactions with the ZA loop while maintaining the key interaction cascade to Asp1071 and Asn1067.

This focused library provided insight into conformational changes upon ligand binding. The same rearrangement of the side chains of the gatekeeper residue Ile1074 and Val1008, as well as the increased size of the binding pocket and slight tilting of the thiazole ring were observed for example for compound **6**. In addition, the phenol moiety engages a superficial solvent-exposed hydrogen bond to Arg1007 (Figure 3A). Interestingly, compound **7**, a close derivative of **6**, lacks the stabilizing interaction with Arg1007 due to the methylation of the hydroxyl group, and its aromatic moiety is disordered in the corresponding co-crystal structure (Fig. S1A).




Figure 3. A-B – Crystal structures of the ATAD2 bromodomain in the complex with compounds **6** and **8**, respectively. C – Different binding modes seen in the all-atom MD simulations of compound **8**-ATAD2 bromodomain complex: left – van der Waals interaction between the side chain of Val1013 and amino-pyridine, right – hydrogen bond between the backbone NH of Asp1014 and nitrogen atom of pyridine ring.

Some of these molecules bound to the enlarged conformation of the pocket but without engaging the ZA loop. In these cases, both the loop and aromatic moiety appear to be

disordered, as for pyridine-containing molecule **8** (Fig. 3B). Significant mobility of the ZA loop is expected, based on previous MD simulations.³¹ To understand the interaction mode of **8** and the higher mobility of the ZA loop, a series of all-atom simulations of the complex were performed, revealing the larger disorder in the ZA loop, as compared to simulations previously reported by us.³¹ While the key interactions of **8** in the binding pocket remained stable for more than 150 ns (e.g. salt bridge to Asp1071), multiple interaction modes of the aromatic substituent with the ZA loop emerge from the simulations (Fig. 3C). These include the direct hydrogen bond between the nitrogen atom of the pyridine and the main chain NH of Asp1014, and the hydrophobic contact between the aromatic ring and the side chain of Val1013. These interactions engage and disengage over the course of the simulation without a single major defined conformation, which is reminiscent of fuzzy binding. Similarly to compound **8**, compound **9** bears a pyridine substituent that cannot be seen in the electron density in the co-crystal structure, along with part of the ZA loop. In contrast, the ZA loop and the R₁ substituent are ordered for **10**, which bears a 3-aminobenzene substituent, and shows a binding mode similar to

1
2
3 the phenol derivative **6** (Figure S1C). Thus, the promiscuous fuzzy binding behavior
4
5
6
7 seems to be induced by the pyridine moiety.
8
9

10 It should be noted that fuzzy or disordered binding is usually avoided in ligand design
11
12 to favor well-defined enthalpy-driven interactions, which provide potency and selectivity.
13
14 To study the relationship between the compound structure and the fuzzy binding mode
15
16
17 further, molecules containing an aminopyridine moiety were extended by merging them
18
19
20
21 with different motifs designed to better engage the Asp1071 residue. As these molecules
22
23
24 might be able to select different conformations of the binding pocket, we envisaged it
25
26
27 would be worth to recapitulate how this affects the fuzzy binding behavior. As seen before,
28
29
30
31 compound **4** selects for enlarged conformation of the pocket and would project the
32
33
34 aromatic group alongside the same vector as in **8**. As expected, the aminopyridine-
35
36
37 bearing derivative **11** exhibits very similar fuzzy binding to that seen for **8** (Figure 4). In
38
39
40
41 contrast, both derivatives of **2** and **3** (**12** and **13**, respectively), show ordered binding of
42
43
44 the aminopyridine group as predicted in the original docking efforts. Importantly, **12** binds
45
46
47 to the same conformation of the pocket as **2**, and introduction of the aromatic moiety does
48
49
50
51 not shift its conformational selectivity, unlike in the case of **1** and **8**. It is clear from these
52
53
54
55
56
57
58
59
60

results that the fuzzy binding behavior arises as a result of very delicate interplay of multiple factors, so that even small modifications to the ligand may disrupt such binding mode, which further complicates the structure-based design against such a highly mobile target. To slightly modulate interactions with Tyr1063, the alanine moiety was modified with a highly strained cyclobutyl ring (**14**), which not only led to ordering of the aminopyridine moiety, but also shifted the conformational equilibrium of the gatekeeper residue Ile1074 and slightly improved binding affinity for this compound.




Figure 4. A-D – Crystal structures of the ATAD2 bromodomain in the complex with compounds **11-14**.

1
2
3 A second AutoCouple campaign was launched to design derivatives that would engage
4
5
6
7 the ZA loop (see Supp. Fig. 2 for co-crystal structures of ATAD2 bromodomain with
8
9
10 compounds **15** and **16**). Unfortunately, no significant gain in affinity was observed, which
11
12
13 showcased the intrinsic difficulty associated with ordering this highly flexible loop (i.e.
14
15
16
17 overcoming a high entropic penalty).
18
19

20
21 Next, we concentrated on further extending these molecules so that they could pick up
22
23
24 interactions with Arg1077, in line with improved affinities observed for compounds
25
26
27 previously reported by others (see supporting information for IC₅₀ in our TR-FRET
28
29
30 assay).²⁴ The proximal nitrogen atom in compound **2**, which does not engage Asp1071
31
32
33
34 was alkylated with an ethyl-linked cyclohexyl ring scaffold yielding ligand **17**. This
35
36
37
38 derivative forms a favorable charge-dipole interaction with Arg1077 through its CF₂ group
39
40
41 (Figure 5). The rigidity of the ethyl difluorocyclohexyl group also allowed to maintain the
42
43
44
45 Ile1074 conformation with a smaller opening of the pocket. This molecule is also the most
46
47
48
49 potent derivative in this series, with IC₅₀ of 15 μM in our TR-FRET assay and K_d of 25 μM
50
51
52
53 in ITC. In contrast to observations made for the more potent previously reported chemical
54
55
56
57 probe,²⁴ further modifications tackling the interactions with the ZA loop by introduction of
58
59
60

an aminopyridine moiety (compound **18**) substantially lowered the potency, thus highlighting how not only the extensions, but the molecule's core need to synergize to achieve substantial gain in affinity for the ATAD2 bromodomain.

In conclusion, we have extensively explored conformational selection of the ATAD2 bromodomain. The MD trajectories and crystal structures allow us to formulate the following conclusions. First, the key hydrophobic residues in the KAc binding site, i.e., the gatekeeper Ile1074, Val1008, and Val1013, can assume multiple orientations for different ligands and even for the same ligand. The local disorder of the hydrophobic side chains involved in binding is observed in the simulations (carried out with compounds **5** and **8**) and the crystal structures with ligands **7**, **8**, **9**, and **11**. Such disorder is congruent with sub-optimal interactions of high micromolar affinity binders. Second, the intrinsic plasticity of the ZA loop results in fuzzy binding or well-defined packing onto the ligand depending on fine details of the latter. As an example, compounds **8** and **10** differ by a single atomic element and show fuzzy binding and single well-defined pose, respectively. Note that this variability is not a consequence of differences in packing between protein molecules in the crystal as all structures belong to the same space group. While the multiple

orientations of the hydrophobic side chains have not been reported previously, the formation of a fuzzy complex and adaptable interface of binding, which is mainly due to the flexibility of the ZA loop, have been observed in atomistic simulations of the histone H4 peptide/ATAD2 bromodomain complex.³¹ The coupling of the ZA-loop rigidification and the rearrangement of the hydrophobic side chains involved in binding can be explored next as a potential strategy to attain potent and selective ligands for this highly mobile target.




Figure 5. Crystal structure of the ATAD2 bromodomain in the complex with compound **17** (A) and the GSK8814 chemical probe (B). C – Superposition of compound **17** and the GSK8814 chemical probe (shown in black) in the binding site. D – 2D structure of GSK8814 and binding affinity determined in house via TR-FRET assay.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website. Experimental procedures, crystallographic and biophysical data, as well as details of the organic compound characterization and molecular dynamics simulations (PDF)

AUTHOR INFORMATION

Corresponding Author

*Phone: +41 44 635 55 21. E-mail: caflisch@bioc.uzh.ch.

*Phone: +41 44 635 39 45. E-mail: cristina.nevado@chem.uzh.

*Phone: +41 44 635 55 87. E-mail: p.sledz@bioc.uzh.ch.

Present Addresses

†Present address: Astex Therapeutics, Cambridge, UK

Author Contributions

‡These authors contributed equally.

Funding Sources

This work has been funded by the Swiss National Science Foundation (Project Funding to AC) and Swiss Cancer League - Krebsliga (to CN and AC). PŠ is grateful to the UZH BioEntrepreneurship program for financial support.

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

The use of beamlines at the Swiss Light Source (Paul Scherrer Institut, Villigen, Switzerland) is gratefully acknowledged. We thank Cassiano Langini for help with

analysis of the MD data, and Maurus Mathis and Johanne Ling for their help with the synthesis and characterization of the compounds on the outset of the project. We are grateful to ChemAxon for providing academic licence for Marvin version 16.2.15.0 (2016), which was used for drawing, displaying and characterizing chemical structures and protonation states, as well as generating conformers (www.chemaxon.com).

ABBREVIATIONS

ATAD2, ATPase family AAA domain-containing 2

PDB CODES

The PDB codes of ATAD2 in complex with the following compounds are listed herein below:

1 5F36; 2 6EPU; 3 6EPX; 4 6HI3; 5 6EPV; 6 6EPJ; 7 6HI4; 8 6HI5; 9 6HI6; 10 6HI7; 11 6HI8; 12 6EPT; 13 6HIA; 14 6HIB; 15 6HIC; 16 6HID; 17 6HIE.

Authors will release the atomic coordinates and experimental data upon article publication.

REFERENCES

(1) Blundell, T. L. Structure-Based Drug Design. *Nature* **1996**, *384*, 23–26.

- (2) Arkin, M. R.; Randal, M.; DeLano, W. L.; Hyde, J.; Luong, T. N.; Oslob, J. D.; Raphael, D. R.; Taylor, L.; Wang, J.; McDowell, R. S.; et al. Binding of Small Molecules to an Adaptive Protein-Protein Interface. *Proc. Natl. Acad. Sci.* **2003**, *100*, 1603–1608.
- (3) Wells, J. A.; McClendon, C. L. Reaching for High-Hanging Fruit in Drug Discovery at Protein-Protein Interfaces. *Nature*. **2007**, *450*, 1001–1009.
- (4) Śledź, P.; Caflisch, A. Protein Structure-Based Drug Design: From Docking to Molecular Dynamics. *Curr. Opin. Struct. Biol.* **2018**, *48*, 93–102.
- (5) Durrant, J. D.; McCammon, J. A. Molecular Dynamics Simulations and Drug Discovery. *BMC Biol.* **2011**, *9*, 71.
- (6) Ganesan, A.; Coote, M. L.; Barakat, K. Molecular Dynamics-Driven Drug Discovery: Leaping Forward with Confidence. *Drug Discovery Today* **2017**, *22*, 249–269.
- (7) Zhao, H.; Caflisch, A. Molecular Dynamics in Drug Design. *Eur. J. Med. Chem.* **2015**, *91*, 4–14.

- (8) Su, J.; Liu, X.; Zhang, S.; Yan, F.; Zhang, Q.; Chen, J. Insight Into Selective Mechanism of Class of I-BRD9 Inhibitors Toward BRD9 Based on Molecular Dynamics Simulations. *Chem. Biol. Drug Des.* **2019**, *93*, 163–176.
- (9) Su, J.; Liu, X.; Zhang, S.; Yan, F.; Zhang, Q.; Chen, J. A Theoretical Insight Into Selectivity of Inhibitors Toward Two Domains of Bromodomain-Containing Protein 4 Using Molecular Dynamics Simulations. *Chem. Biol. Drug Des.* **2018**, *91*, 828–840.
- (10) Huang, D.; Rossini, E.; Steiner, S.; Caflisch, A. Structured Water Molecules in the Binding Site of Bromodomains Can Be Displaced by Cosolvent. *ChemMedChem* **2014**, *9*, 573–579.
- (11) Magno, A.; Steiner, S.; Caflisch, A. Mechanism and Kinetics of Acetyl-Lysine Binding to Bromodomains. *J. Chem. Theory Comput.* **2013**, *9*, 4225–4232.
- (12) Steiner, S.; Magno, A.; Huang, D.; Caflisch, A. Does Bromodomain Flexibility Influence Histone Recognition? *FEBS Letters* **2013**, *587*, 2158–2163.
- (13) Bacci, M.; Langini, C.; Vymětal, J.; Caflisch, A.; Vitalis, A. Focused Conformational Sampling in Proteins. *J. Chem. Phys.* **2017**, *147*, 195102.
- (14) Gay, J. C.; Eckenroth, B. E.; Evans, C. M.; Langini, C.; Carlson, S.; Lloyd, J. T.; Caflisch, A.; Glass, K. C. Disulfide Bridge Formation Influences Ligand Recognition by the ATAD2 Bromodomain. *Proteins* **2019**, *87*, 157–167.
- (15) Jain, A. K.; Barton, M. C. Bromodomain Histone Readers and Cancer. *J. Mol. Biol.* **2017**, *429*, 2003–2010.

- (16) Muller, S.; Filippakopoulos, P.; Knapp, S. Bromodomains as Therapeutic Targets. *Expert Rev. Mol. Med.* **2011**, *13*, e29.
- (17) Ferri, E.; Petosa, C.; McKenna, C. E. Bromodomains: Structure, Function and Pharmacology of Inhibition. *Biochem. Pharmacol.* **2016**, *106*, 1–18.
- (18) Hwang, H. W.; Ha, S. Y.; Bang, H.; Park, C. K. ATAD2 as a Poor Prognostic Marker for Hepatocellular Carcinoma after Curative Resection. *Cancer Res. Treat.* **2015**, *47*, 853–861.
- (19) Morozumi, Y.; Boussouar, F.; Tan, M.; Chaikuad, A.; Jamshidikia, M.; Colak, G.; He, H.; Nie, L.; Petosa, C.; De Dieuleveult, M.; et al. ATAD2 Is a Generalist Facilitator of Chromatin Dynamics in Embryonic Stem Cells. *J. Mol. Cell Biol.* **2016**, *8*, 349–362.
- (20) Ciró, M.; Prosperini, E.; Quarto, M.; Grazini, U.; Walfridsson, J.; McBlane, F.; Nucifero, P.; Pacchiana, G.; Capra, M.; Christensen, J.; et al. ATAD2 Is a Novel Cofactor for MYC, Overexpressed and Amplified in Aggressive Tumors. *Cancer Res.* **2009**, *69*, 8491–8498.

(21) Zhang, M. J.; Zhang, C. Z.; Du, W. J.; Yang, X. Z.; Chen, Z. P. ATAD2 Is

Overexpressed in Gastric Cancer and Serves as an Independent Poor Prognostic

Biomarker. *Clin. Transl. Oncol.* **2016**, *18*, 776–781.

(22) Hou, M.; Huang, R.; Song, Y.; Feng, D.; Jiang, Y.; Liu, M. ATAD2

Overexpression Is Associated with Progression and Prognosis in Colorectal Cancer.

Jpn. J. Clin. Oncol. **2016**, *46*, 222–227.

(23) Caron, C.; Lestrat, C.; Marsal, S.; Escoffier, E.; Curtet, S.; Virolle, V.; Barbry, P.;

Debernardi, A.; Brambilla, C.; Brambilla, E.; et al. Functional Characterization of ATAD2

as a New Cancer/Testis Factor and a Predictor of Poor Prognosis in Breast and Lung

Cancers. *Oncogene* **2010**, *29*, 5171–5181.

(24) Bamborough, P.; Chung, C. wa; Demont, E. H.; Furze, R. C.; Bannister, A. J.;

Che, K. H.; Diallo, H.; Douault, C.; Grandi, P.; Kouzarides, T.; et al. A Chemical Probe

for the ATAD2 Bromodomain. *Angew. Chemie. Int. Ed.* **2016**, *55*, 11382–11386.

(25) Fernández-Montalván, A. E.; Berger, M.; Kuropka, B.; Koo, S. J.; Badock, V.;

Weiske, J.; Puetter, V.; Holton, S. J.; Stöckigt, D.; Ter Laak, A.; et al. Isoform-Selective

ATAD2 Chemical Probe with Novel Chemical Structure and Unusual Mode of Action.

ACS Chem. Biol. **2017**, *12*, 2730–2736.

(26) Bamborough, P.; Chung, C. W.; Furze, R. C.; Grandi, P.; Michon, A. M.; Watson,

R. J.; Mitchell, D. J.; Barnett, H.; Prinjha, R. K.; Rau, C.; et al. Aiming to Miss a Moving

Target: Bromo and Extra Terminal Domain (BET) Selectivity in Constrained ATAD2

Inhibitors. *J. Med. Chem.* **2018**, *61*, 8321–8336.

(27) Bamborough, P.; Chung, C. W.; Demont, E. H.; Bridges, A. M.; Craggs, P. D.;

Dixon, D. P.; Francis, P.; Furze, R. C.; Grandi, P.; Jones, E. J.; et al. A Qualified Success:

Discovery of a New Series of ATAD2 Bromodomain Inhibitors with a Novel Binding Mode

Using High-Throughput Screening and Hit Qualification. *J. Med. Chem.* **2019**, *62*, 7506–

7525.

(28) Sunseri, J.; Koes, D. R. Pharmit: Interactive Exploration of Chemical Space.

Nucleic Acids Res. **2016**, *44*, W442–W448.

(29) Baell, J. B.; Holloway, G. A. New Substructure Filters for Removal of Pan Assay Interference Compounds (PAINS) From Screening Libraries and for Their Exclusion in Bioassays. *J. Med. Chem.* **2010**, *53*, 2719–2740.

(30). A recent study on 2-AT containing fragments and HTS hits revealed that not all 2-AT molecules are problematic and that acylation of the 2-amino group, as in our case, reduces substantially the risk of pan hitting: Devine, S. M.; Mulcair, M. D.; Debono, C. O.; Leung, E. W. W.; Nissink, J. W. M.; Lim, S. S.; Chandrashekar, I. R.; Vazirani, M.; Mohanty, B.; Simpson, J. S.; et al. Promiscuous 2-Aminothiazoles (PrATs): A Frequent Hitting Scaffold. *J. Med. Chem.* **2015**, *58*, 1205–1214.

(31) Langini, C.; Caflisch, A.; Vitalis, A. The ATAD2 Bromodomain Binds Different Acetylation Marks on the Histone H4 in Similar Fuzzy Complexes. *J. Biol. Chem.* **2017**, *292*, 16734–16745.

(32) Crawford, T. D.; Tsui, V.; Flynn, E. M.; Wang, S.; Taylor, A. M.; Côté, A.; Audia, J. E.; Beresini, M. H.; Burdick, D. J.; Cummings, R.; Dakin, L. A.; Duplessis, M.; Good, A. C.; Hewitt, M. C.; Huang, H.-R.; Jayaram, H.; Kiefer, J. R.; Jiang, Y.; Murray, J.;

Nasveschuk, C. G.; Pardo, E.; Poy, F.; Romero, F. A.; Tang Y.; Wang, J.; Xu, Z.;
Zawadzke, L. E.; Zhu, X.; Albrecht, B. K.; Magnuson, S. R.; Bellon, S.; Cochran, A. G.
Diving into the Water: Inducible Binding Conformations for BRD4, TAF1(2), BRD9, and
CECR2 Bromodomains. *J. Med. Chem.* **2016**, *59*, 5391–5402.

(33) Batiste, L.; Unzue, A.; Dolbois, A.; Hassler, F.; Wang, X.; Deerain, N.; Zhu, J.;
Spiliotopoulos, D.; Nevado, C.; Caflisch, A. Chemical Space Expansion of
Bromodomain Ligands Guided by in Silico Virtual Couplings (AutoCouple). *ACS Cent.*
Sci. **2018**, *4*, 180–188.




Table of contents graphics