Nickel-Catalyzed Double Silylation of a Variety of **Carbonyl Compounds with** 1,2-Bis(dimethylsilyl)carborane

Youngjin Kang,[†] Jinsik Kim,[‡] Young Kun Kong,[‡] Jihwa Lee,[§] Soon W. Lee,[§] Sang Ook Kang,^{*,†} and Jaejung Ko^{*,†}

Departments of Chemistry, Korea University, Chochiwon, Chungnam 339-700, Korea, Kyonggi University, Suwon, Kyonggido, 440-760, Korea, and Sungkyunkwan University, Natural Science Campus, Suwon, 440-746, Korea

Received June 21, 2000

The nickel-catalyzed reactions of 1,2-bis(dimethylsilyl)carborane 1 with aldehydes such as isobutyraldehyde, trimethylacetaldehyde, hexanal, and benzaldehyde afforded 5,6carboranylene-2-oxa-1,4-disilacyclohexanes. The dehydrogenative 1,4-double silylation of methacrolein and *trans*-4-phenyl-3-buten-2-one in the presence of a catalytic amount of Ni-(PEt₃)₄ took place under similar conditions. In contrast, the reaction of 1 with α -methyltrans-cinnamaldehyde and trans-cinnamaldehyde under the same reaction conditions yielded the insertion compounds formed via insertion of a carbonyl group into each of the C-Si bonds of 1. The reaction of 1 with diphenylketene gave the 1,2-adduct across the carbonyl group. The structures of compounds 8, 11, and 14 were determined by single-crystal X-ray crystallography.

Introduction

The double silvlation of unsaturated organic substrates catalyzed by group 10 metal complexes is a useful route to disilyl organic compounds in which two Si-C bonds are created by the addition of two silicon units to unsaturated organic substrates such as alkynes,¹ alkenes,² and 1,3-dienes.³ Nickel complexes, in particular, are excellent catalysts for such reactions of siliconcontaining linear compounds⁴ and cyclic disilanes.⁵

Recently, we⁶ reported the double silvlation reactions of [o-(SiMe₂)₂C₂B₁₀H₁₀]Pt(PPh₃)₂, A, containing an ocarboranylene unit with a variety of unsaturated organic substrates. Interestingly, its reaction with transcinnamaldehyde afforded the insertion compound **B** formed through the insertion of a carbonyl ligand into each of the C–Si bonds. Usually, α,β -unsaturated ketones undergo double silvlation to give 1,4-adducts.⁷

 ⁸ Sungkyunkwan University.
 (1) (a) Naka, A.; Okazaki, S.; Hayashi, M.; Ishikawa, M. J. Organomet. Chem. 1995, 499, 35. (b) Ishikawa, M.; Ohshita, J.; Ito, Y. Organometallics 1986, 5, 1518. (c) Kiso, Y.; Tamao, K.; Kumada, M. J. Organomet. Chem. 1974, 76, 105. (d) Ishikawa, M.; Matsuzawa, S.; Hirotsu, K.; Kamitori, S.; Higuchi, T.; Kamitori, S.; Hirotsu, K.

Hirotsu, K.; Kalintori, S.; Higuth, T.; Kalintori, S.; Hirotsu, K. Organometallics 1985, 4, 2040.
(2) (a) Ishikawa, M.; Okazaki, S.; Naka, A.; Tachibana, A.; Kawauchi, S.; Yamabe, T. Organometallics 1995, 14, 114. (b) Ozawa, F.; Sugawara, M.; Hayashi, T. Organometallics 1994, 13, 3237. (c) Hayashi, T.; Kobayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1994, 13, 3237. (c) Hayashi, T.; Kobayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1994, 13, 3237. (c) Hayashi, T.; Kobayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1995, 14, 114. (b) Ozawa, F.; Sugawara, M.; Hayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1995, 14, 114. (b) Ozawa, F.; Sugawara, M.; Hayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1994, 13, 3237. (c) Hayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1994, 13, 3237. (c) Hayashi, T.; Kawamoto, A. M.; Yamashita, H.; Tanaka, K.; Conganometallics 1995, 14, 114. (c) N.; Conganometallics 1995, M. Organometallics 1990, 9, 280.

(3) (a) Obora, Y.; Tsuji, Y.; Kawamura, T. Organometallics 1993, 12, 2853. (b) Tsuji, Y.; Lago, R. M.; Tomohiro, S.; Tsuneishi, H. Organometallics 1992, 11, 2353.
 (4) (a) Tamao, K.; Miyake, N.; Kiso, Y.; Kumada, M. J. Am. Chem. Soc. 1975, 97, 5603. (b) Okinoshima, H.; Yamamoto, K.; Kumada, M. J. Am. Chem. Soc. 1975, 97, 5603. (c) Ishikawa M.; Nishimura, Y.;

J. Am. Chem. Soc. 1972, 94, 9263. (c) Ishikawa, M.; Nishimura, Y.;

(5) Jan. Chem. Soc. 1972, 94, 9205. (c) Islinkawa, W., Wishindia, T.,
Sakamoto, N.; Ono, T.; Ohshita, J. Organometallics 1992, 11, 483.
(5) Jzang, T.; Lee, C.; Liu, C. S. Organometallics 1988, 7, 1265.
(6) (a) Kang, Y.; Kang, S. O.; Ko, J. Organometallics 1999, 18, 1818.
(b) Kang, Y.; Kang, S. O.; Ko, J. Organometallics 2000, 19, 1216.



Recently, Tanaka and co-workers⁸ reported that a metal complex catalyzed double silvlation of transcinnamaldehyde with o-bis(dimethylsilyl)benzene afforded 1,2 and 1,4 adducts C. The peculiar behavior of the 1.2-bis(dimethylsilyl)carborane 1 has allowed novel transformations of some carbonyl compounds. The double silvlation of the carbonyl by the cyclic bis(silvl)platinum complex was a stoichiometric process and thus was limited. Accordingly, we have started a systematic study of the double silvlation of carbonyl compounds catalyzed by a nickel complex. Here we report the double silvlation reactions of carbonyl compounds such as aldehydes, α,β -unsaturated aldehydes, α,β -unsaturated ketones, and a ketene with 1,2-bis(dimethylsilyl)carborane.

Results and Discussion

Nickel-Catalyzed Reactions with Aldehydes. Although the platinum complex (C₂H₄)Pt(PPh₃)₂ is not a catalyst for the 1,2-double silvlation of aldehydes when 1,2-bis(dimethylsilyl)carborane 1 is used, due to its

Korea University.

[‡] Kyonggi University.

^{(7) (}a) Hayashi, T.; Matsumoto, Y.; Ito, Y. J. Am. Chem. Soc. 1988, 110, 5579. (b) Tamao, K.; Okazaki, S.; Kumade, M. J. Organomet. Chem. 1978, 146, 87.

⁽⁸⁾ Uchimaru, Y.; Lautenschlager, H.-J.; Wynd, A. J.; Tanaka, M.; Goto, M. Organometallics 1992, 11, 2639.

Table 1	1. Cryst	al Data d	of Com	pounds	8 , 1	11, ano	114
---------	----------	-----------	--------	--------	--------------	---------	-----

	8	11	14
empirical formula	C ₁₄ H ₂₇ B ₁₀ NOSi ₂	$C_{16}H_{34}B_{10}O_2Si_2$	C20H32B10OSi2
molecular weight	389.65	422.71	452.74
crystal system	orthorhombic	orthorhombic	monoclinic
space group	A_{ba2}	$P2_{1}2_{1}2_{1}$	$P2_1/n$
a, Å	21.704(4)	9.105(2)	14.110(3)
b, Å	17.575(3)	10.750(2)	11.228(15)
<i>c</i> , Å	11.997(4)	26.495	34.201(6)
β , deg			97.025(12)
V, Å ³	4576(2)	2593.3(8)	5377.4(16)
Zvalue	8	4	8
$D_{ m calcd}$, g cm $^{-3}$	1.131	1.083	1.118
F(000)	1632	869	1904
$\mu ({\rm cm}^{-1})$	0.160	0.148	0.144
λ (Mo K α radiation) (Å)	3.5 - 50	3.5 - 50	1.91 - 22.50
scan type	ω	ω	ω
no. of reflns measd	2082	2581	7245
no. of obsd reflns $I > 2\sigma(I)$	1068	2028	6911
R	0.0822	0.0453	0.0458
$R_{ m w}$	0.2009	0.1115	0.1124
goodness of fit	1.013	1.019	1.018
-			

strong Pt-Si bond strength, the Ni complex Ni(PEt₃)₄ was found to be a good catalyst for the dehydrogenative double silvlation of aldehydes. The nickel-catalyzed double silvlation can be carried out either by reacting 1 and the aldehyde in the presence of the $Ni(PEt_3)_4$ catalyst or by reacting 1 and the aldehyde in the presence of a catalytic amount of the cyclic bis(silyl)nickel complex Me₂Si(o-C₂B₁₀H₁₀)SiMe₂Ni(PEt₃)₂ 2.9 Both procedures showed a high catalytic activity. However, the former procedure is preferred due to the easy accessibility of Ni(PEt₃)₄. The results are summarized in Table 2.

Isobutyraldehyde readily reacted at 80 °C with 1 in the presence of a catalytic amount of Ni(PEt₃)₄ to give 5,6-carboranylene-1,1,4,4-tetramethyl-2-oxa-3-isopropyl-1,4-disilacyclohexane 3 in 74% yield (eq 1). In a similar



fashion, the nickel-catalyzed reaction of 1 with other aliphatic aldehydes such as trimethylacetaldehyde, 5-norbornene-2-carboxaldehyde, and hexanal yielded the corresponding six-membered cyclic insertion products. Benzaldehyde also underwent the 1,2-double silylation reaction under mild conditions. Experiments utilizing substituted aromatic aldehydes such as 4-cyanobenzaldehyde followed a similar course. These reactions of 1 with aromatic aldehydes were faster than those of aliphatic aldehydes. Such dehydrogenative 1,2double silvlation reactions have been reported using a platinum complex as the catalyst with o-bis(dimethylsilyl)benzene.⁸ The insertion of an aldehyde carbonyl group into the Si-Si bond promoted by fluoride ion was also noted.¹⁰

Table 2. Nickel-Catalyzed Reactions of Aldehyde with 1,2-Bis(dimethylsilyl)carborane

entry	reactns	products	yield (%)
i)—сно	Me ₂ Si O Si CHMe ₂	74 %
2	сно	Me2 Si O Si CMe3 Me2	66 %
3	СНО	Me ₂ Si o Si o Me ₂	38 %
4	Сно	Si o Si o Si o Me ₂	83 %
5	СНО	Me ₂ Si O Me ₂	85 %
6	NCСНО	Me ₂ SL o Me ₂	92 %

To provide structural information for one of the newly prepared compounds, a single-crystal X-ray diffraction study of the 4-cyanobenzaldehyde addition product 8 was undertaken. The molecular structure of 8 is shown in Figure 1. A summary of cell constants and data collection parameters is included in Table 1. The X-ray crystal structure of 8 confirmed the presence of a sixmembered ring comprising the two carboranylene carbon atoms, two silicon atoms, and a C–O fragment. The heterocyclic ring is best described as a flattened boat configuration. The Si(1)-O(1) bond length (1.621(7) Å) and O(1)-C(7) bond length (1.156(10) Å) are slightly shorter than those in 5,6-benzo-1,1,4,4-tetramethyl-3-(diphenylmethylidene)-1,4-disila-2-oxa-5-cyclohexene⁶ and

^{(9) (}a) Kang, Y.; Lee, J.; Kong, Y. K.; Kang, S. O.; Ko, J. *Chem. Commun.* **1998**, 2343. (b) Kang, Y.; Lee, J.; Kong, Y. K.; Kang, S. O.; Ko, J. *Organometallics* **2000**, *19*, 1722.

⁽¹⁰⁾ Hiyama, T.; Obayashi, M. Tetrahedron Lett. 1983, 24, 4109.



Figure 1. ORTEP drawing of **8** showing the atom-labeling scheme with 50% probability thermal ellipsoids. Selected bond lengths (Å) and angles (deg): Si(1)–O(1) 1.621(7), O(1)–C(7) 1.156(10), Si(2)–C(70) 1.825(10), Si(2)–C(2) 1.902(8), C(1)–C(2) 1.698(10), C(7)–C(8) 1.55(2), N(1)–C(14) 1.130(2), O(1)–Si(1)–C(1) 105.1(3), Si(1)–C(1)–C(2) 118.5(5), C(1)–C(2)–Si(2) 120.5(5), C(7)–Si(2)–C(2) 105.8-(4), Si(2)–C(7)–O(1) 125.8(8).

comparable to those of $Me_2SiOReO_3^{11}$ and $bis[(8'-(di-methylamino)naphth-1'-yl)dimethylsilyl]ether.^{12}$

The ¹H, ¹³C, and ²⁹Si NMR spectra and the mass spectra of compounds 3-8 were consistent with the structure determined for 8. The ¹H NMR spectrum of 8 contained a distinguishing low-field methine resonance (δ 4.79) which was used to monitor its formation. Four singlets (δ 0.49, 0.44, 0.28, 0.03) in the ¹H NMR spectrum and four singlets (-0.64, -1.68, -2.04, -5.42) ppm) in the ¹³C NMR spectrum of 8 could be assigned to the methyl groups on the silicon atoms which indicated that the methyl groups are not equivalent. The ²⁹Si NMR spectrum of 8 exhibited two resonances at 12.59 and 0.95 ppm arising from the nonequivalent silicon atoms. Similar nickel-catalyzed reactions of 1 with ketones such as acetone, acetophenone, and pinacolone afforded a complicated mixture of products which were difficult to separate in pure form.

Ishikawa,¹³ Tanaka,⁸ and their co-workers have suggested mechanistic pathways for the nickel- and platinum-catalyzed reaction of 3,4-benzo-1,1,2,2-tetraethyl-1, 2-disilacyclobut-3-ene or o-bis(dimethylsilyl)benzene with carbonyl compounds. On the basis of their suggestions, a reasonable mechanism for the formation of 3-8 involves the initial formation of bis(silyl)nickel complex 2 as a key intermediate, a species which we had reported earlier.⁹ Intermediate 2 then coordinates to a carbonyl group (D), followed by insertion of the carbonyl bond into a Ni-Si bond to give a seven-membered cyclic intermediate (E). Analogous reactions of a silyl-substituted transition metal complex with carbonyl compounds have been reported previously.8 Reductive elimination of the nickel fragment from the latter intermediate then leads to the product (F).



Nickel-Catalyzed Reaction with α , β -Unsaturated Aldehydes and Ketones. Since platinum complex catalyzed 1,4-double silylation of α , β -unsaturated carbonyl compounds with disilanes is a known process,⁸ we attempted the platinum complex catalyzed double

Table 3.	Nickel-C	atalyzed	Reactions	of Enones
wit	h 1,2-Bis((dimethy	lsilyl)carbo	orane



silylation of prop-2-enal and methacrolein with 1,2-bis-(dimethylsilyl)carborane at high temperature. No reaction was observed in all cases when cyclic bis(silyl)platinum complex was tried as the catalyst. However, α,β -unsaturated carbonyl compounds readily reacted with 1,2-bis(dimethylsilyl)carborane in the presence of a catalytic amount of Ni(PEt₃)₄ to give the 1,4-double silylating products. Thus, reaction of **1** with methacrolein in the presence of a catalytic amount of Ni(PEt₃) yielded 7,8-carboranylene-1,1,6,6-tetramethyl-4-methyl-2-oxa-1,6-disilacyclooct-3-ene (**9**) in 56% yield (eq 2). The



reaction of 1 with the more hindered *trans*-4-phenyl-3buten-2-one afforded 7,8-carboranylene-1,1,6,6-tetramethyl-3-methyl-5-phenyl-2-oxa-1,6-disilacyclooct-3-ene 10 in 28% yield (Table 3). These products, 9 and 10, could readily be isolated by preparative GLC. The ¹H NMR spectrum of 9 contains five resonances at 1.56, 0.42, 0.37, 0.35, and 0.33 ppm, assigned to the five kinds of methyl groups. The four resonances at 0.42, 0.37, 0.35, and 0.33 ppm were assigned to the methyl groups on the silicon atoms and the peak at 1.56 ppm to the methyl group on the olefinic carbon atom. In the olefinic region of the ¹³C NMR spectrum of **9**, two resonances at 136.03 and 107.22 ppm are present. The ²⁹Si NMR spectrum of 9 shows two resonances at 16.42 and -1.36 ppm. The chemical shifts of 9 are in the range of those of the other eight-membered ring compounds. The structure of 10 was further indirectly confirmed by the

⁽¹¹⁾ Sheldrick, G. M.; Sheldrick, W. S.; J. Chem. Soc. A 1969, 2160.

⁽¹²⁾ Spiniello, M.; White, J. M. Organometallics 2000, 19, 1350.

⁽¹³⁾ Ishikawa, M.; Naka, A.; Okazaki, S.; Sakamoto, H. Organometallics 1993, 12, 87.



Figure 2. ORTEP drawing of **11** showing the atomlabeling scheme with 50% probability thermal ellipsoids. Selected bond lengths (Å) and angles (deg): Si(1)–C(7) 1.893(4), Si(1)–C(2) 1.939(4), C(7)–C(8) 1.529(6), C(8)–C(9) 1.501(6), C(9)–O(2) 1.205(5), C(9)–C(10) 1.488(8), Si(2)– O(1) 1.613(3), Si(2)–C(1) 1.912(4), C(1)–C(2) 1.724(5), C(1)–Si(2)–O(1) 104.18(18), Si(2)–C(1)–C(2) 123.9(2), C(2)– Si(1)–C(7) 109.43(16), Si(1)–C(7)–C(8) 113.8(3), C(7)– C(8)-C(9) 113.3(4), C(8)–C(9)–O(2) 121.0(4), O(2)–C(9)– C(10) 121.2(5).

crystal structure of **11**, the hydrolysis product of **10** (eq 3). Such 1,4-double silylation has been observed in the



transition metal catalyzed double silvlation of α,β unsaturated ketones and aldehydes by Ito,14 Kumada,15 Ishkawa,¹⁶ and their respective co-workers. The product **10** is sensitive to hydrolysis; when crystal growth of **10** was attempted, it was 11 which was isolated. In a subsequent experiment, treatment of 10 with 2 equiv of H₂O in benzene- d_6 was monitored by ¹H NMR spectroscopy. The disappearance of the olefinic hydrogen peak at 6.36 ppm and the formation of a new methylene signal at 2.55 ppm and a hydroxy signal at 2.16 ppm were noted. The ¹H NMR spectrum of the product exhibited four resonances at 0.32, 0.22, 0.21, and 0.07 ppm due to the methyl groups on silicon. The IR spectrum of the compound showed new absorptions due to the carbonyl stretch at 1683 cm⁻¹ and ν_{OH} stretch at 3205 cm⁻¹, concomitant with the disappearance of the C–O stretching mode at 987 cm⁻¹. The mass spectrum of the product showed a molecular ion at m/z 422. The molecular structure of 11 as determined by a singlecrystal X-ray diffraction study is shown in Figure 2. The crystallographic data are given in Table 1.

The 1,4-double silvlation reactions are quite sensitive to the substituents on the C=O group. In contrast to the 1,4-double silvlation reactions of 1 with methac-

rolein, when α -methyl-*trans*-cinnamaldehyde was employed in the reaction of **1** under the same conditions, the insertion product, **12**, of a carbonyl group into each of the C–Si bonds was obtained as a colorless solid in 88% yield (eq 4). A key feature in the ¹H NMR spectrum



of **12** includes a singlet at 5.46 ppm assigned to the methine, concomitant with the disappearance of the aldehyde hydrogen peak. *trans*-Cinnamaldehyde reacted similarly to give **13**. As noted in the Introduction, platinum complex **A** had been found⁶ to react with *trans*-cinnamaldehyde to give a product (**B**) in which a C=O bond had inserted into both of the C-Si bonds. The spectral data for **13** were identical to those we had observed for compound **A**. It is of interest that in the case of compound **A** it had been proven that the oxygen atom of the disiloxane unit was derived from the aldehyde.⁶ Similar insertion was observed in the reac-

tions of (CO)FeSiMe₂CH₂CH₂SiMe₂,¹⁷ (CO)₅MnSiMe₃,¹⁸ and MeEt₂SiCo(CO)₃L¹⁹ with benzaldehyde and of 3,4-benzo-1,1,2,2-tetraethyldisilacyclobut-3-ene with benzophenone in cyclohexene.

Such an insertion of the carbonyl functionality into the *o*-carborane C–H bond was observed in Yamamoto's work on the chemoselective addition of *o*-carborane to aldehyde groups²⁰ or in the fluoride-promoted reaction of *o*-carborane with aldehydes.²¹

Diphenylketene also underwent dehydrogenative double silylation with 1 (eq 5) and the product, 1,2-



adduct to the carbonyl group, was obtained in 80% yield. The initial indication of the cyclization to form **14** was given by the observation of a parent ion in the mass spectrum at m/z 468. Although the ¹H and ¹³C NMR spectra of **14** supported the proposed structure, unambiguous confirmation was provided by an X-ray crystallographic analysis. The molecular structure of **14** is shown in Figure 3. The molecule contains one C₂Si₂CO six-membered ring comprising two carboranylene car-

^{(14) (}a) Hayashi, T.; Matsumoto, Y.; Ito, Y. *Tetrahedron Lett.* **1988**, *29*, 4147. (b) Hayashi, T.; Matsumoto, Y.; Ito, Y. *J. Am. Chem. Soc.* **1988**, *110*, 5579.

⁽¹⁵⁾ Tamao, K.; Okazaki, S.; Kumada, M. *J. Organomet. Chem.* **1978**, *146*, 87.

⁽¹⁶⁾ Ishikawa, M.; Sakamoto, H.; Tabuchi, T. Organometallics 1991, 10, 3173.

⁽¹⁷⁾ Nakazawa, H.; Johnson, D. L.; Gladysz, J. A. Organometallics 1983, 2, 1846.

^{(18) (}a) Johnson, D. L.; Gladysz, J. A. J. Am. Chem. Soc. 1979, 101,
6433. (b) Johnson, D. L.; Gladysz, J. A. Inorg. Chem. 1981, 20, 2508.
(19) Murai, S.; Kato, T.; Sonoda, N.; Seki, Y.; Kawamoto, K. Angew.

Chem, Int. Ed. Engl. 1979, 18, 393.
 (20) Nakamura, H.; Aoyagi, K.; Yamamoto, Y. J. Org. Chem. 1997,

⁽²⁰⁾ Nakamura, H., Aoyagi, K., Tamamoto, T. J. Org. Chem. **1997**, 62, 780.

^{(21) (}a) Cai, J.; Nemoto, H.; Nakamura, H.; Singaram, B.; Yamamoto, Y. *Chem. Lett.* **1996**, 792. (b) Nakamura, H.; Aoyagi, K.; Yamamoto, Y. *J. Am. Chem. Soc.* **1998**, *20*, 1167.



Figure 3. ORTEP drawing of **14** showing the atomlabeling scheme with 50% probability thermal ellipsoids. Selected bond lengths (Å) and angles (deg): Si(1)-O(1)1.641(2), O(1)-C(7) 1.414(3), C(7)-Si(2) 1.875(3), Si(2)-C(2) 1.919(3), C(1)-C(2) 1.700(3), C(1)-Si(1) 1.908(3), C(7)-C(8) 1.335(4), C(7)-O(1)-Si(1) 122.3(2), O(1)-Si(1)-C(1) 105.11(11), Si(1)-C(1)-C(2) 118.18(18), C(1)-C(2)-Si(2) 120.39(17), C(2)-Si(2)-C(7) 103.80(11), Si(2)-C(7)-O(1) 107.46(18), C(7)-O(1)-Si(1) 122.29(16), Si(2)-C(7)-C(8) 132.5(2), C(7)-C(8)-C(15) 119.6(2), C(9)-C(8)-C915)115.8(2).

bon atoms, two silicon atoms, and a saturated C–O fragment. The C₂Si₂CO ring is described as a tilted boat configuration. The C7–C8 bond length (1.335(4) Å) is slightly longer than the typical value for the carbon– carbon double bond (1.317 Å) and comparable to that in 5,6-carboranylene-1,1,4,4-tetramethyl-2,3-diphenyl-1,4-disilacyclohex-2-ene (1.33(1) Å). The bond lengthening of C7–C8 may be related to the wideness of the Si(2)–C(7)–C(8) bond angle (132.5(2)°), resulting in a diminishing of the s character of the C7 atom. The two phenyl rings attached to C8 are nearly perpendicular to one another with a dihedral angle of 86.9°.

In conclusion, the cyclic bis(silyl)nickel complex was found to be a good catalyst for the double silylation of a variety of carbonyl compounds. Thus, the nickelcatalyzed reactions of 1,2-bis(dimethylsilyl)carborane with aldehydes such as isobutyraldehyde, trimethylacetaldehyde, hexanal, and benzaldehyde afforded 5,6carboranylene-2-oxa-1,4-disilacyclohexane.

The 1,4-double silvlation of α , β -unsaturated carbonyl compounds such as methacrolein and *trans*-4-phenyl-3-buten-2-one gave the 1,4-double silvlated products. In contrast, the double silvlation reaction of **1** with α -methyl-*trans*-cinnamaldehyde and *trans*-cinnamaldehyde yielded products formed by insertion of a carbonyl group into each of the C–Si bonds of **2**. Thus, the cyclic bis(silyl)nickel complex as a catalyst has been further exploited in a series of novel chemical transformations.

Experimental Section

All experiments were performed under a nitrogen atmosphere in a Vacuum Atmospheres drybox or by standard Schlenk techniques. Benzene, toluene, and THF were distilled from sodium benzophenone. Dichloromethane and hexane were distilled over CaH₂. EtOH was distilled over Linde type 4 Å molecular sieves. ¹H and ¹³C NMR spectra were recorded on a Varian Gemini 200 spectrometer operating at 200.1 and 50.3 MHz, respectively. ²⁹Si NMR spectra were recorded on a JEOL model EX-270 spectrometer. Chemical shifts were referenced relative to TMS. IR spectra were recorded on a Biorad FTS-165 spectrometer. Mass spectra were recorded on a Shimadzu model QP-1000 spectrometer, and elemental analyses were performed with a Carlo Erba Instruments CHNS-O EA 1108 analyzer.

Carborane was purchased from the Callery Chemical Co. and used without purification. The starting materials, Ni-(COD)₂, PEt₃, and ClSiMe₂H, were purchased from Strem Chemicals. All the aldehydes were purchased from Aldrich. 1,2-Bis(dimethylsilyl)carborane and Ni(PEt₃)₄²² were prepared according to previous described procedures.

Nickel-Catalyzed Reaction of 1 with Various Aldehydes. General Procedure. Compounds 3-14 were prepared by the reaction of 1 with the corresponding aldehydes in the presence of a catalytic amount of Ni(PEt₃)₄. In a typical synthesis, a mixture of isobutyraldehyde (0.04 mL, 0.43 mmol), 1 (0.09 g, 0.36 mmol), and Ni(PEt₃)₄ (0.005 g, 0.018 mmol) in toluene (15 mL) was heated to 80 °C for 12 h. The mixture was passed through a short silica gel column to remove the nickel species from the reaction mixture. The solution was chromatographed using benzene/hexane (1:4) as eluent. Recrystallization from ethanol at -10 °C afforded 5,6-carboranylene-1,1,4,4-tetramethyl-3-isopropyl-2-oxa-1,4-disilacyclohexane **3** in 74% yield: mp 82 °C; ¹H NMR (CDCl₃) δ 3.14 (d, 1H, *J*_{HH} = 8.4 Hz, OC*H*), 1.79 (d, sept, 1H, *J*_{HH} = 8.2 Hz, C*H*), 0.93 (d, 6H, $J_{\rm HH} = 8.2$ Hz, CCH₃), 0.36 (s, 3H, SiCH₃), 0.33 (s, 3H, SiCH₃), 0.31 (s, 3H, SiCH₃), 0.28 (s, 3H, SiCH₃); ${}^{13}C{}^{1}H{}$ NMR (CDCl₃) δ 70.78, 31.86, 19.77, -0.97, -1.10, -3.30, -5.09; ²⁹Si NMR (CDCl₃) δ 11.94, -0.56; MS *m*/*z* 330 [M⁺]; IR (KBr pellet; cm⁻¹) 2968 (m), 2908 (w), 2602 (s), 1471 (m), 1416 (w), 1382 (w), 1361 (w), 1288 (w), 1258 (s), 1218 (w), 1092 (m), 1076 (m), 1043 (s), 982 (w), 942 (w), 878 (m), 820 (m), 801 (m), 692 (m), 662 (w). Anal. Calcd for C₁₀H₃₀B₁₀OSi₂: C, 36.32; H, 9.15. Found: C, 35.94; H, 8.86.

4 from the Reaction with Trimethylacetaldehyde. Compound **4** was prepared by using the same procedure as described for **3** in 66% yield. It was purified by sublimation: mp 123 °C; ¹H NMR (CDCl₃) δ 3.32 (s, 1H, OC*H*), 0.90 (s, 9H, CC*H*₃), 0.38 (s, 3H, SiC*H*₃), 0.37 (s, 3H, SiC*H*₃), 0.35 (s, 3H, SiC*H*₃), 0.33 (s, 3H, SiC*H*₃); ¹³C{¹H} NMR (CDCl₃) δ 74.88, 29.38, 21.45, -0.26, -0.48, -0.92, -1.80; ²⁹Si NMR (CDCl₃) δ 11.36, 0.35 ; MS *m*/*z* 344 [M⁺]; IR (KBr pellet; cm⁻¹) 2972 (m), 2911 (w), 2880 (w), 2602 (s), 1476 (w), 1462 (w), 1415 (w), 1400 (w), 1365 (w), 1258 (m), 1202 (w), 1160 (w), 1090 (m), 1076 (m), 1032 (m), 984 (w), 942 (w), 881 (m), 862 (m), 850 (m), 822 (m), 804 (w), 736 (w), 722 (w), 696 (w), 688 (w), 660 (w). Anal. Calcd for C₁₁H₃₂B₁₀OSi₂: C, 38.33; H, 9.35. Found: C, 37.94; H, 9.06.

5 from the Reaction with Hexanal. Compound **5** was prepared by using the same procedure as described for **3** as a liquid in 38% yield: ¹H NMR (CDCl₃) δ 3.46 (t, 1H, $J_{HH} = 6.6$ Hz, OCH), 1.62–1.22 (m, 8H, CH₂), 0.89 (t, 3H, $J_{HH} = 4.3$ Hz, CH₃), 0.37 (s, 3H, SiCH₃), 0.30 (s, 3H, SiCH₃), 0.22 (s, 3H, SiCH₃), 014 (s, 3H, SiCH₃); ¹³C{¹H} NMR (CDCl₃) δ 77.23, 38.24, 31.82, 28.16, 22.68, 11.14, -0.12, -0.98, -3.10, -6.19; ²⁹Si NMR (CDCl₃) δ 13.08, 0.22; MS *m*/*z* 358 [M⁺]. Anal. Calcd for C₁₂H₃₄B₁₀OSi₂: C, 40.19; H, 9.55. Found: C, 39.79; H, 9.22.

6 from the Reaction with 5-Norbornene-2-carboxaldehyde. Compound **6** was prepared by using the same procedure as described for **3** in 83% yield: mp 168 °C; ¹H NMR (CDCl₃) δ 6.18 (dd, 1H, J_{HH} = 8.40 Hz, J_{HH} = 1.04 Hz), 5.83 (dd, 1H, J_{HH} = 8.40 Hz, J_{HH} = 1.04 Hz), 3.41 (d, 1H, J_{HH} = 4.08 Hz, OCH), 2.17 (s, 2H, CH₂), 1.94–1.05 (m, 5H), 0.33 (s, 9H, SiCH₃), 0.31 (s, 3H, SiCH₃), 0.28 (s, 3H, SiCH₃), 0.25 (s, 3H, SiCH₃); ¹³C{¹H} NMR (CDCl₃) δ 138.42, 131.95 (C=*C*H), 77.03 (O*C*H), 52.68, 43.51, 41.92, 39.27, 36.77, -0.97 (Si-*CH₃*), -1.39 (Si-*C*H₃), -3.06 (Si-*C*H₃), -5.76 (Si-*C*H₃); MS *m*/*z* 380 [M⁺]; IR (KBr pellet; cm⁻¹) 3072 (m), 2954 (s), 2877 (m), 2598 (s), 1448 (w), 1402 (w), 1257 (s), 1087 (m), 1022 (s,br), 983 (w), 942 (w), 908 (w), 870 (m), 822 (s), 802 (w), 744 (w), 682 (w). Anal. Calcd for C₁₄H₃₂B₁₀OSi₂: C, 44.20; H, 8.41. Found: C, 43.94; H, 8.22.

7 from the reaction with benzaldehyde: mp 146 °C; ¹H NMR (CDCl₃) δ 7.37–7.12 (m, 5H, *Ph*), 4.74 (s, 1H, OC*H*), 0.47

⁽²²⁾ Browning, J.; Cundy, C. S.; Green, M.; Stone, F. G. A. *J. Chem. Soc. A* **1969**, 20.

(s, 3H, SiC*H*₃), 0.43 (s, 3H, SiC*H*₃), 0.25 (s, 3H, SiC*H*₃), 0.06 (s, 3H, SiC*H*₃); ${}^{13}C{}^{1}H$ } NMR (CDCl₃) δ 139.61, 128.91, 127.85, 126.89, 126.73, 124.61, 67.86, -0.91, -2.78, -3.78, -6.48; ${}^{29}Si$ NMR (CDCl₃) δ 10.46, -0.72; MS *m*/*z* 364 [M⁺]; IR (KBr pellet; cm⁻¹) 30 (62 (w), 3035 (w), 2924 (m), 2852 (w), 2602 (s), 1494 (w), 1450 (w), 1402 (w), 1258 (s), 1162 (w), 1086 (m), 1078 (m), 1040 (m), 1022 (m), 982 (w), 962 (w), 918 (w), 878 (m), 870 (m), 838 (m), 820 (m), 792 (w), 770 (w), 721 (w), 629 (w), 662 (w). Anal. Calcd for C₁₃H₂₈B₁₀OSi₂: C, 42.85; H, 7.68. Found: C, 42.41; H, 7.88.

8 from the reaction with 4-cyanobenzaldehyde: mp 173 °C; ¹H NMR (CDCl₃) δ 7.64–7.23 (m, 4H, *Ph*), 4.79 (s, 1H, OC*H*), 0.49 (s, 3H, SiC*H*₃), 0.44 (s, 3H, SiC*H*₃), 0.28 (s, 3H, SiC*H*₃), 0.03 (s, 3H, SiC*H*₃); ¹³C{¹H} (NMR (CDCl₃) δ 217.74, 137.27, 129.32, 128.77, 127.45, 127.04, 126.82, 68.04, -0.64, -1.68, -2.04, -5.42; ²⁹Si NMR (CDCl₃) δ 12.59, 0.95; MS *m*/*z* 390 [M⁺]; IR (KBr pellet; cm⁻¹) 3046 (w), 2968 (w), 2918 (w), 2852 (w), 2596 (s), 2238 (m), 1604 (m), 1498 (m), 1408 (w), 1335 (w), 1258 (s), 1204 (w), 1108 (w), 1084 (m), 1061 (s), 1020 (w), 982 (w), 882 (s), 863 (s), 832 (s), 805 (w), 764 (m), 722 (w), 682 (w), 664 (w), 652 (w), 562 (m). Anal. Calcd for C₁₄H₂₇B₁₀-NOSi₂: C, 43.18; H, 6.93. Found: C, 43.52; H, 7.08.

9 from the reaction with methacrolein: ¹H NMR (CDCl₃) δ 6.01 (t, 1H, $J_{\text{HH}} = 2.4$ Hz, OC*H*), 1.65 (d, 2H, $J_{\text{HH}} = 2.4$ Hz, C*H*₂), 1.56 (s, 3H, CC*H*₃), 0.42 (s, 3H, SiC*H*₃), 0.37 (s, 3H, SiC*H*₃), 0.35 (s, 3H, SiC*H*₃), 0.33 (s, 3H, SiC*H*₃); ¹³C{¹H} NMR (CDCl₃) δ 136.03, 107.22, 22.70, 18.38, -0.64, -1.09, -2.54, -2.94; ²⁹Si NMR (CDCl₃) δ 16.42, -1.36; MS *m*/*z* 328 [M⁺]; IR (KBr pellet; cm⁻¹) 2968 (m), 2938 (w), 2855 (w), 2602 (s), 1678 (w), 1455 (w), 1258 (s), 1172 (s), 1136 (w), 1082 (s), 1072 (w), 1042 (s), 864 (w), 850 (w), 822 (br), 802 (s). Anal. Calcd for C₁₀H₂₈B₁₀OSi₂: C, 36.58; H, 8.53. Found: C, 36.32; H, 8.38.

10 from the reaction with *trans*-4-phenyl-3-butyn-2one: mp 145 °C; ¹H NMR (CDCl₃) δ 7.43–7.18 (m, 5H, *Ph*), 6.36 (d, 1H, *J*_{HH} = 9.6 Hz, =*CH*), 2.11 (d, 1H, *J*_{HH} = 9.6 Hz, *CHP*h), 1.53 (s, 3H, CC*H*₃), 0.45 (s, 3H, SiC*H*₃), 0.43 (s, 3H, SiC*H*₃), 0.33 (s, 3H, SiC*H*₃), 0.25 (s, 3H, SiC*H*₃); ¹³C{¹H} NMR (CDCl₃) δ 207.17, 136.99, 134.51, 128.87, 128.55, 127.68, 126.75, 126.48, 71.65, 29.93, 26.03, 1.76, 1.25, 0.58, -2.17; ²⁹Si NMR (CDCl₃) δ 24.72, 6.24; MS *mlz* 404 [M⁺]; IR (KBr pellet; cm⁻¹) 3026 (w), 2971 (m), 2932 (s), 2854 (s), 2602 (s), 1598 (w), 1492 (w), 1462 (m), 1442 (w), 1260 (m), 1182 (m), 1104 (m), 1092 (m), 1036 (w), 987 (s), 962 (m), 862 (m), 823 (m), 722 (m). Anal. Calcd for C₁₆H₃₂B₁₀O Si₂: C, 47.51; H, 7.91. Found: C, 47.34; H, 7.68.

Preparation of Carboranylene-1-dimethyl(hydroxy)silyl-2-(1-phenyl-butan-3-one)dimethylsilane (11). To a stirred toluene solution (10 mL) of 10 (0.18 g, 0.43 mmol) was added H_2O (0.02 g, 1.11 mmol) at room temperature. The solution was stirred for 36 h at that temperature. The product was isolated by chromatography using benzene/hexane (1:8) as eluent in 92% yield: mp 139 °C; 1H NMR (CDCl₃) & 7.36-7.08 (m, 5H, *Ph*), 2.55 (d, 2H, $J_{\rm HH} = 6.8$ Hz, CH₂), 2.35 (t, 1H, $J_{\rm HH} = 6.8$ Hz, CH), 2.16 (s, 1H, OH), 1.25 (s, 3H, CH₃), 0.32 (s, 3H, SiCH₃), 0.22 (s, 3H, SiCH₃), 0.21 (s, 3H, SiCH₃), 0.07 (s, 3H, SiCH₃); ¹³C{¹H} NMR (CDCl₃) & 207.06, 128.73, 128.64, 128.07, 127.87, 127.63, 127.04, 31.02, 29.44, 14.20, 2.42, 1.10, 0.59, -0.13; ²⁹Si NMR (CDCl₃) δ 19.22, -2.36; MS $\mathit{m/z}$ 422 [M⁺]; IR (KBr pellet; cm⁻¹) 3205 (m), 2968 (s), 2601 (s), 1683 (m), 1462 (m), 1384 (w), 1364 (w), 1261 (s), 1082 (m), 1052 (m), 1036 (m), 1017 (m), 910 (w), 864 (w), 855 (w), 804 (s). Anal. Calcd for: C₁₆H₃₄B₁₀O₂Si₂: C, 45.49; H, 8.05. Found: C, 45.18; H. 7.86.

12 from the reaction with α -Methyl-*trans*-cinnamaldehyde. Compound 12 was prepared by using the same procedure as described for **3**. The product was isolated by chromatography using benzene/hexane (1:9) as eluent in 88% yield: mp 158 °C; ¹H NMR (CDCl₃) δ 7.47–7.18 (m, 10H, *Ph*), 6.54 (d, 2H, *J*_{HH} = 1.4 Hz, *CH*=), 5.46 (d, 1H, *J*_{HH} = 1.4 Hz, *CH*O), 1.96 (s, 6H, =CC*H*₃), 0.27 (s, 6H, Si–*CH*₃), 0.08 (s, 6H, Si–*CH*₃); ¹³C{¹H} NMR (CDCl₃) δ 136.82, 136.59, 131.80, 129.22, 129.15, 128.45, 127.41, 71.22, 14.58, 1.17, -0.67; ²⁹Si NMR (CDCl₃) δ 13.6; MS(EI) *m*/*z* 566 [M⁺]; IR (KBr pellet; cm⁻¹) 3088 (w), 3055 (w), 3042 (w), 2968 (m), 2625 (s), 1598 (w), 1479 (w), 1442 (w), 1365 (m), 1261 (s), 1182 (w), 1092 (s), 1048 (s), 1030 (s), 954 (w), 918 (m), 892 (m), 808 (s), 782 (m), 722 (m), 666 (w). Anal. Calcd for C₂₆H₄₂B₁₀O₃Si₂: C, 55.12; H, 7.41. Found: C, 54.88; H, 7.28.

13 from the Reaction with *trans*-**Cinnamaldehyde.** Compound **13** was prepared by using the same procedure as described for **3**. The product was isolated by chromatography using benzene/hexane (1:9) as eluent in 53% yield: mp 56 °C; ¹H NMR (CDCl₃) δ 7.52–6.18 (m, 10H, *Ph*), 6.88 (dd, 2H, *J*_{HH} = 12.3 Hz, *J*_{HH} = 5.8 Hz, =C*H*), 6.74 (dd, 2H, *J*_{HH} = 12.3 Hz, *J*_{HH} = 4.4 Hz, =C*H*), 4.89 (dd, 2H, *J*_{HH} = 5.8 Hz, *J*_{HH} = 4.4 Hz, *CH*), 0.23 (s, 6H, Si–*CH*₃), 0.11 (s, 6H, Si–*CH*₃); ¹³C{¹H} NMR (CDCl₃) δ 162.2, 156.6, 137.4, 133.6, 132.1, 129.1, 128.7, 31.1, -0.46 (Si–*CH*₃), -0.68 (Si–*CH*₃); ²⁹Si NMR (CDCl₃) δ 17.4; MS(EI) *m*/*z* 538 [M⁺]; Anal. Calcd for C₂₄H₃₈B₁₀O₃Si₂: C, 53.52; H, 7.05. Found: C, 53.22; H, 6.83.

14 from the Reaction with Diphenylketene. Compound **14** was prepared by using the same procedure as described for **3**. The product was isolated by chromatography using benzene/hexane (1:1) as eluent in 80% yield: mp 98 °C; ¹H NMR (CDCl₃) δ 7.44–7.12 (m, 10H, *Ph*), 0.19 (s, 6H, Si–*CH*₃), -0.02 (s, 6H, Si–*CH*₃); ¹³C{¹H} NMR (CDCl₃) δ 143.28, 141.18, 136.25, 130.57, 129.24, 128.45, 128.36, 128.25, 127.95, 127.68, 127.43, 127.28, 71.53, -1/26, -1.53; ²⁹Si NMR (CDCl₃) δ 11.97, -6.47; MS(EI) *m*/*z* 468 [M⁺]; IR (KBr pellet; cm⁻¹) 3001 (w), 2950 (w), 2925 (m), 2600 (s), 1460 (m), 1260 (s), 1080 (m), 950 (m), 860 (m), 710 (w). Anal. Calcd for C₂₀H₃₂B₁₀OSi₂: C, 53.09; H, 5.90. Found: C, 52.86; H, 5.72.

X-ray Crystallography. Details of the crystal data and a summary of intensity data collection parameters for **8**, **11**, and **14** are given in Table 1. Crystals of **8**, **11**, and **14** were grown from hexane at -5 °C. Crystals of **8**, **11**, and **14** were mounted in thin-walled glass capillaries and sealed under argon. The data sets of three crystals were collected on an Enraf CAD4 automated diffractometer. Mo K α radiation ($\lambda = 0.7107$ Å) was used for all structure. Each structure was solved by the application of direct methods using the SHELX-96 program and least-squares refinement using SHELX-97. All nonhydrogen atoms in compounds **8**, **11**, and **14** were refined anisotropically. All other hydrogen atoms were included in calculated positions.

Acknowledgment. J.K. wishes to acknowledge the financial support of the Korea Research Foundation (H0181000) made in the program year of 1998.

Supporting Information Available: Tables listing crystallographic information, atomic coordinates and B_{eq} values, anisotropic thermal parameters, and intramolecular bond distances, angles, and torsion angles for **8**, **11**, and **14**. This material is available free of charge via the Internet at http://pubs.acs.org.

OM000527K