

This article was downloaded by: [Chulalongkorn University]

On: 03 January 2015, At: 06:14

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn19>

Synthesis and Biological Activity of 4'-Thio-2'-deoxy Purine Nucleosides

Lea Messini^a, Kamal N. Tiwari^a, John A. Montgomery^a & John A. Secrist III^a

^a Southern Research Institute, P.O. Box 55305, Birmingham, AL, 35255-5305, USA

Published online: 04 Oct 2006.

To cite this article: Lea Messini, Kamal N. Tiwari, John A. Montgomery & John A. Secrist III (1999) Synthesis and Biological Activity of 4'-Thio-2'-deoxy Purine Nucleosides, *Nucleosides and Nucleotides*, 18:4-5, 683-685, DOI: [10.1080/15257779908041540](https://doi.org/10.1080/15257779908041540)

To link to this article: <http://dx.doi.org/10.1080/15257779908041540>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.


This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS AND BIOLOGICAL ACTIVITY OF 4'-THIO-2'-DEOXY PURINE NUCLEOSIDES

Lea Messini, Kamal N. Tiwari, John A. Montgomery, and John A. Secrist III
Southern Research Institute, P.O. Box 55305, Birmingham, AL 35255-5305, USA

ABSTRACT: Coupling of 1-*O*-acetyl-2-deoxy-3,5-di-*O*-toluoyl-4-thio-D-ribofuranose with 6-chloropurine and 2,6-dichloropurine gave a mixture of 9 α and 9 β anomers as major products. These anomers were separated and converted to 2'-deoxy-4'-thio analogues of adenosine, inosine, guanosine, 2-amino-adenosine, and 2-chloro adenosine as well as their α -anomers.

Some years ago we reported our initial efforts on the synthesis of 2'-deoxy-4'-thio purine nucleosides. In that work, we found that formation of the α anomer was strongly favored (1 β :9 α), making the preparation of significant amounts of desired β nucleosides difficult. Herein we present a new coupling method for the synthesis of 2'-deoxy-4'-thio purine nucleosides, one that allows us to obtain reasonable quantities of both α and β anomers of the target nucleosides, which we then evaluated for their anticancer activity.²



Coupling of sugar 1 with 6-chloropurine was carried out using a mixture of SnCl₄ and TMSOTf in the following manner. In a typical experiment, to sugar 1 (1.1 mmol) and 6-chloropurine (1.37 mmol) in 20 mL of CH₂Cl₂ was added consecutively HMDS (1.3

mmol), TMSCl (3.77 mmol). The resulting mixture was stirred at R.T. for 0.5 h and then cooled to 0°C; TMSOTf (0.65 mmol) was added and the reaction mixture was stirred for 1 h at 0°C; then CH₃CN (20 mL) and SnCl₄ (0.65 mmol) were added to it. The reaction mixture was stirred at R.T. for 2 h and concentrated at reduced pressure. After standard work-up and flash chromatography (silica gel) **2α** and **2β** were obtained in 55% yield with a ratio of **2α:2β** (7:3). Treatment of **2α** and **2β** with ethanolic ammonia at 130°C in a bomb for 12 h gave compounds **3α** and **3β** respectively, which were purified on an XAD-4 resin column. Similarly, coupling of sugar **1** with 2,6-dichloropurine using the method described for 6-chloropurine gave **4α** and **4β** in 66% total yield ($\alpha:\beta$ ratio 7:3). Compounds **4α** and **4β** were treated with ethanolic ammonia at 130°C for 24 h in a pressure bomb to produce **5α** and **5β** respectively in approximately 85% yields. Compound **5β** was similar to the authentic sample made earlier in our laboratory by an alternative route.³ Compounds **4α** and **4β** were converted to diazido compounds **6α** and **6β** using sodium azide in 95% ethanol in quantitative yields, which upon reaction with SnCl₂ in MeOH:CH₂Cl₂ (98:2) gave **7α** and **7β** respectively in about 80% yield after purification. Both **7α** and **7β** were deblocked with NaOMe in MeOH at R.T. for 6 h and after purification on an XAD-4 resin column⁴ gave **8α** (85% yield) and **8β** (79% yield). Compound **3β** and **8β** upon treatment with adenosine deaminase at R.T. 16 h gave after purification **9β** (75% yield) and **10β** (80% yield) respectively. Compound **10β** has similar spectral data as reported earlier.⁴ Similarly, **3α** and **8α** were converted to **9α** and **10α** (65% yield) respectively by adenosine deaminase at 37°C for 14 days. All the compounds were characterized by mass spectra, ¹H and ¹³C NMR, UV spectra, and elemental analysis. Anomeric configuration and point of attachment of sugars to purines were derived from NOE experiments on compounds **2α**, **2β** and **5α**, **5β**.

The cell culture cytotoxicity of all ten target compounds was determined against four different human cell lines. Significant cytotoxicity was seen with **5β**, the 2-chloroadenine analog with the β configuration, as reported earlier.³ This compound did not show any significant inhibition of tumor growth at the maximum tolerated doses using the murine colon 36 animal model.

REFERENCES

1. Secrist III, J. A.; Parker, W. B.; Tiwari, K. N.; Messini, L.; Shaddix, S. C.; Rose, L. M.; Bennett Jr., L. L.; Montgomery, J. A. The Synthesis and Biological Activity of Certain 4'-Thio Nucleosides. *Nucleosides Nucleotides*, **1995**, *14*(3-5), 675-686.
2. We wish to thank Prof. Grahame Mackenzie of the University of Hull for sharing with us several years ago a procedure from his laboratory those resulted in a β/α ratio comparable to that reported herein. His procedure employed a carbohydrate similar to **1** (with benzoyl groups at *O*-3 and *O*-5), used an excess of trimethylsilyl triflate, and was successful with 6-chloropurine. We were forced to search for an alternate method when that procedure did not work satisfactorily with 2,6-dichloropurine.
3. Tiwari, K. N.; Secrist III, J. A.; Montgomery, J. A. Synthesis and Biological Activity of 4'-Thio Nucleosides of 2-Chloro Adenine. *Nucleosides Nucleotides*, **1994**, *13*, 1819-1828.
4. Van Draanen, N.A.; Freeman, G. A.; Short, S. A.; Harvey, R.; Jansen, R.; Szczech, G.; Koszalka, G. W. Synthesis and Antiviral Activity of 2'-Deoxy-4'-thio Purine Nucleosides. *J. Med. Chem.* **1996**, *39*(2), 538-542.