


Trans-Selective Asymmetric Aziridination of Diazoacetamides and *N*-Boc Imines Catalyzed by Axially Chiral Dicarboxylic Acid

Takuya Hashimoto, Nanase Uchiyama, and Keiji Maruoka*

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo, Kyoto 606-8502, Japan

Received July 19, 2008; E-mail: maruoka@kuchem.kyoto-u.ac.jp

Enantiomerically pure aziridines constitute an important class of chiral synthons, since they can be easily converted to various types of important building blocks, such as functionalized α - or β -amino acid derivatives.¹ Hence, catalytic asymmetric aziridinations have drawn attention as a means to access these valuable compounds in a straightforward manner. One prevailing strategy for this purpose is the asymmetric nitrogen transfer to alkene by transition-metal catalysis² or organocatalysis.³ Another attractive but less-studied approach to this end is the use of chiral Lewis acid-catalyzed asymmetric aziridination using imine and diazoacetate pioneered by Wulff et al.^{4–6} It is well-known that this sort of aziridination including achiral acid catalysis^{7,8} almost has always dominantly provided *cis* aziridine. In this context, we report herein that axially chiral dicarboxylic acid-catalyzed reaction of *N*-Boc imines and diazoacetamides offers a novel organocatalytic way for the asymmetric aziridination with remarkably high enantioselectivity and unique *trans* selectivity, hitherto unattainable using this strategy.



Chiral Brønsted acid catalysis recently introduced a new paradigm into the acid-catalyzed reaction of imine and diazoacetate. Terada and co-workers reported that chiral phosphoric acid facilitated the reaction of *N*-acyl imines and *tert*-butyl diazoacetate to furnish the Friedel–Craft-type adduct with an excellent level of enantioselectivity (path a).⁹ We revealed the catalytic activity of axially chiral dicarboxylic acid (*R*)-**1e** in the similar reaction using *N*-Boc imines and diazo compounds.¹⁰ The mode of these reactions is considered to involve the hydrogen abstraction of the putative intermediate **A** by the basic site of the catalyst as suggested by Terada. Here, we envisaged that the course of the reaction might be switched in favor of the formation of aziridine by lowering the acidity of the α -proton of diazo carbonyls (path b), thereby realizing unprecedented Brønsted-acid-catalyzed asymmetric aziridination.⁸

As diazo carbonyl derivatives capable of resisting the hydrogen abstraction, we chose diazoacetamides in consideration of the lower acidity of α -proton of amides compared to that of esters. Accordingly, dicarboxylic acid (*rac*)-**1**-catalyzed reaction of several diazoacetamides having different *N,N*-substituents and benzaldehyde *N*-Boc imine was initially investigated (Scheme 1). Although the use of *N,N*-disubstituted diazoacetamides was not found fruitful, we were encouraged by the result of the reaction using *N*-monosubstituted diazoacetamides. Thus, the reaction of *N*-phenyldiazoacetamide with benzaldehyde *N*-Boc imine provided the desired aziridine as a major product. Surprisingly, the *trans* isomer was formed exclusively, in sharp contrast to the previous acid-catalyzed aziridination which gives *cis* aziridine dominantly starting

Scheme 1. Preliminary Investigation of Dicarboxylic-Acid-Catalyzed Aziridination of Diazoacetamides and *N*-Boc Imine




Table 1. Screening of Catalysts (*R*)-**1**^a


entry	catalyst	solvent	2:3 ^c	2 yield ^d (%)	2 ee ^e (%)
1	(<i>R</i>)- 1a	CH ₂ Cl ₂	66:34	15	–19
2	(<i>R</i>)- 1b	CH ₂ Cl ₂	79:21	54	68
3	(<i>R</i>)- 1c	CH ₂ Cl ₂	77:23	34	66
4	(<i>R</i>)- 1d	CH ₂ Cl ₂	79:21	70	92
5	(<i>R</i>)- 1e	CH ₂ Cl ₂	65:35	24	72
6	(<i>R</i>)- 1d	CHCl ₃	78:22	69	95
7	(<i>R</i>)- 1d	toluene	90:10	61	97

^a Reactions were performed with benzaldehyde *N*-Boc imine (0.10 mmol) and *N*-phenyldiazoacetamide (0.12 mmol) in the presence of 5 mol % (*R*)-**1** (0.005 mmol). ^b Determined by ¹H NMR analysis of the crude material. ^c Determined by ¹H NMR analysis of the crude material. ^d Isolated yield. ^e Determined by chiral HPLC analysis.

from diazoacetates and *N*-aryl or *N*-alkyl imines.¹¹ The stereochemical relationship was unambiguously established by X-ray crystallographic analysis (vide infra). Scrutiny of the byproduct led to confirmation of the major component as the 1,2-aryl shift product **3**, whereas a trace amount of 1,2-hydride shift product was also observed (not shown). The aziridination also proceeded with *N*-benzyldiazoacetamide, albeit in diminished yield.


We then set out to investigate the viability of enantiomerically pure dicarboxylic acid (*R*)-**1** in this novel *trans*-selective aziridination as summarized in Table 1. Screening of the reaction with benzaldehyde *N*-Boc imine and *N*-phenyldiazoacetamide in CH₂Cl₂ using 5 mol % of (*R*)-**1** having 3,3'-diaryl substituents led to the identification of 3,3'-dimesityl substituted dicarboxylic acid (*R*)-**1d** as optimal catalyst, giving the *trans* aziridine exclusively in 70% yield with 92% ee (entry 4). Further optimization revealed the use of toluene as solvent of choice (entries 7 vs 6).

The scope of this *trans*-selective asymmetric aziridination was then examined using 5 mol % of (*R*)-**1d** as summarized in Table

Table 2. Scope of Trans-Selective Asymmetric Aziridination^a


entry	Ar ¹	Ar ²	2:3 ^c	2 yield ^d (%)	2 ee ^e (%)
1	Ph	Ph	90:10	61	97
2	4-tolyl	Ph	73:27	51	99
3	3-tolyl	Ph	81:19	52	98
4	2-Np	Ph	82:18	66	99
5	4-MeOC ₆ H ₄	Ph	56:44	<20	N.D.
6	4-PivOC ₆ H ₄	Ph	77:23	49	96
7	3-MeOC ₆ H ₄	Ph	72:28	55	96
8	4-FC ₆ H ₄	Ph	72:28	31	89
9	3-ClC ₆ H ₄	Ph	77:23	50	91
10	Ph	4-MeOC ₆ H ₄	87:13	61	97
11	2-Np	4-MeOC ₆ H ₄	88:12	71	99
12	4-PivOC ₆ H ₄	4-MeOC ₆ H ₄	74:26	57	97
13	Ph	4-ClC ₆ H ₄	85:15	60	97
14	2-Np	4-ClC ₆ H ₄	81:19	70	99


^a Reactions were performed with arylaldehyde *N*-Boc imine (0.10 mmol) and *N*-aryldiazoacetamide (0.12 mmol) in the presence of 5 mol % (*R*)-**1d** (0.005 mmol). ^b Determined by ¹H NMR analysis of the crude material. ^c Determined by ¹H NMR analysis of the crude material. ^d Isolated yield. ^e Determined by chiral HPLC analysis. N.D. = Not determined

**Figure 1.** Ortep representation of **2** (Ar¹ = 3-ClC₆H₄, Ar² = Ph) with ellipsoids shown at 50% probability level. Hydrogen atoms are omitted for clarity.

2. The reaction of 3- and 4-tolualdehyde *N*-Boc imine provided the trans aziridine in moderate yields with high enantioselectivities (entries 2 and 3). 2-Naphthaldehyde *N*-Boc imine was converted to the aziridine in 66% yield with 99% ee (entry 4). Although the use of *N*-Boc imine bearing the electron-donating 4-methoxyphenyl moiety led to poor conversion (entry 5), use of 4-pivaloyloxybenzaldehyde *N*-Boc imine as an alternative gave the aziridine having 4-oxygenated aryl group in 49% yield with 96% ee (entry 6). 3-Methoxybenzaldehyde *N*-Boc imine could be utilized uneventfully (entry 7). As for arylaldehyde *N*-Boc imines bearing electron-withdrawing group, substitution at para position led to the diminished yield regardless of the prolonged reaction time of 8 h (entry 8). The reaction of 3-chlorobenzaldehyde *N*-Boc imine was completed within 5 h, giving the trans aziridine in 50% yield with 91% ee (entry 9). Both electron-donating and withdrawing groups were tolerated as *N*-aryl substituents of diazoacetamide in the reaction with a range of *N*-Boc imines (entries 10–14).

While (*R*)-**1d** catalyzed Friedel–Craft-type reaction proceeded via the attack of diazoacetate from the *si*-face of *N*-Boc imines,¹⁰ this *trans*-selective aziridination furnished the compound stemming from the *re*-face selective nucleophilic addition of diazoacetamide as determined by X-ray crystallographic analysis of **2** (Ar¹ = 3-ClC₆H₄, Ar² = Ph) (Figure 1). This inverted enantiofacial selectivity is indicative of the intervention of a different mechanism as suggested (*vide supra*).

We speculated the *trans* selectivity arises from the preference of a rotamer wherein the carboxamide group and the aryl group of

**Figure 2.** Possible explanation of the observed *trans* selectivity.

N-Boc imine adopt antiperiplanar orientation. Synclinal orientation would be destabilized by the steric repulsion (Figure 2).¹² The hydrogen-bonding between amide *N*–*H* bond and Boc group might act as a secondary factor.¹³

Acknowledgment. This work was partially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan. T.H. thanks a Grant-in-Aid for Young Scientists (Start-up) and GCOE program “Integrated Materials Science” for financial support.

Supporting Information Available: Experimental details and characterization data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) *Aziridines and Epoxides in Organic Synthesis*, Yudin, A. K., Ed.; Wiley-VCH: Weinheim, Germany, 2006. (b) Watson, I. D. G.; Yu, L.; Yudin, A. K. *Acc. Chem. Res.* **2006**, *39*, 194.
- (a) Jacobsen, E. N. in *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer, Berlin, 1999, Vol. 2, p. 607. (b) Müller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905.
- (a) Shen, Y.-M.; Zhao, M.-X.; Xu, J.; Shi, Y. *Angew. Chem., Int. Ed.* **2006**, *45*, 8005. (b) Vesely, J.; Ibrahim, I.; Zhao, G.-L.; Rios, R.; Córdova, A. *Angew. Chem., Int. Ed.* **2007**, *46*, 778.
- (a) Antilla, J. C.; Wulff, W. D. *J. Am. Chem. Soc.* **1999**, *121*, 5099. (b) Antilla, J. C.; Wulff, W. D. *Angew. Chem., Int. Ed.* **2000**, *39*, 4518. (c) Loncaric, C.; Wulff, W. D. *Org. Lett.* **2001**, *3*, 3675. (d) Patwardhan, A. P.; Pulgam, V. R.; Zhang, Y.; Wulff, W. D. *Angew. Chem., Int. Ed.* **2005**, *44*, 6169. (e) Lu, Z.; Zhang, Y.; Wulff, W. D. *J. Am. Chem. Soc.* **2007**, *129*, 7185. (f) Zhang, Y.; Desai, A.; Lu, Z.; Hu, G.; Ding, Z.; Wulff, W. D. *Chem.–Eur. J.* **2008**, *14*, 3785.
- For other chiral Lewis acid catalyzed reactions, see: (a) Juhl, K.; Hazell, R. G.; Jørgensen, K. A. *J. Chem. Soc., Perkin Trans. 1* **1999**, 2293. (b) Redlich, M.; Hossain, M. M. *Tetrahedron Lett.* **2004**, *45*, 8987. (c) Wipf, P.; Lyon, M. A. *ARKIVOC* **2007**, xii, 91.
- For asymmetric aziridination with chiral sulfur ylides, see: (a) Aggarwal, V. K.; Thompson, A.; Jones, R. V. H.; Standen, M. C. H. *J. Org. Chem.* **1996**, *61*, 8368. (b) Aggarwal, V. K.; Alonso, E.; Fang, G.; Ferrara, M.; Hynd, G.; Porcelloni, M. *Angew. Chem., Int. Ed.* **2001**, *40*, 1433.
- For achiral Lewis acid catalyzed reactions, see: (a) Casarrubios, L.; Pérez, J. A.; Brookhart, M.; Templeton, J. L. *J. Org. Chem.* **1996**, *61*, 8358. (b) Gunnoe, T. B.; White, P. S.; Templeton, J. L.; Casarrubios, L. *J. Am. Chem. Soc.* **1997**, *119*, 3171. (c) Gunnoe, T. B.; Surgen, M.; White, P. S.; Templeton, J. L.; Casarrubios, L. *Organometallics* **1997**, *16*, 4865. (d) Rasmussen, K. G.; Jørgensen, K. A. *J. Chem. Soc., Perkin Trans. 1* **1997**, 1287. (e) Rasmussen, K. G.; Juhl, K.; Hazell, R. G.; Jørgensen, K. A. *J. Chem. Soc., Perkin Trans. 2* **1998**, 1347. (f) Ha, H.-J.; Suh, J.-M.; Kang, K.-H.; Ahn, Y.-G.; Han, O. *Tetrahedron* **1998**, *54*, 851. (g) Nagayama, S.; Kobayashi, S. *Chem. Lett.* **1998**, 27, 685. (h) Mayer, M. F.; Hossain, M. M. *J. Org. Chem.* **1998**, *63*, 6839. (i) Mayer, M. F.; Wang, Q.; Hossain, M. M. *J. Organomet. Chem.* **2001**, *630*, 78. (j) Xie, W.; Fang, J.; Li, J.; Wang, P. G. *Tetrahedron* **1999**, *55*, 12929. (k) Sengupta, S.; Mondal, S. *Tetrahedron Lett.* **2000**, *41*, 6245. (l) Kubo, T.; Sakaguchi, S.; Ishii, Y. *Chem. Commun.* **2000**, 625. (m) Akiyama, T.; Ogi, S.; Fuchibe, K. *Tetrahedron Lett.* **2003**, *44*, 4011.
- For achiral Brønsted acid catalyzed *cis*-aziridination using benzhydrylimines, see: Williams, A. L.; Johnston, J. N. *J. Am. Chem. Soc.* **2004**, *126*, 1612. Use of benzhydrylimine in our catalytic system led to the poor conversion and low *trans/cis* selectivity (~1:1).
- Uraguchi, D.; Sorimachi, K.; Terada, M. *J. Am. Chem. Soc.* **2005**, *127*, 9360.
- Hashimoto, T.; Maruoka, K. *J. Am. Chem. Soc.* **2007**, *129*, 10054.
- Trans/cis* ratio was >20/1, as determined by ¹H NMR analysis of the crude material.
- Diazoacetates having a bulky ester moiety are known to give the *trans* aziridine favorably (see ref 7m).
- In the aziridination using sulfur ylides and sulfonylimines, similar internal CH–O hydrogen bonding was indicated by a computational study, see: Robiette, R. *J. Org. Chem.* **2006**, *71*, 2726.

JA805635C