

Heterogeneous & Homogeneous & Bio- & Nano-

CHEM**CAT**CHEM

CATALYSIS

Accepted Article

Title: Bicataltytic multistep reactions en route to the one-pot total synthesis of complex molecules: easy access to chromene and 1,2-dihydroquinoline derivatives from simple substrates

Authors: Pascal D. Giorgi, Peter Miedziak, Jennifer Edwards, Graham Hutchings, and Sylvain Antoniotti

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *ChemCatChem* 10.1002/cctc.201600925

Link to VoR: <http://dx.doi.org/10.1002/cctc.201600925>

WILEY-VCH

www.chemcatchem.org



Bicatalytic multistep reactions en route to the one-pot total synthesis of complex molecules: easy access to chromene and 1,2-dihydroquinoline derivatives from simple substrates.

Pascal D. Giorgi,^[a] Peter J. Miedziak,^[b] Jennifer K. Edwards,^[b] Graham J. Hutchings,^[b] and Sylvain Antoniotti^{[a]*}

Abstract: By combining nanocatalysis and base-catalysis, a novel one-pot multistep process was found for the synthesis of substituted heterocycles of biological relevance from simple substrates. It is based on an initial Au/O₂ oxidation of allylic alcohols followed by a base-catalysed tandem hetero-Michael/aldolisation/crotonisation with *ortho*-hydroxy or *ortho*-amino benzaldehydes. The flexibility of the reaction even allowed the benzaldehyde partner to be prepared in situ in an example of one-pot/5-steps process.

After decades of research in organic synthesis methods dominated by the chemical yield as the only metric to evaluate chemical efficiency, new concepts and paradigms have emerged in the 90's introducing atom economy, waste prevention, and safety among others, as mandatory aspects to consider in the development of sustainable chemical processes.^[1] Catalysis has thus played a prominent role not only because of the amount of waste prevented when switching from a stoichiometric to a catalytic process, but also because of indirect advantages such as generally lower energy demand (both for temperature and pressure settings), higher degree of selectivity, and new reactions design. In tandem reactions, several chemical steps are achieved in the same pot either upon the sequential introduction of reagents and reactants or upon the use of one or more catalysts operating in both synchronous and asynchronous fashion. The step-economy results in a large amount of waste production avoided, because one or more isolation and purification steps are skipped. These approaches were made possible in modern catalytic methods development either with dual catalysis, cooperative catalysis, bifunctional catalysis, or orthogonal multicatalysis.^[2]

Multicatalysis has thus attracted an increasing interest recently in a quest for highly selective and efficient synthetic methodologies with a limited impact in terms of waste generation and resource consumption. Examples of combinations of catalysts of the same type have been reported with two organocatalysts,^[3] or two or more metal catalysts, for example.^[4] More sophisticated cases of orthogonal catalysis were reported by mixing transition metals catalysis with biocatalysis in tandem metathesis/epoxidation reaction,^[5] transition metals and organocatalysis with cross-metathesis/iminium/enamine catalysis,^[6] organo- and N-heterocyclic carbene catalysis,^[7] and more recently photoredox catalysis with organocatalysis^[8] and

with transition metal catalysis,^[9] for example. Conveniently, tandem/domino processes could be triggered the introduction of an electrophilic carbonyl function. Multicatalytic processes based on an oxidation as the first step involved for example a Cu-TEMPO/peptide catalysis,^[10] a Pd NPs-catalysed oxidation/supported amine-catalysed aldolisation,^[11] TPAP/NMO-catalysed oxidation/prolinol derivative-catalysed conjugated addition, oxidation/Knoevenagel condensation catalysed by octahedral MnO₂ molecular sieve,^[12] or oxidation/Biginelli condensation in mechanochemical conditions,^[13] for example. Recently, the one-pot synthesis of flavones was proposed catalysed by Au NPs supported on LDH (Layered Double Hydroxide) for which transient alcohols formed upon an aldol reaction were oxidised and further cyclised.^[14] In different approaches, a whole carbonyl group could be installed for example by hydroformylation of olefins followed an aldol^[15] or acyloin^[16] reaction, or Pd(II)-catalysed aminochloro-carbonylation/In(III)-catalysed Friedel-Crafts reaction of unsaturated amines.^[17] The strategy consisting in the Au NPs/O₂ oxidation of alcohols to generate in situ aldehydes further used as electrophiles has been efficiently used in processes leading to esters, amides, and imines.^[18-20] Oxidation of benzyl alcohols to aldehydes under Au/O₂ catalysis is well established,^[21] but our strategy required moving towards less studied substrates such as allylic alcohols, which were usually oxidised by more sophisticated nanocatalysts and/or pressurised O₂, and simple nanocatalysts.^[20] The use of such selective Au NPs/O₂ oxidative system was envisioned as the best choice to avoid undesired oxidation reactions that should further occur on advanced intermediates or on the final product during tandem processes.

The development of multicatalytic reactions has for prerequisite a good understanding of the individual reactions of concern as well as of their operating conditions (temperature, solvents, rate, conversion, yield, substrates scope). We thus initially focused on the oxidation of benzyl alcohol **1a** as a model substrate, as it is considered a benchmark for oxidation by nanoparticles in the literature.^[18, 22, 23] Selected results of the preliminary screening of the reaction conditions and catalysts are summarised in Table 1 (additional data could be found in SI). The use of small amounts of H₂O₂ or TBHP (5 mol%) as initiators was beneficial to the oxidation.^[24] The best results were obtained with small gold nanoparticles (Au NPs, size below 3 nm) supported on TiO₂, commercially available under the trade name AUROLite®. In particular, these small supported NPs were much more active than Au NPs prepared by impregnation (entries 3, 6, 7). This is likely to be due to the fact that impregnation method often results in bimodal distribution with small (<10 nm) and large (40-70 nm) NPs.^[25] The latter are catalytically inert in our conditions while accounting for a large fraction of the gold content. Reactions in toluene and THF allowed the best results in terms of conversion and selectivity in favour of the formation of

[a] M. P. D. Giorgi, Dr S. Antoniotti
Université Côte d'Azur, CNRS, ICN, Parc Valrose, 06108 Nice
cedex 2, FRANCE.
E-mail: sylvain.antoniotti@unice.fr

[b] Dr P. J. Miedziak, Dr. J. K. Edwards, Prof. Dr G. J. Hutchings
Cardiff Catalysis Institute, School of Chemistry, Cardiff University,
Park Place, Cardiff, CF10 3AT, UNITED KINGDOM.

benzaldehyde **1b**, which was the sole product in our optimised conditions (entry 6). Control experiments were performed to rule out possibilities of hidden homogeneous catalysis by leaching of gold species (entry 8) and thermal reactions (entry 9). We also found that small amounts of **1b** (less than 1%) could be formed by oxidation of toluene used as the solvent (entry 10), but one has to consider that toluene could also be formed at the surface of Au NPs by reduction of benzyl alcohol **1a**.




Table 1. Catalysts screening for the oxidation of benzyl alcohol **1a**.

Entry	Cat. ^a	Additives (x mol%)	Solvent, T °C, Time (h)	Conv [%] ^b
1	AuPd NPs/TiO ₂ ^c	-	THF, 65, 16	0
2	Au NPs/TiO ₂ ^c	-	THF, 65, 16	2
3	Au NPs/TiO ₂ ^c	TBHP (5 mol%)	Toluene, 90, 6	3
4	AuPd NPs/TiO ₂ ^c	TBHP (5 mol%)	Toluene, 90, 6	4
5	Au NPs/Mg(OH) ₂ ^c	TBHP (5 mol%)	Toluene, 90, 6	12
6	AU NPs/TiO ₂ ^d	TBHP (5 mol%)	THF, 65, 16	78
7	AU NPs/TiO ₂ ^d	TBHP (5 mol%)	Toluene, 90, 16	51
8	AuCl ₃	TBHP (5 mol%)	Toluene, 90, 5	0
9	-	TBHP (5 mol%)	Toluene, 90, 16	14
10 ^e	-	-	Toluene, 90, 16	N.A.

^a AuPd/TiO₂: 5% w/w. Au/TiO₂: 5% w/w, 1.94 nm. Au/Mg(OH)₂: 1% w/w. Au/TiO₂ (AURO/ite): 1% w/w, 1.56 nm. Used in 1 mol% catalyst/substrate ratio. ^b Based on GC-TCD quantification by external calibration. ^c Prepared by impregnation. ^d Commercially available under the trade name AURO/ite. ^e Reaction performed in the absence of substrate **1a**. Traces of **1b** were observed (less than 1%), formed upon oxidation of toluene.

We further evaluated these reaction conditions in the selective oxidation of benzylic and allylic alcohols to the corresponding aldehydes by Au NPs supported on metal oxides (only the best results are presented) as shown in Table 2.

Benzylic alcohol **1a** was converted with excellent conversion (99%) and selectivity in **1b** in 87% isolated yields (entry 1). Aliphatic allylic alcohols **2-4a** were similarly oxidised with high selectivities, but with lower conversions (68-71%) and 66-68% isolated yields (entries 2-4). Cinnamyl alcohol **5a** and some derivatives **6-7a** with different substitution patterns on the benzenic ring were efficiently converted to cinnamaldehyde **5b** and related products **8-7b** with excellent conversions (100%) and almost quantitative isolated yields (entries 5-7). A sufficient O₂ supply was obtained by saturation of the solvent by bubbling before use and running the reaction under O₂ (1 atm).

Table 2. Conversion and selectivity in the oxidation of various alcohols.

Entry	Substrate	Conditions ^a	Conv [%]	Product, Isolated yield
1		Au NPs/TiO ₂ H ₂ O ₂ (5 mol%) Toluene, 90 °C	99	 1b , 87%
2		Au NPs/ZnO TBHP (10 mol%) THF, 60 °C	70	 2b , 68%
3		Au NPs/Al ₂ O ₃ TBHP (10 mol%) Toluene, 80 °C	71	 3b , 66%
4		Au NPs/Al ₂ O ₃ TBHP (10 mol%) Toluene, 80 °C	68	 4b , 68%
5		Au NPs/Al ₂ O ₃ TBHP (10 mol%) THF, 60 °C	100	 5b , 99%
6		Au NPs/Al ₂ O ₃ TBHP (10 mol%) THF, 60 °C	100	 6b , 99%
7		Au NPs/Al ₂ O ₃ TBHP (10 mol%) THF, 60 °C	100	 7b , 99%

^a Catalysts were used in 1 mol% catalyst/substrate ratio for reaction times of 1 hrs. Reactions were conducted under an O₂ atmosphere (1 atm).

TEM analysis of the Au NPs used revealed average particle sizes of 1.56 nm for Au/TiO₂, 2.63 nm for Au/Al₂O₃, and 1.54 nm for Au/ZnO with relatively narrow dispersities (Figure 1 and SI). In most instances, optimal results were obtained with Au NPs/Al₂O₃ as catalyst. In previous reports, it was shown that the presence of mineral bases or fluoride ions during the oxidation could be beneficial,^[22] but in our conditions, catalytic testing with K₂CO₃ and KF showed no significant effect. An effect of the basic character of the support could not thus be confirmed. More likely, the physical features of the support, i.e. the specific surface area (200-260 m²/g for Al₂O₃, 40-50 m²/g for TiO₂, 40-50 m²/g for ZnO) or the internal structure (pores, channels ...) could be of concern herein.

With these promising results in hand for allylic alcohols obtained thanks to a relatively mild and flexible protocol, we could further develop a bicatalytic reaction leading to chromenes by mixing with a second catalytic system.





Figure 1. TEM images of Au NPs supported on TiO₂ (a), Al₂O₃ (b) and ZnO (c).

Chromenes are motifs occurring in a large number of natural products exhibiting bioactivity. Daurichromene and confluentin have been isolated from *Rhododendron dauricum* used in traditional Chinese medicine against acute bronchitis^[26, 27] Cordiachromene, isolated from ascidians, is a naturally occurring chromene studied for anti-inflammatory and anticancer properties for which multistep synthesis have been proposed.^[28] Other chromenes such as eulatachromene,^[29] eneccalol angelate and eneccalol methyl ether,^[30] or precocene I and II^[31] have been studied for their diverse biological activities. Chromenes synthesis thus constitutes an important topic of research for which various routes have been described using unsaturated aldehydes as Michael acceptors^[27, 32, 33] or in base catalysed/promoted reactions or involving salicylaldehyde as electrophile.^[34] Organocatalytic asymmetric versions of the reaction have been reported with prolinol derivatives such as Jorgensen's catalyst.^[35] Alternatively, prenylated phenols could also be used as substrates in biomimetic cyclisation^[36] or 2-(1-hydroxyallyl)phenols in Fe(III)-catalysed allylic cyclisation.^[37] Recently, tandem reactions involving a Rh-catalyzed intermolecular hydroacylation of alkynes with salicylaldehydes followed by intramolecular oxo-Michael addition were described for the synthesis of chroman-4-ones.^[38]


Our plan was to use the Au NPs/O₂ catalytic system to generate in situ α,β -unsaturated aldehydes further undergoing a base-catalysed oxa-Michael addition of salicylaldehydes to yield chromenes substituted on positions 2, 3 and 5 in a sequential bicatalytic one-pot/4-steps procedure (Scheme 1). The availability of both unsaturated alcohols and substituted salicylaldehydes^[39] granted the applicability of the approach to access various substitution patterns on the resulting chromenes, potentially useful for structure-activity relationship studies and/or synthesis of bioactive compounds. Replacing salicylaldehyde by *ortho*-aminobenzaldehyde, substituted dihydroquinolines could be similarly obtained efficiently. For substrate **2a**, the use of triazabicyclodecene (TBD) as the base at a catalyst ratio of 30 mol% allowed the formation of chromene **2c** in 64% yield. For nerol (**3a**) and (*E,E*)-farnesol (**4a**), activation with K₂CO₃ (1.1 equiv., added as a methanolic solution) was sufficient and chromenes **3c** and **4c** were obtained in 67 and 59% yields, respectively. With cinnamyl alcohol derivatives **5a–7a**, the use of pyrrolidine (30 mol%) in THF in the presence of molecular sieves (3 Å) was necessary for the reaction to proceed efficiently, and the expected chromenes **5c–7c** were obtained in 86–93% yields. By replacing salicylaldehyde with *ortho*-aminobenzaldehyde, dihydroquinolines **5d** and **6d** could be obtained similarly in 93 and 79% yield, respectively. Variation on

the salicylaldehyde partner was also possible with **3a–5a** as the starting alcohols. Salicylaldehydes substituted on the benzene ring with both electron-donating and withdrawing groups could be used and allowed the formation of the corresponding chromenes in 52–79% isolated yields. It is worth noting that in all these examples, the second step, started upon addition of benzaldehyde derivatives, was performed at room temperature.



Scheme 1. Bicatalytic one-pot/4-steps synthesis of substituted chromenes/dihydroquinolines from allyl alcohols and aromatic aldehydes. Method A: TBD (30 mol%), MS 3 Å, toluene, 60 °C then rt; Method B: K₂CO₃ (1.1 equiv. in MeOH), toluene, 80 °C then rt; Method C: pyrrolidine (30 mol%), MS 3 Å, THF, 60 °C then rt. See supporting information for details.

Since the preparation of *ortho*-aminobenzaldehyde was found to be moderately efficient,^[40] we reasoned that *ortho*-aminobenzyl alcohol could be used as a substrate too and oxidised in situ. Upon sequential oxidation of cinnamyl alcohol and *ortho*-aminobenzyl alcohol, dihydroquinoline **5d** was thus obtained in 85% yield, in an example fitting with the quest for ideal synthesis without purification (Scheme 2).^[41]




Scheme 2. Synthesis of dihydroquinoline **5d** in one-pot from simple alcohols.

In terms of reaction mechanism, and taking into account existing literature on the two core reactions taken separately,^[22, 32] we focused on the possible influence of Au NPs on the hetero-Michael/aldolisation process, the base being added after the oxidation process took place.

First of all, when the reaction depicted in Scheme 1 was performed with **3a** in the absence of base, **3c** was obtained in only 5% yield. In the absence of Au NPs, no oxidation reaction occurred.

With in situ formed Michael acceptors such as citral **3b**, K₂CO₃ was offering a sufficient activation, while for substrates such a cinnamaldehyde **5b**, the formation of a covalent activated intermediate, i.e. an iminium ion, was necessary for the reaction to proceed through iminium/enamine mechanism. A working mechanism for this bicatalytic process is summarised in Scheme 3.



Scheme 3. Working mechanism of the bicatalytic tandem oxidation/hetero-Michael addition/aldolisation/crotonisation reaction

Allylic alcohol **A** could be first deprotonated to the corresponding alcoholate coordinated to the surface of the NP. A redox process would then take place to yield Michael acceptor **B** and an Au NP hydride, further re-oxidised by O₂ to yield HO⁻. Upon activation by K₂CO₃, deprotonated salicylaldehyde would attack intermediate **B** to yield potassium enolate **D**. The latter would cyclise by aldolisation reaction to potassium alcoholate **E**. Upon re-protonation to **F** and dehydration, either thermally-promoted or base-catalysed, chromene **C** would be obtained. In the organocatalytic version, i.e. with pyrrolidine, the reaction would proceed through iminium **B'** and enamine **D'** to form intermediate **E'** hydrolysed to **F** with the recovery of the catalyst. In summary, we have developed a step- and atom-economical bicatalytic tandem process allowing a simplified access to

biologically relevant chromene and 1,2-dihydroquinoline scaffolds with a large space of substitution possibilities. We have shown that up to five reactions could proceed in the same pot to convert simple starting materials into complex molecules by combining nanocatalysis and base-catalysis. These approaches could be further developed in the future toward an ideal goal of one-pot total synthesis.

Acknowledgements

This work was supported by the University Nice Sophia Antipolis, the CNRS, and the ANR program CD2I (Nanocausys, grant 12-

CDII-0010-02). The French MESR is acknowledged for a PhD fellowship to P.D.G. G.J.H., P.J.M. and J.K.E. thank the EPSRC for funding (EP/K014854/1).

Keywords: nanocatalysis • tandem reactions • oxidation • sustainable chemistry • step-economy

- [1] a) P. Anastas, J. Warner, *Green chemistry: theory and practice*, Oxford University Press, Oxford, **1998**, 135. b) M. Poliakoff, P. Anastas, *Nature* **2001**, *413*, 257. c) R. A. Sheldon, *Chem. Commun.* **2008**, 3352-3365. d) B. M. Trost, *Science* **1991**, *254*, 1471-1477.
- [2] T. L. Lohr, T. J. Marks, *Nat. Chem.* **2015**, *7*, 477-482.
- [3] a) X. Li, L. Yang, C. Peng, X. Xie, H.-J. Leng, B. Wang, Z.-W. Tang, G. He, L. Ouyang, W. Huang, B. Han, *Chem. Commun.* **2013**, *49*, 8692-8694. b) R. C. Wende, P. R. Schreiner, *Green Chem.* **2012**, *14*, 1821-1849.
- [4] a) Y. Jeong, Y. Moon, S. Hong, *Org. Lett.* **2015**, asap. b) L. Li, X.-H. Xia, Y. Wang, P. P. Bora, Q. Kang, *Adv. Synth. Catal.* **2015**, *357*, 2089-2097. c) Y. Oonishi, A. Gomez-Suarez, A. R. Martin, Y. Makida, A. M. Z. Slawin, S. P. Nolan, *Chem. - Eur. J.* **2014**, *20*, 13507-13510.
- [5] C. A. Denard, M. J. Bartlett, Y. Wang, L. Lu, J. F. Hartwig, H. Zhao, *ACS Catalysis* **2015**, *5*, 3817-3822.
- [6] B. Simmons, A. M. Walji, D. W. C. MacMillan, *Angew. Chem.* **2009**, *121*, 4413-4417; *Angew. Chem. Int. Ed.* **2009**, *48*, 4349-4353.
- [7] a) G. He, F. Wu, W. Huang, R. Zhou, L. Ouyang, B. Han, *Adv. Synth. Catal.* **2014**, *356*, 2311-2319. b) C. B. Jacobsen, L. Albrecht, J. Udmark, K. A. Joergensen, *Org. Lett.* **2012**, *14*, 5526-5529. c) Y. Liu, M. Nappi, E. C. Escudero-Adan, P. Melchiorre, *Org. Lett.* **2012**, *14*, 1310-1313.
- [8] D. Hager, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2014**, *136*, 16986-16989.
- [9] a) A. Noble, S. J. McCarver, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2015**, *137*, 624-627. b) C. Vila, *ChemCatChem* **2015**, *7*, 1790-1793. c) L. Chu, J. M. Lipshultz, D. W. C. MacMillan, *Angew. Chem.* **2015**, *127*, 8040-8044; *Angew. Chem. Int. Ed.* **2015**, *54*, 7929-7933.
- [10] K. Akagawa, S. Takigawa, E. Mano, K. Kudo, *Tetrahedron Lett.* **2011**, *52*, 770-773.
- [11] N. Nelson, U. Chaudhary, K. Kandel, I. Slowing, *Top. Catal.* **2014**, *57*, 1000-1006.
- [12] B. Sarmah, R. Srivastava, P. Manjunathan, G. V. Shanbhag, *ACS Sustain. Chem. Eng.* **2015**, *3*, 2933-2943.
- [13] P. K. Sahoo, A. Bose, P. Mal, *Eur. J. Org. Chem.* **2015**, *2015*, 6994-6998.
- [14] T. Yatabe, X. Jin, K. Yamaguchi, N. Mizuno, *Angew. Chem.* **2015**, *127*, 13500-13504; *Angew. Chem. Int. Ed.* **2015**, *54*, 13302-13306.
- [15] X. Fang, R. Jackstell, A. Börner, M. Beller, *Chem. Eur. J.* **2014**, *20*, 15692-15696.
- [16] a) K. A. Ostrowski, T. A. Faßbach, A. J. Vorholt, *Adv. Synth. Catal.* **2015**, *357*, 1374-1380. b) K. A. Ostrowski, T. A. Faßbach, D. Vogelsang, A. J. Vorholt, *ChemCatChem* **2015**, *7*, 2607-2613. c) K. Dong, R. Sang, J.-F. Soule, C. Bruneau, R. Franke, R. Jackstell, M. Beller, *Chem. Eur. J.* **2015**, *21*, 18033-18037.
- [17] L. M. Ambrosini, T. A. Cernak, T. H. Lambert, *Tetrahedron* **2010**, *66*, 4882-4887.
- [18] G. L. Brett, P. J. Miedziak, Q. He, D. W. Knight, J. K. Edwards, S. H. Taylor, C. J. Kiely, G. J. Hutchings, *ChemSusChem* **2013**, *6*, 1952-1958.
- [19] a) L. L. Chng, J. Yang, J. Y. Ying, *ChemSusChem* **2015**, *8*, 1916-1925. b) H. Miyamura, H. Min, J.-F. Soulé, S. Kobayashi, *Angew. Chem. Int. Ed.* **2015**, *54*, 7564-7567. c) J.-F. Soulé, H. Miyamura, S. Kobayashi, *Chem. Asian. J.* **2013**, *8*, 2614-2626. d) G. T. Whiting, S. A. Kondrat, C. Hammond, N. Dimitratos, Q. He, D. J. Morgan, N. F. Dummer, J. K. Bartley, C. J. Kiely, S. H. Taylor, G. J. Hutchings, *ACS Catalysis* **2015**, *5*, 637-644.
- [20] W.-J. Yoo, H. Miyamura, S. Kobayashi, *J. Am. Chem. Soc.* **2011**, *133*, 3095-3103.
- [21] A. S. K. Hashmi, G. J. Hutchings, *Angew. Chem.* **2006**, *118*, 8064-8105; *Angew. Chem. Int. Ed.* **2006**, *45*, 7896-7936.
- [22] E. Skupien, R. Berger, V. Santos, J. Gascon, M. Makkee, M. Kreutzer, P. Kooyman, J. Moulijn, F. Kapteijn, *Catalysts* **2014**, *4*, 89-115.
- [23] H. Zou, R. Wang, J. Dai, Y. Wang, X. Wang, Z. Zhang, S. Qiu, *Chem. Commun.* **2015**, *51*, 14601-14604.
- [24] D. I. Enache, J. K. Edwards, P. Landon, B. Solsona-Espriu, A. F. Carley, A. A. Herzing, M. Watanabe, C. J. Kiely, D. W. Knight, G. J. Hutchings, *Science* **2006**, *311*, 362-365.
- [25] a) N. Dimitratos, J. A. Lopez-Sanchez, J. M. Anthonykutti, G. Brett, A. F. Carley, R. C. Tiruvalam, A. A. Herzing, C. J. Kiely, D. W. Knight, G. J. Hutchings, *Phys. Chem. Chem. Phys.* **2009**, *11*, 4952-4961. b) J. K. Edwards, B. E. Solsona, P. Landon, A. F. Carley, A. Herzing, C. J. Kiely, G. J. Hutchings, *J. Catal.* **2005**, *236*, 69-79.
- [26] a) B. Lesch, J. Toröng, M. Nieger, S. Bräse, *Synthesis* **2005**, *2005*, 1888-1900. b) N. Iwata, N. Wang, X. Yao, S. Kitanaka, *J. Nat. Prod.* **2004**, *67*, 1106-1109.
- [27] Y. R. Lee, J. H. Choi, S. H. Yoon, *Tetrahedron Lett.* **2005**, *46*, 7539-7543.
- [28] a) M. Menna, A. Aiello, F. D'Aniello, C. Imperatore, P. Luciano, R. Vitalone, C. Irace, R. Santamaria, *Eur. J. Org. Chem.* **2013**, *2013*, 3241-3246. b) Y. Noda, M. Yasuda, *Helv. Chim. Acta* **2012**, *95*, 1946-1952.
- [29] J. H. Kim, N. Mahoney, K. L. Chan, R. J. Molyneux, B. C. Campbell, *Cur. Microbiol.* **2004**, *49*, 282-287.
- [30] D. Harel, S. A. Khalid, M. Kaiser, R. Brun, B. Wünsch, T. J. Schmidt, *J. Ethnopharmacol.* **2011**, *137*, 620-625.
- [31] M. Szczepanik, R. Obara, A. Szumny, B. Gabrys, A. Halarewicz-Pacan, J. Nawrot, C. Wawrzęczyk, *J. Agric. Food Chem.* **2005**, *53*, 5905-5910.
- [32] D. Lanari, O. Rosati, M. Curini, *Tetrahedron Lett.* **2014**, *55*, 1752-1755.
- [33] M. C. Bröhmer, N. Volz, S. Bräse, *Synlett* **2009**, *2009*, 1383-1386.
- [34] B. Sarmah, G. Baishya, N. Hazarika, P. J. Das, *Synlett* **2014**, *26*, 2151-2155.
- [35] a) T. Govender, L. Hojabri, F. M. Moghaddam, P. I. Arvidsson, *Tetrahedron-Asymmetr.* **2006**, *17*, 1763-1767. b) K. Liu, X. Jiang, *Eur. J. Org. Chem.* **2015**, *2015*, 6423-6428. c) H. Sundén, I. Ibrahim, G.-L. Zhao, L. Eriksson, A. Córdova, *Chem. Eur. J.* **2007**, *13*, 574-581.
- [36] a) K. Takenaka, Y. Tanigaki, M. L. Patil, C. V. L. Rao, S. Takizawa, T. Suzuki, H. Sasai, *Tetrahedron-Asymmetr.* **2010**, *21*, 767-770. b) M. Bobin, I. J. Day, S. M. Roe, E. M. E. Vieux, *Dalton Trans.* **2013**, *42*, 6592-6602.
- [37] L. Calmus, A. Corbu, J. Cossy, *Adv. Synth. Catal.* **2015**, *357*, 1381-1386.
- [38] X.-W. Du, L. M. Stanley, *Org. Lett.* **2015**, *17*, 3276-3279.
- [39] D. Reinhard, L. Schöttner, V. Brosius, F. Rominger, M. Mastalerz, *Eur. J. Org. Chem.* **2015**, *2015*, 3274-3285.
- [40] PCC: 0%; TEMPO/Cu: ~20%; AUROLite/O₂: ~25% isolated yield of ortho-aminobenzaldehyde
- [41] S. Balieu, G. E. Hallett, M. Burns, T. Bootwicha, J. Studley, V. K. Aggarwal, *J. Am. Chem. Soc.* **2015**, *137*, 4398-4403.