

Synthesis of 5'-Methylthio Coformycins: Specific Inhibitors for Malarial Adenosine Deaminase

Peter C. Tyler,^{*,†} Erika A. Taylor,[‡] Richard F. G. Fröhlich,[†] and Vern L. Schramm^{*,‡}

Contribution from the Carbohydrate Chemistry Team, Industrial Research Ltd., Lower Hutt, New Zealand, and Department of Biochemistry, Albert Einstein College of Medicine at Yeshiva University, 1300 Morris Park Avenue, Bronx, New York 10461

Received February 5, 2007; E-mail: p.tyler@irl.cri.nz; vern@aecom.yu.edu

Abstract: Transition state theory suggests that enzymatic rate acceleration ($k_{\text{cat}}/k_{\text{non}}$) is related to the stabilization of the transition state for a given reaction. Chemically stable analogues of a transition state complex are predicted to convert catalytic energy into binding energy. Because transition state stabilization is a function of catalytic efficiency, differences in substrate specificity can be exploited in the design of tight-binding transition state analogue inhibitors. Coformycin and 2'-deoxycoformycin are natural product transition state analogue inhibitors of adenosine deaminases (ADAs). These compounds mimic the tetrahedral geometry of the ADA transition state and bind with picomolar dissociation constants to enzymes from bovine, human, and protozoan sources. The purine salvage pathway in malaria parasites is unique in that *Plasmodium falciparum* ADA (PfADA) catalyzes the deamination of both adenosine and 5'-methylthioadenosine. In contrast, neither human adenosine deaminase (HsADA) nor the bovine enzyme (BtADA) can deaminate 5'-methylthioadenosine. 5'-Methylthio coformycin and 5'-methylthio-2'-deoxycoformycin were synthesized to be specific transition state mimics of the *P. falciparum* enzyme. These analogues inhibited PfADA with dissociation constants of 430 and 790 pM, respectively. Remarkably, they gave no detectable inhibition of the human and bovine enzymes. Adenosine deamination is involved in the essential pathway of purine salvage in *P. falciparum*, and prior studies have shown that inhibition of purine salvage results in parasite death. Inhibitors of HsADA are known to be toxic to humans, and the availability of parasite-specific ADA inhibitors may prevent this side-effect. The potent and *P. falciparum*-specific inhibitors described here have potential for development as antimalarials without inhibition of host ADA.

Introduction

Adenosine deaminase (ADA) deaminates adenosine and deoxyadenosine to form the respective inosines and thereby plays an essential role in purine metabolism.¹ Genetic deficiency of ADA in humans causes severe combined immunodeficiency disease (SCID), which results from the accumulation of dATP in B- and T-cells.^{2–4} Adenosine deaminase is also important in purine salvage pathways in microorganisms and is thought to be essential for salvage pathways in purine auxotrophs like *Plasmodium falciparum*.^{5,6} The adenosine deaminase inhibitors coformycin and 2'-deoxycoformycin (also known as Pentostatin) are natural product transition state analogue inhibitors that bind with picomolar affinity to adenosine deaminases.^{1,7–9} Inhibition of adenosine deaminase results in the accumulation of adenosine

and deoxyadenosine. Production of excess dATP from accumulated deoxyadenosine unbalances the deoxynucleotide pool, inhibits ribonucleotide reductase, and initiates apoptosis in B- and T-cells.¹⁰ Deoxycoformycin has been approved by the FDA for the treatment of hairy cell leukemia since 1991.¹¹ However, it causes renal, liver, pulmonary, and central nervous system toxicities at high doses and is therefore limited in its therapeutic profile.¹²

Recently, the ADA from *P. falciparum* (PfADA) has been characterized and found to catalyze the deamination of adenosine, as well as the conversion of 5'-methylthioadenosine (MTA) to 5'-methylthioinosine (MTI) and ammonia (Figure 1).¹³ 5'-Methylthioinosine was not previously reported as a metabolite in any biological system. The ability of PfADA to utilize MTA

[†] Industrial Research Ltd.

[‡] Albert Einstein College of Medicine at Yeshiva University.

- (1) Agarwal, R. P. *Pharmacol. Ther.* **1982**, *17*, 399–429.
- (2) Giblett, E. R.; Anderson, J. E.; Cohen, F.; Pollara, B.; Meuwissen, H. J. *Lancet* **1972**, *2*, 1067–1069.
- (3) Agarwal, R. P.; Crabtree, G. W.; Parks, R. E., Jr.; Nelson, J. A.; Keightley, R.; Parkman, R.; Rosen, F. S.; Stern, R. C.; Polmar, S. H. *J. Clin. Invest.* **1976**, *57*, 1025–1035.
- (4) Ullman, B.; Gudas, L. J.; Cohen, A.; Martin, D. W., Jr. *Cell* **1978**, *14*, 365–375.
- (5) Webster, H. K.; Weismann, W. P.; Pavia, C. S. *Adv. Exp. Med. Biol.* **1984**, *165*, 225–229.
- (6) Wiesmann, W. P.; Webster, H. K.; Lambros, C.; Kelly, W. N.; Daddona, P. E. *Prog. Clin. Biol. Res.* **1984**, *165*, 325–342.

- (7) Agarwal, R. P.; Spector, T.; Parks, J. R. E. *Biochem. Pharmacol.* **1977**, *26*, 359–367.
- (8) Cha, S.; Agarwal, R. P.; Parks, J. R. E. *Biochem. Pharmacol.* **1975**, *24*, 2187–2197.
- (9) Schramm, V. L. *Arch. Biochem. Biophys.* **2005**, *433*, 13–26.
- (10) Blackburn, M. R.; Kellems, R. E. *Adv. Immunol.* **2005**, *86*, 1–41.
- (11) Published online at: <http://www.accessdata.fda.gov/scripts/cder/onctools/summary.cfm?ID=111>.
- (12) Johnson, S. A. *Clin. Pharmacokinet.* **2000**, *39*, 5–26. Margolis, J.; Grever, M. R. *Semin. Oncol.* **2000**, *27*, 9–14.
- (13) Ting, L. M.; Shi, W.; Lewandowicz, A.; Singh, V.; Mwakwingwe, A.; Birck, M. R.; Taylor Ringia, E. A.; Bench, G.; Madrid, D. C.; Tyler, P. C.; Evans, G. B.; Fumeaux, R. H.; Schramm, V. L.; Kim, K. J. *Biol. Chem.* **2005**, *280*, 9547–9554.




Figure 1. The reaction catalyzed by adenosine deaminase involves nucleophilic attack on adenosine by a hydroxide ion formed by a catalytic site zinc atom. A short-lived Meisenheimer complex intermediate is protonated at the exocyclic amino group, followed by loss of NH₃ to produce inosine. PfADA has substrate specificity that includes both adenosine and 5'-MeS-adenosine, an activity not shared by mammalian ADAs.

is specific to the parasite, and human ADA does not catalyze this reaction. Prior studies have shown that inhibition of malarial purine salvage enzymes results in parasite death,¹⁴ however, potent ADA inhibitors with specificity for the enzymes from malaria parasites have not been described.

5'-Methylthioformycin, 5'-methylthio-2'-deoxycoformycin, and other 5'-functionalized 2'-deoxycoformycin molecules were synthesized and investigated for their ability to inhibit PfADA, as compared to the ADAs from human (HsADA) and bovine (BtADA) sources. The comparative enzymology and inhibitor specificity of malarial and human ADAs highlight the importance of examining substrate specificity in addition to transition state analysis in the design of species-specific inhibitors for use as therapeutic agents.

Experimental Section

Reagents, Bacterial Strains, and Transformation. Calf spleen ADA, human erythrocyte ADA, adenosine, potassium phosphate, and EDTA were purchased from Sigma-Aldrich Chemical Co. LR Clonase II and pDEST-14 vector were both obtained from Invitrogen (Carlsbad, CA). Ni Sepharose High Performance and Source 15Q media were obtained from GE Healthcare (Piscataway, NJ). A diastereomeric mixture of 2'-deoxy-8-(*R/S*)-coformycin was a gift from Dr. David C. Baker. Concentrations of inhibitors were determined spectrophotometrically using the extinction coefficient of 8250 M⁻¹ cm⁻¹ at 282 nm.¹⁵ Enzyme concentrations for BtADA and HsADA were determined by the manufacturer, and enzyme concentrations for PfADA was determined spectrophotometrically using the extinction coefficient of 32340 M⁻¹ cm⁻¹ at 280 nm, as determined by the ExPASy - ProtParam tool. Adenosine concentrations were also determined spectrophotometrically using the extinction coefficient of 14900 M⁻¹ cm⁻¹ at 260 nm. *Escherichia coli* cells BL21 Star-(DE3) and BL21 Star cells (Invitrogen, Carlsbad) were grown in Luria broth (LB) with 100 µg/mL ampicillin (AMP) at 37 °C.

PfADA Gene Construction. A chemically synthesized gene encoding the adenosine deaminase from *P. falciparum* was synthesized, cloned into a pDONR221 vector, and sequenced by DNA 2.0 (Menlo Park). The gene was synthesized using optimized codons for *E. coli* protein expression and the addition of an N-terminal thrombin cleavable (histidine)₆ tag.

Protein Expression and Purification. The designed gene was transferred to the pDEST-14 vector with LR Clonase II. The pDEST-14-PfADA construct was transformed into BL21 Star competent cells, and the protein was expressed without induction with shaking at 37 °C overnight. The cells were ruptured by passage through a French press, the cell debris was pelleted by centrifugation, and the remaining supernatant was purified over a 25 mL Ni Sepharose High Performance His-tag affinity column with elution by a 0–250 mM imidazole gradient. The enzyme was then dialyzed into 100 mM Tris-HCl, 10

µM ZnCl₂, pH 8.0 and frozen in 20 µL aliquots at –80 °C. The protein was equally active with or without the (histidine)₆ tag, which could be removed by combining 1 unit thrombin (GE Healthcare, Piscataway) for every 1 mg of protein in PBS at 22 °C, incubation for 1 h, followed by purification over an 8 mL Source 15Q column with elution by a 0–1 M NaCl gradient in the dialysis buffer.

Enzyme Inhibition Assay. Adenosine deaminase activity was determined by monitoring the conversion of adenosine to inosine via the change in absorbance at 267 nm in assay mixtures containing 20 mM potassium phosphate, pH 7.0, 100 µM adenosine, and 1 µM EDTA at 30 °C.¹⁵ The kinetics for slow-onset inhibition and *K_i* measurement were performed using inhibitor concentrations ranging from 10 µM to 2 nM and an enzyme concentration of 1 nM. The *K_i* values were determined by fitting the initial rate and inhibitor concentrations to the following expression of competitive inhibition:^{16,17}

$$(V_o'/V_o) = (K_M + [S]) / (K_M + [S] + (K_M[I]/K_i))$$

where *V_o'* is the initial rate in the presence of inhibitor, *V_o* is the initial rate in the absence of inhibitor, [I] is the inhibitor concentration, and [S] is the substrate concentration. This expression is valid only under the condition where the inhibitor concentration is 10 times greater than the enzyme concentration. In conditions where inhibitor concentration does not exceed 10 times the enzyme concentration, the effective inhibitor concentration was obtained by the expression:

$$I' = I - (1 - V_o'/V_o)E_t$$

where *I'* is the effective inhibitor concentration, *V_o'* and *V_o* are the initial rates in the presence and absence of inhibitor, and *E_t* is the total enzyme concentration.

In cases where slow onset inhibition was observed, where the inhibitor reached a tighter binding thermodynamic equilibrium with the enzyme, the equilibrium dissociation constant (*K_i'*) was obtained by fitting the rates to the following equation for competitive inhibition:

$$(V_o'/V_o) = (K_M + [S]) / (K_M + [S] + (K_M[I]/K_i'))$$

with [I] concentrations being corrected as above.¹⁸ The *K_M* values for

- (14) Kicska, G. A.; Tyler, P. C.; Evans, G. B.; Furneaux, R. H.; Schramm, V. L.; Kim, K. *J. Biol. Chem.* **2002**, 277, 3226–3231.
- (15) Schramm, V. L.; Baker, D. C. *Biochemistry* **1985**, 24, 641–646.
- (16) Merkler, D. J.; Brenowitz, M.; Schramm, V. L. *Biochemistry* **1990**, 29, 8358–8364.
- (17) Singh, V.; Shi, W.; Evans, G. B.; Tyler, P. C.; Furneaux, R. H.; Almo, S. C.; Schramm, V. L. *Biochemistry* **2004**, 43, 9–18.
- (18) Singh, V.; Evans, G. B.; Lenz, D. H.; Mason, J. M.; Clinch, K.; Mee, S.; Painter, G. F.; Tyler, P. C.; Furneaux, R. H.; Lee, J. E.; Howell, P. L.; Schramm, V. L. *J. Biol. Chem.* **2005**, 280, 18265–18273.

adenosine with the enzymes are $16 \pm 2 \mu\text{M}$ for the bovine enzyme,¹⁵ $29 \pm 3 \mu\text{M}$ for the malarial enzyme,¹³ and $22 \pm 2 \mu\text{M}$ for the human enzyme.¹⁹

(8R)-6-(tert-Butoxycarbonyl)-8-(tert-butyldimethylsilyloxy)-3-(2'-deoxy-3',5'-di-O-toluoyl- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepine (3). (8R)-6-(tert-Butoxycarbonyl)-8-(tert-butyldimethylsilyloxy)-3-(2'-cyanoethyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepine **2a** (0.76 g, 1.81 mmol) was dissolved in dry THF (15 mL), and potassium *tert*-butoxide (1.0 M in THF, 3.6 mL) was added at room temperature. The mixture turned brown immediately and was quenched by addition of glacial acetic acid (206 μL , 3.6 mmol) and coevaporated with toluene (30 mL). The residue was suspended in chloroform/ethyl acetate (5 mL, 1:2 v/v) by ultrasonication and applied on a chromatography column (50 g of silica, chloroform/ethyl acetate = 1:2 v/v, then ethyl acetate), which gave (8R)-6-(tert-butoxycarbonyl)-8-(tert-butyldimethylsilyloxy)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepine **2b** as an amorphous yellow solid (0.51 g, 1.39 mmol, 77%).

2b was dissolved in dry acetonitrile (10 mL) and evaporated in vacuo, flushed with argon, and dissolved again in dry acetonitrile (15 mL). Addition of sodium hydride (60% in mineral oil, 72 mg) gave a visible gas formation. After 30 min, 2-deoxy-3,5-di-O-(*p*-toluoyl)- α -D-erythro-pentofuranosyl chloride **1**²⁰ (0.70 g, 1.81 mmol) was added to the reaction mixture, and a thick heterogeneous slurry was formed. After further 40 min, the reaction mixture was filtered through flux calcined diatomaceous earth and rinsed thoroughly with ethyl acetate. The filtrate was evaporated to dryness in vacuo. Purification of the residue by chromatography (60 g of silica, petroleum ether/ethyl acetate = 2:1 v/v) gave **3** as yellowish oil (0.85 g, 1.18 mmol, 85% calcd. from **2b**). R_f = 0.31 (petrol ether/EtOAc = 3:2 v/v), $[\alpha]_D^{20}$ = -17.4 (c 11.8, chloroform). ¹H NMR: (CDCl₃) δ 0.05 (s, 3H, SiMe), 0.19 (s, 3H, SiMe), 0.88 (s, 9H, Si^tBu), 1.54 (s, 9H, O^tBu), 2.40 (s, 3H, ArMe), 2.43 (s, 3H, ArMe), 2.70 (ddd, J = 2.2, 5.9, 14.2 Hz, 1H, H-2'), 2.84 (ddd, J = 6.4, 8.4, 14.2 Hz, 1H, H-2'), 3.08 (br d, J = 13.6 Hz, 1H, H-7), 4.50 (dd, J = 4.4, 13.6 Hz, 1H, H-7), 4.55–4.60 (m, 1H, H-4'), 4.61–4.65 (m, 2H, H-5'), 5.13 (br d, J = 4.4 Hz, H-8), 5.70 (ddd, J = 2.2, 2.3, 6.0 Hz, 1H, H-3'), 6.42 (dd, J = 5.9, 8.4 Hz, 1H, H-1'), 7.19–7.30 (m, 4H, Ar), 7.57 (s, 1H, H-2), 7.85–7.99 (m, 5H, 4 \times Ar and H-5). ¹³C NMR: (CDCl₃) δ -4.6 (SiMe), -4.2 (SiMe), 18.6 (SiC(CH₃)₃), 22.0 (ArCH₃), 26.2 (SiC(CH₃)₃), 28.5 (OC(CH₃)₃), 38.9 (C-2'), 47.2 (C-7), 64.6 (C-5'), 67.6 (C-8), 75.6 (C-3'), 82.7 (C-4'), 83.7 (OC(CH₃)₃), 84.4 (C-1'), 127.0, 127.2, 129.6, 130.1, 130.2, 132.3 (C-2), 133.1, 134.8, 142.7 (C-5), 144.4, 144.7, 152.8, 166.3, 166.6. HRMS: (MH⁺) calcd for C₃₈H₅₁N₄O₈Si⁺, 719.3476; found, 719.3492.

(8R)-6-(tert-Butoxycarbonyl)-8-(tert-butyldimethylsilyloxy)-3-(2'-deoxy- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d]-[1,3]diazepine (4). In a rubber sealed round-bottom flask, **3** (0.79 g, 1.10 mmol) was treated with ammonia (7 M in methanol, 75 mL) at room temperature. After 17 h, the mixture was evaporated in vacuo, the residue was transferred into an ace pressure tube, fresh ammonia (7 M in methanol, 50 mL) was added, and the tube was sealed and warmed to 45 °C for another 4 h. After evaporation in vacuo, the residue was purified by chromatography (41 g of silica, chloroform/methanol = 15:1 v/v), which gave **4** as colorless oil (375 mg, 71%). R_f = 0.34 (chloroform/methanol = 10:1 v/v), $[\alpha]_D^{20}$ = $+10.0$ (c 5.23, chloroform). ¹H NMR: (CDCl₃) δ 0.06 (s, 3H, SiMe), 0.19 (s, 3H, SiMe), 0.88 (s, 9H, Si^tBu), 1.53 (s, 9H, O^tBu), 2.25 (ddd, J = 1.1, 5.9, 14 Hz, 1H, H-2'), 2.94 (ddd, J = 5.3, 8.7, 14 Hz, 1H, H-2'), 3.08 (br d, J = 13.6 Hz, 1H, H-7), 3.74 (dd, J = 1.3, 12.2 Hz, 1H, H-5'), 3.89 (dd, J = 1.8, 12.2 Hz, 1H, H-5'), 4.10–4.15 (m, 1H, H-4'), 4.54 (dd, J = 4.7, 13.6 Hz, 1H, H-7), 4.65–4.71 (m, 1H, H-3'), 5.14 (d, J = 4.7 Hz, 1H, H-8), 6.20 (dd, J = 5.9, 8.7 Hz, 1H, H-1'), 7.49 (s, 1H, H-2), 7.85 (s, 1H,

H-5). ¹³C NMR: (CDCl₃) δ -4.6 (SiMe), -4.3 (SiMe), 18.7 (SiC(CH₃)₃), 26.2 (SiC(CH₃)₃), 28.5 (OC(CH₃)₃), 41.6 (C-2'), 47.0 (C-7), 63.5 (C-5'), 67.6 (C-8), 73.3 (C-3'), 84.3 (OC(CH₃)₃), 87.3 (C-1'), 88.7 (C-4'), 132.7, 134.9 (C-2), 136.2, 143.3 (C-5), 152.6. HRMS: (MH⁺) calcd for C₂₂H₃₉N₄O₆Si⁺, 483.2639; found, 483.2643.

(8R)-6-(tert-Butoxycarbonyl)-8-(tert-butyldimethylsilyloxy)-3-(2'-deoxy-5'-O-tosyl- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepine (5). **4** (0.375 g, 0.777 mmol) was dissolved in dry pyridine (7 mL, 87 mmol), and tosyl chloride (0.222 g, 1.17 mmol) was added at 0 °C. After 20 min, the mixture was warmed to room temperature and stirred overnight. The reaction mixture was diluted with chloroform (30 mL) and consecutively washed with water (40 mL), citric acid (5% w/w, 3 \times 50 mL), and saturated NaHCO₃ (60 mL), dried (MgSO₄), and evaporated in vacuo. The residue was purified by chromatography (46 g of silica, chloroform/methanol = 20:1 v/v), which gave the title compound as a glass (313 mg, 63%). R_f = 0.39 (chloroform/methanol = 10:1 v/v), $[\alpha]_D^{20}$ = $+25.8$ (c 3.62, CHCl₃). ¹H NMR: (CDCl₃) δ 0.07 (s, 3H), 0.20 (s, 3H), 0.88 (s, 9H), 1.54 (s, 9H), 2.43 (m, 4H), 2.34–2.46 (m, 4H), 2.55–2.66 (m, 2H, 1H D₂O exchangeable), 3.07 (br d, J = 13.5 Hz, 1H), 4.06–4.16 (m, 2H), 4.20–4.29 (m, 1H), 4.48 (dd, J = 4.5, 13.5 Hz, 1H), 4.60–4.70 (m, 1H), 5.12 (br d, J = 4.5 Hz, 1H), 6.30 (t, J = 6.7 Hz, 1H), 7.28–7.34 (m, 2H), 7.48 (s, 1H), 7.70–7.75 (m, 2H), 7.83 (s, 1H). ¹³C NMR: (CDCl₃) δ -4.6 , -4.1 , 18.7, 22.0, 26.2, 28.5, 40.6, 47.2, 67.6, 69.1, 72.2, 83.8, 83.9, 128.4, 130.4, 132.4, 132.6, 133.0, 134.6, 142.6, 145.6, 152.8. HRMS: (MH⁺) calcd for C₂₉H₄₅N₄O₈SSi⁺, 637.2727; found, 637.2753.

(8R)-6-(tert-Butoxycarbonyl)-3-(2'-deoxy-5'-O-tosyl- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepin-8-ol (6). **5** (0.31 g, 0.49 mmol) was dissolved in dry methanol (12 mL), ammonium fluoride (0.54 g, 14.6 mmol) was added, and the mixture was heated to reflux for 6.5 h. After the mixture was cooled to room temperature, stirring was continued overnight. The next morning the mixture was refluxed for an additional 6 h. The reaction mixture was evaporated in vacuo, and the residue was purified by chromatography (27 g of silica, chloroform/methanol = 10:1 v/v), which gave **6** as a hard foam (147 mg, 58%). R_f = 0.23 (chloroform/methanol = 10:1 v/v), $[\alpha]_D^{20}$ = $+30.3$ (c 2.15, chloroform). ¹H NMR: (CDCl₃) δ 1.53 (s, 9H), 2.37–2.64 (m, 6H), 3.59 (br d, J = 13.5 Hz, 1H), 4.03 (dd, J = 5.9, 13.5 Hz, 1H), 4.08–4.17 (m, 3H), 4.18–4.26 (m, 1H), 4.58–4.67 (m, 1H), 5.01 (dd, J = 2.0, 5.9 Hz, 1H), 6.31 (t, J = 6.6 Hz, 1H), 7.27–7.32 (m, 2H), 7.59 (s, 1H), 7.67–7.73 (m, 2H), 7.80 (s, 1H). ¹³C NMR: (CDCl₃) δ 22.0, 28.4, 40.8, 46.8, 65.8, 69.4, 71.6, 84.0, 84.2, 128.3, 130.4, 132.3, 132.6, 133.0, 134.2, 142.9, 145.6, 152.9. HRMS: (MH⁺) calcd for C₂₃H₃₁N₄O₈S⁺, 523.1863; found, 523.1848.

(8R)-3-(2'-Deoxy-5'-methylthio- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepin-8-ol (7). Under argon, **6** (35 mg, 0.067 mmol) was dissolved in dry methanol (3 mL) and treated with sodium thiomethoxide (95% w/w, 0.30 g) at room temperature. After 2 h, the reaction mixture was evaporated in vacuo. Purification of the residue by chromatography (12 g of silica, chloroform/methanolic ammonia (7 M) = 5:1 v/v, 120 mL, 4:1 v/v 100 mL) gave **7** as a white solid (14.5 mg, 73%). R_f = 0.35 (chloroform/methanolic ammonia (7 M) = 4:1 v/v), $[\alpha]_D^{20}$ = $+89$ (c 0.41, methanol). ¹H NMR: (D₂O) δ 2.09 (s, 3H, SMe), 2.48 (ddd, J = 4.1, 6.4, 14.1 Hz, 1H, H-2'), 2.68 (ddd, J = 6.7, 7.2, 14.1 Hz, 1H, H-2'), 2.74 (dd, J = 6.6, 14.1 Hz, 1H, H-5'), 2.82 (dd, J = 5.7, 14.1 Hz, 1H, H-5'), 3.34 (br d, J = 13.5 Hz, 1H, H-7), 3.49 (dd, J = 4.4, 13.5 Hz, 1H, H-7), 4.16 (ddd, J = 3.5, 6.0, 6.2 Hz, 1H, H-4'), 4.52 (m, 1H, H-3'), 5.12 (br d, J = 3.5 Hz, 1H, H-8), 6.25 (t, J = 6.8 Hz, 1H, H-1'), 7.20 (s, 1H), 7.75 (s, 1H). ¹³C NMR: (D₂O) δ 15.8 (SMe), 36.7 (C-5'), 39.1 (C-2'), 47.7 (C-7), 67.2 (C-8), 73.8 (C-3'), 83.2 (C-1'), 85.8 (C-4'), 128.4, 131.4 (C-2), 135.9, 151.0 (C-5). HRMS: (MH⁺) calcd for C₁₂H₁₉N₄O₃S⁺, 299.1178; found, 299.1187. Anal. Calcd for C₁₂H₁₉N₄O₃S \cdot 1.4H₂O: C, 44.54; H, 6.48; N, 17.31; S, 9.91. Found: C, 44.42; H, 6.13; N, 17.19; S, 9.69.

(8R)-3-(2'-Deoxy-5'-propylthio- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepin-8-ol (8). 1-Propanethiol (0.42


(19) The human enzyme sample received from Sigma was characterized to have a k_{cat} of $36 \pm 1 \text{ s}^{-1}$ and a K_M of $22 \pm 2 \mu\text{M}$.

(20) Hoffer, M. *Chem. Ber.* **1960**, *93*, 2777–2781.

mL, 4.64 mmol) was diluted in dry methanol (3 mL), and sodium methoxide (30% w/w in methanol, 0.72 mL) was added. After 5 min at room temperature, **6** (40 mg, 0.077 mmol) was added, and the mixture was stirred overnight at room temperature. The next day the solution was evaporated in vacuo and purified by chromatography (15 g of silica, chloroform/methanolic ammonia (7 M) = 5:1 v/v), which gave the product as a white soapy material (17 mg, 68%). R_f = 0.31 (chloroform/methanolic ammonia (7 M) = 4:1 v/v), $[\alpha]^{20}_D$ = +76 (c 0.85, methanol). ^1H NMR: (CD_3OD) δ 0.97 (t, J = 7.3 Hz, 3H), 1.60 (sext, J = 7.3 Hz, 2H), 2.35 (ddd, J = 3.9, 6.3, 13.5 Hz, 1H, H-2'), 2.47 (dd, J = 6.9, 13.5 Hz, 1H, H-2'), 2.55 (t, J = 7.3 Hz, 2H), 2.76 (dd, J = 5.9, 13.8 Hz, 1H, H-5'), 2.81 (dd, J = 5.9, 13.8 Hz, 1H, H-5'), 3.27–3.35 (m, 1H, H-7), 3.39 (dd, J = 4.4, 13.1 Hz, 1H, H-7), 4.02 (ddd, J = 3.5, 5.9, 5.9 Hz, 1H, H-4'), 4.35–4.43 (m, 1H, H-3'), 4.98–5.05 (m, 1H, H-8), 6.27 (t, J = 6.8 Hz, 1H, H-1'), 7.08 (s, 1H), 7.66 (s, 1H). ^{13}C NMR: (CD_3OD) δ 13.7, 24.0, 35.7, 35.9, 41.4 (C-2'), 49.1 (C-7), 68.4 (C-8), 74.5 (C-3'), 84.4 (C-1'), 87.6 (C-4'), 130.3 (C-2), 131.4, 136.8, 149.8 (C-5). HRMS: (MH^+) calcd for $\text{C}_{14}\text{H}_{23}\text{N}_4\text{O}_3\text{S}^+$, 327.1491; found, 327.1487. Anal. Calcd for $\text{C}_{14}\text{H}_{22}\text{N}_4\text{O}_3\text{S}$: C, 51.51; H, 6.79; N, 17.16. Found: C, 51.49; H, 6.77; N, 16.94.

(8*R*)-3-(2'-Deoxy-5'-phenylthio- β -D-erythro-pentofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-*d*][1,3]diazepin-8-ol (9**).** Under argon, sodium methoxide (30% w/w in methanol, 0.72 mL) was added to a solution of thiophenol (0.47 mL, 4.6 mmol) in dry methanol (3 mL). After 5 min, **6** (39 mg, 0.075 mmol) was added, and the solution was stirred at room temperature for 24 h. Sodium thiomethoxide (95%, 0.32 g) was added to the mixture and stirred for further 3 h 20 min. The reaction mixture was concentrated in vacuo, redissolved in chloroform/methanol (4:1 v/v), and silica (1 g) was added and again concentrated to dryness in vacuo and applied to a column (14 g of silica, chloroform/methanol = 10:1 v/v 200 mL which washed off excess thiophenol, then 4:1 v/v 100 mL eluted the product), which gave compound **9** as an oil (24 mg, 90%). R_f = 0.21 (chloroform/methanol = 4:1 v/v), $[\alpha]^{20}_D$ = +83 (c 0.49, methanol). ^1H NMR: (CD_3OD) δ 2.35 (ddd, J = 3.4, 6.2, 13.6 Hz, 1H, H-2'), 2.51 (ddd, J = 6.2, 7.4, 13.6 Hz, 1H, H-2'), 3.19 (dd, J = 6.0, 13.8 Hz, 1H, H-5'), 3.24 (dd, J = 6.1, 13.8 Hz, 1H, H-5'), 3.27–3.34 (m, 1H, H-7), 3.39 (dd, J = 4.4, 13.2 Hz, 1H, H-7), 4.05 (ddd, J = 3.1, 6.0, 6.1 Hz, 1H, H-4'), 4.41 (ddd, J = 3.0, 3.2, 6.2 Hz, 1H, H-3'), 5.01 (dd, J = 1.4, 4.4 Hz, 1H, H-8), 6.26 (dd, J = 6.2, 7.4 Hz, 1H, H-1'), 7.07 (s, 1H, H-5), 7.14–7.22 (m, 1H), 7.23–7.32 (m, 2H), 7.36–7.43 (m, 2H), 7.59 (s, 1H, H-2). ^{13}C NMR: (CD_3OD) δ 37.7 (C-5'), 41.3 (C-2'), 49.1 (C-7), 68.4 (C-8), 74.6 (C-3'), 84.7 (C-1'), 86.6 (C-4'), 127.3, 130.1, 130.4, 130.6, 131.5 (C-2), 136.8, 137.5, 149.8 (C-5). HRMS: (MH^+) calcd for $\text{C}_{17}\text{H}_{21}\text{N}_4\text{O}_3\text{S}^+$, 361.1334; found, 361.1324. Anal. Calcd for $\text{C}_{17}\text{H}_{20}\text{N}_4\text{O}_3\text{S} \cdot 1.7\text{H}_2\text{O}$: C, 52.21; H, 6.03; N, 14.33. Found: C, 52.01; H, 5.75; N, 14.50.

Allyl 2,3-Di-O-acetyl-5-S-methyl-5-thio- α,β -D-ribofuranoside (10**).** Acetyl chloride (2.0 mL, 28 mmol) was added to a stirred suspension of D-ribose (6.0 g, 40 mmol) in allyl alcohol (100 mL). After 0.5 h, pyridine (5 mL, 61.8 mmol) was added to the clear solution, and it was concentrated to dryness. Chromatography ($\text{CHCl}_3/\text{EtOAc}/\text{MeOH}$ = 5:2:1 v/v) afforded allyl α,β -D-ribofuranoside (7.47 g, 39.3 mmol, 98%) as a syrup. A solution of this material in dry pyridine (60 mL) was treated with 2,4,6-triisopropylbenzenesulfonyl chloride (14.2 g, 46.9 mmol) and then allowed to stand for 2 days. It was diluted with water (1 L) and extracted two times with chloroform. The combined extracts were washed consecutively with water, diluted HCl, and aqueous NaHCO_3 , dried, and concentrated. Chromatography ($\text{EtOAc}/\text{Pet Ether}$ = 1:3 v/v) afforded 9.77 g of major product. A stirred solution of 8.84 g of this material in DMF (30 mL) was treated with sodium thiomethoxide (4.1 g, 58.6 mmol). After 2 h, pyridine (50 mL) and acetic anhydride (50 mL) were added, and the solution was stirred for 16 h before being partitioned between toluene and water. The organic phase was washed with water, diluted HCl, and aqueous NaHCO_3 , dried,

Scheme 1^a

^a Reagents: (i) KOtBu, THF, 77%; (ii) NaH, CH_3CN , 85%; (iii) NH_3 , MeOH, 71%; (iv) TsCl, pyridine, 60%; (v) NH_4F , MeOH, reflux, 58%; (vi) NaSMe, MeOH, 73%; (vii) PrSH, NaOMe, MeOH, 68%; (viii) PhSH, NaOMe, MeOH, then NaSMe, 90%.


and concentrated. Chromatography ($\text{EtOAc}/\text{Pet Ether}$ = 1:4 v/v) gave separately the two anomers of the title compound: first the major compound (4.16 g, 13.7 mmol) followed by the minor (0.73 g, 2.4 mmol). Total yield 40% overall. For the major anomer, ^1H NMR: (CDCl_3) δ 5.94–5.81 (1H, m), 5.34–5.12 (4H, m), 5.03 (1H, s), 4.30–4.19 (2H, m), 4.04–3.97 (1H, m), 2.77 (2H, m), 2.18 (3H, s), 2.10 (3H, s), 2.06 (3H, s). ^{13}C NMR: δ 170.1, 170.0, 133.9, 117.9, 104.7, 80.8, 75.5, 74.8, 68.9, 38.8, 20.9, 16.6. For the minor anomer, ^1H NMR: (CDCl_3) δ 5.95–5.84 (1H, m), 5.34–5.02 (5H, m), 4.35–4.21 (2H, m), 4.12–4.05 (1H, m), 2.87 (1H, dd, J = 14.0, 4.6 Hz), 2.78 (1H, dd, J = 14.0, 5.5 Hz), 2.18 (3H, s), 2.12 and 2.11 (3H each, s). ^{13}C NMR: δ 170.8, 170.2, 134.4, 117.5, 99.8, 81.6, 72.4, 71.3, 68.9, 37.1, 21.1, 20.9, 17.4. HRMS: (MH^+) calcd for $\text{C}_{13}\text{H}_{21}\text{O}_6\text{S}$, 305.1059; found, 305.1072.

8-(*R*)-3-(2,3-Di-O-acetyl-5-S-methyl-5-thio- β -D-ribofuranosyl)-6-tert-butoxycarbonyl-8-(tert-butyldimethylsilyloxy)-3,6,7,8-tetrahydroimidazo[4,5-*d*][1,3]diazepine (12**).** To a solution of **10** (3.48 g, 11.5 mmol) in acetic acid (70 mL) was added tetrakis(triphenylphosphine)palladium(0) (5.3 g, 4.59 mmol), and the solution was stirred under argon at 70 °C. After 1.5 h, more tetrakis(triphenylphosphine)palladium(0) (1.0 g, 0.87 mmol) was added, and after another 0.5 h, the solution was concentrated to dryness. Chromatography ($\text{EtOAc}/\text{Pet Ether}$ = 1:1 v/v) afforded de-allylated material as a syrup (1.9 g, 7.2 mmol, 63%). Trichloroacetonitrile (0.75 mL, 7.5 mmol) and then DBU (4 drops) were added to a stirred solution of this material (0.31 g, 1.17 mmol) in dichloromethane (3 mL). After 0.5 h, the solution was diluted with hexanes and chromatographed ($\text{EtOAc}/\text{hexanes}$ 1:3 with 0.5% Et_3N) to give 0.40 g (0.98 mmol) of syrupy trichloroacetimidate **11**. A solution of this material and **2b** (0.12 g, 0.328 mmol) in dry dichloromethane (5 mL) was cooled in an ice bath, trimethylsilyl trifluoromethanesulfonate (0.10 mL, 0.55 mmol) was added, and the solution was allowed to warm to room temperature. After 1 h, the solution was washed with aqueous NaHCO_3 , dried, and concentrated. Chromatography (1% Et_3N in $\text{EtOAc}/\text{CHCl}_3/\text{hexanes}$ 1:2:4) gave **12**

Table 1. Inhibition Constants of Transition State Analogue Inhibitors of Adenosine Deaminase

inhibitor	BiADA		HsADA		PfADA	
	K_i (nM)	K_i^* (nM)	K_i (nM)	K_i^* (nM)	K_i (nM)	K_i^* (nM)
coformycin	1.1 ± 0.4	0.06 ± 0.01	13.9 ± 3.4	0.11 ± 0.02	0.68 ± 0.07	0.08 ± 0.02
5'-MeS-coformycin (13)	$> 10\,000^a$	ND ^b	$> 10\,000$	ND	2.66 ± 0.13	0.43 ± 0.12
2'-deoxycorformycin	0.39 ± 0.12	0.027 ± 0.004	0.5 ± 0.1	0.026 ± 0.005	8.2 ± 2.9	0.038 ± 0.009
5'-MeS-2'-deoxycorformycin (7)	$> 10\,000$	ND	$> 10\,000$	ND	2.3 ± 0.8	0.73 ± 0.22
5'-PrS-2'-deoxycorformycin (8)	$> 10\,000$	ND	$> 10\,000$	ND	12 ± 1	ND
5'-PhS-2'-deoxycorformycin (9)	$> 10\,000$	ND	$> 10\,000$	ND	61 ± 11	ND

^a K_i values of $> 10\,000$ indicate that assays with $10\ \mu\text{M}$ of the indicated inhibitor exhibited no inhibition under the conditions of the assay. ^b No slow onset phase was detected.

Scheme 2^a

^a Reagents: (i) HCl, AlOH; (ii) triisopropylbenzenesulfonyl chloride, py; (iii) NaSMc, DMF, then py/Ac₂O; (iv) (Ph₃P)₄Pd(0), HOAc; (v) Cl₃CCN, DBU, CH₂Cl₂; (vi) **2b**, TMSOTf, CH₂Cl₂; (vii) Bu₄NF, HOAc, THF; (viii) NaSMc, MeOH.

as a syrup (0.109 g, 0.178 mmol, 54% based on the acceptor). ¹H NMR: (CDCl₃) δ 7.88 (1H, s), 7.60 (1H, s), 6.03 (1H, d, $J = 5.0$ Hz), 5.80 (1H, t, $J = 5.4$ Hz), 5.50 (1H, t, $J = 5.5$ Hz), 5.13 (1H, d, $J = 4.4$ Hz), 4.48 (1H, dd, $J = 13.5, 4.6$ Hz), 4.34 (1H, dd, $J = 13.5, 5.2$ Hz), 3.13 (1H, d, $J = 13.5$ Hz), 2.87 (2H, m), 2.14, 2.10, 2.08 (3H each, s), 1.54 (9H, s), 0.88 (9H, s), 0.19, 0.05 (3H each, s). ¹³C NMR: δ 169.9, 169.7, 152.8, 142.8, 135.0, 133.2, 86.4, 83.9, 82.0, 73.9, 72.8, 67.6, 47.2, 36.9, 28.5, 26.2, 21.0, 20.8, 18.7, 17.4, -4.2, -4.6. HRMS: (MH⁺) calcd for C₂₇H₄₅N₄O₈Si, 613.2727; found, 613.2739.

8-(R)-Hydroxy-3-(5-S-methyl-5-thio- β -D-ribofuranosyl)-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepine (13**).** Tetrabutylammonium fluoride (2 mL, 1 M in THF) was added to a solution of **12** (0.25 g, 0.408 mmol) and acetic acid (0.05 mL) in THF (2 mL), and the solution was allowed to stand for 2 days. The solution was evaporated, and chromatography (5% MeOH in EtOAc v/v) afforded 8-(R)-3-(2,3-di-O-acetyl-5-S-methyl-5-thio- β -D-ribofuranosyl)-6-tert-butoxycarbonyl-8-hydroxy-3,6,7,8-tetrahydroimidazo[4,5-d][1,3]diazepine as a foam (0.143 g, 0.287 mmol, 70%). ¹H NMR: (CDCl₃) δ 7.86 (1H, s), 7.68 (1H, s), 6.06 (1H, d, $J = 5.2$ Hz), 5.78 (1H, t, $J = 5.4$ Hz), 5.48 (1H, t, $J = 5.4$ Hz), 4.98 (1H, br s), 4.35 (1H, dd, $J = 10.4, 5.2$ Hz), 3.94–3.75 (2H, m), 2.94–2.82 (2H, m), 2.16, 2.11, 2.09 (3H each, s), 1.55 (9H, s). ¹³C NMR: δ 170.0, 169.7, 152.7, 143.3, 133.6, 86.4, 84.4, 82.1, 73.9, 72.7, 65.7, 46.7, 36.9, 28.5, 20.9, 20.8, 17.4. Sodium thiomethoxide (0.20 g, 2.85 mmol) was added to a stirred solution of this material (0.14 g, 0.28 mmol) in methanol (3 mL). After 4 h, the solution was concentrated, and chromatography of the residue gave **13** as a white solid (0.062 g, 0.197 mmol, 70%). ¹H NMR: (CD₃OD) δ 8.00 (1H, s), 7.16 (1H, s), 5.93 (1H, d, $J = 4.3$ Hz), 5.01 (1H, dd, $J = 4.5, 2.2$ Hz), 4.43 (1H, t, $J = 4.5$ Hz), 4.23–4.16 (2H, m), 3.46–3.34 (2H, m), 2.94–2.79 (2H, m), 2.15 (3H, s). ¹³C NMR: δ 150.9, 137.5, 131.5, 128.4, 90.2, 85.3, 76.4, 74.1, 67.7, 37.9, 17.1. HRMS: (MH⁺) calcd for C₁₂H₁₈N₄O₄S, 315.1127; found, 315.1130.

Results and Discussion

The recent determination that the adenosine deaminase from *P. falciparum* has catalytic activity for both adenosine and 5'-methylthioadenosine as physiological substrates¹³ led to the investigation of adenosine deaminase inhibitors with 5'-methylthio and related groups. Two potent inhibitors of human adenosine deaminase, coformycin and 2'-deoxycorformycin, were targeted as parent compounds for this investigation. Both inhibitors have picomolar dissociation constants for human adenosine deaminase,^{7–9} and 2'-deoxycorformycin (Pentostatin) was approved by the FDA for treatment of hairy cell leukemia in 1991. The use of Pentostatin as a therapeutic agent has been limited due to its toxicity related to the inhibition of ADA and its incorporation into DNA.¹² Modification of ADA inhibitors to eliminate binding to the human enzyme and to eliminate 5'-phosphorylation for entry into DNA pathways, while retaining powerful inhibition of the malarial enzyme, would allow for therapeutic efficacy against the parasite while eliminating the toxicity of the parent compounds.

Synthesis of 5'-Functionalized Coformycins. Previous syntheses of coformycin and 2'-deoxycorformycin have focused on the preparation of 8-keto-diazepine derivatives followed by reduction of the ketone and separation of the diastereoisomeric 8-(R/S)-alcohols.^{21–23} The chromatographic separation of these isomers is a synthetically inefficient and difficult route. Consequently, we have concentrated on methods to synthesize the

- (21) Chan, E.; Putt, S. R.; Showalter, H. D. H.; Baker, D. C. *J. Org. Chem.* **1982**, *47*, 3457–3464.
- (22) Hawkins, L. D.; Hanvey, J. C.; Boyd, F. L., Jr.; Baker, D. C.; Showalter, H. D. H. *Nucleosides Nucleotides* **1983**, *2*, 479–494.
- (23) Jeanette, T. H.; Riordan, J. M.; Montgomery, J. A. *Nucleosides Nucleotides* **1986**, *5*, 431–439.




Figure 2. Dissociation constants (K_d) for inhibition of ADAs from bovine, human, and *P. falciparum* sources.

desired 8-(*R*)-diazepine specifically. Synthesis of the chiral aglycone **2b** has been reported,²⁴ and we have developed an improved synthesis of this material, the details of which will be published elsewhere. The aglycone was conveniently stored as the *N*-protected **2a**, as **2b** does not survive storage for even a few days.

There have been no reports of the successful glycosylation of **2b**, but we found that a number of 5'-substituted thio-2'-deoxycycoformycin derivatives (**7**, **8**, and **9**) were accessible by using 2-deoxy-di-*O*-toluoylribosyl chloride **1** as the glycosylating agent, which was made from 2-deoxy-D-ribose (Scheme 1). The aglycone **2a** was deprotected at *N*-3 with potassium *tert*-butoxide in THF to give free amine **2b**. This was deprotonated by treatment with sodium hydride followed by coupling with the α -glycosyl chloride **1** under S_N2 conditions.^{25–27} That the isolated anomerically pure nucleoside analogue **3** was the β -anomer depicted was confirmed by NOESY NMR experiments on compounds **7**, **8**, and **9**, which showed interactions between H-1' \leftrightarrow H-4', H-3' \leftrightarrow H-2, and H-5' \leftrightarrow H-2. This product was deacetylated with ammonia in methanol to give diol

Table 2. Kinetic Constants for HsADA, BtADA, and PfADA with Adenosine and 5'-Methylthioadenosine as Substrates^a

organism	adenosine			5'-methylthioadenosine		
	k_{cat} (s ⁻¹)	K_M (μ M)	k_{cat}/K_M (M ⁻¹ s ⁻¹)	k_{cat} (s ⁻¹)	K_M (μ M)	k_{cat}/K_M (M ⁻¹ s ⁻¹)
<i>B. taurus</i>	65	16	1.1×10^6	<0.02	NA ^b	NA
<i>H. sapiens</i>	36.3 ± 1.3	22 ± 3	1.6×10^6	<0.02	NA	NA
<i>P. falciparum</i>	1.8	29	6.2×10^5	15	170	9.6×10^4

^a *H. sapiens* values were determined here; *B. taurus* and *P. falciparum* values and associated errors reported in references 13 and 15. ^b Not applicable because no significant reaction rate is observed.

4. Selective tosylation at the primary hydroxy group afforded tosylate **5** in moderate yield (60%). Desilylation²⁸ then gave diol **6**, the treatment of which with sodium thiomethoxide simultaneously displaced the tosylate and, somewhat surprisingly, cleaved the Boc carbamate to provide methylthio ether **7** in 73% yield from compound **6**. The propylthio analogue **8** was prepared in the same fashion in a similar yield (68%). However, when sodium thiophenoxide was used, the tosylate was displaced, but the Boc group remained unaffected. The addition of sodium thiomethoxide to this reaction mixture removed the Boc group and provided the phenylthio compound **9**, which was isolated in 90% yield in a one-pot procedure.

The successful synthesis of the β -nucleoside derivative **3** above depends on the use of crystalline 2-deoxy- α -ribosyl chloride **1** so that essentially a single anomer of ribosyl chloride is available in solution for the S_N2 reaction. There are no readily available crystalline α -ribosyl halides available, so the 5'-methylthioformycin derivative was prepared using a ribosyl donor with a participating group at *O*-2 to control the anomeric stereochemistry. We found it convenient to use a 5-methylthio-ribose donor and after some experimentation settled on a trichloroacetimidate donor as the most effective. Thus, D-ribose was converted to the allyl 5-*S*-methylribofuranosides **10** (Scheme 2) in 40% overall yield. Deallylation and formation of the trichloroacetimidate donor **11** were followed directly by glycosylation of **2b** in the presence of TmsOTf to give the protected coformycin derivative **12** (54%). The stereochemistry of **12** was confirmed by NMR NOESY experiments. Desilylation with Bu₄NF was complicated by degradation, but when buffered with HOAc gave the desired alcohol in 70% yield. Since we have found that sodium thiomethoxide will remove the Boc carbamate from diazepines such as **12**, it was convenient to use these conditions (NaSMe, MeOH) to effect deacetylation and the Boc removal to give 5'-methylthioformycin **13** in 70% yield.

Inhibition of Adenosine Deaminase by Coformycin Analogues. The inhibition of human (HsADA), bovine (BtADA), and malarial (PfADA) adenosine deaminases was examined with the coformycins described above (Table 1, Figure 2). Both coformycin and 2'-deoxycycoformycin exhibit slow onset inhibition of all three enzymes with the dissociation constants ranging from 27 to 110 pM. None of the 5'-functionalized analogues inhibited the bovine or human enzymes, whereas they all inhibited PfADA. The 5'-methylthio analogues were the most effective inhibitors of PfADA, with slow onset inhibition and dissociation constants in the range of 430–730 pM. The propylthio- and phenylthio-analogues also inhibited

(24) Van Truong, T.; Rapoport, H. *J. Org. Chem.* **1993**, 58, 6090–6096.

(25) Kayzimierzuk, Z.; Binding, U.; Seela, F. *Helv. Chim. Acta* **1989**, 72, 1527–1536.

(26) Kayzimierzuk, Z.; Cotton, H. B.; Revankar, G. R.; Robins, R. K. *J. Am. Chem. Soc.* **1984**, 106, 6379–6382.

(27) Kondo, K. S. H.; Ishibashi, M.; Kobayashi, J. *Tetrahedron* **1992**, 48, 7145–7148.

(28) Zhang, W. R. M. *J. Tetrahedron Lett.* **1992**, 33, 1177–1180.




Figure 3. Amino acid sequence alignment for adenosine deaminases from human, bovine, mouse and two plasmodium sources. Residues highlighted in gray are completely conserved in sequence, those in light and dark blue are similar.³¹ Residues located in the active site (defined on the basis of structural analysis from a superposition of the mouse and *P. yoelii* structures, PDB ID 1ADD and 2AMX, respectively, and being located within 10 Å of bound 1-deazaadenosine) are indicated in red. Those active site residues within 5 Å of the 5'-hydroxyl of 1-deazaadenosine, where identity is not conserved between the mammalian and *Plasmodium* sequences, are highlighted in yellow. The insertion in the mammalian enzymes corresponding to residues 109–119 (MmADA) causes an alteration in the structural orientation of residues 101–127, relative to the malarial enzyme, which is indicated by a black bar.

PfADA, but with no detectable slow onset phase, and these compounds gave dissociation constants²⁹ of 12 and 61 nM, respectively.

The substrate specificity of the ADA enzymes with adenosine and 5'-methylthioadenosine as reactants reveals that the mammalian enzymes, although more proficient at utilization of adenosine than PfADA, are unable to utilize 5'-methylthioadenosine (Table 2). Comparison of the amino acid sequences of PfADA, BtADA, and HsADA with the structurally characterized ADAs from *Mus musculus* (MmADA) and *Plasmodium yoelii* (PyADA) reveals a high level of sequence

conservation, especially of active site residues (Figure 3).³⁰ An exception is in the amino acids that are near the 5'-position, where residues Cys153 and Met155 in MmADA structurally and sequentially align with residues Ile171 and Asp173 in PyADA. Additionally, Tyr129 of PyADA, whose identity is maintained with respect to the mammalian homologue (corresponds to Tyr102 in MmADA), is 1 Å removed from the active site as compared to Tyr102 (corresponding to distances of 5.6 and 4.7 Å, respectively; data not shown). These differences cause an increased pocket size for the 5'-group in PfADA, which

(29) Dissociation constants are defined as the final equilibrium dissociation constant, which corresponds to K_i^* , or K_i if no slow-onset inhibition is observed.

(30) The PDB ID's for the structurally characterized ADAs from *Mus musculus* and *Plasmodium yoelii* are 1ADD and 2AMX, respectively.

(31) Sequences were aligned with the CLUSTALW algorithm using the Gonnet series for amino acid alignment.

is likely to be the structural change that allows for the binding and deamination of the 5'-functionalized compounds.

Conclusions

The substrate specificity differences between mammalian and malarial ADAs have been exploited for the generation of inhibitors specific for ADA from *P. falciparum*. The 5'-methylthio group results in a 5-fold difference in the binding affinity of coformycin and 5'-methylthiocoformycin to PfADA. However, this same change causes a >20 000-fold preference in inhibitor binding to PfADA as compared to the human

and bovine enzymes. The ADA isozyme specificity highlights the importance of examining substrate specificity in addition to transition state features in the design of species-specific inhibitors.

Acknowledgment. This work was supported by research grant AI49512 from the National Institutes of Health and contract C08X0209 from the New Zealand Foundation for Research Science and Technology.

JA0708363