

- (15) A. M. Felix, D. P. Winter, S. S. Wang, I. D. Kulesha, W. R. Pool, D. L. Hane, and H. Sheppard, *J. Med. Chem.*, **17**, 422 (1974).
- (16) (a) M. Brenner and W. Huber, *Helv. Chim. Acta*, **36**, 1109 (1953); (b) R. P. Patel and S. Price, *J. Org. Chem.*, **30**, 3575 (1965).
- (17) J. C. Sheehan and D. D. H. Yang, *J. Am. Chem. Soc.*, **80**, 1154 (1958).
- (18) G. Losse, A. Barth, and K. Jasche, *J. Prakt. Chem.*, **21**, 32 (1963).
- (19) G. Losse, A. Barth, and W. Langenbeck, *Chem. Ber.*, **95**, 918 (1962).
- (20) K. Nishimura, M. Arai, K. Sasahara, and T. Nitani, U.S. Patent 3961041 (June 1, 1976).
- (21) G. M. Tyce, M. D. Muneter, and C. A. Owen, *Mayo Clin. Proc.*, **45**, 438 (1970).
- (22) R. Lavery and K. M. Taylor, *Anal. Biochem.*, **22**, 269 (1968).
- (23) D. Shepard and G. B. West, *J. Physiol.*, **120**, 15 (1953).
- (24) H. J. Schumann, *Arch. Exp. Pharmacol.*, **227**, 566 (1956); **234**, 17 (1958); **236**, 474 (1959).
- (25) K. Imai, M. Sugiura, and Z. Tamura, *Chem. Pharm. Bull.*, **18**, 2134 (1970).
- (26) K. Imai, M. Sugiura, H. Kubo, Z. Tamura, K. Ohya, N. Tsunakawa, K. Hirayama, and H. Narabayashi, *Chem. Pharm. Bull.*, **20**, 759 (1972).
- (27) M. Horiguchi, K. Shimizu, and A. Saita, presented at the 91st Meeting of Pharmaceutical Society of Japan, Fukuoka, April 1971.


Antibacterial Halogenoacetyl Derivatives of Amino Acids and Simple Peptides

Jennifer Goodacre, Leonard Jeffries, John H. C. Naylor,* Roger J. Ponsford, and Irene Stirling

Beecham Pharmaceuticals, Research Division, Brockham Park, Betchworth, Surrey, England. Received May 4, 1977

The vital role of D-alanine and L-lysine in the peptidoglycan crosslinking process in the bacterial cell wall prompted preparation of various small peptides incorporating these amino acids. *N*-Iodoacetyl or -bromoacetyl derivatives of the peptides were then prepared in the hope that they would serve as active-site-directed irreversible inhibitors of cell wall transpeptidases. Certain of the halogenoacetyl dipeptide esters, but not the corresponding free acids, showed slight antistaphylococcal activity. Subsequent structural variation showed that inclusion of D-alanine or L-lysine was not necessary, since antibacterial activity was at least as good when the dipeptide unit was replaced by glycylglycine or by an ω -aminoalkanoic acid. It was concluded that the observed antibacterial activity was probably not due to specific inhibition of a cell wall transpeptidase.

A vital process in the growth and division of bacterial cells is the construction of new peptidoglycan, a macromolecular framework which determines the integrity and shape of the cell wall.¹ The mucopeptide units which comprise the peptidoglycan are elaborated in the interior of the cells, but their assembly involves a peptide cross-linking reaction in the exterior situation. Penicillins and other β -lactam antibiotics are believed to exert their antibacterial action by inhibiting the membrane-bound transpeptidase enzyme or enzymes responsible for this reaction, probably by acylation at or near the active site.² In all bacteria so far studied cross-linking involves a C-terminal D-alanyl-D-alanine component of one peptide chain, the terminal D-alanine being displaced by the amino group of a second peptide chain, the following reaction in staphylococci being typical.



Certain small peptides structurally related to the terminal sequences involved in the cross-linking reaction act as competitive inhibitors of the DD-carboxypeptidase/transpeptidase of certain bacteria, but none of these peptides proved toxic to intact bacteria.³ It occurred to us that such peptides might be converted into effective antibacterial agents by the attachment of a chemically reactive group, which could facilitate irreversible combination with the target enzyme according to Baker's concept⁴ of active-site-directed irreversible enzyme inhibition. Other enzymes which have been irreversibly inhibited by chemically reactive peptide derivatives include elastase,⁵ neurohypophyseal hormone-stimulated adenylate cyclase,⁶ trypsin, and chymotrypsin.⁴ The reactive groups which have proved valuable in designing effective inhibitors include *N*-bromoacetyl or -iodoacetyl groups. In

this paper we describe the incorporation of such groups in certain amino acids or di- and tripeptides. In constructing the initial range of peptides, either D-alanine or L-lysine or both were included, since these two amino acids appear to play a central role in the cross-linking process in the staphylococcal cell wall. Later use was also made of D-glutamic acid, which had proved a useful C-terminus in the previously mentioned³ reversible inhibitors, and of glycylglycine and various ω -aminoalkanoic acids as possible substitutes for a dipeptide unit.

Chemistry. Abbreviations used are those recommended by the IUPAC-IUB Commission on Biochemical Nomenclature Symbols for Amino Acid and Peptides: DCC, dicyclohexylcarbodiimide; Cbz, benzyloxycarbonyl; Bzl, benzyl; Boc, *tert*-butoxycarbonyl; Bu, *tert*-butyl; Phth, phthalimidyl; Suc, succinyl; Ac, acetyl; TosOH, *p*-toluenesulfonic acid.

The protected peptide esters listed in Table I were synthesized by a standard coupling technique using DCC in methylene chloride. They were then converted into the corresponding compounds containing a single free amino group (Table II). For this purpose Cbz groups were removed by catalytic hydrogenation over 10% palladium on charcoal. Boc groups were removed by means of 1 equiv of *p*-toluenesulfonic acid, a procedure⁷ which does not affect *tert*-butyl or benzyl ester functions, and in these cases (18, 22–26, 30–32, 34, 35) the product was frequently isolated as the TosOH salt.

The free amines were then treated with mixed anhydrides prepared from iodoacetic acid or bromoacetic acid to yield the *N*-halogenoacetyl derivatives listed in Table III. Finally the free acids (Table IV) were liberated by treating the *tert*-butyl esters with trifluoroacetic acid. In the case of certain lysine-containing peptides the TFA treatment also served to remove a Boc group and gave the peptides as TFA salts (63–66 and 72).

Table I. N-Protected Peptide Esters

No.	Structure	Yield, %	Mp, °C (solvent) ^a	$[\alpha]^{22}_D$, deg ^b	Formula ^c
1	Cbz-D-Ala-D-Ala-OBu	71	60 (EA-PE)	+54.0	C ₁₈ H ₂₆ N ₂ O ₅
2	Boc-D-Ala-D-Ala-OBzl	68	69 (B-PE)	+53.1	C ₁₈ H ₂₆ N ₂ O ₅
3	Cbz-D-Ala-D-Glu-(OBu) ₂	68	Oil	+27.2	C ₂₄ H ₃₆ N ₂ O ₇
4	Cbz-L-Lys(Boc)-D-Ala-OBu	66	64-65 (B-PE)	+7.3	C ₂₆ H ₄₁ N ₃ O ₇
5	Cbz-L-Lys(Boc)-D-Ala-D-Ala-OBu	54	146-147 (B)	+28.1	C ₂₉ H ₄₆ N ₄ O ₈
6	Boc-L-Lys(Cbz)-D-Ala-D-Ala-OBu	85	95 (EA-PE)	+23.7	C ₂₉ H ₄₆ N ₄ O ₈
7	Boc-L-Lys(Ac)-D-Ala-D-Ala-OBu	82	115-116 (EA-PE)	+31.5	C ₂₃ H ₄₂ N ₃ O ₇
8	Boc-L-Lys(Suc)-D-Ala-D-Ala-OBu	80	69-70 (EA-PE)	+22.3	C ₂₅ H ₄₂ N ₄ O ₈
9	Boc-L-Lys(Phth)-D-Ala-D-Ala-OBu	52	132-133 (EA)	+26.6	C ₂₉ H ₄₂ N ₄ O ₈
10	Boc-L-Lys(Cbz)-D-Ala-D-Glu-(OBu) ₂	70	48	+14.5	C ₃₅ H ₅₆ N ₄ O ₁₀
11	Boc-L-Lys(Phth)-D-Ala-D-Glu-(OBu) ₂	41	130-131 (ET-PE)	+22.7	C ₃₅ H ₅₂ N ₄ O ₁₀
12	Boc-L-Lys(Cbz)-NH(CH ₂) ₄ CO ₂ Bu	97	Oil	+3.1	C ₂₈ H ₄₅ N ₃ O ₇
13	Boc-L-Lys(Phth)-NH(CH ₂) ₄ CO ₂ Bu	64	173-174 (ET-PE)	+1.2	C ₂₈ H ₄₁ N ₃ O ₇
14	Boc-Gly-Gly-OBzl	88	77-79 (B-PE)		C ₁₆ H ₂₂ N ₂ O ₅
15	Cbz-NH(CH ₂) ₄ CO ₂ Me	60	Oil		C ₁₄ H ₁₉ NO ₄
16	Cbz-NH(CH ₂) ₄ CO ₂ Bu	87	Oil		C ₁₇ H ₂₅ NO ₄

^a B, benzene; A, ethyl acetate; ET, ether; PE, petroleum ether (bp 60-80 °C). ^b Rotations were measured at concentrations of 0.9-3.0% in methanol. ^c Analyses (C, H, and N) were within ±0.4% of theory.

Table II. Deprotected Peptide Esters

No.	Structure	Yield, %	Mp, °C (solvent) ^a	$[\alpha]^{22}_D$, deg ^b	Formula ^c
17	D-Ala-D-Ala-OBu	97	Oil	+48.0	C ₁₀ H ₂₀ N ₂ O ₃
18	D-Ala-D-Ala-OBzl·TosOH	82	165-166 (E-PE)	+21.8	C ₂₀ H ₂₆ N ₂ O ₅ S
19	D-Ala-D-Glu-(OBu) ₂	94	Oil	+25.3	C ₁₅ H ₂₈ N ₂ O ₅
20	L-Lys(Boc)-D-Ala-OBu	98	Oil	+25.2	C ₁₈ H ₃₅ N ₃ O ₅
21	L-Lys(Boc)-D-Ala-D-Ala-OBu	91	Oil	+44.5	C ₂₁ H ₄₀ N ₄ O ₆
22	L-Lys(Ac)-D-Ala-D-Ala-OBu·TosOH	90	93-95 (ET*)	+35.5	C ₂₅ H ₄₂ N ₄ O ₅ S
23	L-Lys(Suc)-D-Ala-D-Ala-OBu·TosOH·H ₂ O	65	80 (ET*)	+35.3	C ₂₇ H ₄₀ N ₄ O ₈ S·H ₂ O
24	L-Lys(Phth)-D-Ala-D-Ala-OBu·TosOH·0.5H ₂ O	81	158-160 (E)	+39.1	C ₃₁ H ₄₂ N ₄ O ₅ S·0.5H ₂ O
25	L-Lys(Phth)-D-Ala-D-Glu-(OBu) ₂ ·TosOH	44	97 (ET*)	+27.8	C ₃₇ H ₅₂ N ₄ O ₁₁ S
26	L-Lys(Phth)-NH(CH ₂) ₄ CO ₂ Bu·TosOH	95	131 (EA-PE)	+7.4	C ₃₀ H ₄₁ N ₃ O ₅ S
27	Boc-L-Lys(NH ₂)-D-Ala-D-Ala-OBu·0.5H ₂ O	95	79-80 (ET*)	+25.9	C ₂₁ H ₄₀ N ₄ O ₅ ·0.5H ₂ O
28	Boc-L-Lys(NH ₂)-D-Ala-D-Glu-(OBu) ₂	99	58	+17.4	C ₂₇ H ₅₀ N ₄ O ₅
29	Boc-L-Lys(NH ₂)-NH(CH ₂) ₄ CO ₂ Bu	100	Oil	+3.9	C ₂₀ H ₃₉ N ₃ O ₅
30	Gly-Gly-OBzl·TosOH	57	145-148 (E-PE)		C ₁₈ H ₂₂ N ₂ O ₅ S
31	H ₂ N(CH ₂) ₃ CO ₂ Bzl·TosOH	75	88-89 (EA)		C ₁₈ H ₂₃ NO ₅ S
32	H ₂ N(CH ₂) ₄ CO ₂ Me·TosOH	81	86-87 (EA-PE)		C ₁₃ H ₂₁ NO ₅ S
33	H ₂ N(CH ₂) ₄ CO ₂ Bu	99	Oil		C ₉ H ₁₉ NO ₂
34	H ₂ N(CH ₂) ₄ CO ₂ Bzl·TosOH	60	98-99 (EA-PE)		C ₁₉ H ₂₅ NO ₅ S
35	H ₂ N(CH ₂) ₅ CO ₂ Bzl·TosOH	55	94-95 (EA-PE)		C ₂₀ H ₂₇ NO ₅ S

^a E, ethanol; EA, ethyl acetate; PE, petroleum ether (bp 60-80 °C); ET*, trituration with ether. ^b Rotations were measured at concentrations of 0.5-3.0% in methanol. ^c Analyses were within ±0.4% of theory for C, H, and N and within ±0.5% for S.

Biology. All compounds listed in Tables III and IV were tested for their ability to inhibit growth of *Staphylococcus aureus* and *Escherichia coli* in Sensitivity Test Broth (Oxoid). None of the end products (Table IV) showed significant activity. Surprisingly, however, several of the intermediate esters (Table III) showed appreciable activity, especially against *S. aureus*. On the whole, the dipeptide esters tended to be more active than the tripeptide esters. The few bromoacetyl esters tested (39, 42, and 44) were slightly less active than the iodoacetyl analogues (37, 41, and 43). A further surprising feature of the antibacterial activities listed in Table III is that the constituent amino acids appear to have little importance. Indeed, central portions of the molecules of the most active compounds (52, 56, 57, and 59) are simple nonchiral chains.

The above findings led us to doubt whether the compounds owed their activity to the mechanism originally postulated, i.e., inhibition of cell-wall transpeptidases. We therefore submitted a selection of the compounds comprising one of the most active iodoacetyl dipeptide esters (52), two iodoacetyl tripeptide esters (47 and 48), and seven free iodoacetyl peptides (62, 67-71, and 73) to Dr. P. E. Reynolds (Department of Biochemistry, University of Cambridge, England) who kindly tested their ability to inhibit the action of three DD-carboxypeptidase/transpeptidases in releasing C-terminal D-alanine (estimated

using D-amino acid oxidase) from suitable peptide substrates. Ac₂-L-Lys-D-Ala-D-Ala served as substrate for the *Streptomyces* R61 enzyme and UDP-*N*-acetylmuramyl-L-alanyl-meso-diaminopimeloyl-D-alanyl-D-alanine for the enzymes of *Bacillus stearothermophilus* and *Salmonella typhimurium*. The compounds were tested at concentrations of 10 and 100 µg/mL but in no case was any significant inhibition observed. We are consequently unable to explain the biochemical basis for the observed antibacterial activity of certain of the compounds in Table III.

Experimental Section

Materials and Methods. All amino acids were purchased from Sigma Chemical Co. or Fluka A.G. Basle. Melting points were determined on a Büchi Tottoli melting point apparatus and are uncorrected. All new compounds were characterized by infrared and NMR spectra. Elemental microanalyses were determined to within ±0.4% of theoretical values for C, H, and N; for S and halogen the limits were ±0.5%. Optical rotations were determined on a Perkin-Elmer Model 141 polarimeter. Thin-layer chromatography was performed on silica gel plates in MeOH-DMF-NH₄OH (12:3:5) or MeOH and developed with ninhydrin, bromocresol green (0.05% in methanol), or iodine (1% solution in methanol).

The following preparative procedures are representative. Details of analogues are given in Tables I-IV.

Table III. Halogenoacetyl Peptide Esters

No.	Structure	Yield, %	Mp, °C (solvent) ^a	[α] _D ²² , deg ^b	Formula ^c	Antibacterial act. ^d (MIC in μg/mL)	
						<i>S. aureus</i>	<i>E. coli</i>
36	ICH ₂ CO-D-Ala-OBu	90	Oil	+39.6	C ₈ H ₁₆ INO ₃	31	>500
37	ICH ₂ CO-D-Ala-D-Ala-OBu	45	158-160 (B-PE)	+64.9	C ₁₂ H ₂₁ INO ₄	31	>500
38	ICH ₂ CO-D-Ala-D-Ala-OBzl	76	176 (EA-PE)	+67.3	C ₁₃ H ₂₁ INO ₄	7.5	>500
39	BrCH ₂ CO-D-Ala-D-Ala-OBu	44	148-150 (B-PE)	+67.8	C ₁₂ H ₂₁ BrN ₂ O ₄	62	>500
40	ICH ₂ CO-D-Ala-D-Glu-(OBu) ₂	60	41-42	+41.2	C ₁₃ H ₂₁ INO ₆	31	>500
41	ICH ₂ CO-L-Lys(Boc)-D-Ala-OBu	59	150-151 (B-PE)	+0.5	C ₂₀ H ₃₆ INO ₆	15	>500
42	BrCH ₂ CO-L-Lys(Boc)-D-Ala-OBu	61	140-141 (B-PE)	+2.2	C ₂₀ H ₃₆ BrN ₂ O ₆	62	>500
43	ICH ₂ CO-L-Lys(Boc)-D-Ala-D-Ala-OBu	40	136 (B-PE)	+16.3	C ₂₃ H ₄₁ INO ₇	250	500
44	BrCH ₂ CO-L-Lys(Boc)-D-Ala-D-Ala-OBu	61	125-127 (B)	+25.8	C ₂₃ H ₄₁ BrN ₂ O ₇	500	500
45	ICH ₂ CO-L-Lys(Ac)-D-Ala-D-Ala-OBu	55	162-164 (EA-PE)	+31.3	C ₂₀ H ₃₅ INO ₆	500	>500
46	ICH ₂ CO-L-Lys(NH ₂)-D-Ala-D-Ala-OBu·TosOH	86	103-104 (E-EA)	+12.0	C ₂₂ H ₄₁ INO ₆ S	250	>500
47	ICH ₂ CO-L-Lys(Suc)-D-Ala-D-Ala-OBu	80	167-169 (EA-PE)	+17.2	C ₂₂ H ₃₅ INO ₇	>500	>500
48	ICH ₂ CO-L-Lys(Phth)-D-Ala-D-Ala-OBu	85	180-182 (EA)	+9.7 (HOAc)	C ₂₆ H ₃₃ INO ₇	62	>500
49	ICH ₂ CO-L-Lys(Phth)-D-Ala-D-Glu-(OBu) ₂	50	152-153 (EA)	+11.4	C ₃₃ H ₄₅ INO ₉	62	>500
50	ICH ₂ CO-L-Lys(Phth)-NH(CH ₂) ₄ CO ₂ Bu·0.5H ₂ O	60	138 (EA)	-13.5	C ₂₃ H ₃₄ INO ₆	>500	>500
51	Boc-L-Lys(COCH ₂)-D-Ala-D-Ala-OBu	65	85 (EA-PE)	+21.4	C ₂₃ H ₄₁ INO ₇	125	>500
52	ICH ₂ CO-Gly-Gly-OBzl	75	171-172 (E-EA)		C ₁₃ H ₁₉ INO ₄	0.9	62
53	ICH ₂ CO-Gly-OBzl	75	111 (B)		C ₁₁ H ₁₇ INO ₃	7.5	125
54	ICH ₂ CO-NH(CH ₂) ₂ CO ₂ Bzl	60	36-38 (PE*)		C ₁₂ H ₁₉ INO ₃	7.5	500
55	ICH ₂ CO-NH(CH ₂) ₃ CO ₂ Bzl	40	50 (EA-PE)		C ₁₃ H ₂₁ INO ₃	7.5	250
56	ICH ₂ CO-NH(CH ₂) ₄ CO ₂ Bzl	40	45-46 (EA-PE)		C ₁₄ H ₂₃ INO ₃	<0.4	62
57	ICH ₂ CO-NH(CH ₂) ₄ CO ₂ Me	75	42 (EA-PE)		C ₈ H ₁₅ INO ₃	3.7	7.5
58	ICH ₂ CO-NH(CH ₂) ₄ CO ₂ Bu	70	58 (EA-PE)		C ₁₁ H ₂₀ INO ₃	7.5	500
59	ICH ₂ CO-NH(CH ₂) ₅ CO ₂ Bzl	60	53-54 (EA-PE)		C ₁₃ H ₂₃ INO ₃	3.7	62

^a B, benzene; E, ethanol; EA, ethyl acetate; PE, petroleum ether (bp 60-80 °C); PE*, trituration with petroleum ether. ^b Rotations were measured at concentrations of 0.5-3.0% in methanol unless indicated otherwise. ^c Analyses were within ±0.4% of theory for C, H, and N and within ±0.5% for halogens. ^d 0.5-mL vol of Sensitivity Test Broth (Oxoid Ltd., Basingstoke, England) inoculated with 0.2 mL of 10⁻⁴ dilution of 18-h broth culture. Minimum inhibitory concentration (MIC) recorded after 18-h incubation.

Table IV. Halogenoacetyl Peptides

No.	Structure	Yield, %	Mp, °C (solvent) ^a	[α] _D ²² , deg ^b	Formula ^c	Antibacterial act. ^d (MIC in μg/mL) ^e	
						<i>S. aureus</i>	<i>E. coli</i>
60	ICH ₂ CO-D-Ala-D-Ala	90	175 (ET*)	+60.0	C ₈ H ₁₃ INO ₄	500	>500
61	BrCH ₂ CO-D-Ala-D-Ala	85	164 (ET*)	+68.0	C ₈ H ₁₃ BrN ₂ O ₄	500	>500
62	ICH ₂ CO-D-Ala-D-Glu	80	Oil	+46.9	C ₁₀ H ₁₅ INO ₆	500	>500
63	ICH ₂ CO-L-Lys(NH ₂)-D-Ala-CF ₃ CO ₂ H	62	55 (ET*)	-12.0	C ₁₂ H ₂₁ F ₃ INO ₆	>500	>500
64	BrCH ₂ CO-L-Lys(NH ₂)-D-Ala-CF ₃ CO ₂ H	78	51-53 (ET*)	-11.3	C ₁₃ H ₂₁ BrF ₃ N ₂ O ₆	>500	>500
65	ICH ₂ CO-L-Lys(NH ₂)-D-Ala-D-Ala-CF ₃ CO ₂ H	90	190 (ET*)	+2.0 (HOAc)	C ₁₆ H ₂₆ F ₃ INO ₇	>500	>500
66	BrCH ₂ CO-L-Lys(NH ₂)-D-Ala-D-Ala-CF ₃ CO ₂ H	100	100 (ET*)	-26.6 (HOAc)	C ₁₆ H ₂₆ BrF ₃ N ₂ O ₇	>500	>500
67	ICH ₂ CO-L-Lys(Ac)-D-Ala-D-Ala	98	212 (E-ET)	+4.1	C ₁₆ H ₂₇ INO ₆	>500	>500
68	ICH ₂ CO-L-Lys(Suc)-D-Ala-D-Ala	90	125-127 (ET*)	+3.6 (HOAc)	C ₁₈ H ₂₇ INO ₇	500	>500
69	ICH ₂ CO-L-Lys(Phth)-D-Ala-D-Ala	98	189-190 (EA)	+5.9 (HOAc)	C ₂₄ H ₃₇ INO ₇	500	>500
70	ICH ₂ CO-L-Lys(Phth)-D-Ala-D-Glu	90	135 (ET*)	+6.8	C ₂₄ H ₃₉ INO ₉	250	>500
71	ICH ₂ CO-L-Lys(Phth)-NH(CH ₂) ₄ CO ₂ H	85	124 (ET*)	-13.6	C ₃₁ H ₄₆ INO ₆	>500	>500
72	H ₂ N-L-Lys(COCH ₂)-D-Ala-D-Ala-CF ₃ CO ₂ H	80	110 (E-ET)	+31.9 (HOAc)	C ₁₆ H ₂₆ F ₃ INO ₇	500	>500
73	ICH ₂ CONH(CH ₂) ₅ CO ₂ H	60	94-95 (ET)		C ₇ H ₁₂ INO ₃	125	500

^a E, ethanol; EA, ethyl acetate; ET, ether; ET*, trituration with ether. ^b Rotations were measured at concentrations of 0.5-3.0% in methanol unless indicated otherwise. ^c Analyses were within ±0.4% of theory for C, H, and N and within ±0.5% for halogens. ^d 0.5-mL vol of Sensitivity Test Broth (Oxoid Ltd., Basingstoke, England) inoculated with 0.2 mL of 10⁻⁴ dilution of 18-h broth culture. Minimum inhibitory concentration (MIC) recorded after 18-h incubation.

N-Protected Peptide Esters (Table I). The N-protected amino acid (0.01 mol) was dissolved in dry CH_2Cl_2 (50 mL) and a solution of amino acid ester *p*-tosylate (0.01 mol) in dry CH_2Cl_2 (25 mL) containing N-methylmorpholine (0.01 mol) was added. The solution was cooled to 0 °C and dicyclohexylcarbodiimide (DCC) (0.01 mol) was added. The solution was allowed to warm to room temperature, stirred overnight, and filtered, and the filtrate was washed with 20% citric acid solution (2 × 50 mL) and 3% sodium bicarbonate (3 × 50 mL). The organic phase was dried over MgSO_4 , the solvent evaporated, and the residual oil crystallized from the appropriate solvent (Table I).

Removal of N-Protecting Groups (Table II). (a) The N-Cbz peptide *tert*-butyl ester (0.01 mol) was dissolved in MeOH (10 mL) and hydrogenated at ambient temperature and pressure over 10% Pd/C (500 mg). When the theoretical amount of hydrogen had been adsorbed, the solution was filtered through Kieselguhr and the solvent evaporated to leave the free amine.

(b) The N-Boc peptide ester (0.01 mol) in EtOH (50 mL) was cooled to 0 °C and treated dropwise with *p*-toluenesulfonic acid (0.01 mol) in EtOH (10 mL). The solution was allowed to warm to room temperature; then the solvent was evaporated and the residual oil triturated with ether to give the TosOH salt of the amine.

Introduction of N-Halogenoacetyl Groups (Table III). Iodo- or bromoacetic acid (0.01 mol) in dry CH_2Cl_2 (50 mL) containing N-methylmorpholine (0.01 mol) was cooled to -5 °C

and isobutyl chloroformate (0.01 mol) was added over 15 min. A solution of the amine (0.01 mol) (or of the amine TosOH salt plus N-methylmorpholine) in CH_2Cl_2 was added portionwise at -5 °C and the mixture was stirred thus for 30 min, then allowed to warm to room temperature, and washed with 20% citric acid followed by 3% sodium bicarbonate solution. The organic phase was dried (MgSO_4) and evaporated to leave the desired amide.

Halogenoacetyl Peptides (Table IV). The N-halogenoacetyl peptide *tert*-butyl ester (0.01 mol) was stirred with trifluoroacetic acid (2 mL) at room temperature for 1 h. Dry ether (50 mL) was added, the suspension was stirred for a further hour, and the solid product was collected by filtration.

References and Notes

- (1) M. J. Osborn, *Annu. Rev. Biochem.*, **38**, 501 (1969).
- (2) P. M. Blumberg and J. L. Strominger, *Bacteriol. Rev.*, **38**, 291 (1974).
- (3) N. Nieto, H. R. Perkins, M. Leyh-Bouille, J.-M. Frere, and J.-M. Ghysen, *Biochem. J.*, **131**, 163, 707 (1973).
- (4) B. R. Baker, "Design of Active-Site-Directed Irreversible Enzyme Inhibitors", Wiley, New York, N.Y., 1967.
- (5) R. C. Thompson and E. R. Blout, *Biochem. J.*, **12**, 44 (1973).
- (6) R. Walter, I. L. Schwartz, O. Hechter, T. Dousa, and P. L. Hoffman, *Endocrinology*, **91**, 39 (1972).
- (7) J. Goodacre, R. J. Ponsford, and I. Stirling, *Tetrahedron Lett.*, **42**, 3609 (1975).

Synthesis of β -Spiro[pyrrolidinoindolines], Their Binding to the Glycine Receptor, and in Vivo Biological Activity

Fred M. Hershenson,* Kathleen A. Prodan, Ronald L. Kochman, James L. Bloss, and Carl R. Mackerer

Searle Laboratories, Chicago, Illinois 60680. Received May 4, 1977


A series of β -spiro[pyrrolidinoindolines], **3a-d**, was prepared and evaluated for their ability to bind to the glycine receptor. These compounds were also tested in vivo to determine if they would produce convulsant or anxiolytic effects. The target indolines were chosen because they represent rings A, B, E, and a portion of ring C of strychnine. Results of this study indicate that, in this series, an acetylindoline in the endo configuration and a tertiary amine, such as that of the pyrrolidine ring nitrogen, are required for biological activity. In all of the cases studied, the activity was of a convulsant rather than a relaxant nature. Excellent correlation was found to exist between the binding affinities to the strychnine site of the glycine receptor and clonic convulsions (ED_{50}) and death (LD_{50}) in the mouse.

The excitatory actions of strychnine on the central nervous system (CNS) have been attributed to its ability to interact with postsynaptic receptors which are sensitive to glycine, thereby blocking the inhibitory effects of this amino acid neurotransmitter.¹ Snyder et al. have demonstrated that [^3H]strychnine binds to synaptosomal membrane fragments obtained from the rat brain stem and spinal cord² and that the regional location of this binding within the CNS correlates with endogenous glycine concentrations.³

The inhibition of [^3H]strychnine binding to synaptosomal preparations has thus far been reported for glycine and β -alanine,² a group of strychnine alkaloids,⁴ and a series of 1,4-benzodiazepine derivatives.⁵ The convulsant activities of the series of strychnos alkaloids studied were found to be highly correlated with their respective binding affinities. Likewise, Snyder was able to show that the rank order of potency of a series of 21 1,4-benzodiazepines in a variety of animal and human pharmacological and behavioral tests correlated with the ability of these compounds to displace bound [^3H]strychnine. This observation led to the hypothesis that the 1,4-benzodiazepines exert their anxiolytic, muscle-relaxant, and anticonvulsant effects by mimicking glycine at its receptor site.⁵

The purpose of this study was twofold. First, we were interested in determining what structural features within

the strychnine molecule were responsible for the high binding affinity to the glycine receptor and, secondly, to establish if a portion of the strychnine skeleton would bind to this receptor and elicit anxiolytic, rather than convulsant, effects. Although the likelihood of finding a strychnine antagonist (glycine agonist) within the framework of the strychnine molecule might seem remote, such an approach has been successful elsewhere. The tetrahydropyridine, **1**, which constitutes a portion of the LSD molecule, **2**, was recently reported to bind with high affinity to the LSD receptor and, more importantly, was found to antagonize the behavioral effects of LSD.⁶



Furthermore, Rees and Smith⁷ reported that certain seco derivatives of strychnine, strychnidine, and brucidine,