

Nucleosides and Nucleotides

Publication details, including instructions for authors and
subscription information:

<http://www.tandfonline.com/loi/Incn19>

Inhibitory Effects of 1-Deazaadenosine Analogues on HIV Replication and Adenosine Deaminase

Sauro Vittori ^a, Rosaria Volpini ^a, Emidio Camaioni ^b, Giorgio Palùg ^a
& Giorgio Cristalli ^a

^a Dipartimento di Scienze Chimiche, University of Camerino, 62032, Camerino, (MC)

^b Istituto di Microbiologia, University of Padua, Italy

Published online: 16 Feb 2007.

To cite this article: Sauro Vittori, Rosaria Volpini, Emidio Camaioni, Giorgio Palùg & Giorgio Cristalli (1995) Inhibitory Effects of 1-Deazaadenosine Analogues on HIV Replication and Adenosine Deaminase, *Nucleosides and Nucleotides*, 14:3-5, 603-606

To link to this article: <http://dx.doi.org/10.1080/15257779508012435>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

INHIBITORY EFFECTS OF 1-DEAZAADENOSINE ANALOGUES ON HIV REPLICATION AND ADENOSINE DEAMINASE

Sauro Vittori, Rosaria Volpini, Emidio Camaioni, §Giorgio Palù, Gloria Cristalli*

Dipartimento di Scienze Chimiche, University of Camerino, 62032 Camerino (MC)
and §Istituto di Microbiologia, University of Padua, ITALY.


Abstract. A series of 2',3'-dideoxy-N⁶-(cyclo)alkyl-1-deazaadenosine derivatives were synthesized starting from 2,6-dichloro-1-deazapurine (**9**). The new nucleosides proved to be good inhibitors of HIV-1 replication, the most active being the 2',3'-dideoxy-2-chloro-N⁶-cycloctyl-1-deazaadenosine (**14h**, ED₅₀ = 0.4 μM).

1-Deazaadenosine derivatives have been shown to possess cytotoxic activity,^{1,2} adenosine receptor affinity,³ and to inhibit adenosine deaminase^{4,5} and platelet aggregation.^{3,6} We have already reported the coupling of 7-nitro-3H-imidazo[4,5-b]pyridine (**1**)¹ and of 5-chloro-7-nitro-3H-imidazo[4,5-b]pyridine⁷ with ribose and 2-deoxyribose derivatives to obtain 6-amino-¹ and 6-hydroxylamino-1-deazapurine nucleosides.^{2,8} Moreover, 2',3'-dideoxynucleosides are good tools for inhibition of HIV replication.


On this basis we attempted the synthesis of 2',3'-dideoxy-1-deazaadenosine by reaction of compound **1** with the benzyl derivative of dideoxy sugar **2** (Scheme 1). A mixture of α and β anomers (**3**) was obtained in 40% yield but attempts of deprotection failed leading to N-3 (**4**) and N-1 (**5**) benzyl derivatives. Catalytic reduction of compound **3** gave the still protected nucleoside **6** in low yield. Coupling of **1** with the *t*-butyldimethylsilyl derivative **7** led to the β anomer **8** in very low yield. Alternatively, we used the more versatile base 5,7-dichloro-3H-imidazo[4,5-b]pyridine, (**9**)^{7,9} which was glycosylated using first the sugar **7**, to give small quantity of **10**, and then the toluoyl derivative **11**,¹⁰ to give mixture of α and β anomers (**12**) in 70% yield (Scheme 2).

The anomeric and isomeric configuration of the new nucleosides was assigned on the basis of NMR and UV spectra.¹¹

Substitution of chlorine in 6 position from compound **12** or from the deprotected derivative **13** with ammonia, methylamine and several



Scheme 1



Scheme 2

TABLE 1. Biological activity of 2',3'-dideoxy-N⁶-(cyclo)alkyl-1-deazaadenosines.

Cp	R	EC ₅₀ (μM) HIV-1	CC ₅₀ (μM) C8166 cells	K _i (μM) ADA calf intest.
14a	H	>1000	>1000	>100
15a	H	500	>1000	2.5
14b	CH ₃	400	800	11
15b	CH ₃	50	100	2.2
14c	cC ₃ H ₅	200	400	>100
15c	cC ₃ H ₅	100	750	78
14d	cC ₄ H ₇	4	250	>100
15d	cC ₄ H ₇	50	400	92
14e	cC ₅ H ₉	20	50	>100
15e	cC ₅ H ₉	40	200	>100
14f	cC ₆ H ₁₁	4	20	>100
15f	cC ₆ H ₁₁	40	50	69
14g	cC ₇ H ₁₃	0.8	9	>100
15g	cC ₇ H ₁₃	8	40	53
14h	cC ₈ H ₁₅	0.4	10	>100
15h	cC ₈ H ₁₅	0.8	10	>100

cycloalkylamines gave nucleosides **14a-h**, whose reduction with Pd/C led to the dechlorinated products (**15a-h**).

BIOLOGICAL EVALUATION

The anti HIV-1 activities and toxicities of synthesized compounds were assessed in C8166 cells infected with HIV-1_{IIIIB}. The results are reported on TABLE 1. While the parent compound **15a** was inactive, the presence of cycloalkyl substituents on N⁶ brought about a good anti HIV activity, which increased with the dimension of the substituent; on the other hand, the presence of a chlorine atom in 2-position improved both activity and therapeutic index, the most active compound being 2',3'-dideoxy-2-chloro-N⁶-cyclooctyl-1-deazaadenosine (**14h**, EC₅₀ = 0.4 μM).

The compounds were also tested for their ability to inhibit calf intestine adenosine deaminase (ADA). Results (TABLE 1) showed that none of the tested derivatives were substrate of the enzyme, and some of them were good inhibitors, with 2',3'-dideoxy-1-deazaadenosine (**15a**) and 2',3'-dideoxy-N⁶-methyl-1-deazaadenosine (**15b**) being the most active compounds, with a K_i of 2.5 and 2.2 μM, respectively. A considerable decrease in activity is produced by the presence of a chlorine atom

in the 2-position and of a cycloalkyl substituent in N⁶. This is in agreement with our hypothesis that 1-deazaadenosine derivatives interact directly with the enzyme at the catalytic site, since the presence of a bulk, hydrophobic substituent on the exocyclic nitrogen is detrimental for the hydrogen bonding of the molecules.

REFERENCES

- 1 Cristalli, G.; Franchetti, P.; Grifantini, M.; Vittori, S.; Bordoni, T.; Geroni, C. *J. Med. Chem.* **1987**, *30*, 1686.
- 2 Cristalli, G.; Vittori, S.; Eleuteri, A.; Grifantini, M.; Volpini, R.; Lupidi, G.; Capolongo, L.; Pesenti, E. *J. Med. Chem.* **1991**, *34*, 2226.
- 3 Cristalli, G.; Franchetti, P.; Grifantini, M.; Vittori, S.; Klotz, K.-N.; Lohse, M. J. *J. Med. Chem.* **1988**, *31*, 1179.
- 4 Lupidi, G.; Cristalli, G.; Marmocchi, F.; Riva, F.; Grifantini, M. *J. Enzyme Inhib.* **1985**, *1*, 67.
- 5 Cristalli, G.; Eleuteri, A.; Vittori, S.; Volpini, R.; Camaioni, E.; Lupidi, G. *Drug Dev. Res.* **1993**, *28*, 253.
- 6 Antonini, I.; Cristalli, G.; Franchetti, P.; Grifantini, M.; Martelli, S.; Petrelli, F. *J. Pharm. Sci.* **1984**, *73*, 366.
- 7 Cristalli, G.; Grifantini, M.; Vittori, S.; Balduini, W.; Cattabeni, F. *Nucleosides & Nucleotides* **1985**, *4* (5), 625.
- 8 Cristalli, G.; Vittori, S.; Eleuteri, A.; Grifantini, M.; Lupidi, G.; Capolongo, L.; Pesenti, E. *Nucleosides & Nucleotides* **1991**, *10*(1-3), 253.
- 9 Cristalli, G.; Vittori, S.; Eleuteri, A.; Volpini, R.; Camaioni, E.; Lupidi, G. *Nucleosides & Nucleotides* **1994**, *13*(1-3), 835.
- 10 Abdel-Megied, A. E.-S.; Pedersen, E.B.; Nielsen, C.M. *Monatsh. Chem.* **1991**, *122*, 59.
- 11 Cristalli, G.; Vittori, S.; Camaioni, E.; Eleuteri, A.; Volpini, R.; Lupidi, G.; Palù, G.; Bevilacqua, F. *J. Med. Chem.* **1994** in press