



A Journal of



## Accepted Article

**Title:** Visible-Light-Promoted Efficient Aerobic Dehydrogenation of N-Heterocycles by Tiny Organic Semiconductor under Ambient Conditions

**Authors:** Kunyi Yu, Hanjie Zhang, Chenliang Su, and Yongfa Zhu

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Eur. J. Org. Chem.* 10.1002/ejoc.202000170

**Link to VoR:** <http://dx.doi.org/10.1002/ejoc.202000170>

Supported by



WILEY-VCH

## COMMUNICATION

# Visible-Light-Promoted Efficient Aerobic Dehydrogenation of N-Heterocycles by Tiny Organic Semiconductor under Ambient Conditions

Kunyi Yu,<sup>[a]</sup> Hanjie Zhang,<sup>[a]</sup> Chenliang Su,<sup>\*[b]</sup> Yongfa Zhu<sup>\*[a]</sup>

[a] K. Y. Yu, H. J. Zhang, Pro Y. F. Zhu  
Department of Chemistry  
Tsinghua University  
Beijing 100084, PR China  
E-mail: zhuyf@tsinghua.edu.cn  
URL: <http://www.zhuyfgroup.com>

[b] Pro C. L. Su  
International Collaborative Laboratory of 2D Materials for Optoelectronics Science and Technology of Ministry of Education, Engineering Technology Research Center for 2D Materials Information Functional Devices and Systems of Guangdong Province, Institute of Microscale Optoelectronics, Shenzhen University  
Shenzhen 518060, PR China  
Email: chmsuc@szu.edu.cn  
URL: <http://2dchem.szu.edu.cn/>

Supporting information for this article is given via a link at the end of the document.

**Abstract:** An efficient reusable catalytic system has been developed based on perylene diimide (PDI) organic semiconductor for the aerobic dehydrogenation of N-heterocycles with visible light. This practical catalytic system without any additives proceeds under ambient conditions. The minute aggregates of PDI molecules on the surface of SiO<sub>2</sub> nanospheres form tiny organic semiconductors, resulting in high-efficiency photo-oxidative activity. Notably, the robustness of this method is demonstrated by the synthesis of a wide range of N-heteroarenes, gram-scale experiments as well as reusability tests.

N-heteroarenes are considered privileged structure in nature products, medicinal chemistry and material science.<sup>[1]</sup> Therefore, the development of mild methods for synthesis of N-heteroarenes has attracted the interest of organic synthetic chemists.<sup>[2]</sup> The site-selective introduction of substituents onto N-heteroarenes is a formidable challenge, while this is relatively straightforward to achieve for the corresponding N-heterocycles. For example, tetrahydroquinoline can easily undergo 6-site electrophilic substitution functionalization, and the corresponding substituted quinoline can be obtained by dehydrogenation.<sup>[3]</sup> Therefore, the dehydrogenation of N-heterocycles to N-heteroarenes is considered a viable method.

When compared to thermally promoted organic reaction, the utilisation of visible-light energy as a driving force for organic transformation is appealing, as it is a mild, safe, and easy-to-handle energy resource<sup>[4]</sup>. In general, there are two subclasses of visible-light-promoted dehydrogenation of N-heterocycles: acceptorless<sup>[5, 6]</sup> and acceptor assisted,<sup>[3, 6]</sup> depending on whether hydrogen acceptors are involved. Acceptor assisted dehydrogenation, especially aerobic oxidative dehydrogenation using oxygen as a cheap and readily available oxidant, has the advantages of scalability and operational simplicity. From the point of view of sustainable development, heterogeneous catalysts are preferable due to facile product separation, catalyst

reusability.<sup>[1k]</sup> In recent years, semiconductor photocatalysis with heterogeneous advantages has been widely used in photocatalytic splitting water into H<sub>2</sub> and O<sub>2</sub>, the degradation of organic pollutants and organic synthesis.<sup>[7]</sup> To the best of our knowledge, only two cases of aerobic dehydrogenation of N-heterocycles by semiconductors have been reported. Operation temperatures up to 80 °C and pure oxygen was required in the mpg-C<sub>3</sub>N<sub>4</sub> catalytic system;<sup>[8]</sup> TEMPO as a co-catalyst was required in the TiO<sub>2</sub> catalytic system.<sup>[9]</sup> Therefore, an efficient semiconductor photocatalytic protocol that functions under ambient conditions is highly desired.

Perylene diimide (PDI) has been widely used as a basic structural unit for fabricating organic photo-functional materials in recent years due to its unique optical and electronic properties.<sup>[10]</sup> Compared to some semiconductor photocatalysts (TiO<sub>2</sub> and C<sub>3</sub>N<sub>4</sub>), easy chemically modified PDIs can be used as functional photocatalysts in various photo-promoted chemical transformation. In 2014, König et al. reported a monomolecular PDI-AN, which can reductively cleave inert C–X bond due to accumulating the energy of two photons.<sup>[11]</sup> In 2016, Duan et al. reported a metal-organic framework Zn-PDI, which can achieve the photo-oxidation of benzyl alcohol and benzyl amine.<sup>[12]</sup> In the same year, our group reported a self-assembled supramolecule PDINH, capable of photo-oxidative degradation of organic pollutants.<sup>[13]</sup> In 2017, Chen et al. reported a covalent immobilized PDI-SN, which could be used for photo-reductive degradation of halogenated organic pollutants.<sup>[14]</sup> PDIs exhibit a variety of attractive photocatalytic activities in various phase states, prompting us to develop a new photocatalytic system based on PDI for the photocatalytic aerobic dehydrogenation of N-heterocycles.

Firstly, we examined three heterogeneous photocatalysts based on PDI (Figure 1): PDINH (self-assembly supramolecular), PDI-SN (covalent immobilized to SiO<sub>2</sub> nanospheres) and PDI-U<sup>[15]</sup>

## COMMUNICATION




Figure 1. PDIs in different phase states.

(polymerisation with urea) in the aerobic dehydrogenation of N-heterocyclic amines under ambient conditions (see Figure S1, b in the Supporting Information), utilising tetrahydroquinoline (THQ) as the model substrate. To our delight, good catalytic activity was exhibited by all three catalysts (Table 1, entries 1-3). The heterogeneous catalysts remained stable following the catalytic process, apart from supramolecular PDINH, which was structurally destroyed. In particular, the highest photo-oxidative activity was exhibited by PDI-SN, which had a larger specific surface area than PDI-U (27.7 m<sup>2</sup>/g for PDI-SN and 12.1 m<sup>2</sup>/g for PDI-U). Interestingly, the free monomer PDI-AN (condensation of PDI with 2, 6-diisopropyl aniline as showed in Figure 1) did not display any photo-oxidative activity (Table 1, entry 4); the significant differences in catalytic activity between the free monomer PDI-AN and PDI-SN prompted us to examine the phase state of the PDI molecules in PDI-SN. Compared with the free monomer PDI-AN, the wide peak in the UV-Vis absorption spectrum of PDI-SN implied that the PDI molecules exist in aggregation state in PDI-SN (see Figure S8 in the Supporting

Table 1. Optimization of the reaction conditions.<sup>[a][b]</sup>


Entry	Catalyst	Yield (%)
1	PDINH	83
2	PDI-U	54
3	PDI-SN	94(91) <sup>[c]</sup>
4	PDI-AN	trace
5	none	trace
6	PDI-SN	trace <sup>[d]</sup>
7	PDI-SN	14 <sup>[e]</sup>
8	TiO <sub>2</sub>	8
9	C <sub>3</sub> N <sub>4</sub>	18

[a]. Reaction conditions: THQ (**1a**, 0.2 mmol), catalyst (5 mg), DMA (1 mL), ambient air atmosphere under 18 W 400 nm LED irradiation overnight. [b]. Yield determined by GC analysis with an internal standard (dodecane). [c]. Isolated yield. [d]. In the absence of light. [e]. under Ar atmosphere.

Information). In addition, the fluorescence emission spectra of the supramolecular PDINH and PDI-SN supported the hypothesis mentioned above (see Figure S8 in the Supporting Information). Unlike the thick layer of the polymer PDI-U semiconductors, PDI-SN consists of minute aggregates of PDI molecules on the surface of SiO<sub>2</sub> nanospheres, which form thin layers of PDI semiconductors, resulting in a higher photogenic carrier transport efficiency, and superior photo-oxidative activity. In other words, the particular aggregation of PDI molecules in PDI-SN ensured both photo-oxidative activity and the high catalytic efficiency.

Next, we carried out solvent optimization experiments. The current catalyst has catalytic activity in many polar solvents such as DMA, DMF, DMSO, MeCN, and water. However, no product was obtained when using methanol, which probably acted as a role of electron donor and quenched the photo-generated holes (see Table S3 in the Supporting Information). In order to determine the necessity of each component, namely the catalyst, LED irradiation and air, a number of control experiments were conducted and the results indicated that they were all indispensable components (Table 1, entries 5-7). In addition, wavelengths of LED light sources were studied, with 400 nm leading to the highest catalytic activity (see Table S2 in the Supporting Information). Comparatively, the inorganic semiconductors TiO<sub>2</sub> and organic semiconductor bulky g-C<sub>3</sub>N<sub>4</sub> exhibited poor activity under standard reaction conditions (Table 1, entries 8-9).

Table 2. Aerobic dehydrogenation of cyclic amines.<sup>[a][b]</sup>




[a]. Reaction conditions: cyclic amines (**1**, 0.4 mmol), PDI-SN (10 mg), DMA (2 mL), ambient air atmosphere under 18 W 400 nm LED irradiation overnight. [b]. Isolated yield.

With the optimal reaction conditions in hand, this practical catalytic protocol was then applied to the aerobic dehydrogenation of diverse cyclic amines. As shown in Table 2,

## COMMUNICATION

various substituted tetrahydroquinolines and indolines were compatible and the expected dehydrogenated products were obtained in good to excellent yields. Tetrahydroquinolines bearing a methyl, substituted at 2-, 3-, 4- and 6- positions were well tolerated; the 2-position substitution afforded slightly lower yields. It is probably due to the higher degree of steric hindrance at the 2-position compared to that at the other positions (Table 2, **2b-2e**). A variety of 6-substituted tetrahydroquinolines were successfully subjected to this practical catalytic system; electron-deficient substitutions gave superior results (Table 2, **2f-2h**). The readily oxidised phenol, as well as aniline fragments were likewise compatible, thus the 8-substituted hydroxyquinoline and aminoquinoline being produced in 93% and 88% yield, respectively (Table 2, **2i-2j**). Similarly, various substituted indolines gave the corresponding indoles in good yields (Table 2, **2k-2q**), except for 2-indoleic acid (Table 2, **2r**). The ternary heterocyclic dihydroacridine likewise performed successful (Table 2, **2s**). Finally, a heterocyclic substrate containing two nitrogen atoms were investigated and the corresponding quinoxaline was obtained in good yield (Table 2, **2t**).


**Table 3.** Aerobic dehydrogenation of cyclic amines generated *in situ*.<sup>[a][b]</sup>



[a]. Reaction conditions: *o*-diaminobenzene (**3**, 0.4 mmol), Benzylamine or aldehyde (**4**, 0.44 mmol), PDI-SN (10 mg), and DMA (2 mL), ambient air atmosphere under 18 W 400 nm LED irradiation overnight. [b]. Isolated yield.


Encouraged by these results, we attempted to apply this practical catalytic protocol to cyclic amines generated *in situ*. As shown in Table 3, we investigated a strategy whereby benzylamine was the precursor for the imine; it reacted with *o*-phenylenediamine to form a cyclic amine intermediate, after which dehydrogenative aromatization afforded 2-phenylbenzimidazole (Table 3, **5a**) in an isolated yield of 91%. Benzylamines with varying electronic properties were compatible, including those substituted with electron-donating group, such as a methoxy (Table 3, **5b**), and electron-withdrawing group, such as a nitro group (Table 3, **5c**); the corresponding products were obtained in 87% and 93% yield, respectively. In addition, the combinations of *o*-hydroxyaniline and *o*-mercaptoaniline with benzylamines were studied, and the corresponding 2-phenylbenzoxazole (Table 3, **5d**) and 2-thiophene benzoxazole (Table 3, **5e**) were generated in 69–73% yield under standard reaction condition. Finally, vanillic aldehyde was not oxidized to vanillic acid in this photo-oxidative system, and the desired product benzimidazole derivative (Table 3, **5f**) was obtained in a yield of 78%.

As agglomeration of heterogeneous nanocatalysts is ubiquitous deficient,<sup>[2k]</sup> we examined the reusability of the present catalyst by performing six consecutive experimental runs (see the Supporting Information for details of the recycling procedure). To our delight, no significant reduction (Figure 2) were observed in yield (83 - 91%). In addition, no significant changes were seen in the UV and IR spectra of the catalyst isolated by centrifugal filtration, compared with the fresh one (see Figure S3 in the Supporting Information).



**Figure 2.** Recycling test of the PDI-SN catalyst. Reaction conditions: THQ (**1a**, 0.4 mmol), PDI-SN (20 mg), DMA (2 mL), ambient air atmosphere under 18 W 400 nm LED irradiation overnight.

To evaluate the scalability of the aerobic dehydrogenation protocol, a gram-scale experiment was performed (see the Supporting Information for details of the gram-scale reaction procedure). We found that the reactivity was no significant altered in this case, and the gram-scale reaction proceeded smoothly, giving the product in an isolated yield of 82% by a prolonged reaction time of up to 72 h, and with a combination light source (Scheme 1).




**Scheme 1.** Gram-scale preparation for the aerobic dehydrogenation of THQ. Reaction conditions: THQ (**1a**, 10 mmol), PDI-SN (500 mg), DMA (50 mL), ambient air atmosphere under 4x30 W 400 nm LED irradiation 72 h.

In order to determine the possible intermediates and reaction mechanism, a series of quenching and trapping experiments were performed (Figure 3).<sup>[16]</sup> The detection of reactive oxygen species was conducted firstly. The reaction proceeded unaffectedly in the presence of TBA,<sup>[17]</sup> a well-known hydroxyl radical ( $\text{HO}\cdot$ ) quenching agent, which indicated that no  $\text{HO}\cdot$  was involved. Similarly, the reaction is unaffected by the presence of singlet oxygen quenching ( $^1\text{O}_2$ ) agent DBACO,<sup>[18]</sup> which indicated that  $^1\text{O}_2$  was not the reactive oxygen species. In the presence of superoxide radical ( $\text{O}_2^{\cdot-}$ ) quenching agent BHT,<sup>[19]</sup> a significantly reduction in yield suggested that the reaction mechanism entail a  $\text{O}_2^{\cdot-}$ , and analogous results were obtained in nitrotetrazolium blue chloride (NBT) control experiments.<sup>[20]</sup> The reduction in reactivity in the presence of the photo-generated electron quenching agent AgOTf,<sup>[21]</sup> likewise supported the existence of  $\text{O}_2^{\cdot-}$  intermediates. The reaction was severely hampered in the presence of photo-generated holes quenching agent EG.<sup>[22]</sup> Reduction quenching


## COMMUNICATION

generally required participation by the photo-generated holes and fluorescence quenching experiments indicated electrons transfer from tetrahydroquinoline to organic semiconductor (see Figure S9 in the Supporting Information). The presence of TEMPO reduced the reaction yield, which supported a free radical pathway.<sup>[23]</sup> Finally, H<sub>2</sub>O<sub>2</sub> as a possible by-product was also detected in a typical horseradish peroxidase experiment.<sup>[24]</sup>



**Figure 3.** Experimental results of mechanism research. Reaction conditions: THQ (**1a**, 0.2 mmol), quenching agent (0.4 mmol), PDI-SN (5 mg), DMA (1 mL), ambient air atmosphere under 18 W 400 nm LED irradiation overnight. TBA = *tert*-butyl alcohol; BHT = butylated hydroxytoluene; DABCO = bicyclo[2.2.2]-1,4-diazaoctane; AgOTf = silver triflate; EG = ethylene glycol; TEMPO = 2,2,6,6-tetramethylpiperidiny-N-oxide radical.

Based on the above experimental results and known literatures,<sup>[3, 5a, 5b, 5d, 8]</sup> we proposed a possible reaction mechanism (Scheme 2). Firstly, oxidative photo-generated holes and reductive photo-generated electrons are produced by excitation of the organic semiconductor PDI-SN, respectively. Then, the substrate **1a** is oxidized by photo-generated holes through single electron transfer (SET),<sup>[25]</sup> thereby producing the corresponding cation radical **A**; O<sub>2</sub> from the air is reduced by photo-generated electrons, thereby producing a O<sub>2</sub><sup>•-</sup>, respectively. Coherently, the radical cation **A** is oxidized by O<sub>2</sub><sup>•-</sup> to form an imine cation intermediate **B**, via the so-called hydrogen-atom transfer (HAT) mechanism.<sup>[3]</sup> Consequently, Imine intermediate **C** is obtained from the imine cation **B** by the release of one proton (H<sup>+</sup>) with a concomitant generation of H<sub>2</sub>O<sub>2</sub>. The imine **C** can be rapidly isomerized to intermediate **D**.<sup>[5a, 26]</sup> The second dehydrogenation step could be completed by another catalytic cycle or aromatization,<sup>[3, 5a]</sup> providing the final dehydrogenation product **2a**.



**Scheme 2.** Proposed mechanism for the aerobic dehydrogenation of THQ.

In summary, we have reported an efficient semiconductor photocatalytic system for the aerobic dehydrogenation of N-heterocycle under ambient conditions. This reusable photocatalytic system proceeds under very mild conditions: in the ambient air atmosphere, without additional heat sources and additives. The minute aggregates of PDI molecules on the surface of SiO<sub>2</sub> nanospheres form tiny PDI semiconductors, resulting in high-efficiency photo-oxidative activity. This practical catalytic protocol is attractive, and offers a wide scope related to the synthesis of quinoline, indole, benzimidazole, benzothiazole and benzoxazole derivatives. The mechanistic studies show that photo-generated holes and O<sub>2</sub><sup>•-</sup> are pivotal components in the reaction pathway.

## Acknowledgements

This work was partly supported by Chinese National Science Foundation (21437003, 21673126, 21761142017, 21621003, and 21972094), Beijing Municipal Science and Technology Project (Z181100005118007), Collaborative Innovation Center for Regional Environmental Quality, Shenzhen Peacock Plan (KQJSCX20170727100802505 and KQTD2016053112042971), Guangdong Special Support Program, and Pengcheng Scholar program. The authors thank Dr. Chuntian Qiu, Shenzhen University for providing important advice in preparing manuscripts.

## Conflict of interest

The authors declare no conflict of interest.

**Keywords:** aerobic dehydrogenation • N-heterocyclic amines • perylene diimide • organic semiconductor • visible light

- [1] O. Afzal, S. Kumar, M. R. Haider, M. R. Ali, R. Kumar, M. Jaggi, S. Bawa, *Eur. J. Med. Chem.* **2015**, *97*, 871-910
- [2] a) S. Chakraborty, W. W. Brennessel, W. D. Jones, *J. Am. Chem. Soc.* **2014**, *136*, 8564-8567; b) A. E. Wendlandt, S. S. Stahl, *J. Am. Chem. Soc.* **2014**, *136*, 506-512; c) A. E. Wendlandt, S. S. Stahl, *J. Am. Chem. Soc.* **2014**, *136*, 506-512; d) X. Cui, Y. Li, S. Bachmann, M. Scalone, A. E. Surkus, K. Junge, C. Topf, M. Beller, *J. Am. Chem. Soc.* **2015**, *137*, 10652-10658; e) S. Chen, Q. Wan, A. K. Badu-Tawiah, *Angew. Chem.* **2016**, *128*, 9491-9495; *Angew. Chem. Int. Ed.* **2016**, *55*, 9345-9349; f) D. Forberg, T. Schwob, M. Zaheer, M. Friedrich, N. Miyajima, R. Kempe, *Nat. Commun.* **2016**, *7*, 13201; g) D. Jung, M. H. Kim, J. Kim, *Org. Lett.* **2016**, *18*, 6300-6303; h) M. Kojima, M. Kanai, *Angew. Chem.* **2016**, *128*, 12412-12415; *Angew. Chem. Int. Ed.* **2016**, *55*, 12224-12227; i) Y. Han, Z. Wang, R. Xu, W. Zhang, W. Chen, L. Zheng, J. Zhang, J. Luo, K. Wu, Y. Zhu, C. Chen, Q. Peng, Q. Liu, P. Hu, D. Wang, Y. Li, *Angew. Chem.* **2018**, *130*, 11432-11436; *Angew. Chem. Int. Ed.* **2018**, *57*, 11262-11266; j) D. Xu, H. Zhao, Z. Dong, J. Ma, *ChemCatChem* **2019**, *11*, 5475-5486; k) X. Bi, T. Tang, X. Meng, M. Gou, X. Liu, P. Zhao, *Catal. Sci. Technol.* **2020**, *10*, 360-371.
- [3] M. K. Sahoo, G. Jaiswal, J. Rana, E. Balaraman, *Chem. Eur. J.* **2017**, *23*, 14167-14172.
- [4] a) X. Li, J. Yu, M. Jaroniec, *Chem. Soc. Rev.* **2016**, *45*, 2603-2636; b) S. Lazzaroni, D. Ravelli, S. Protti, M. Fagnoni, A. Albini, *C. R. Chim.* **2017**, *20*, 261-271.
- [5] For selected examples on the visible-light-promoted acceptorless dehydrogenation of N-heterocycles, see: a) K. H. He, F. F. Tan, C. Z. Zhou, G. J. Zhou, X. L. Yang, Y. Li, *Angew. Chem.* **2017**, *129*, 3126-3130; *Angew. Chem. Int. Ed.* **2017**, *56*, 3080-3084; b) M. Zheng, J. Shi,

## COMMUNICATION

- T. Yuan, X. Wang, *Angew. Chem.* **2018**, *130*, 5585-5589; *Angew. Chem. Int. Ed.* **2018**, *57*, 5487-5491; c) M. Hao, X. Deng, L. Xu, Z. Li, *Appl. Catal., B* **2019**, *252*, 18-23; d) M. K. Sahoo, E. Balaraman, *Green Chem.* **2019**, *21*, 2119-2128.
- [6] For selected examples on the visible-light-promoted acceptor assisted dehydrogenation of N-heterocycles, see: a) K. Suzuki, F. Tang, Y. Kikukawa, K. Yamaguchi, N. Mizuno, *Angew. Chem.* **2014**, *126*, 5460-5464; *Angew. Chem. Int. Ed.* **2014**, *53*, 5356-5360; b) T. Shao, Y. Yin, R. Lee, X. Zhao, G. Chai, Z. Jiang, *Adv. Synth. Catal.* **2018**, *360*, 1754-1760.
- [7] Recent reviews on semiconductor photocatalysis, see: a) N. Serpone, A. V. Emeline, *J. Phys. Chem. Lett.* **2012**, *3*, 673-677; b) S. Perathoner, C. Ampelli, S. Chen, R. Passalacqua, D. Su, G. Centi, *J. Energy Chem.* **2017**, *26*, 207-218; c) A. Savateev, I. Ghosh, B. König, M. Antonietti, *Angew. Chem.* **2018**, *130*, 16164-16176; *Angew. Chem. Int. Ed.* **2018**, *57*, 15936-15947; d) I. Ghosh, J. Khamrai, A. Savateev, N. Shlapakov, M. Antonietti, B. König, *Science* **2019**, *365*, 360-366.
- [8] F. Su, S. C. Mathew, L. Mohlmann, M. Antonietti, X. Wang, S. Blechert, *Angew. Chem.* **2011**, *123*, 683-686; *Angew. Chem. Int. Ed.* **2011**, *50*, 657-660.
- [9] For selected examples on PDI fabricating organic photo-functional materials, see: a) D. Görl, X. Zhang, F. Werthner, *Angew. Chem.* **2012**, *124*, 6434-6455; *Angew. Chem. Int. Ed.* **2012**, *51*, 6328-6348; b) D. Schmidt, M. Son, J. M. Lim, M.-J. Lin, I. Krummenacher, H. Braunschweig, D. Kim, F. Werthner, *Angew. Chem.* **2015**, *127*, 14186-14190; *Angew. Chem. Int. Ed.* **2015**, *54*, 13980-13984; c) P. Spent, F. Werthner, *Angew. Chem.* **2015**, *127*, 10303-10306; *Angew. Chem. Int. Ed.* **2015**, *54*, 10165-10168; d) A. H. Endres, M. Schaffroth, F. Paulus, H. Reiss, H. Wadepohl, F. Rominger, R. Krämer, U. H. F. Bunz, *J. Am. Chem. Soc.* **2016**, *138*, 1792-1795; e) D. Meng, D. Sun, C. Zhong, T. Liu, B. Fan, L. Huo, Y. Li, W. Jiang, H. Choi, T. Kim, J. Y. Kim, Y. Sun, Z. Wang, A. J. Heeger, *J. Am. Chem. Soc.* **2016**, *138*, 375-380; f) A. Zhang, C. Li, F. Yang, J. Zhang, Z. Wang, Z. Wei, W. Li, *Angew. Chem. Int. Ed.* **2017**, *56*, 2694-2698; *Angew. Chem.* **2017**, *129*, 2738-2742.
- [10] N. O. Balayeva, N. Zheng, R. Dillert, D. W. Bahnemann, *ACS Catal.* **2019**, *9*, 10694-10704.
- [11] I. Ghosh, T. Ghosh, J. I. Bardagi, B. König, *Science* **2014**, *346*, 725-728.
- [12] L. Zeng, T. Liu, C. He, D. Shi, F. Zhang, C. Duan, *J. Am. Chem. Soc.* **2016**, *138*, 3958-3961.
- [13] D. Liu, J. Wang, X. Bai, R. Zong, Y. Zhu, *Adv. Mater.* **2016**, *28*, 7284-7290.
- [14] J. Shang, H. Tang, H. Ji, W. Ma, C. Chen, J. Zhao, *Chin. J. Catal.* **2017**, *38*, 2094-2101.
- [15] P. Sharma, D. Damien, K. Nagarajan, M. M. Shaijumon, M. Hariharan, *J. Phys. Chem. Lett.* **2013**, *4*, 3192-3197.
- [16] For detailed information, see the Supporting Information.
- [17] H. Che, C. Liu, W. Hu, H. Hu, J. Li, J. Dou, W. Shi, C. Li, H. Dong, *Catal. Sci. Technol.* **2018**, *8*, 622-631.
- [18] M. Klaper, T. Linker, *J. Am. Chem. Soc.* **2015**, *137*, 13744-13747.
- [19] F. Su, S. C. Mathew, G. Lipner, X. Fu, M. Antonietti, S. Blechert, X. Wang, *J. Am. Chem. Soc.* **2010**, *132*, 16299-16301.
- [20] A. M. Czoska, S. Livraghi, M. Chiesa, E. Giamello, S. Agnoli, G. Granozzi, E. Finazzi, C. D. Valentin, G. Pacchioni, *J. Phys. Chem. C* **2008**, *112*, 8951-8956.
- [21] J. Zhang, Y. Hu, J. Qin, Z. Yang, M. Fu, *Chem. Eng. J.* **2020**, *385*, 123814.
- [22] R. A. Geioushy, S. M. El-Sheikh, A. B. Azzam, B. A. Salah, F. M. Eldars, *J. Hazard. Mater.* **2020**, *381*, 120955.
- [23] D. R. Chambers, A. Juneau, C. T. Ludwig, M. Frenette, D. B. C. Martin, *Organometallics* **2019**, *38*, 4570-4577.
- [24] A. M. Azevedo, V. C. Martins, D. M. F. Prazeres, V. Vojinović, J. M. S. Cabral, L. P. Fonseca, *Biotechnol. Annu. Rev.* **2003**, *9*, 199-247.
- [25] The results of Stern-Volmer luminescence quenching experiment can support the single electron transfer (SET) from tetrahydroquinoline to photo-induced holes (see supporting information).
- [26] H. Li, J. Jiang, G. Lu, F. Huang, Z. Wang, *Organometallics* **2011**, *30*, 3131-3141.

WILEY-VCH

Accepted Manuscript