to superpose the three-dimensional structures of teleocidin and TPA using computers.^{22,23} We have also superposed them by a new rational method which is designed to superpose molecules in terms of physical and chemical properties related to receptor binding, but not in terms of superficial chemical structures or positions of heteroatoms.²⁴ In our study, the superpositions of both the twist and the sofa form conformers of teleocidin onto the TPA molecule were examined independently, whereas the other groups examined the superposition only for the twist conformation. The results showed that the TPA molecule and the teleocidin sofa form could interact with the common receptor through three hydrogen bonds, and the sofa form could be superposed onto TPA much better than the twist form. In the superposed structures, the spatial position of the alkyl group at C-12 in teleocidin corresponded to that of the methyl group at C-2 in the TPA molecule.

In this study, the biological activity of the four indolactams can be reasonably interpreted in terms of the existence ratio of the sofa form. This finding is consistent with the results from the superposition of teleocidin and TPA molecules. The coincidence of the results from two independent approaches favors the hypothesis that the sofa form is very close to the active conformation for tumorpromoting activity of teleocidins.

This work has also shown that MD calculations are very useful for searching for the stable conformations in highly strained cyclic compounds. They are especially useful in molecules whose stability is strongly influenced by the conformations of the substituent groups on them.

Conclusion

The importance of the sofa form for the tumor-promoting activity of teleocidins and indolactams was indicated by conformation analyses of four indolactam congeners using high-temperature MD calculations.

Experimental Section

Registry No. 1, 11032-05-6; 2, 90365-57-4; 3, 84590-50-1; 4, 110073-31-9; 5, 110073-28-4.

Nucleosides and Nucleotides. 107. 2-(Cycloalkylalkynyl)adenosines: Adenosine A_2 Receptor Agonists with Potent Antihypertensive Effects¹

Toichi Abiru,^{*,†} Takanori Miyashita,[†] Yohko Watanabe,[†] Toyofumi Yamaguchi,^{†,§} Haruhiko Machida,[†] and Akira Matsuda[‡]

Biology Laboratory, R & D Division, Yamasa Shoyu Co., Ltd., Araoicho 2-10-1, Choshi, Chiba 288, Japan, and Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060, Japan. Received October 4, 1991

Adenosine receptor-binding profiles in rat brain tissues and antihypertensive effects in spontaneously hypertensive rats (SHR) of a series of 2-(cycloalkylakynyl)adenosines (2-CAAs) and their congeners are described. The structure-activity relationship of this series of compounds is discussed, focusing on the length of the alkynyl side chain and bulkiness of the terminal cycloalkyl substituents in terms of binding activity and cardiovascular effects. All the 2-CAAs had a preferential affinity for A₂ receptors. Of these derivatives, 2-(3-cyclopentyl-1-propyn-1yl)adenosine (10b) exhibited the most selective affinity for A₂ receptors (K_1 ratio: A₁/A₂ = 70) on the basis of receptor binding. In the C-2 binding region of adenosine, compounds often have potent and/or selective A₂ activity from introduction of an acetylenic group at the C-2 position followed by one methylene residue further followed by a hydrophobic substituent such as a cycloalkyl ring at the terminal position of the alkynyl side chain. Intravenous injection of 10b up to 100 μ g/kg had a potent hypotensive effect without a marked decrease in heart rate in anesthetized SHR. Compounds 10j-s, with a hydroxyl group in the C-3" position of the alkynyl side chain, had a potent affinity for both A₁ and A₂ receptors, but they were not highly selective for A₂ receptors. These compounds caused a marked bradycardia upon intravenous administration in anesthetized SHR. Oral administration of 10b (0.1-1 mg/kg) had a potent and long-lasting antihypertensive effect in conscious SHR.

Adenosine receptors in cell membranes have been classified into A_1 and A_2 receptors on the basis of receptor-mediated inhibition (A_1 receptors) or stimulation (A_2 receptors) of adenylate cyclase.² Some adenosine ana-

⁽²²⁾ Jeffrey, A. M.; Liskamp, R. M. J. Computer-Assisted Molecular Modeling of Tumor Promoters: Rationale for the Activity of Phorbol Esters, Teleocidin B, and Aplysiatoxin. Proc. Natl. Acad. Sci. U.S.A. 1986, 83, 241-245.

⁽²³⁾ Wender, P. A.; Koehler, K. F.; Sharkey, N. A.; Dell'Aquila, M. L.; Blumberg, P. M. Analysis of the Phorbol Ester Pharmaco-phore on Protein Kinase C as a Guide to the Rational Design of New Classes of Analogs. Proc. Natl. Acad. Sci. U.S.A. 1986, 83, 4214-4218.

⁽²⁴⁾ Itai, A.; Kato, Y.; Tomioka, N.; Iitaka, Y.; Endo, Y.; Hasegawa, M.; Shudo, K.; Fujiki, H.; Sakai, S. A Receptor Model for Tumor Promoters: Rational Superposition of Teleocidins and Phorbol Esters. Proc. Natl. Acad. Sci. U.S.A. 1988, 85, 3688-3692.

^{(±)-}Indolactam-G (3). The preparation method, combustion elemental analysis, and ¹H NMR at room temperature have been published.¹⁴ The 400-MHz ¹H NMR spectrum of indolactam-G at -30 °C was measured with a JEOL GX400 spectrometer. The chemical shifts at -30 °C are as follows: The fold conformer, 2.78 (dd, 1 H, J = 15.4, 8.3, 8-CH₂), 2.86 (s, 3 H, NCH₃), 3.17 (dd, 1 H, J = 15.4, 6.9, 8-CH₂), 3.51 (d, 1 H, J = 13.5, 12-CH₂), 3.58 (dd, 1 H, J = 11.3, 8.2, 14-CH₂), 3.69 (dd, 1 H, J = 11.3, 4.1, 14-CH₂), 3.94 (d, 1 H, J = 13.5, 12-CH₂), 5.05 (m, 1 H, J = 7.6, 6-CH), 7.09 (d, 1 H, J = 7.6, 7-CH); The S8 conformer, 2.94 (s, 3 H, NCH₃), 3.06 (d, 1 H, J = 13.8, 12-CH₂), 4.18 (d, 1 H, J = 13.8, 12-CH₂), 6.78 (d, 1 H, J = 7.6, 5-CH); other peaks could not be assigned since they overlapped with the peaks of the fold form. The signal ratio of these two conformers is 1:0.2.

[†]Yamasa Shoyu Co., Ltd.

[†]Hokkaido University.

^{*}Author to whom correspondence should be addressed.

¹Present address: Department of Biological Sciences, The Nishi Tokyo University, Uenohara, Yamanashi 409-01, Japan.

Part 106: Matsuda, A.; Nakajima, Y.; Ueda, T. Synthesis and biological activity of 1-(2-deoxy-2-hydroxyimino- or methoxyimino-β-D-erythro-pentofuranosyl)-thymine and -cytosine. Nucleosides Nucleotides, in press.



logues, such as N⁶-substituted adenosines, with a preferential affinity for A_1 receptors cause cardiac depression, while analogues with a high affinity for A2 receptors cause vasodilation.³ Recently, adenosine has been reported to be clinically useful for the treatment of arrhythmia.⁴ However, the therapeutic potential of adenosine agonists as antihypertensives and vasodilators have not been established due to their lack of selectivity for adenosine receptors. Nonselective adenosine agonists produce hypotension and vasodilation accompanied by detrimental effects such as atrio-ventricular block and angina pain.⁵ Therefore, selective A_2 receptor agonists have the potential of being useful agents for the treatment of cardiovascular diseases with minimized toxic effects. To improve and separate the adenosine-induced action, a number of adenosine derivatives have been synthesized. Of these analogues, N^6 -cyclohexyladenosine (CHA)⁶ and N^6 -cyclopentyladenosine (CPA)⁷ are highly selective A_1 receptor

- (2) Van Calker, D.; Muller, M.; Hamprecht, B. Adenosine regulates via two different types of receptors, the accumulation of cyclic AMP in cultured brain cells. J. Neurochem. 1979, 33, 999–1005.
- (3) Haleen, S. J.; Evans, D. B. Selective effects of adenosine receptor agonists upon coronary resistance and heart rate in isolated working rabbit hearts. *Life Sci.* 1985, 36, 127-137.
- (4) DiMarco, J. P.; Sellers, D. T.; Lerman, B. B.; Greenberg, M. L.; Berne, R. M.; Belardinelli, L. Diagnostic and therapeutic use of adenosine in patients with supraventricular tachyarrhythmias. J. Am. Coll. Cardiol. 1985, 6, 417-424.
- (5) Sylven, C.; Jonzon, B.; Brandt, R.; Beermann, B. Adenosineprovoked angina pectoris-like pain-time characteristics, influence of autonomic blockade and naloxone. *Eur. Heart J.* 1987, 8, 738-743.
- (6) Bruns, R. F.; Daly, J. W.; Snyder, S. H. Adenosine receptors in brain membranes: Binding of N⁶-cyclohexyl[³H]adenosine and 1,3-diethyl-8-[³H]phenylxanthine. Proc. Natl. Acad. Sci. U.S.A. 1980, 77, 5547-5551.



^a Reaction conditions: (a) HC=CR, (PPh₃)₂PdCl₂, Cul, Et₃N in DMF, 80° C; (b) HC=CR, (PPh₃)₂PdCl₂, Cul, Et₃N in dioxane, room temperature; (c) concentrated NH₄OH/dioxane, 70 °C.

agonists. However, only a few potent and/or selective A_2 receptor agonists have been reported. 2-(Phenylamino)adenosine (CV-1808) (1) was reported as a potent coronary vasodilator and confirmed as a selective A_2 agonist.⁸ NECA (*N*-ethyladenosin-5'-uronamide) (2) was also found to be a coronary vasodilator and a highly potent A_2 agonist but not selective. Recently, more selective A_2 agonists such as DPMA [N^6 -[2-(3,5-dimethoxyphenyl)-2-(2-methylphenyl)ethyl]adenosine] (3),⁹ CGS 21680 [2-[[4-(2carboxyethyl)phenethyl]amino]-*N*-ethyladenosin-5'uronamide hydrochloride] (4),^{10,11} MPEA [2-[2-(4methylphenyl)ethoxy]adenosine] (5),^{12,13} and CGS 22492 [2-[(2-cyclohexylethyl)amino]adenosine] (6)¹⁴ shown in Chart I have been reported.

Previously, we synthesized a series of 2-alkynyladenosines (2-AAs), some of which were potent inhibitors

- (7) Williams, M.; Braunwalder, A.; Erickson, T. E. Evaluation of the binding of the A-1 selective adenosine radioligand, cyclohexyladenosine (CPA), to rat brain tissues. Naunyn-Schmiedebergs Arc. Pharmacol. 1986, 332, 179–183.
- (8) Daly, J. W.; Jacobson, K. A.; Ukena, D. Adenosine receptors: Development of selective agonists and antagonists. In: Cardiac Electrophysiology and Pharmacology of Adenosine and ATP: Basic and Clinical Aspects; Pellerg, A., Michelson, E. L., Dreifs, L. S., Eds.; Alan R. Liss, Inc.: New York, 1987; p 41-63.
- (9) Bridges, A. J.; Bruns, R. F.; Ortwine, D. F.; Priebe, S. R.; Szotek, D. L.; Trivedi, B. K. N⁶-[2-(3,5-Dimethoxyphenyl)-2-(2-methylphenyl)ethyl]adenosine and its uronamide derivatives. Novel adenosine agonists with both high affinity and high selectivity for adenosine A₂ receptor. J. Med. Chem. 1988, 31, 1282-1285.
- (10) Hutchison, A. J.; Webb, R. L.; Oei, H. H.; Ghai, G. R.; Zimmerman, M. B.; Williams, M. CGS 21680C, an A₂ selective adenosine receptor agonist with preferential hypotensive activity. J. Pharmacol. Exp. Ther. 1989, 251, 47-55.
- (11) Hutchison, A. J.; Williams, M.; de Jesus, R.; Yokoyama, R.; Oei, H. H.; Ghai, G. R.; Webb, R. L.; Zoganas, H. C.; Stone, G. A.; Jarvis, M. F. 2-(Arylalkylamino)adenosin-5'-uronamides: A new class of highly selective adenosine A₂ receptor ligands. J. Med. Chem. 1990, 33, 1919–1924.
- (12) Ueeda, M.; Thompson, R. D.; Arroyo, L. H.; Olsson, R. A. 2-Alkoxyadenosines; Potent and selective agonists at the coronary artery A₂ adenosine receptor. J. Med. Chem. 1991, 34, 1334-1339.
- (13) Ueeda, M.; Thompson, R. D.; Arroyo, L. H.; Olsson, R. A. Aralkoxyadenosines: Potent and selective agonists at the coronary artery A₂ adenosine receptor. J. Med. Chem. 1991, 34, 1340-1344.
- (14) Francis, J. E.; Webb, R. L.; Ghai, G. R.; Hutchison, A. J.; Moskal, M. A.; deJesus, R.; Yokoyama, R.; Rovinski, S. L.; Contardo, N.; Dotson, R.; Barclay, B.; Stone, G. A.; Jarvis, M. F. Highly selective adenosine A₂ receptor agonists in a series of N-alkylated 2-aminoadenosines. J. Med. Chem. 1991, 34, 2570-2579.

Table I. A₁ and A₂ Receptor Binding Activities of Adenosine Analogues in Rat Brain Tissues and Their Cardiovascular Effects in SHR

		K _i , ^a nM		selectivity	BP ED ₃₀ , ^b	HR ED ₁₀ .°
no.	R	A1	A ₂	A_1/A_2	μg/kg	μg/kg
10a	c-C ₅ H ₉	88 ± 5.2	12 ± 2.5	7	0.29 ± 0.03	>100
1 0b	$c-C_5H_9CH_2$	162 ± 36	2.3 ± 0.6	70	0.05 ± 0.001	>100
10c	$c-C_5H_9(CH_2)_2$	136 ± 15	3.4 🌨 1.1	40	0.16 ± 0.005	>100
10 d	c-C ₆ H ₁₁	138 ± 18	10 ± 1.9	14	0.36 ± 0.06	>100
10e	$c-C_6H_{11}CH_2$	208 ± 26	6.5 ± 1.8	32	0.15 ± 0.03	>100
10 f	$c-C_6H_{11}(CH_2)_2$	313 ± 45	26 ± 4.2	12	0.48 ± 0.09	>100
10g	$c-C_{6}H_{11}(CH_{2})_{3}$	>229	46 ± 7.6	>5	1.64 ± 0.56	>100
10 h	Ph	701 ± 13	109 ± 28	6	10.1 ± 2.4	>100
10i	$Ph(CH_2)_2$	400 ± 39	37 ± 4.3	11	1.21 ± 0.28	>100
10j	$c-C_5H_8(OH)$	11 ± 1.8	3.3 ± 0.5	3	0.07 ± 0.01	1.6 ± 0.2
10k	$c-C_6H_{10}(OH)$	21 ± 3.3	0.9 ± 0.2	23	0.02 ± 0.004	3.5 ± 1.2
101	$c-C_7H_{12}(OH)$	27 ± 2.5	1.1 ± 0.3	25	0.02 ± 0.001	5.6 ± 0.9
10m	$c-C_8H_{14}(OH)$	56 ± 6.7	1.9 ± 0.5	29	0.01 ± 0.003	13 ± 2.6
10 n	CH ₃ (OH)CH	15 ± 2.1	18 ± 1.9	0.8	0.15 ± 0.01	0.8 ± 0.1
1 0o	$(CH_3)_2(OH)C$	32 ± 4.2	19 ± 2.1	2	0.14 ± 0.01	1.4 ± 0.3
10p	CH ₃ (CH ₂) ₂ (OH)CH	16 ± 1.9	1.8 ± 0.2	9	0.04 ± 0.01	0.83 ± 0.2
10 q	CH ₃ (CH ₂) ₄ (OH)CH	11 ± 1.5	2 ± 0.3	7	0.06 ± 0.004	0.45 ± 0.12
10r	Ph(OH)ČH	3.4 ± 0.5	1.9 ± 0.2	2	0.05 ± 0.003	1.3 ± 0.03
10s	HOCH ₂	8.3 ± 1.0	20 ± 2.2	0.4	0.36 ± 0.14	0.74 ± 0.1
10t	$CH_3(CH_2)_3$	126 ± 8.8	2.8 ± 0.3	45	0.11 ± 0.03	67 ± 6.8
10 u	$CH_3(CH_2)_5$	133 ± 20	7.9 ± 1.8	17	0.20 ± 0.02	>100
3	J	108 ± 10	6.9 ± 2.3	15	12.3 ± 1.6	>100
4		1232 ± 95	8.8 ± 1.2	140	0.97 ± 0.2	>100

^a Inhibition constant for A_1 (rat brain membranes, [³H]CHA) or A_2 (rat striatal membranes, [³H]NECA receptor binding activities of agonists. Affinities for A_1 and A_2 receptors are means \pm SE of three separate experiments in triplicate. The K_d values for the binding of [³H]CHA and [³H]NECA were 1.24 \pm 0.11 and 4.47 \pm 0.53 nM, respectively. ^bDose of compound which produced a 30% decrease in blood pressure of anesthetized SHR. ^cDose of compound which produced a 10% decrease in heart rate of anesthetized SHR. ED₃₀ and ED₁₀ values are means \pm SE of four animals.

of the passive cutaneous anaphylaxis reaction and had cardiovascular effects in normotensive rats.¹⁵ We also reported that 2-(1-hexyn-1-yl)adenosine (10t) and 2-(1octyn-1-yl)adenosine (YT-146) (10u) are A_2 selective agonists that have a potent and long-lasting antihypertensive effect in spontaneous hypertensive rats (SHR).¹⁶ Furthermore, we examined the SAR of a series of 2-AAs and identified the optimal chain length at the C-2 position for selective A₂ binding affinity.¹⁷ Recently, Ueeda et al.^{12,13} proposed a model of the A_2 receptor where pharmacological effects of a series of 2-(alkoxy)adenosines were examined. The experiment using Langendorf guinea pig heart preparations showed that 5 was highly selective for A_2 receptors. Francis et al. have reported the SAR of N-alkylated 2-aminoadenosines, in which 6 is extremely A_2 selective compound (530-fold).¹⁴ However, the nature of regions in the A₂ receptors has not been fully understood. This paper expands the mapping of C-2 regions of adenosine receptors by examining the SAR of 2-CAAs with respect to potency and selectivity of adenosine A_1/A_2 receptor binding affinity and cardiovascular effects in SHR and describes an improved method for the synthesis of 2-AAs, especially 2-CAAs and their hydroxyl derivatives.

Results and Discussion

Chemistry. Some of 2-AAs 10h,i,o-q were synthesized following published procedures.^{15,17} Treatment of 2-





^aReaction conditions: (a) concentrated $H_2SO_4/48\%$ HBr, reflux; (b) LiC=CH \cdot $H_2NCH_2CH_2NH_2$ in DMSO, room temperature.

iodoadenosine (8) with terminal alkynes, using catalytic amounts of bis(triphenylphosphine)palladium dichloride and cuprous iodide (CuI) in N,N-dimethylformamide (DMF) and triethylamine at 80 °C for 1 h under argon atmosphere gave 2-AAs and their hydroxyl derivatives 10 (method A). However, to obtain the pure products, treatment of the reaction mixture with hydrogen sulfide gas (H₂S) with subsequent purification on silica gel column chromatography was required because of chelation of metals between 6-NH₂ and N⁷ of the adenine ring. To avoid such processes, we devised an improved procedure, which is outlined in Scheme I.

Palladium-catalyzed cross-coupling reaction of 9-(2,3,5-tri-O-acetyl-1-β-D-ribofuranosyl)-6-chloro-2-iodopurine (7), which is readily available in three steps from guanosine,¹⁷ with 1-octyne in dioxane at room temperature gave the corresponding 6-chloro-2-octyn-1-ylpurine nucleoside 9u in high isolated yield. This reaction proceeded smoothly at low temperature with little coloration, which is frequently observed in such reactions in DMF at high temperature. Cuprous iodide was easily removed by washing with aqueous solution of ethylenediaminetetraacetic acid disodium salt without using H_2S . Amination and deacetylation of 9u using concentrated NH₄OH in dioxane in sealed tube at 70 °C gave 10u in 86% yield from 7 (method B). The IR, ¹H NMR, and UV data of 10u were identical to those described previously.¹⁷ A number of 2-CAAs was synthesized in two steps by this new route in 75-85% yield (Table II). The synthesis of the cyclo-

⁽¹⁵⁾ Matsuda, A.; Shinozaki, M.; Miyasaka, T.; Machida, H.; Abiru, T. Palladium-catalyzed cross-coupling of 2-iodoadenosine with terminal alkynes: synthesis and biological activities of 2-alkynyladenosines. *Chem. Pharm. Bull.* 1985, 33, 1766–1769.

⁽¹⁶⁾ Abiru, T.; Yamaguchi, T.; Watanabe, Y.; Kogi, K.; Aihara, K.; Matsuda, A. The antihypertensive effect of 2-alkynyladenosines and their selective affinity for adenosine A₂ receptors. *Eur. J. Pharmacol.* 1991, 196, 69-76.

⁽¹⁷⁾ Matsuda, A.; Shinozaki, M.; Yamaguchi, T.; Homma, H.; Nomoto, R.; Miyasaka, T.; Watanabe, Y.; Abiru, T. Nucleosides and Nucleotides. 103. 2-Alkynyladenosines: a novel class of selective adenosine A₂ receptor agonists with potent antihypertensive effects. J. Med. Chem. 1992, 35, 241-252.

alkylalkynes such as 4-cyclopentyl-1-butyne, 4-cyclohexyl-1-butyne, and 5-cyclohexyl-1-pentyne (13a-c) was done by treatment of a lithium acetylide-ethylenediamine complex with proper cycloalkyl bromides (12a-c) (Scheme II).

Adenosine Receptor Binding Activity. The A_1 binding assay was done in adenosine deaminase (ADA)pretreated rat brain membranes (without cerebellum and brainstem) by measuring the ability of test compounds to displace [³H]CHA binding according to a previously described procedure,¹⁸ with modifications.¹⁶ The A_2 binding assay was done in ADA-pretreated rat striatal membranes using [³H]NECA,¹⁸ with modifications.¹⁶ The results are summarized in Table I.

A series of 2-CAAs 10a-g showed a substantial A₂ selectivity. Of these analogues, 2-(3-cyclopentyl-1-propyn-1-yl)adenosine (10b) had the highest selectivity (70-fold) for A_2 receptors with K_i values of 162 and 2.3 nM for A_1 and A_2 receptors, respectively. These values are essentially similar to those of 2-(1-hexyn-1-yl)adenosine (10t), which we recently reported.¹⁶ Affinities of these 2-CAAs for receptor subtypes varied according to the number of methylene residues between the terminal cycloalkyl ring and the acetylenic bond in the side chain. Among compounds 10a-c, which have a cyclopentyl ring at the terminal position of the alkynyl side chain, 10b had the most potent A2 activity, while it showed modest activity for A1 receptors. Similarly, among the cyclohexyl analogues 10d-g, 10e had the most A_2 selectivity. In both series the maximum binding activity and selectivity for A₂ receptors was observed when one methylene group separated the cycloalkyl ring from the acetylenic bond. Increases or decreases of this number reduced the activity. Similar effects on activity were observed when a bulky substituent was adjacent to the acetylenic bond as shown by 2-[2-(trimethylsilyl)-1-ethyn-1-yl]adenosine¹⁷ and the phenylethynyl derivative 10h. Similarly, it was observed in the SAR of 2-alkoxyadenosines¹² and N-alkylated 2-aminoadenosines¹⁴ that decrease or increase in the number of methylene group varied the affinity and selectivity for A_2 receptors. Therefore, a number of methylene groups is

essential for potent A_2 activity. Recently, Ueeda et al.^{12,13} reported models of C-2 binding regions of adenosine A_1 and A_2 receptors in SAR studies of 2-(alkoxy)adenosines. They proposed at least three subregions near the C-2 position for A_2 receptors: (a) an X subregion that accommodates the oxygen atom that links the alkyl substituent to the adenine C-2, (b) an alkyl subregion, which follows the X subregion and is of very limited bulk tolerance, and (c) a hydrophobic subregion, which is a prominent feature only for A_2 receptors, that accommodates cycloalkyl, bicycloalkyl, or phenyl rings. From our results together with previous observations, it is clear that there is the X subregion that accommodates -O-, -NH-, and -C=C- groups at the C-2 position. The nature of the region together with the N¹ and N³ positions of the adenine ring has polar characteristics including hydrogen-bonding. The alkyl region can be tolerant of a linear alkyl substituent with one or two methylene groups. Two methylene groups are required for maximum affinity to the A_2 receptor for the C-2 alkoxy¹² and alkylamino^{11,14} derivatives but only one methylene for 2-CAAs. Therefore the total number of atoms or distance between the C-2 position and the terminal substituent is most important. Although the presence of a hydrophobic subregion was

proposed, adjacent to the alkyl subregion, it was not clear from our previous studies of a series of 2-AAs having linear alkyl substituents. However, it is now obvious that a region that accommodates the cyclopentyl or cyclohexyl rings is related to the hydrophobic subregion proposed. The cyclopentyl derivative 10b had higher affinities to A₂ receptors than the cyclohexyl derivative 10e, and these relations are inverse to the potencies of 2-(cyclopentylethoxy)adenosine and 2-(cyclohexylethoxy)adenosine as seen in the report by Ueeda et al.^{12,13} This inverse relations were also observed in the SAR of N-alkylated 2-aminoadenosines.¹⁴ These differences can be explained as a consequence of the differences between the position of the distal groups at the C-2 position, since the side chain attached to the oxygen atom or nitrogen atoms is bent at the heteroatom in the 2-alkoxyadenosines and N-alkylated 2-aminoadenosines, respectively, while the acetylenic bond in the 2-AAs is linear. The lower affinities of the phenyl derivatives 10h and 10i than those of the phenylalkoxy derivatives could be explained in a similar way. We previously proposed that the shape of the C-2 substituent of 2-alkynyladenosines may be linear.¹⁷ but this study implies that the shape of the alkyl substituent could be like cycloalkanes when they bind to A_2 receptors, because cyclopentyl derivatives 10b have more potent A_2 binding activity than 10u, with n-octynyl side chain, although they have a C-2 side chain with the same number of carbon atoms. This is supported by the SAR study of N-alkylated 2-aminoadenosines in which a lipophilic side chain alone is not sufficient to produce strong binding affinity and the ring attached to the side chain plays an important role in the binding.¹⁴ Essentially, our results of 2-AAs and 2-CAAs on A_2 affinity support the receptor models proposed.

We next investigated propargyl alcohol derivatives 10j-s for further structural requirements for the affinity and the subtype selectivity. Previously, we disclosed that the propargyl alcohol derivative 10s and its ethers could be inversely important in binding to the A_1 receptors.¹⁷ As can be seen from Table I, introduction of a hydroxyl group at the C-3" position potentiates both A_1 and A_2 binding affinities compared to the corresponding 2-CAAs 10a and 10d. Although we described above how the alkyl subregion is of limited bulk tolerance, a series of 2-AAs and 2-CAAs having a hydroxyl group at the C-3" position instead favors steric bulk for binding to both receptors. Especially, 10r showed about 200- and 60-fold more potent affinities to A_1 and A_2 receptors than those of 10h, respectively. From the viewpoint of increase in the selectivity to the receptor subtypes, introduction of the hydroxyl group is not adequate, but 10k is one of the most potent agonists so far known. These suggest that in addition to the three subregions described above there is another polar subregion adjacent to the alkyl subregions in both receptors.

Antihypertensive Activity. The potency of test compounds to decrease blood pressure (BP) and heart rate (HR) were compared in SHR. The relative potency to decrease BP was estimated on the basis of the ED_{30} value, the mean dose that produced a 30% decrease in BP. Similarly, the relative potency to decrease HR was estimated on the basis of the ED_{10} value, the mean dose that produced a 10% decrease in HR. Some compounds were examined for antihypertensive effects by oral administration in conscious SHR.

A series of 2-CAAs 10a-g showed a potent hypotensive effect without showing a marked HR decrease in doses lower than 100 μ g/kg in intravenous (iv) administration (Figures 1 and 2). These compounds produced a weak tachycardia at the doses lower than 10 μ g/kg. However,

⁽¹⁸⁾ Bruns, R. F.; Lu, G. H.; Pugsley, T. A. Characterization of the A₂ adenosine receptor labeled by [³H]NECA in rat striatal membranes. *Mol. Pharmacol.* 1986, 29, 331-346.



Figure 1. Effects of 2-(cyclopentylalkynyl)adenosines and other 2-substituted adenosines on mean blood pressure (MBP) in anesthetized male SHR. Each compound was administered iv in a cumulative manner at 5-min intervals. Values are means \pm SE of four animals.



Figure 2. Effects of 2-(cyclohexylalkynyl)adenosines on MBP in anesthetized male SHR. Each compound was administered iv in a cumulative manner at 5-min intervals. Values are means \pm SE of four animals.

the increase and decrease in HR caused by these compounds were within 6% (data not shown). The ED_{30} value for BP and ED₁₀ value for HR of each compound are shown in Table I. Compound 10b, which was the most A_2 selective in the series, had the most potent hypotensive effect and was 17- and 240-fold more potent than those of 4 and 3, respectively, which are currently the gold standards as A2 agonists. Compound 6 has been reported to be an extremely A_2 selective agent and showed a hypotensive effect accompanied by a marked tachycardia, which may result from its extremely low A₁ affinity.¹⁴ On the other hand, the tachycardia caused by 2-CAAs were weaker than that of 6, since A_1 affinities of 2-CAAs were much higher than that of 6. Generally, hypotensive effects of these nucleosides paralleled affinities for the A2 receptors

Although 2-CAAs with a hydroxyl group at the C-3''position 10j-s showed a highly potent hypotensive effect in iv administration, a marked HR decrease was also observed at doses more than $1 \mu g/kg$ as shown in Figure 3. Dose-response curves of these compounds for decreases in BP showed two phases; i.e., the first phase reached a plateau at a dose as high as 0.3 μ g/kg with a 50-60% decrease in BP and the second phase showed further decreases in BP, up to 70–80% at doses more than $3 \mu g/kg$. The decrease in BP observed in the second phase is thought to be caused by the depression of the heart via the A_1 receptor activation, since 10j-m had potent affinities for A_1 receptors. It has been known that the adenosine-induced decrease in BP and HR are mediated via A_2 and A_1 receptors, respectively. In this study, there was a positive correlation between the A_2 activity and decreases in the BP of tested compounds (r = 0.79, p < 0.001) and



Figure 3. Effects of 2-[2-(1-hydroxycycloalkyl)-1-ethyn-1-yl]adenosines and 10u on MBP and HR in anesthetized male SHR. Each compound was administered iv in a cumulative manner at 5-min intervals. Values are means \pm SE of four animals.



Figure 4. Antihypertensive effects of a single oral administration of 10b (0.1 mg/kg, \odot ; 0.3 mg/kg, \bigstar ; 1 mg/kg, \Box) on systolic blood pressure and heart rate in conscious male SHR. Values are means \pm SE of five animals. Significantly different from corresponding values of predose at p < 0.05 (*).

between the A_1 activity and decreases in the HR (r = 0.62, p < 0.05) among compounds of which the ED₁₀ is lower than 100 $\mu g/kg$. The potencies of 3 and 4 for the hypotensive effects were weaker than those of 2-CAAs, even though they had similar degrees of A_2 affinities to those nucleosides 10a, 10d, and 10i. Hence, it is possible that hypotensive effects induced by 2-CAAs may involve some other mechanism than direct A_2 receptor activation.

In the next experiment, the antihypertensive effect of 10b, which had a potent hypotensive effect in iv admin-

Table II. 2-CAAs and Their Congeners

		synthetic			crystn	IR (KBr)	
no.	R	method	yield," %	mp, °C	solvent	$(\nu_{\rm C=C}), {\rm cm}^{-1}$	formula ^b
10a	c-C ₅ H ₉	В	77	127-133	EtOH/H ₂ O	2230	C ₁₇ H ₂₁ N ₅ O ₄ ·H ₂ O
10b	c-C ₅ H ₉ CH ₂	В	80	125-127	$EtOH/H_2O$	2230	$C_{18}H_{28}N_5O_4 \cdot 2/_3H_2O$
10c	$c-C_5H_9(CH_2)_2$	В	75	108-114	EtOH/H ₂ O	2235	$C_{19}H_{25}N_5O_4 \cdot 2/_3H_2O$
10 d	$c-C_{6}H_{11}$	В	77	135-141	EtOH/H ₂ O	2220	C ₁₈ H ₂₃ N ₅ O ₄ ·H ₂ O
10e	$c-C_{6}H_{11}CH_{2}$	В	75	97-103	EtOH/H ₂ O	2235	$C_{19}H_{25}N_5O_4 \cdot 1/_2H_2O$
1 0f	$c-C_{6}H_{11}(CH_{2})_{2}$	В	77	104-111	EtOH/H ₂ O	2240	$C_{20}H_{27}N_5O_4\cdot^2/_3H_2O$
10g	$c-C_{6}H_{11}(CH_{2})_{3}$	В	77	117-127	EtOH/H ₂ O	2230	C ₂₁ H ₂₉ N ₅ O ₄ ·H ₂ O
10 h	Ph	Α	87	146-147	CHCl ₃	2210	C ₁₈ H ₁₇ N ₅ O ₄ ·H ₂ O
10 i	$Ph(CH_2)_2$	Α	55	115-120	EtOH/H ₂ O	2230	$C_{20}H_{21}N_5O_4 \cdot 1/_2H_2O$
10j	$c-C_5H_8(OH)$	В	80	138144	$EtOH/H_2O$	2230	C ₁₇ H ₂₁ N ₅ O ₅ ·H ₂ O
10k	$c-C_{6}H_{10}(OH)$	В	79	142-147	EtOH/H ₂ O	2230	C ₁₈ H ₂₃ N ₅ O ₅ ·H ₂ O
101	$c-C_7H_{12}(OH)$	В	79	foam	, -	2230	C ₁₉ H ₂₅ N ₅ O ₅ ·H ₂ O
10m	$c-C_8H_{14}(OH)$	В	79	foam		2230	C ₂₀ H ₂₇ N ₅ O ₅ - ³ / ₂ H ₂ O
10o	(CH ₃) ₂ (OH)C	Α	50	142-147	MeOH	2250	C ₁₅ H ₁₉ N ₅ O ₅ ·H ₂ O
10p	CH ₃ (CH ₂) ₂ (OH)CH	Α	58	167-169	MeOH	2230	$C_{16}H_{21}N_5O_5$
10g	CH ₃ (CH ₂) ₄ (OH)CH	Α	60	177-179	EtOAc	2230	$C_{18}H_{25}N_5O_5$
10u	$CH_3(CH_2)_5$	В	86	101-103	EtOH/H ₂ O	2230	$C_{18}H_{25}N_5O_4 \cdot 1/_5H_2O$

^a Overall yields from 7. ^bAll compounds had satisfactory C, H, and N microanalytical data within ±0.4% of the theoretical value.

istration, was investigated in oral administration in conscious SHR. Oral administration of 10b at a dose of 0.1, 0.3, or 1 mg/kg decreased systolic blood pressure of SHR dose-dependently as shown in Figure 4. The duration of the antihypertensive effect of this compound was maintained for at least 7 h after administration. Compound 10b caused a significant tachycardia at 1 or 3 h after the oral administration, but it recovered thereafter. Thus, 2-CAAs are promising as antihypertensive agents that should be considered for further detailed preclinical evaluation.

In conclusion, 2-CAAs were A_2 selective adenosine agonists. In the C-2 binding region of adenosine, compounds often have potent and/or selective A_2 activity from introduction of an acetylenic group at the C-2 position followed by one methylene residue further followed by a hydrophobic substituent such as a cycloalkyl ring at the terminal position of the alkynyl side chain. Above all, 2-(3-cyclopentyl-1-propyn-1-yl)adenosine (10b) showed 70-fold A_2 selectivity and potent antihypertensive activity in both iv and oral administration. Introduction of a hydroxyl group into the C-3" position of the alkynyl side chain increased the affinity for both receptor subtypes, while it also produced a potent HR decreasing effect by the increase in the A_1 affinity.

Experimental Section

Melting points were measured on a Yamato MP-21 melting point apparatus and are uncorrected. Elemental analyses were done at Yanaco MT-5. The ¹H NMR spectra were recorded on a JEOL GSX-400 (400 MHz) spectrometer with tetramethylsilane as an internal standard. Chemical shifts are reported in parts per million (δ), and signals are expressed as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), or br (broad). All exchangeable protons were detected by addition of D₂O. UV absorption spectra were recorded with a Shimadzu UV-160A spectrophotometer. IR spectra was recorded with Hitachi 260-50 spectrometer with KBr pellets. TLC was done on Merck Kieselgel F254 precoated plates. The silica gel used for column chromatography was Merck Kieselgel 60 (70-230 mesh). All acetylene compounds except for 4-cyclopentyl-1-butyne (13a), 4-cyclohexyl-1-butyne (13b), and 5-cyclohexyl-1-pentyne (13c) were purchased from Hydrus Chemical Co. Physical and analytical data for the compounds 10a-u were shown in Table II.

General Method for the Preparation of 2-AAs (10h,i,o-q). Compound 8 (393 mg, 1 mmol), CuI (9.5 mg, 0.05 mmol), bis-(triphenylphosphine)palladium dichloride (36 mg, 10 mol %), triethylamine (0.16 mL, 1.2 mmol), and terminal alkyne (1.2 equiv of phenylacetylene, 4-phenyl-1-butyne, 2-methyl-3-butyn-2-ol, 1-hexyn-3-ol, or 1-octyn-3-ol) in DMF (7 mL) was heated at 80 °C for several hours under an argon atmosphere. After the starting material was completely consumed, judged by TLC (CHCl₃/EtOH = 10:1, v/v), the reaction mixture was concentrated to dryness under reduced pressure. The residue was dissolved in CHCl₃, and H₂S gas was introduced to the solution (\sim 30 s) followed by N₂ gas. The suspension was filtered through a Celite pad and washed with CHCl₃. The combined filtrate and washings were concentrated to dryness in vacuo, and the residue was purified by a silica gel column using an appropriately mixed MeOH-CHCl₃ solvent system.

2-(Phenylethyn-1-yl)adenosine (10h). Compound 10h (373 mg, 97%) was obtained from 8 with phenylacetylene: UV (H₂O) λ_{max} 261 nm (23 000), 285 (20 800), 304 (19 000), λ_{min} 239 (17 100), 275 (19 400), 293 (18 800); UV (0.05 N HCl) λ_{max} 274 nm (16 600), 320 (21 300), λ_{min} (10 800), 291 (12 400); NMR (DMSO-d₆) δ 3.67 (2 H, m, H-5'a,b), 4.03 (1 H, m, H-4'), 4.19 (1 H, dd, H-3'), 4.58 (1 H, t, H-2'), 5.93 (1 H, d, H-1'), 7.42 (2 H, br s, NH₂), 7.45–7.67 (5 H, m, Ph), 8.49 (1 H, s, H-8).

2-(4-Phenyl-1-butyn-1-yl)adenosine (10i). Compound **10i** (240 mg, 61%) was obtained from 8 with 4-phenyl-1-butyne: UV (H₂O) λ_{max} 270 nm (14 500), 287 sh (9900), λ_{min} 247 nm (7300); UV (0.05 N HCl) λ_{max} 271 nm (16000), 295 (11700), λ_{min} 247 (7300), 284 (10700); NMR (DMSO-d₆) δ 2.71 (2 H, t, C=CH₂), 2.87 (2 H, t, PhCH₂), 3.54-3.69 (2 H, m, H-5'a,b), 3.95 (1 H, m, H-4'), 4.13 (1 H, dd, H-3'), 4.54 (1 H, t, H-2'), 5.85 (1 H, d, H-1'), 7.20-7.33 (5 H, m, Ph), 7.42 (2 H, br s, NH₂), 8.40 (1 H, s, H-8).

2-(3-Hydroxy-3-methyl-1-butyn-1-yl)adenosine (100). Compound 100 (190 mg, 55%) was obtained from 8 with 2methyl-3-butyn-2-ol: UV (H₂O) λ_{max} 270 nm (14 300), 289 (9800), λ_{min} 245 (6200), 281 (9500); UV (0.05 N HCl) λ_{max} 271 nm (15 500), 294 (10 700), λ_{min} 246 (6600), 283 (9800); NMR (DMSO-d₆) δ 1.46 (6 H, s, Me), 3.40–3.69 (2 H, m, H-5'a,b), 3.94 (1 H, m, H-4'), 4.12 (1 H, dd, H-3'), 4.48 (1 H, t, H-2'), 5.13–5.17 (2 H, m, OH × 2), 5.45 (1 H, d, OH), 5.55 (1 H, s, C=COOH), 5.87 (1 H, d, H-1'), 7.44 (2 H, br s, NH₂), 8.41 (1 H, s, H-8).

2-(3-Hydroxy-1-hexyn-1-yl)adenosine (10**p**). Compound 10**p** (235 mg, 64%) was obtained from 8 with 1-hexyn-3-ol: UV (H₂O) λ_{max} 265 sh nm (14 100), 270 (14 500), 289 (9900), λ_{min} 246 (7600), 282 (9700); UV (0.05 N HCl) λ_{max} 265 sh nm (12 500), 272 (14 000), 297 (9900), λ_{min} 247 (5700), 283 (8800); NMR (DMSO- d_{e}) δ 0.92 (3 H, t, Me), 1.33–1.76 (4 H, m, CH₂), 3.58 2 H, m, H-5'a,b), 3.95 (1 H, m, H-4'), 4.12 (1 H, dd, H-3'), 4.48 (1 H, t, H-2'), 5.86 (1 H, d, H-1'), 7.44 (2 H, br s, NH₂), 8.41 (1 H, s, H-8).

2-(3-Hydroxy-1-octyn-1-yl)adenosine (10q). Compound 10q (260 mg, 67%) was obtained from 8 with 1-octyn-3-ol: UV (H₂O) λ_{max} 270 nm (13 800), 289 (9500), λ_{min} 245 (6500), 281 (9400); UV (0.05 N HCl) λ_{max} 271 nm (15 000), 294 (10 900), λ_{min} 246 (6700), 282 (9900); NMR (DMSO- d_e) δ 0.88 (3 H, t, Me), 1.24–1.67 (8 H, m, CH₂), 3.53–3.69 (2 H, m, H-5'a,b), 3.95 (1 H, m, H-4'), 4.12 (1 H, dd, H-3'), 4.51 (1 H, dd, H-2'), 5.15–5.17 (2 H, m, OH × 2), 5.45 (1 H, d, OH), 5.50 (1 H, d, OH), 5.86 (1 H, d H-1'), 7.44 (2 H, br s, NH₂), 8.41 (1 H, s, H-8).

Synthesis of Bromoalkylcycloalkanes. A mixture of 2cyclopentylethanol (11a) (5.0 g, 44 mmol), concentrated H_2SO_4 (1.9 mL), and HBr (48%, 11.2 g) was heated under reflux for 6 h. The cooled mixture was diluted with CHCl₃, and the whole was washed several times with H₂O. The separated organic phase was dried (Na₂SO₄), and the solvent was removed in vacuo. The resulting oil was distilled under vacuum to give 12a (5.6 g, 72%): bp 77 °C (16 mmHg); NMR (DMSO-d₆) δ 1.67-1.91 (11 H, m, cyclopentyl-CH₂), 3.51 (2 H, t, CH₂Br). In the case of 2-cyclohexaneethanol (11b) (10 g, 78 mmol) or 1-cyclohexyl-3-propanol (11c) (10 g, 70.3 mmol), a mixture with concentrated H₂SO₄ (3.8 mL) and HBr (48%, 22.3 g) was heated under reflux for 7 h. A similar workup and distillation gave 12b (10.4 g, 70%) [bp 78-79 °C (3 mmHg); NMR (DMSO-d₆) δ 0.87-1.72 (13 H, m, cyclohexyl-CH₂), 3.54 (2 H, t, CH₂Br)] or 12c (10.8 g, 75%) [bp 114-124 °C (3 mmHg); NMR (DMSO-d₆) δ 0.85-1.83 (15 H, m, cyclohexyl-CH₂CH₂), 3.50 (2 H, t, CH₂Br)], respectively.

Synthesis of Cycloalkylalkylacetylenes. Compound 12a (5.2 g, 29.4 mmol) was added to a solution of lithium acetylideethylenediamine complex (1.1 equiv) in DMSO (10 mL) at 0 °C under argon. After the mixture was stirred for 5 h at room temperature, H₂O was added. The mixture was extracted with CHCl₃, washed with H₂O, and dried (Na₂SO₄). The solvent was removed in vacuo and the oil was distilled under vacuum to give 13a (2.1 g, 59%): bp 56-59 °C (20 mmHg); IR (KBr) 2115 cm⁻¹; NMR (DMSO- d_6) δ 1.02-2.16 (13 H, m, cyclopentyl-CH₂CH₂), 2.71 (1 H, t, C=CH). 4-Cyclohexyl-1-butyne (13b) and 5-cyclohexyl-1-pentyne (13c) were obtained similarly from 12b and 12c, respectively. 13b (4.3 g, 60%): bp 30 °C (3 mmHg); IR (KBr) 2115 cm⁻¹; NMR (DMSO-d₆) δ 0.83-2.17 (15 H, m, cyclohexyl-CH₂CH₂), 2.69 (1 H, t, C=CH). 13c (4.3 g, 59%): bp 50 °C (3 mmHg); IR (KBr) 2115 cm⁻¹; NMR (DMSO- d_6) δ 0.83–2.14 (17 H, m, cyclohexyl-CH₂CH₂CH₂), 2.71 (1 H, t, C=CH).

General Method for the Preparation of 9a-g,j-m,u. A mixture of 7 (540 mg, 1 mmol), CuI (9.5 mg, 0.05 mmol), bis-(triphenylphosphine)palladium dichloride (18 mg, 5 mol %), triethylamine (0.16 mL, 1.2 mmol), and 1-octyne (0.18 mL, 1.2 mmol) in 1,4-dioxane (10 mL) was stirred for 2 h at room temperature under argon (judged by TLC; $CHCl_{9}/EtOAc = 4:1, v/v$). The solvent was removed in vacuo and the residue dissolved in CHCl₃ (50 mL) and triethylamine (0.1 mL), which was washed with saturated aqueous ethylenediaminetetraacetic acid disodium salt $(2 \times 10 \text{ mL})$ and brine (20 mL). The separated organic phase was dried (Na₂SO₄) and the solvent was removed in vacuo. The residue was purified by a silica gel column with CHCl₃/EtOAc (4:1) to give 9u (490 mg, 95%) as an oil: IR (KBr) 2235 cm⁻¹; NMR (CDCl₃) δ 0.90 (3 H, t, Me), 1.30–1.71 (8 H, m, CH₂), 2.08 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.18 (3 H, s, Ac), 2.49 (2 H, t, C=CCH₂), 4.41 (2 H, m, H-5'a,b), 4.47 (1 H, dd, H-4'), 5.57 (1 H, dd, H-3'), 5.80 (1 H, t, H-2'), 6.33 (1 H, d, H-1'), 8.31 (1 H, s, H-8). Compounds 9a-g,j-m were prepared similarly.

9-(2,3,5-Tri-O-acetyl-1- β -D-ribofuranosyl)-6-chloro-2-(2cyclopentyl-1-ethyn-1-yl)purine (9a). From 7 with cyclopentylacetylene (0.14 mL, 1.2 mmol) for 1 h, 430 mg of 9a (85% as an oil) was obtained: IR (KBr) 2240 cm⁻¹; NMR (CDCl₃) δ 1.61-2.14 (8 H, m, c-C₅H₉), 2.08 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.90 (1 H, t, C=CCH), 4.40-4.42 (2 H, m, H-5'a,b), 4.46 (1 H, dd, H-4'), 5.58 (1 H, dd, H-3'), 5.80 (1 H, t, H-2'), 6.32 (1 H, d, H-1'), 8.30 (1 H, s, H-8).

9-(2,3,5-Tri-O-acetyl-1- β -D-ribofuranosyl)-6-chloro-2-(3cyclopentyl-1-propyn-1-yl)purine (9b). From 7 with 3cyclopentyl-1-propyne (0.16 mL, 1.2 mmol) for 1 h, 480 mg of 9b (92% as an oil) was obtained: IR (KBr) 2240 cm⁻¹; NMR (CDCl₃) δ 1.33–2.23 (11 H, m, c-C₅H₉), 2.08 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.49 (2 H, t, C=CCH₂), 4.41 (2 H, m, H-5'a,b), 4.47 (1 H, dd, H-4'), 5.58 (1 H, dd, H-3'), 5.81 (1 H, t, H-2'), 6.31 (1 H, d, H-1'), 8.30 (1 H, s, H-8).

9-(2,3,5-Tri-*O*-acetyl-1-β-D-ribofuranosyl)-6-chloro-2-(4cyclopentyl-1-butyn-1-yl)purine (9c). From 7 with 4-cyclopentyl-1-butyne (13a) (0.15 mL, 1.2 mmol) for 1 h, 460 mg of 9c (86% as an oil) was obtained: IR (KBr) 2240 cm⁻¹; NMR (CDCl₃) δ 1.10–1.97 (11 H, m, c-C₅H₉), 2.08 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.50 (2 H, t, C=CCH₂), 4.41 (2 H, m, H-5'a,b), 4.47 (1 H, dd, H-4'), 5.57 (1 H, dd, H-3'), 5.81 (1 H, t, H-2'), 6.32 (1 H, d, H-1'), 8.31 (1 H, s, H-8).

9-(2,3,5-Tri-O-acetyl-1- β -D-ribofuranosyl)-6-chloro-2-(2cyclohexyl-1-ethyn-1-yl)purine (9d). From 7 with cyclohexylacetylene (0.15 mL, 1.2 mmol) for 1 h, 480 mg of 9d (92% as an oil) was obtained: IR (KBr) 2240 cm⁻¹; NMR (CDCl₃) δ 1.36–1.97 (10 H, m, c-C₆H₁₁), 2.09 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.66 (1 H, t, C=CCH), 4.41 (2 H, m, H-5'a,b), 4.46 (1 H, dd, H-4'), 5.59 (1 H, dd, H-3'), 5.81 (1 H, t, H-2'), 6.31 (1 H, d, H-1'), 8.30 (1 H, s, H-8).

9-(2,3,5-Tri-*O*-acetyl-1-β-D-ribofuranosyl)-6-chloro-2-(3cyclohexyl-1-propyn-1-yl)purine (9e). From 7 with 3-cyclohexyl-1-propyne (0.17 mL, 1.2 mmol) for 1 h, 450 mg of 9e (84% as an oil) was obtained: IR (KBr) 2240 cm⁻¹; NMR (CDCl₃) δ 1.0-2.0 (11 H, m, c-C₆H₁₁), 2.08 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.38 (2 H, t, C=CCH₂), 4.41-4.42 (2 H, m, H-5'a,b), 4.47 (1 H, dd, H-4'), 5.58 (1 H, dd, H-3'), 5.81 (1 H, t, H-2'), 6.31 (1 H, d, H-1'), 8.30 (1 H, s, H-8).

9-(2,3,5-Tri-*O***-acetyl-1**- β -D-ribofuranosyl)-6-chloro-2-(4cyclohexyl-1-butyn-1-yl)purine (9f). From 7 with 3-cyclohexyl-1-butyne (13b) (0.16 g, 1.2 mmol) for 1 h, 490 mg of 9f (89% as an oil) was obtained: IR (KBr) 2240 cm⁻¹; NMR (CDCl₃) δ 0.88–1.77 (13 H, m, c-C₆H₁₁), 2.08 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.49 (2 H, t, C=CCH₂), 4.41 (2 H, m, H-5'a,b), 4.47 (1 H, dd, H-4'), 5.57 (1 H, dd, H-3'), 5.80 (1 H, t, H-2'), 6.32 (1 H, d, H-1'), 8.31 (1 H, s, H-8).

9-(2,3,5-Tri-O-acetyl-1-β-D-ribofuranosyl)-6-chloro-2-(5cyclohexyl-1-pentyn-1-yl)purine (9g). From 7 with 5-cyclohexyl-1-pentyne (13c) (0.18 g, 1.2 mmol) for 1 h, 510 mg of 9g (91% as an oil) was obtained: IR (KBr) 2235 cm⁻¹; NMR (CDCl₃) δ 0.84–1.72 (15 H, m, c-C₆H₁₁CH₂CH₂), 2.08 (3 H, s, Ac), 2.16 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.46 (2 H, t, C=CCH₂), 4.41 (2 H, m, H-5'a,b), 4.47 (1 H, dd, H-4'), 5.57 (1 H, dd, H-3'), 5.80 (1 H, t, H-2'), 6.33 (1 H, d, H-1'), 8.31 (1 H, s, H-8).

9-(2,3,5-Tri-O-acetyl-1- β -D-ribofuranosyl)-6-chloro-2-[2-(1-hydroxycyclopentyl)-1-ethyn-1-yl]purine (9j). From 7 with 1-ethynyl-1-cyclopentanol (0.14 mL, 1.2 mmol) for 1 h, 450 mg of 9j (86% as an oil) was obtained: IR (KBr) 2235 cm⁻¹; NMR (CDCl₃) δ 1.77–1.94 (8 H, m, c-C₅H₈), 2.09 (3 H, s, Ac), 2.13 (3 H, s, Ac), 2.18 (3 H, s, Ac), 2.85 (1 H, s, OH), 4.43–4.52 (3 H, m, H-4', 5'a,b), 5.72 (1 H, dd, H-3'), 5.90 (1 H, t, H-2'), 6.22 (1 H, d, H-1'), 8.28 (1 H, s, H-8).

9-(2,3,5-Tri-*O*-acetyl-1- β -D-ribofuranosyl)-6-chloro-2-[2-(1-hydroxycyclohexyl)-1-ethyn-1-yl]purine (9k). From 7 with 1-ethynyl-1-cyclohexanol (0.15 g, 1.2 mmol) for 1 h, 500 mg of 9k (93% as an oil) was obtained: IR (KBr) 2230 cm⁻¹; NMR (CDCl₃) δ 1.64–2.13 (10 H, m, c-C₆H₁₀), 2.10 (3 H, s, Ac), 2.12 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.90 (1 H, s, OH), 4.46–4.51 (3 H, m, H-4', 5'a,b), 5.75 (1 H, dd, H-3'), 5.89 (1 H, t, H-2'), 6.21 (1 H, d, H-1'), 8.26 (1 H, s, H-8).

9-(2,3,5-Tri-O-acetyl-1- β -D-ribofuranosyl)-6-chloro-2-[2-(1-hydroxycycloheptyl)-1-ethyn-1-yl]purine (91). From 7 with 1-ethynyl-1-cycloheptanol (0.17 g, 1.2 mmol) for 1 h, 500 mg of 91 (91% as an oil) was obtained: IR (KBr) 2230 cm⁻¹; NMR (CDCl₃) δ 1.64–2.16 (12 H, m, c-C₇H₁₂), 2.10 (3 H, s, Ac), 2.12 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.82 (1 H, s, OH), 4.44–4.52 (3 H, m, H-4', 5'a,b), 5.75 (1 H, dd, H-3'), 5.90 (1 H, t, H-2'), 6.21 (1 H, d, H-1'), 8.27 (1 H, s, H-8).

9-(2,3,5-Tri-*O***-acetyl-**1- β -D-**ribofuranosyl**)-**6-chloro-2-[2-(1-hydroxycyclooctyl**)-1-ethyn-1-yl]purine (**9m**). From 7 with 1-ethynyl-1-cyclooctanol (0.18 g, 1.2 mmol) for 1 h, 500 mg of **9m** (89% as an oil) was obtained: IR (KBr) 2230 cm⁻¹; NMR (CDCl₃) δ 1.52–2.14 (14 H, m, c-C₈H₁₄), 2.10 (3 H, s, Ac), 2.12 (3 H, s, Ac), 2.17 (3 H, s, Ac), 2.65 (1 H, s, OH), 4.44–4.51 (3 H, m, H-4', 5'a,b), 5.75 (1 H, dd, H-3'), 5.89 (1 H, t, H-2'), 6.20 (1 H, d, H-1'), 8.25 (1 H, s, H-8).

Conversion of 9 into 10. A solution of 9u (490 mg, 0.95 mmol) in concentrated NH₄OH (30 mL) and 1,4-dioxane (60 mL) in a steel sealed tube was heated at 70 °C for 20 h. The solvent was removed in vacuo and the residue was purified by a silica gel column with CHCl₃/MeOH (5:1) to give 2-(1-octyn-1-yl)adenosine (10u, 320 mg, 90%): mp 101-103 °C (lit.¹⁷ mp 101-103 °C).

Similarly 9a-g,j-m were converted into 10a-g,j-m.

2-(2-Cyclopentyl-1-ethyn-1-yl)adenosine (10a). From 9a (430 mg, 0.85 mmol), 10a (280 mg, 91%) was obtained: UV (H₂O) λ_{max} 270 nm (15 900), 286 sh (11 000), λ_{min} 247 (8000); UV (0.05 N HCl) λ_{max} 272 nm (17 000), 290 sh (12 000), λ_{min} 247 (7500); NMR (DMSO-d₆) δ 1.56–1.99 and 2.82–2.86 (9 H, m, c-C₅H₉), 3.52–3.70 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.52 (1 H, dd, H-2'), 5.18 (1 H, d, 2'- or 3'-OH), 5.22 (1 H, d, H-3'), 5.45 (1 H, d, 2'- or 3'-OH), 7.41 (2 H, br s, NH₂), 8.38 (1 H, s, H-8).

2-(3-Cyclopentyl-1-propyn-1-yl)adenosine (10b). From 9b (480 mg, 0.92 mmol), 10b (300 mg) was obtained: UV (H₂O) λ_{max} 271 nm (14900), 286 sh (10500), λ_{min} 246 (6800); UV (0.05 N HCl) λ_{max} 272 (16700), 293 (12000), λ_{min} 248 (6600), 284 (11600); NMR (DMSO-d₆) δ 1.29–2.10 (9 H, m, c-C₅H₉), 2.40 (2 H, d, C=CCH₂), 4.12 (1 H, dd, H-3'), 4.53 (1 H, dd, H-2'), 5.12 (1 H, d, 2' or 3'-OH), 5.25 (1 H, t, 5'-OH), 5.41 (1 H, d, 2' or 3'-OH), 5.88 (1 H, d, H-1'), 7.36 (2 H, br s, NH₂), 8.39 (1 H, s, H-8).

2-(4-Cyclopentyl-1-butyn-1-yl)adenosine (10c). From 9c (460 mg, 0.86 mmol), 10c (290 mg, 87%) was obtained: UV (H₂O) λ_{max} 270 nm (14 300), 286 sh (9400), λ_{min} 246 (7200); UV (0.05 N HCl) λ_{max} 271 nm (16 300), 290 (10 300), λ_{min} 245 nm (7700); NMR (DMSO- d_{θ}) δ 1.09–1.95 (11 H, m, c-C₅H₉CH₂), 2.40 (2 H, t, C= CCH₂), 3.53–3.68 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.15 (1 H, dd, H-3'), 4.53 (1 H, dd, H-2'), 5.16 (1 H, d, 2' - or 3'-OH), 5.21 (1 H, t, 5'-OH), 5.43 (1 H, d, 2' - or 3'-OH), 5.85 (1 H, d, H-1'), 7.41 (2 H, br s, NH₂), 8.38 (1 H, s, H-8).

2-(2-Cyclohexyl-1-ethyn-1-yl)adenosine (10d). From **9d** (480 mg, 0.92 mmol), **10d** (290 mg, 84%) was obtained: UV (H₂O) λ_{max} 270 nm (14700), 285 sh (10500), λ_{min} 246 (6900); UV (0.05 N HCl) λ_{max} 271 nm (15800), 294 (12000), λ_{min} 247 (6600), 284 (11300); NMR (DMSO-d₆) δ 1.29–1.84 and 2.59–2.65 (11 H, m, c-C₆H₁₁), 3.54–3.68 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.51 (1 H, dd, H-2'), 5.86 (1 H, d, H-1'), 7.43 (2 H, br s, NH₂), 8.39 (1 H, s, H-8).

2-(3-Cyclohexyl-1-propyn-1-yl)adenosine (10e). From 9e (450 mg, 0.84 mmol), 10e (290 mg, 89%) was obtained: UV (H₂O) λ_{max} 270 nm (13 700), 286 sh (9500), λ_{min} 246 (6200); UV (0.05 N HCl) λ_{max} 272 nm (15 400), 292 (10 700), λ_{min} 247 (6200), 285 (10 600); NMR (DMSO- d_6) δ 1.02–1.83 (11 H, m, c-C₆H₁₁), 2.31 (2 H, d, C=CCH₂), 3.53–3.68 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.11 (1 H, dd, H-3'), 4.53 (1 H, dd, H-2'), 5.18 (1 H, d, 2'- or 3'-OH), 5.23 (1 H, t, 5'-OH), 5.45 (1 H, d, 2'- or 3'-OH), 5.85 (1 H, d, H-1'), 7.41 (2 H, br s, NH₂), 8.38 (1 H, s, H-8).

2-(4-Cyclohexyl-1-butyn-1-yl)adenosine (10f). From **9f** (490 mg, 0.89 mmol), **10f** (310 mg, 86%) was obtained: UV (H₂O) λ_{max} 270 nm (14600), 285 sh (10200), λ_{min} 246 (6800); UV (0.05 N HCl) λ_{max} 272 nm (17000), 288 sh (11500), λ_{min} 247 (7200); NMR (DMSO-d₆) δ 0.87–1.76 (13 H, m, c-C₆H₁₁CH₂), 2.41 (2 H, t, C=CCH₂), 3.53–3.68 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.53 (1 H, dd, H-2'), 5.16 (1 H, d, 2'- or 3'-OH), 5.22 (1 H, t, 5'-OH), 5.44 (1 H, d, 2'- or 3'-OH), 5.85 (1 H, d, H-1'), 7.41 (2 H, br s, NH₂), 8.38 (1 H, s, H-8).

2-(5-Cyclohexyl-1-pentyn-1-yl)adenosine (10g). From 9g (510 mg, 0.91 mmol), 10g (320 mg, 85%) was obtained: UV (H₂O) λ_{max} 270 nm (13 800), 285 sh (9700), λ_{min} 246 (6200); UV (0.05 N HCl) λ_{max} 272 nm (16900), 290 sh (11600), λ_{min} 246 (6900); NMR (DMSO- d_6) δ 0.86–1.71 (15 H, m, c-C₆H₁₁CH₂CH₂), 2.38 (2 H, t, C=CCH₂), 3.55–3.68 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.53 (1 H, dd, H-2'), 5.18 (1 H, d, 2'- or 3'-OH), 5.24 (1 H, t, 5'-OH), 5.45 (1 H, d, 2'- or 3'-OH), 5.85 (1 H, d, H-1'), 7.42 (2 H, br s, NH₂), 8.39 (1 H, s, H-8).

 $\begin{array}{l} \textbf{2-[2-(1-Hydroxycyclopentyl)-1-ethyn-1-yl]adenosine (10j).}\\ From 9j (450 mg, 0.86 mmol), 10j (300 mg, 93\%) was obtained:\\ UV (H_2O) \lambda_{max} 270 nm (7500), 289 (9100), \lambda_{min} 247 (6600), 281 (9100); UV (0.05 N HCl) \lambda_{max} 272 nm (14\,100), 295 (10\,300), \lambda_{min} 248 (6600), 283 (9500); NMR (DMSO-d_6) \delta 1.66-1.94 (8 H, m, c-C_8H_8-H), 3.53-3.69 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.49 (1 H, dd, H-2'), 5.16-5.18 (2 H, m, OH <math display="inline">\times$ 2), 5.42 (1 H, br s, OH), 5.46 (1 H, d, OH), 5.87 (1 H, d, H-1'), 7.43 (2 H, br s, NH_2), 8.41 (1 H, s, H-8). \end{array}

2-[2-(1-Hydroxycyclohexyl)-1-ethyn-1-yl]adenosine (10k). From **9k** (500 mg, 0.93 mmol), **10k** (310 mg, 85%) was obtained: UV (H₂O) λ_{max} 270 nm (14 400), 287 sh (9600), λ_{min} 246 (7100); UV (0.05 N HCl) λ_{max} 271 nm (15600), 294 (10800), λ_{min} 247 (7300), 284 (10300); NMR (DMSO-d₆) δ 1.25–1.87 (10 H, m, c-C₆H₁₀-H), 3.56–3.71 (2 H, m, H-5'a,b), 3.97 (1 H, dd, H-4'), 4.15 (1 H, dd, H-3'), 4.50 (1 H, dd, H-2'), 5.11 (1 H, d, OH), 5.20 (1 H, t, 5'-OH), 5.44 (1 H, d, OH), 5.50 (1 H, d, OH), 5.89 (1 H, d, H-1'), 7.41 (2 H, br s, NH₂), 8.39 (1 H, s, H-8).

2-[2-(1-Hydroxycycloheptyl)-1-ethyn-1-yl]adenosine (101). From **91** (500 mg, 0.91 mmol), **101** (320 mg) was obtained: UV (H₂O) λ_{max} 270 nm (13100), 289 (9000), λ_{min} 246 (6100), 283 (8800); UV (0.05 N HCl) λ_{max} 272 (14 300), 296 (10 300), λ_{min} 247 (6300), 284 (9500); NMR (DMSO-d₆) δ 1.49–2.00 (12 H, m, c-C₇H₁₂-H), 3.53–3.69 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.50 (1 H, dd, H-2'), 5.15–5.17 (2 H, m, OH × 2), 5.41 (1 H, br s, OH), 5.45 (1 H, d, OH), 5.88 (1 H, d, H-1'), 7.44 (2 H, br s, NH₂), 8.41 (1 H, s, H-8).

2-[2-(1-Hydroxycyclooctyl)-1-ethyn-1-yl]adenosine (10m). From **9m** (500 mg, 0.89 mmol), **10m** (330 mg, 89%) was obtained: UV (H₂O) λ_{max} 271 nm (13000), 289 (9100), λ_{min} 246 (6000), 282 (8900); UV (0.05 N HCl) λ_{max} 272 nm (13900), 295 (10 200), λ_{min} 248 (6200), 283 (9400); NMR (DMSO-d₆) δ 1.45–1.93 (14 H, m, c-C_gH₁₄-H), 3.54–3.68 (2 H, m, H-5'a,b), 3.95 (1 H, dd, H-4'), 4.12 (1 H, dd, H-3'), 4.50 (1 H, t, H-2'), 5.87 (1 H, d, H-1'), 7.45 (2 H, br s, NH₂), 8.41 (1 H, s, H-8).

Adenosine Receptor Binding Assay. The A_1 and A_2 receptor binding assays were done by previously described methods.¹⁶ Briefly, the A_1 receptor binding assay was done in rat brain membranes using [³H]CHA as a radioligand. The reaction mixture containing [³H]CHA, adenosine deaminase, a test compound solution, and the A₁ receptor preparation was incubated at 23 °C for 2 h. After the incubation, the reaction mixture was filtrated with a glass fiber filter under reduced pressure. Nonbound radioactivity was removed by washing the filter with ice-cold buffer. The radioactivity of the glass fiber filter was counted in a liquid scintillation counter. The A2 receptor binding assay was done in rat striatal membranes using [³H]NECA in the presence of 50 nM CPA. Nonspecific binding of [³H]CHA and [³H]NECA were defined as binding in the presence of 10 mM (R)-phenyliso-propyladenosine and 100 μ M CPA, respectively. IC₅₀ values were calculated from a nonlinear, log transformation of specific binding data from the ligand binding assay and converted to K_i values as described previously.¹⁶

Blood Pressure and Heart Rate. Effects of the compound on BP and HR in iv administration were measured in anesthetized male SHR as described previously.¹⁶

The solution of the test compound $(0.03-100 \ \mu g/mL)$ was administered iv at 5-min intervals in a cumulative manner. Changes in BP and HR were expressed in terms of percent changes of their control values. The relative potency to decrease BP was estimated on the basis of the ED₃₀ value, the mean dose that produced a 30% decrease in BP of SHR. Similarly, the relative potency to decrease HR was estimated on the basis of the ED₁₀ value, the mean dose that produced a 10% decrease in HR of SHR. Antihypertensive effect of the compound in oral administration was also examined in conscious SHR as described previously.¹⁶

Acknowledgment. The authors thank Yoshinori Sekimoto for his skillful technical assistance. This investigation was supported in part by Grants-in-Aid for Developmental Scientific Research (A.M.) and Scientific Research on Priority Areas (A.M.) from the Ministry of Education, Science, and Culture of Japan.

Registry No. 8, 35109-88-7; 9a, 141345-24-6; 9b, 141345-25-7; 9c, 141345-26-8; 9d, 141345-27-9; 9e, 141345-28-0; 9f, 141345-29-1; 9g, 141345-30-4; 9j, 141345-31-5; 9k, 141345-32-6; 9l, 141345-33-7; 9m, 141345-34-8; 9u, 133560-14-2; 10a, 141345-09-7; 10b, 141345-10-0; 10c, 141345-11-1; 10d, 141345-12-2; 10e, 141345-13-3; 10f, 141345-14-4; 10g, 141345-15-5; 10h, 90596-70-6; 10i, 141345-16-6; 10j, 141345-17-7; 10k, 141345-18-8; 10l, 141345-19-9; 10m, 141345-20-2; 10o, 141345-21-3; 10p, 141345-22-4; 10q, 141345-23-5; 10u, 90596-75-1; 11a, 766-00-7; 11b, 4442-79-9; 11c, 1124-63-6; 12a, 18928-94-4; 12b, 1647-26-3; 12c, 34094-21-8; 13a, 141345-07-5; 13b, 141345-08-6; 13c, 5963-75-7; c-C₅H₉C==CH, 930-51-8; c-C₅H₉CH₂C=CH, 116279-08-4; c-C₆H₁₁C=CH, 931-48-6; c-C₆H₁₁CH₂C=CH, 17715-00-3; PhC=CH, 536-74-3; Ph-(CH₂)₂C=CH, 16520-62-0; c-C₅H₈(OH)C=CH, 17356-19-3; c-C₆H₁₀(OH)C=CH, 78-27-3; c-C₇H₁₂(OH)C=CH, 2809-78-1; c-C₈H₁₄(OH)C=CH, 55373-76-7; (CH₈)₂(OH)CC=CH, 115-19-5; $CH_{3}(CH_{2})_{2}(OH)CHC = CH, 15352-98-4; CH_{3}(CH_{2})_{4}(OH)CHC =$ CH, 37911-28-7; CH₃(CH₂)₅C=CH, 629-05-0; LiC=CH·H₂NC-H₂CH₂NH₂, 50475-76-8.