ORIGINAL RESEARCH



Synthesis and biological evaluation of novel 2',4',5'trimethoxyflavonol derivatives as anti-inflammatory and antimicrobial agents

Girish D. Hatnapure · Ashish P. Keche · Atish H. Rodge · Rajesh H. Tale · Satish S. Birajdar · Mahendra J. Pawar · Vandana M. Kamble

Received: 28 February 2013 / Accepted: 25 May 2013 © Springer Science+Business Media New York 2013

Abstract A series of novel 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (flavonol) derivatives (2a-u) of biological interest have been prepared via CLAISEN-SCHMIDT condensation followed by ALGAR-FLYNN-OYAMADA reaction and to search for the potent nonsteroidal anti-inflammatory agents from this novel series. All the synthesized compounds have been screened for their in vitro proinflammatory cytokines tumor necrosis factor (TNF- α) and interleukin-6 (IL-6) inhibitory activity along with antimicrobial activity. As many as three compounds viz. 2h, 2l, and 2q from this novel series were found to be potent TNF- α and IL-6 inhibitor (up to 72–81 % TNF- α and 86-92 % IL-6 inhibitory activity) but at 10 µM concentration as compared with the standard dexamethasone (71 %TNF-a and 84 % IL-6 inhibitory activities at 1 µM concentration). While the compounds 2d, 2m, 2n, and 2s were found to be potent antimicrobial agent showing even 2-2.5-fold more potency than that of standard ciprofloxacin and miconazole at the same MIC value of 10 µg/mL.

G. D. Hatnapure · A. P. Keche · S. S. Birajdar Department of Chemistry, Maharashtra Udayagiri Mahavidyalaya, Udgir, India

V. M. Kamble (⊠) Department of Chemistry, The Institute of Science, Mumbai, India e-mail: m.vandana248@gmail.com

A. H. Rodge · R. H. Tale Dendrimer and Bio-organic Chemistry Laboratory, School of Chemical Science, S. R. T. M. University, Nanded, India

M. J. Pawar

Department of Nanotechnology, Art, Commerce and Science College, Amravati, India

Keywords Flavonoid · NSAIDs · ALGAR–FLYNN–OYAMADA anti-inflammatory and antimicrobial activities

Introduction

Nonsteroidal anti-inflammatory drugs (NSAIDs) have been recognized as a vital therapeutic agents for the alleviation of pain and inflammation associated with a numerous pathologic conditions viz. arthritis, bursitis, and tendinitis. However, chronic administration of NSAIDs has been associated with clinically significant complications such as gastrointestinal (GI) symptoms including mucosal damage, bleeding, nausea, heartburn, dyspepsia, abdominal pain, and renal toxicity, etc. Nowadays available polytherapy for inflammatory conditions associated with microbial infections increases the risk for developing NSAID-related complications especially in elderly, patients with prior history of peptic ulcer disease, patients with impaired liver or kidney functions, and patients taking anticoagulants, corticosteroids, etc. Hence, there is a pressing need for the drugs having both anti-inflammatory and antimicrobial activities, both from the pharmacoeconomic as well as the patient compliance point of view. (Allison et al., 1992; Flower, 2003; Bekhit and Abdel-Azeim, 2004).

The flavonoids are important bioactive molecules having a group of natural products present in a wide variety of plants reported to have potent anti-inflammatory activity in both (in vitro and in vivo) although not fully understood (Harborne and Williams, 2000; Middleton *et al.*, 2000). The several action mechanisms are proposed to explain in vivo anti-inflammatory action of flavonoids depending on their chemical structures. The unique action mechanisms and significant in vivo activity make flavonoids a reasonable candidate for new anti-inflammatory drugs. Flavonoids exhibit a broad range of biological activities, including antiviral, anti-inflammatory, antioxidant, antiallergic, hepatoprotective, antithrombotic and antitumoral actions (Manthey *et al.*, 2001; Rice-Evans and Miller, 1996; Hollman *et al.*, 1996).

However, the naturally found flavonoids like baicalein, quercetin, and myricetin have the -OH group in their structure, but in recent studies it is noteworthy that the O-alkylation of these phenolic -OH has considerable attention. Because of converting the trihydroxyl groups to the alkoxy groups on the "A" ring of baicalein resulting in increased activity on P-gp 170 inhibition (Lee et al., 2004). Fernández et al. analyzed inhibition of aldose reductase by 33 baicalein derivatives with modifications on the "A" and "B" rings, leading to the conclusion that compounds with either three hydroxyl or methoxyl functionalities on the "A" ring are effective inhibitors of aldose reductase (Fernandez et al., 2005). Furthermore, Morita et al. showed that the increased lipophilicity of phenolic hydroxyls via alkylation may enhance the therapeutic potential to X-linked adrenoleukodystrophy (X-ALD) (Morita et al., 2005).

Based on the aforementioned information, the combination of the biologically effective –O alkylation with a flavonol skeleton in continuation with our previous study (Hatnapure *et al.*, 2012), herein, we reporting a novel series of 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4one derivatives and have been screened for their in vitro proinflammatory cytokines TNF- α and IL-6 inhibitory activity.

The cytokines are intercellular messengers responsible for host defense mechanisms as inflammatory, immune and hematogenic responses. One of the key proinflammatory cytokine tumor necrosis factor- α (TNF- α) is mainly produced by the activated macrophages and monocytes, which further induces the production of the several inflammatory cytokines such as interleukin-1 β (IL-1 β), interleukin-6 (IL-6), and granulocyte-macrophage colony-stimulating factor (GM-CSF). It is also a multitude of biological activities linked to pathology of autoimmune diseases such as rheumatoid arthritis (RA) (Macnaul et al., 1990; Brennan et al., 1992), Crohn's disease (van Dullemen et al., 1995), systemic lupus erythematosus (Maury and Teppo, 1989), multiple sclerosis (Sharief and Hentges, 1991; Beck et al., 1988; Gallo et al., 1989), septic shock (Lechner et al., 1992), and AIDS (Makonkawkeyoon et al., 1993). On the other hand, cytokine IL-6 (from the series of cytokine-signaling pathway) contributes to the initiation and extension of the inflammatory process and considered as a central mediator in a range of inflammatory diseases but has not received the desired attention in drug discovery (Dominic and Raj, 2009). TNF- α and IL-6 are thus pharmaceutically important molecular targets for the treatment of the above-mentioned diseases. The available biopharmaceuticals [TNF-soluble receptor (Enbrel TM) and TNF antibody (Remicade TM)] are expensive, difficult to administer orally and have major side effects on prolonged clinical use. Therefore, there is an urgent medical need to discover small-molecule agents to deal with higher levels production of TNF- α .

On going through the literature, we found some flavonoids such as luteolin, galangin, etc., inflammatory inhibitors, when their structural activity relationships were compared (Fig. 1), the important moieties are the C-2, 3-double bond, A-ring 5, 7-hydroxyl groups, and B-ring 4'or 3',4'-hydroxyl groups. The C-3 hydroxyl group as in flavonol is favorable for lipoxygenase (LOX) inhibition and oral anti-inflammatory activity (Bauman *et al.*, 1980; Landolfi *et al.*, 1984).

In flavones and flavonol of the same type, flavonol showed greater inhibition than flavones. The tri methoxy flavonol derivatives have not hitherto tested as antiinflammatory agents, in particular against the TNF- α and IL-6 and also their antimicrobial activities. In order to improve the anti-inflammatory and antimicrobial activity of the compound based on quercetin and myricetin core (Cho *et al.*, 2003). We envisage the new scaffold by incorporating O-alkylation on ring "B" and different substituents on ring "A" hitherto remained untested (Fig. 2). We set out to test our notion by design, synthesis, and biological evaluation of novel flavonol derivatives.

Nowadays, our research group also engaged in antiinflammatory activity study program of novel urea and thiourea derivatives (Keche *et al.*, 2012a, b; Tale *et al.*, 2011). Herein, we presenting results on the synthesis and discovery of potent TNF- α and IL-6 inhibitor-based 2',4',5'-trimethoxyflavonol scaffold. The libraries of 21 different compounds (**2a–u**) are obtained according to Scheme 1 with good yield.

Our synthetic strategy for 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one derivatives have been planned by employing the general synthetic path of reaction. The 21 different 3-hydroxyflavones have been prepared by CLAISEN–SCHMIDT condensation followed by ALGAR–FLYNN–OYAMADA reaction (Scheme 1) (Li *et al.*, 2008). The key intermediates 1-(2-hydroxyphenyl)-



Fig. 1 Structure of luteolin and galangin

Fig. 2 New design scaffold



Reagents and conditions:- (a) EtOH,60% KOH, 0-rt 6-8h: (b) 15% aq NaOH/ 16% H2O2 (v/ v 1:1) 5-6h.

Scheme 1 Synthesis of 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one derivatives

3-(2,4,5-trimethoxyphenyl) prop-2-en-1-one (1) derivatives have been synthesized by the CLAISEN–SCHMIDT condensation between substituted 2-hydroxyacetophenones and 2,4,5-trimethoxybenzaldehyde in moderate to good yields (60–70 %). The treatment of 1 with hydrogen peroxide and base in ethanol under ALGAR–FLYNN–OYA-MADA conditions afforded the corresponding flavonol (**2a–u**) in (50–70 % yields). The desired products were purified by column chromatography (silica gel with 20 % ethyl acetate/hexane). The purity of the compounds was checked by TLC and HPLC. Spectral data ¹H, C¹³ NMR, and MS of the newly synthesized compounds have been in full agreement with their proposed structures.

Having secured the structurally diverse novel 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one derivatives, next in order to search for their proinflammatory cytokines, TNF- α , and IL-6 production by lipopolysaccharide (LPS) in THP-1 cells was measured according to the method described by Hwang *et al.* (1993). The results of the TNF- α and IL-6 inhibitory activity are shown in Table 1. As many as **3** compounds from this library have been found to be potent TNF- α and IL-6 inhibitor with reference standard drug dexamethasone. While the compounds **2d**, **2m**, **2n**, and **2s** found to be potent antimicrobial agent showing even 2–2.5-fold more potency than that of standard ciprofloxacin and miconazole at the same MIC value of 10 μ g/mL by using agar-well diffusion method (Sridhar *et al.*, 2004).

Thus the compounds **2h**, **2l**, and **2q** from the series found to be promising TNF- α and IL-6 inhibitors (up to 72–81 % TNF- α and 82–92 % IL-6 inhibitory activity but at 10 μ M concentration), when compared with standard dexamethasone (71 % TNF- α and 84 % IL-6 inhibitory activities at 1 μ M concentration). Notably, the compound **2h** exhibited much higher TNF- α or IL-6 (81 and 92 %) inhibitory activity. The compounds **2c**, **2e**, **2i**, **2j**, and **2k** exhibited moderate activity even as remaining compounds found to have low, very low, or no activity at the same level of concentration against the proinflammatory cytokines TNF- α and IL-6. As a result, the most potent

 Table 1
 Anti-inflammatory activity of 3-hydroxy-2-(2,4,5-trime-thoxyphenyl)-4H-chromen-4-one derivatives



| Compound 2(a-u) | R | % inhibition at 10 µM | |
|-----------------|---------------------------------------|-----------------------|------|
| | | TNF-α | IL-6 |
| 2a | 5-Cl | 45 | 39 |
| 2b | 5-Cl, 7-CH ₃ | 52 | 46 |
| 2c | 8-Cl | 60 | 65 |
| 2d | 6-Cl, 8-Cl | 30 | 28 |
| 2e | 6-CN, 8-Cl | 65 | 78 |
| 2f | 5-Br | 17 | 22 |
| 2g | 6-Br, 8-NO ₂ | - | - |
| 2h | 5-F | 81 | 92 |
| 2i | 7-F | 67 | 74 |
| 2j | 8-F | 60 | 75 |
| 2k | 6-F, 8-F | 64 | 76 |
| 21 | 5-F, 7-F | 72 | 86 |
| 2m | 6-NO ₂ , 8-F | 28 | 41 |
| 2n | 6-CN, 8-F | 27 | 49 |
| 20 | 7-CH ₃ | 15 | 21 |
| 2p | 6-CH ₃ , 8-NO ₂ | _ | - |
| 2q | 6-CF ₃ | 75 | 82 |
| 2r | 6-NO ₂ | - | - |
| 2s | 6-SO ₂ Me | 16 | 30 |
| 2t | 8-CN | 08 | 11 |
| 2u | 5-CN | _ | _ |
| Ref-1 | Dexamethasone (1 μ M) | 71 | 84 |

compound **2h** could prove to be a promising candidate for drug discovery.

In order to search for the potent antimicrobial agent from these newly synthesized compounds (2a-u) have been evaluated for in vitro antibacterial and antifungal activity against various Gram-positive, Gram-negative bacteria, and fungal strains (Tables 2, 3). As can be seen from obtained results, many compounds from the newly synthesized series found to be potent antibacterial and antifungal agents. Thus the compounds 2d, 2m, 2n, and 2s exhibited comparable to or even higher antimicrobial activity than the standard ciprofloxacin and miconazole, respectively, almost against all the tested bacteria or fungi. The remaining compounds of this series have been found to be moderate, low, or no activity. Moreover, the compounds 2d and 2s have more potent antibacterial agents than the standard drug against some bacteria viz. Staphylococcus aureus, Bacillus subtilis, and Salmonella typhimurium. Also the compound 2d found to be better antifungal agents as compared with standard miconazole.

The SAR of 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4Hchromen-4-one derivatives are shown in Tables 1, 2, and 3, some interesting trend has been observed as the effect of substituent present on ring "A" along with the O-methylation of "B" ring on taken activities that is the nature of the substituent affecting the biological activity of the synthesized analogues. From the TNF- α and IL-6 inhibitory activity data (Table 1), it is observed that a majority of the analogues of this series found to be active as IL-6 inhibitor while very few exhibited TNF- α inhibitory activity. So it is clear from the results that the position of the substituent on "A" ring of flavonol moiety has profound effect on the activity.

The 5-, 6-, and 7-positions are favorable sites for the higher potency. Evidently, the compounds 2h and 2l with F at 5 and 7 positions or both as well as CF₃ at 6-positions exhibiting highest TNF- α and IL-6 inhibitory activity, while presence of Cl, Br, CH₃, CN, and NO₂ at same position (compounds 2a, 2g, 2p, 2r, 2o, and 2u) exhibits no TNF- α or IL-6 inhibitory activity. However, surprisingly the compounds 2c and 2e have shown moderate activity with chloro and cyano groups at 8- and 6- or 8-positions of "A" ring, whereas, the fluoro group at 7- and 6-, 8-positions is moderately potent with (64-67 % TNF- α and 76-77 % IL-6 inhibitory activity). The compound 2h shows better TNF- α (81 % and IL-6 92 %) activity at 10 µM concentration. None of the compounds exhibits no or very low anti-inflammatory activity. It clears from the SAR fluoro group at 5-, 6-, and 7-positions tolerate the procytokine activity.

The antimicrobial activity data are presented in Tables 2 and 3. As shown in our results, some analogues of this series have been found to be more potent than the standard drug

Table 2Antibacterial activity of 3-hydroxy-2-(2,4,5-trimethoxy-phenyl)-4H-chromen-4-one derivatives

| Compounds | Gram-positive | | Gram-negative | |
|---------------|--------------------------|----------------------|---------------------|---------------------------|
| | Staphylococcus aureus | Bacillus subtilis | Escherichia coli | Salmonella typhimurium |
| 2a | 35 | 30 | 25 | 40 |
| 2b | 30 | 45 | 35 | 50 |
| 2c | 85 | - | 80 | 90 |
| 2d | 10 | 10 | 15 | 10 |
| 2e | 90 | 80 | 75 | 80 |
| 2f | 80 | 65 | - | 75 |
| 2g | 40 | 35 | 45 | 35 |
| 2h | 90 | _ | 75 | 80 |
| 2i | 65 | 55 | 45 | 50 |
| 2j | _ | 80 | 60 | 90 |
| 2k | 75 | 85 | - | 80 |
| 21 | 80 | 90 | 85 | - |
| 2m | 20 | 15 | 15 | 20 |
| 2n | 25 | 15 | 10 | 25 |
| 20 | 90 | 85 | - | 80 |
| 2p | _ | _ | 85 | - |
| 2q | 85 | 70 | 90 | 90 |
| 2r | 85 | 90 | 90 | 90 |
| 2s | 15 | 10 | 20 | 10 |
| 2t | 40 | 30 | 25 | 30 |
| 2u | 35 | 45 | 60 | 40 |
| Ciprofloxacin | 25 | 20 | 15 | 20 |

MIC values are the average of three reading ($\mu g/mL)$

- No activity was observed up to 200 µg/mL

ciprofloxacin, while some of them have comparable potency. Interestingly none of the compounds with high anti-inflammatory activity found to be potent antibacterial or antifungal agents. The compounds 2d, 2m, 2n, and 2s bearing Cl, NO₂, CN, and SO₂Me groups, respectively, at 6-position with Cl, F, at 8-position or at both have higher potency than the compounds bearing such a group at 5- or 7-positions or at both. It is clear from our results that 6- or 8-position is the favorable site for high-antibacterial activity. The high potency of 2d, 2m, 2n, and 2s may be attributed to the presence of lipophilic or H-bond acceptor type group's placement such as F, Cl, CN, SO₂Me, and NO₂ at 6- or 8-positions, but unfortunately this functional group presence on 5, 7 or both shows moderate or no activity with respect to standard drug against the tested strains. No activity has been observed in case of compounds 20-2r up to concentration of 200 µg/mL against some bacteria and fungi.

It is clear from activity data (Table 2 vs. Table 3) that the SAR of 3-hydroxy-2-(2,4,5-dimethoxyphenyl)-4H-

chromen-4-one derivatives of antibacterial activity strongly correlates with their SAR of antifungal activity. Again the positions 6 and 8 of ring "A" are favorable sites for high activity. The compounds 2d and 2s were found to be 2–2.5fold more potent than the standard drug miconazole, while 2m and 2n exhibited comparable antifungal activity similar to the antibacterial activity trend, polar lipophilic groups such as Cl, SO₂Me, NO₂, F at 6- and 8-position or both at same level of concentration 10 μ g/mL. Apart from the compounds 2a, 2b, 2g, 2t, and 2u have no major effect on the antifungal activity.

It reveals from our SAR studied, the presence of lipophilic F and CF_3 groups with trimethoxy (O-alkylation) tolerates the procytokine activity because of fluorine imparts the special characteristics that enhance therapeutic efficacy and improved pharmacological properties in bioactive molecules. While the fluoro is only slightly more lipophilic than hydrogen and trifluoromethyl is much more lipophilic than methyl or chloro, this factor is often the most significant in improving the pharmacological activity.

In conclusion, we have developed the novel approach for the synthesis of structurally diverse 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one derivatives (**2a–u**) using simple reaction sequence under mild conditions and their TNF- α and IL-6 inhibitory and antimicrobial activity have been evaluated. This study on the novel flavonol derivative bearing lipophilic fluoro group at 5-, 6-, and 7-positions and presence of trimethoxy group on "B" ring has lead to the discovery of novel and potent TNF- α and IL-6 inhibitors viz. **2h**, **2l**, and **2q** and could prove to be the valuable platform for the search of nonsteroidal anti inflammatory agents based on such a scaffold in vivo. While the presence of chloro, cyano or sulfonyl groups on 6- and 8-positions or both on ring "A" found to be an effective potent antimicrobial agents viz. compounds **2d**, **2m**, **2n**, and **2s**.

Experimental

Chemistry

General techniques

All commercial chemicals and solvents are of reagents grade and have been used without further purification. The thin layer chromatography was performed on Merck precoated silica gel 60 F254 plates, with visualization under UV light. ¹H NMR spectra have been recorded with Bruker 300 MHz AVANCE instrument and *J* values are in Hertz and chemical shifts (δ) are reported in ppm relative to internal tetramethylsilane. The mass spectra have been measured with Thermo Finnigan-TSQ Quarter Ultra (triple Quad). The purity of all compounds was determined by

 Table 3
 Antifungal
 activity
 of
 3-hydroxy-2-(2,4,5-trimethoxy-phenyl)-4H-chromen-4-one
 derivatives

| Compounds | Candida albicans | Aspergillus niger | Fusarium solani | Aspergillus flavus |
|------------|---------------------|----------------------|--------------------|-----------------------|
| 2a | 25 | 40 | 35 | 45 |
| 2b | 30 | 25 | 20 | 35 |
| 2c | - | 80 | 90 | 80 |
| 2d | 10 | 10 | 15 | 10 |
| 2e | 75 | 85 | - | 90 |
| 2f | - | - | 90 | - |
| 2g | 45 | 30 | 25 | 40 |
| 2h | - | 90 | 85 | 70 |
| 2i | 70 | 65 | 85 | 90 |
| 2j | 90 | - | 80 | 95 |
| 2k | - | 80 | 75 | 80 |
| 21 | 65 | 80 | - | - |
| 2m | 20 | 15 | 20 | 10 |
| 2n | 25 | 20 | 15 | 25 |
| 20 | - | 80 | 90 | 85 |
| 2p | - | 90 | - | - |
| 2q | 60 | 70 | 75 | 85 |
| 2r | 90 | - | 80 | - |
| 2s | 15 | 10 | 10 | 20 |
| 2t | 40 | 30 | 25 | 30 |
| 2u | 25 | 30 | 15 | 25 |
| Miconazole | 20 | 20 | 15 | 20 |

MIC values are the average of three reading (μ g/mL)

- No activity was observed up to 200 µg/mL

HPLC (Waters 2695 Alliance) using column Kromasil C18, solvent acetonitrile and buffer (0.01 M ammonium acetate + 0.5 % triethylamine, pH 5.0 with acetic acid). Melting points were determined with (PEW-340 MP) melting point apparatus and are uncorrected.

General procedure for the synthesis of 1-(2hydroxyphenyl)-3-(2,4,5-trimethoxyphenyl) prop-2-en-1one derivatives (**1***a*–*u*)

Different substitutes (2-hydroxyacetophenone) (1 eqi.) and 2,4,5-trimethoxy benzaldehyde (1 eqi.) in EtOH (15 mL) were added and a 60 % of KOH (5 mL) solution was added drop wise at 0 °C. The reaction mixture was stirred at room temperature for 6–8 h. The reaction mixture was cooled and poured into ice water, neutralized using 1 N HCl. The light-yellow solid thus obtained was filtered and purified using silica gel column chromatography (with 1:9 ethyl acetate in hexane) or recrystallized from EtOH, yielded the desired 2,4,5-trimethoxy chalcone analogues (1a-u) with an average yield of 60–70 %.

General procedure for the synthesis of 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one derivatives (2a–u)

To a well-stirred solution of 2-hydroxychalcones **1** (1 eqi.) in MeOH (20 mL) and aq. KOH (10 mL, 20 %), cooled at 0–5 °C, 30 % H_2O_2 (10 mL) was added drop wise for 30 min. The reaction mixture was further stirred for 4–5 h. The resulting light-yellow reaction mixture was poured on crushed ice and neutralized with dil. HCl. The light-yellow solid thus obtained was filtered, washed with water and dried. The crude product was purified by chromatography on SiO₂ (100–200 silica gel) (EtOAc/petroleum ether v/v = 2:8) to give the desired compounds as solids. The physical, analytical, and spectral data of new 3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4-H-chromen-4-one analogues are given below.

5-*Chloro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (2a)* White solid: yield 66 %; mp: 191–193 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 7.77 (d, J = 8.5 Hz, 1H), 7.51 (d, J = 9 Hz, 1H), 7.35 (m, 1H), 7.10 (s, 1H), 6.61 (s, 1H), 3.90 (s, 3H), 3.83 (s, 3H), 3.81 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 172.49(C=O), 165.71(C-2), 164.14 (C-9), 155.19(C-2'), 152.29(C-4'), 151.19(C-5'), 144.25(C-5), 143.31(-OH/C-3), 139.12(C-7), 127. 2 3(C-10), 118.16(C-6), 113.35(C-8), 111.14 (C-6'), 104.54(C-1'), 97.72 (C-3'), 57.16, 56.28, 56.23 (OMe-C); MS (APCI); *m/z* 363.1 [M+H]⁺; HPLC: 95.96 %.

5-*Chloro-3-hydroxy-7-methyl-2-(2,4,5-trimethoxyphenyl)*-4*H-chromen-4-one (2b)* Off white solid: yield 67 %; mp: 187–189 °C .¹H NMR (CDCl₃, 300 MHz): δ 9.25 (bs, 1H), 8.00 (s, 1H), 7.63 (s, 1H), 7.05 (s, 1H), 6.79 (s, 1H), 3.84 (s, 3H), 3.77 (s, 3H), 3.69 (s, 3H), 2.96 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 173.21(C=O), 157.10(C-2), 152.91 (C-9), 150.31(C-2'), 148.63(C-4'), 147.37(C-5'),143.10(C-7), 141.21(C-5), 140.31(-OH/C-3), 125.41(C-10), 120.51(C-6), 118.12(C-8), 112.33(C-6'), 110.10(C-1'), 98.10(C-3'), 57.33, 56.46, 56.23 (OMe-C), 22.23 (Me/C); MS (APCI); *m/z* 377.1 [M+H]⁺; HPLC: 96.89 %.

8-*Chloro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (2c)* White solid: yield 64 %; mp: 196–198 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 7.77 (d, J = 8.5 Hz, 1H), 7.51 (d, J = 9 Hz, 1H), 7.35 (m, 1H), 7.10 (s, 1H), 6.61 (s, 1H), 3.90 (s, 3H), 3.83 (s, 3H), 3.81 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 172.59(C=O), 165.71(C-2), 164.14 (C-9), 155.19(C-2'), 152.29(C-4'), 151.19(C-5'), 143.25(C-8), 142.31(-OH/C-3), 138.12(C-7), 127.33(C-10), 117.16(C-6), 115.35(C-5), 112.14 (C-6'), 105.54(C-1'), 98.72 (C-3'), 56.30, 56.28, 56.23 (OMe-C); MS (APCI); m/z 363.1 [M+H]⁺; HPLC: 93.96 %.

6,8-Dichloro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (2d) Pale yellow solid: yield 68 %; mp: 199– 201 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.19 (bs, 1H), 8.13 (s, 1H), 7.73 (s, 1H), 7.23 (s, 1H), 6.65 (s, 1H), 3.97 (s, 3H), 3.92 (s, 3H), 3.89 (s, 3H); (CDCl₃, 300 MHz, δ ppm): 173.03(C=O), 158.13(C-2), 155.47(C-9), 150.86(C-2'), 145.37(C-4'), 144.12 (C-5'), 142.21(-OH/C-3), 136.42(C-8), 133.05(C-6), 130.20(C-10), 123.20(C-7), 121.14(C-5), 117.02 (C6'), 112.32 (C-1'), 97.11(C-3'), 57.22, 56.15, 56.10 (OMe-C); MS (APCI); *m*/*z* 398.1 [M+H]⁺; HPLC: 96.06 %.

8-*Chloro-3-hydroxy-4-oxo-2-(2,4,5-trimethoxyphenyl)-4Hchromene-6-carbonitrile (2e)* Off white solid: yield 57 %; mp: 203–205 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.27 (bs, 1H), 8.31 (s, 1H), 8.26 (s, 1H), 7.15 (s, 1H), 6.86 (s, 1H), 3.89 (s, 3H), 3.86 (s, 3H), 3.83 (s, 3H); (CDCl₃, 300 MHz, δ ppm): 172.13(C=O), 157.14(C-2), 154.57(C-9), 150.66 (C-2'), 146.42(C-4'), 143.32(C-5'), 142.31(-OH/C-3), 139.16 (C-8), 132.20(C-10), 124.15(C-5), 123.36(C-7), 119.13(CN), 117.12 (C-6'), 113.16(C-6), 111.33(C-1'), 97.11(C-1'), 57.23, 56.25, 56.14 (OMe-C); MS (APCI); *m/z* 388.1 [M+1]⁺; HPLC: 94.39 %.

5-Bromo-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (**2f**) Off white solid: yield 63 %; mp: 193– 195 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 8.14 (s, 1H), 7.85 (d, J = 8.5 Hz, 1H), 7.57 (d, J = 9 Hz 1H), 7.04 (s, 1H), 6.78 (s, 1H), 3.82 (s, 3H), 3.74 (s, 3H), 3.66 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 172.49(C=O), 163.72(C-2), 164.14(C-9), 155.19(C-2'), 152.29(C-4'), 151.19(C-5'), 143.25(C-5), 142.31(-OH/C-3), 138.12(C-7), 127. 2 3(C-10), 120.16(C-6), 115.35(C-8), 110.14 (C-6'), 105.54(C-1'), 99.72 (C-3'), 57.17, 56.30, 56.29 (OMe-C); MS (APCI); m/z 409.0 [M+2H]⁺; HPLC: 96.65 %.

6-*Bromo-3-hydroxy-8-nitro-2-(2,4,5-trimethoxyphenyl)-*4*H-chromen-4-one* (**2***g*) Yellow solid: yield 70 %; mp: 197 –199 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.27 (BS, 1H), 8.31 (s, 1H), 8.26 (s, 1H), 7.15 (s, 1H), 6.84 (s, 1H), 3.89 (s, 3H), 3.83 (s, 3H), 3.78 (s, 3H); (CDCl₃, 300 MHz, δ ppm): 174.03(C=O), 157.13(C-2), 155.47(C-9), 150.86(C-2'), 146.37(C-4'), 144.12(C-5'), 14 2.21(–OH/C-3), 141.42(C-8), 133.05(C-6), 129.20(C-10), 123.20(C-7), 121.14(C-5), 117.02 (C-6'), 1 12.32 (C-1'), 97.11(C-3'), 57.24, 56.21, 56.18 (OMe-C); MS (APCI); *m/z* 453.1 [M+H]⁺; HPLC: 95.03 %.

5-*Fluoro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one* (*2h*) Brown solid: yield 71 %; mp: 186–188 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 7.70 (d, *J* = 9 Hz 3H), 7.07 (s, 1H), 6.80 (s, 1H), 3.84 (s, 3H), 3.76 (s, 3H), 3.68 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm):172.60(C=O), 163.71(C-2), 164.14 (C-9), 155.19(C-2'), 152.29(C-4'), 151.19(C-5'), 143.25(C-5), 141.31(-OH/C-3), 139.12(C-7), 127. 2 3(C-10), 118.16(C-6), 113.35(C-8), 111.14 (C-6'), 102.54(C-1'), 95.72(C-3'), 57.28, 56.27, 56. 23 (OMe-C); MS (APCI); *m/z* 347.1 [M+2H]⁺; HPLC: 97.39 %.

7-*Fluoro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one* (2*i*) Off white solid: yield 69 %; mp: 190–192 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 8.26 (d, *J* = 8.6 Hz 1H), 7.21 (d, *J* = 9 Hz, 1H), 7.17 (m, 1H), 7.12 (s, 1H), 6.65 (s, 1H), 3.96 (s, 3H), 3.90 (s, 3H), 3.90 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 173.60(C=O), 162.71(C-2), 160.14(C-9), 155.19(C-2'), 152.29(C-4'), 151.19(C-5'), 144.12(C-7), 141.31(–OH/C-3), 137.25(C-5), 129.23(C-10), 118.16(C-6), 113.35(C-8), 111.14 (C-6'), 102.54(C-1'), 95.72 (C-3'), 57.30, 56.28, 56. 23 (OMe-C); MS (APCI); *m/z* 347.1 [M+2H]⁺; HPLC: 95.33 %.

8-*Fluoro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (2j)* Off white solid: yield 65 %; mp: 189– 191 °. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 8.04 (d, *J* = 8.6 Hz, 1H), 7.44 (d, *J* = 9 Hz, 1H), 7.33 (m, 1H), 7.15 (s, 1H), 6.66 (s, 1H), 3.96(s, 3H), 3.90 (s, 3H), 3.89 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 175.60(C=O), 163.71(C-2), 160.14(C-9), 155.19(C-2'), 152.29(C-4'), 151.19(C-5'), 146.12(C-8), 141.31(-OH/C-3), 137.25(C-10), 129.23(C-5), 118.16(C-6), 115.35(C-7), 113.14 (C-6'), 102.54(C-1'), 95.72 (C-3'), 57.27, 56.21, 56. 20 (OMe-C); MS (APCI); *m/z* 347.1 [M+H]⁺; HPLC: 94.39 %.

6,8-Difluoro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4Hchromen-4-one (**2k**) Pale yellow solid: yield 60 %; mp: 206–208 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.19 (bs, 1H), 7.90 (dd, 1H), 7.65 (dd, 1H), 7.10 (s, 1H), 6.85 (s, 1H), 3.88 (s, 3H), 3.80 (s, 3H), 3.72 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 173.03(C=O), 158.13(C-2), 155.47(C-9), 150.86(C-2'), 145.37(C-4'), 144.12(C-5'), 142.21(-OH/ C-3), 141.42(C-8), 140.05(C-6), 132.20(C-10), 125.20(C-7), 122.14(C-5), 119.02(C-6'), 113.32(C-1'), 97.15(C-3'), 57.23, 56.15, 56.12 (OMe-C); MS (APCI); *m/z* 365.1 [M+H]⁺; HPLC: 98.57 %.

5,7-Difluoro-3-hydroxy-2-(2,4,5-trimethoxyphenyl)-4Hchromen-4-one (**2***l*) Pale yellow solid: yield 66 %; mp: 200–202 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.19 (bs, 1H), 7.09 (s, 1H), 6.76 (dd, 1H), 6.64 (s, 1H), 6.52 (dd, 1H), 3.96 (s, 3H), 3.88 (s, 3H), 3.86 (s, 3H); (CDCl₃, 300 MHz, δ ppm): ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 174.03(C=O), 159.14(C-2), 157.47(C-9), 152.86(C-2'), 146.37(C-4'), 145.12(C-5'), 143.21 (–OH/C-3), 142.52(C-7), 141.25(C-5), 130.20(C-10), 127.20 (C-6), 124.14(C-8), 118.02(C-6'), 114.32(C-1'), 98.15(C-3'), 57.21, 56.17, 56. 14 (OMe-C); MS (APCI); *m*/*z* 365.1 [M+1]⁺; HPLC: 96.34 %. 8-*Fluoro-3-hydroxy-6-nitro-2-(2,4,5-trimethoxyphenyl)-*4*H-chromen-4-one* (**2m**) Off white solid: yield 57 %; mp: 211–213 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.27 (bs, 1H), 8.31 (s, 1H), 8.27(s, 1H), 7.17 (s, 1H), 6.89 (s, 1H), 3.91 (s, 3H), 3.89 (s, 3H), 3.86 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 176.03(C=O), 159.13(C-2), 157.47(C-9), 152.86(C-2'), 146.37(C-4'), 145.12(C-5'), 143.21(-OH/ C-3), 142.42(C-6), 137.05(C-8), 130.20(C-10), 124.20(C-7), 122.14(C-5), 118.02 (C6'), 113.32 (C-1'), 97.25(C-3'), 57.27, 56.25, 56.21 (OMe-C); MS (APCI); *m/z* 392.1 [M+H]⁺; HPLC: 95.54 %.

8-*Fluoro-3-hydroxy-4-oxo-2-(2,4,5-trimethoxyphenyl)-4Hchromene-6-carbonitrile* (**2n**) Brown solid: yield 64 %; mp: 209–211 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.27 (bs, 1H), 8.31 (s, 1H), 8.26 (s, 1H), 7.15 (s, 1H), 6.84 (s, 1H), 3.89 (s, 3H), 3.83 (s, 3H), 3.78 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 172.13(C=O), 157.14(C-2), 154.57(C-9), 151.66(C-2'), 146.27 (C-4'), 144.32(C-5'), 143.16(C-8), 142.31(-OH/C-3), 132.20(C-10), 124.15(C-5), 123.36(C-7), 120.13(CN), 117.12(C-6'), 115.16(C-6), 112.33(C-1'), 97.20 (C-1'), 57.23, 56.24, 56.17 (OMe-C); MS (APCI); *m/z* 372.1 [M+H]⁺; HPLC: 93.67 %.

3-Hydroxy-7-methyl-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (**2o**) Brown solid: yield 68 %; mp: 189–191 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.25 (bs, 1H), 8.13 (s, J = 8 Hz, 1H), 7.31 (s, 1H),7.23 (d, J = 8.5 Hz, 1H), 7.11 (s, 1H), 6.65 (s, 1H), 3.96 (s, 3H), 3.92 (s, 3H), 3.88 (s, 3H), 2.49 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 174.21(C=O), 158.10(C-2), 155.91(C-9), 151.31(C-2'), 148.55(C-4'), 147.37(C-5'), 141.10(C-7), 140.31(-OH/C-3), 130.21(C-5), 125.41(C-10), 120.51(C-6), 118.12(C-8), 113.33(C-6'), 111.10(C-1'), 98.30(C-3'), 57.35, 56.53, 56.42 (OMe-C), 21.23 (Me/C); MS (APCI); *m*/z 343.1 [M+H]⁺; HPLC: 98.64 %.

3-Hydroxy-6-methyl-8-nitro-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (**2p**) Yellow solid: yield 70 %; mp: 181–183 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.25 (bs, 1H), 8.34 (s, 1H), 8.27 (s, 1H), 7.14 (s, 1H), 6.85 (s, 1H), 3.89 (s, 3H), 3.81 (s, 3H), 3.72 (s, 3H), 2.51 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 173.03(C=O), 157.13(C-2), 155.47(C-9), 151.86(C-2'), 145.37(C-4'), 143.12(C-5'), 142.21(-OH/C-3), 141.42(C-8), 139.05(C-6), 129.20(C10), 123.20(C-7), 121.14(C-5), 117.02 (C-6'), 114.32(C-1'), 97.14(C-3'), 57.21, 56.23, 56.18 (OMe-C), 21.27 (Me/C); MS (APCI); *m/z* 388.1 [M+H]⁺; HPLC: 95.04 %.

3-Hydroxy-6-(trifluoromethyl)-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (**2q**) Off white solid: yield 66 %; mp: 216–218 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 8.59 (s, 1H), 7.85 (dd, 1H), 7.63 (d, *J* = 9 Hz, 1H), 7.12 (s, 1H), 6.66 (s, 1H), 3.83 (s, 3H), 3.74 (s, 3H), 3.71 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 175.01(C=O), 159.13(C-2), 157.47(C-9), 150.86(C-2'), 147.37(C-4'), 145.12(C-5'), 141.21(-OH/C-3), 131.42(C-7), 129.05(C-6), 127.17 (C/CF₃), 130.20(C-10), 123.20(C-8), 121.14(C-5), 118.02(C-6'), 112.32(C-1'), 98.11(C-3'), 57.24, 56.16, 56.12 (OMe-C); MS (APCI); *m*/*z* 397.1 [M+H]⁺; HPLC: 95.03 %.

3-Hydroxy-6-nitro-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (**2r**) Yellow solid: yield 66.5 %; mp: 196–198 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.35 (bs, 1H), 8.82 (s, 1H), 8.50 (d, J = 8.5 Hz, 1H), 7.87 (d, J = 9 Hz, 1H), 7.11 (s, 1H), 6.83 (s, 1H), 3.89 (s, 3H), 3.81 (s, 3H), 3.72 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 173.23(C=O), 159.51(C-2), 158.47(C-9), 151.85(C-2'), 146.37(C-4'), 145.30 (C-5'), 142.10(C-6), 141.21(-OH/C-3), 131.42(C-7), 130.20 (C-10), 123.20(C-8), 121.14(C-5), 119.02(C-6'), 112.32(C-1'), 98.20(C-3'), 57.24, 56.18, 56.15 (OMe-C); MS (APCI); m/ z 374.1 [M+H]⁺; HPLC: 97.58 %.

3-Hydroxy-6-(methylsulfonyl)-2-(2,4,5-trimethoxyphenyl)-4H-chromen-4-one (2s) Off white solid: yield 66 %; mp: 220–222 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.24 (bs, 1H), 8.58 (s, 1H), 8.20 (d, J = 9 Hz, 1H), 7.86 (d, J = 8.6 Hz, 1H), 7.10 (s, 1H), 6.83 (s, 1H), 3.86 (s, 3H), 3.82 (s, 3H), 3.78 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 172.45 (C=O), 158.51(C-2), 157.47(C-9), 151.81(C-2'), 146.37(C-4'), 145.30(C-5'), 141.21(-OH/C3), 137.42.(C-6), 131.42 (C-7), 131.20(C-10), 123.20(C-8), 120.14(C-5), 119.02(C-6'), 112.32(C-1'), 96.20(C-3'), 57.25, 56.19, 56.16 (OMe-C), 39.89 (Me/SO₂); MS (APCI); *m*/*z* 407.1 [M+H]⁺; HPLC: 94.95 %.

3-Hydroxy-4-oxo-2-(2,4,5-trimethoxyphenyl)-4H-chromene-8-carbonitrile (2t) Off white solid: yield 58 %; mp: 199–201 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 8.22 (d, *J* = 8.5 Hz, 1H), 8.06 (d, *J* = 9 Hz, 1H), 7.41 (m, 1H), 7.21 (s, 1H), 6.75 (s, 1H), 3.81 (s, 3H), 3.77 (s, 3H), 3.67 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 172.60(C=O), 163.71(C-2), 161.14(C-9), 156.19(C-2'), 153.29(C-4'), 151.19 (C-5'), 142.31(–OH/C-3), 130.33(C-10), 129.12(C-7), 125.25 (C-8), 117.16(C-6), 115.35(C-5), 112.14 (C-6'), 107.54(C-1'), 98.72 (C-3'), 56.27, 56.25, 56.24 (OMe-C); MS (APCI); *m*/ *z* 372.1 [M+H]⁺; HPLC: 95.36 %.

3-Hydroxy-4-oxo-2-(2,4,5-trimethoxyphenyl)-4H-chromene-5-carbonitrile (**2u**) Off white solid: yield 60 %; mp: 207–209 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.05 (bs, 1H), 8.67 (s, 1H), 8.18 (d, J = 8.5 Hz, 1H), 7.59 (d, J = 9 Hz 1H), 7.59 (s, 1H), 6.78 (s, 1H), 3.83 (s, 3H), 3.76 (s, 3H), 3.68 (s, 3H); ¹³C NMR (CDCl₃, 300 MHz, δ ppm): 173.16(C=O), 162.20(C-2), 160.10(C-9), 156.21(C-2'), 154.29(C-4'), 151.19(C-5'), 142.31(-OH/C-3), 129.79(C-10), 129.12(C-7), 126.25(C-5), 118.16(C-6), 115.35(C-8), 113.14 (C-6'), 107.54(C-1'), 97.70 (C-3'), 56.22, 56.21, 56.19 (OMe-C); MS (APCI); *m*/*z* 372.1 [M+H]⁺; HPLC: 96.76 %.

Biological assay

Anti-inflammatory assay

Proinflammatory cytokine production by LPS in THP-1 cells has been measured according to the method described by Hwang in 1993. During assay, THP-1 cells were cultured in RPMI 1640 culture medium (Gibco BRL, Pasley, UK) containing 100 U/mL penicillin and 100 mg/mL streptomycin containing 10 % fetal bovine serum (FBS, JRH). Cells have been differentiated with phorbol myristate acetate (PMA, Sigma). Following cell plating, the test compounds in 0.5 % DMSO have been added to each well separately and the plate has been incubated for 30 min at 37 °C. Finally, LPS (E. coli 0127:B8, Sigma Chemical Co., St. Louis, MO, USA) has been added at a final concentration of 1 µg/mL in each well. Plates have been further incubated at 37 °C for 24 h in 5 % CO₂. After incubation, supernatants have been harvested, and assayed for TNF- α and IL-6 by ELISA as described by the manufacturer (BD Biosciences).

Antibacterial assay

All the newly synthesized compounds have been screened for their antibacterial activity against selected Gram-positive organism's viz. Bacillus subtilis (MTCC 441) and Staphylococcus aureus (MTCC 96), and Gram-negative organism's viz. Salmonella typhimurium (MTCC 98) and Escherichia coli (MTCC 443) bacterial strains by agar-well diffusion method with little modification. Different concentrations (10-200 µg/mL) of test compounds have been prepared in DMSO. The bacterial suspension has been spread over nutrient agar plates and the well with of 6 mm diameter has been punched with sterile cork borer. The sample (50 μ L) has been added to the well and the plates have been incubated at 36 °C for 24 h. DMSO used as solvent control and ciprofloxacin has been used as standard antibacterial agent. The lowest concentration of compound which completely inhibits the bacterial growth has been taken as minimum inhibitory concentration (MIC).

Antifungal assay

All the newly synthesized compounds have been screened for their antifungal activity against *Candida albicans* (MTCC 227), *Aspergillus niger* (MTCC 281), *Aspergillus* *flavus* (MTCC 277), and *Fusarium solani* (MTCC 350) by agar-well diffusion method with little modification. Normal saline has been used to make a suspension of spores of fungal strain, prepared as same as antibacterial assay. Miconazole has been used as standard antifungal agent.

Acknowledgments We are grateful to the Department of Chemistry, Maharashtra Udayagiri Mahavidyalaya, Udgir for providing necessary facilities and would like to thank Mr. R. B. Ingle, Department of Microbiology, Shri Shivaji College, Akola for biological screening studies of the newly synthesized compounds.

References

- Allison MC, Howatson AG, Torrance CJ, Lee FD, Russell RI (1992) Gastrointestinal damage associated with the use of nonsteroidal anti-inflammatory drugs. N Engl J Med 327:749–754
- Bauman J, Bruchhausen FV, Wurm G (1980) Flavonoids and related compounds as inhibitors of arachidonic acid peroxidation. Prostaglandins 20:627–639
- Beck J, Rondot P, Catinot L, Falcoff E, Kirchner H, Wietzerbin J (1988) Increased production of interferon gamma and tumor necrosis factor precedes clinical manifestation in multiple sclerosis: do cytokines trigger off exacerbations? Acta Neurol Scand 78:318–323
- Bekhit AA, Abdel-Azeim T (2004) Design, synthesis and biological evaluation of some pyrazole derivatives as anti-inflammatory antimicrobial agents. Bioorg Med Chem 12:1935–1945
- Brennan FM, Gibbons DL, Mitchell T, Cope AP, Maini RN, Feldmann M (1992) Enhanced expression of tumor necrosis factor receptor mRNA and protein in mononuclear cells isolated from rheumatoid arthritis synovial joints. Eur J Immunol 22:1907–1912
- Cho SY, Park SJ, Kwon MJ, Jeong TS, Bok SH, Cho WY (2003) Quercetin suppresses proinflammatory cytokines production through MAP kinases and NF-kappa B pathway in lipopolysaccharide-stimulated macrophage. Mol Cell Biochem 243:153–160
- Dominic SC, Raj MD (2009) Role of interleukin-6 in the anemia of chronic disease. Semin in Arthritis Rheum 38:382
- Fernandez M, Caballero J, Helguera AM, Castro EA, Gonzalez MP (2005) Quantitative structure-activity relationship to predict differential inhibition of aldose reductase by flavonoid compounds. Bioorg Med Chem 13:3269–3277
- Flower RJ (2003) The development of COX-2 inhibitors. Nat Rev Drug Discov 2:179–191
- Gallo P, Piccinno MG, Krzalic L, Tavolato B (1989) Tumor necrosis factor alpha (TNF alpha) and neurological diseases. Failure in detecting TNF- α in the cerebrospinal fluid from patients with multiple sclerosis, AIDS dementia complex, and brain tumours. J Neuroimmunol 23:41–44
- Harborne JB, Williams CA (2000) Advances in flavonoid research since 1992. Phytochemistry 55:481–504
- Hatnapure GD, Keche AP, Rodge AH, Birajdar SS, Tale RH, Kamble VM (2012) Synthesis and biological evaluation of novel piperazine derivatives of flavone as potent anti-inflammatory and antimicrobial agent. Bioorg Med Chem Lett 22:6385–6390
- Hollman PCH, Hertog MGL, Katan MB (1996) Role of dietary flavonoids in protection against cancer and coronary heart disease. Biochem Soc Trans 24:785–789
- Hwang C, Gatanaga M, Granger GA, Gatanaga T (1993) Mechanism of release of soluble forms of tumor necrosis factor/lymphotoxin receptors by phorbol myristate acetate-stimulated human THP-1 cells in vitro. J Immunol 151:5631–5638

- Keche AP, Hatnapure GD, Tale RH, Rodge AH, Birajdar SS, Kamble VM (2012a) Novel pyrimidine derivatives with aryl urea, thiourea and sulfonamide moieties: synthesis, anti-inflammatory and antimicrobial evaluation. Bioorg Med Chem Lett 22:3445–3448
- Keche AP, Hatnapure GD, Tale RH, Rodge AH, Kamble VM (2012b) Synthesis, anti-inflammatory and antimicrobial evaluation of novel 1-acetyl-3,5-diaryl-4,5-dihydro (1H) pyrazole derivatives bearing urea, thiourea and sulfonamide moieties. Bioorg Med Chem Lett 22:6611–6615
- Landolfi R, Mower RL, Steiner M (1984) Modification of platelet function and arachidonic acid metabolism by bioflavonoid. Structure-activity relations. Biochem Pharmacol 33:1525–1530
- Lechner AJ, Tredway TL, Brink DS, Klein CA, Matuschak GM (1992) Differential systemic and intrapulmonary TNF-alpha production in *Candida sepsis* during immunosuppression. Am J Physiol 263:526–535
- Lee Y, Yeo H, Liu SH, Jiang Z, Savizky RM, Austin DJ, Cheng YC (2004) Increased anti-P-glycoprotein activity of baicalein by alkylation on the A ring. J Med Chem 47:5555–5566
- Li Z, Ngojeh G, DeWitt P, Zheng Z, Min Chen, Lainhart B, Li V, Felpo P (2008) Synthesis of a library of glycosylated flavonols. Tetrahedron Lett 49:7243–7245
- Macnaul KL, Hutchinson NI, Parsons JN, Bayne EK, Tocci MJ (1990) Analysis of IL-1 and TNF-alpha gene expression in human rheumatoid synoviocytes and normal monocytes by in situ hybridization. J Immunol 145:4154–4166
- Makonkawkeyoon S, Limson-Pombre RNR, Moreira AL, Schauf V, Kaplan G (1993) Thalidomide inhibits the replication of human immunodeficiency virus type 1. Proc Natl Acad Sci USA 90:5974–5978
- Manthey JA, Grohmann K, Guthrie N (2001) Biological properties of citrus flavonoids pertaining to cancer and inflammation. Curr Med Chem 8:135–153

- Maury CP, Teppo AM (1989) Tumor necrosis factor in the serum of patients with systemic lupus erythematosus. Arthritis Rheum 32:146–150
- Middleton EJ, Kandaswami C, Theoharides TC (2000) The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. Pharmacol Rev 52:673–751
- Morita M, Takahashi I, Kanai M, Okafuji F, Iwashima M, Hayashi T, Watanabe S, Hamazaki T, Shimozawa N, Suzuki Y, Furuya H, Yamada T, Imanaka T (2005) Baicalein 5,6,7-trimethyl ether, a flavonoid derivative, stimulates fatty acid beta-oxidation in skin fibroblasts of X-linked adrenoleukodystrophy. FEBS Lett 579:409–414
- Rice-Evans CA, Miller NJ (1996) Antioxidant activities of flavonoids as bioactive components of food. Biochem Soc Trans 24:790–795
- Sharief MK, Hentges R (1991) Association between tumor necrosis factor-alpha and disease progression in patients with multiple sclerosis. N Engl J Med 325:467–472
- Sridhar R, Perumal PT, Etti S, Shanmugam G, Ponnuswamy MN, Prabavathyc VR, Mathivanan N (2004) Design, synthesis and anti-microbial activity of 1H-pyrazole carboxylates. Bioorg Med Chem Lett 14:6035
- Tale RH, Rodge AH, Hatnapure GD, Keche AP (2011) The novel 3,4dihydropyrimidin-2(1H)-one urea derivatives of *N*-aryl urea: synthesis, anti-inflammatory, antibacterial and antifungal activity evaluation. Bioorg Med Chem Lett 21:4648–4651
- van Dullemen HM, van Deventer SJ, Hommes DW, Bijl HA, Jansen J, Tytgat GN, Woody J (1995) Treatment of Crohn's disease with anti-tumor necrosis factor chimeric monoclonal antibody (cA2). Gastroenterology 109:129–135