

Green Chemistry

Cutting-edge research for a greener sustainable future

Accepted Manuscript

View Article Online
View Journal

This article can be cited before page numbers have been issued, to do this please use: Z. Fang, M. G. Flynn, J. E. Jackson and E. L. Hegg, *Green Chem.*, 2021, DOI: 10.1039/D0GC03597A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Thio-assisted reductive electrolytic cleavage of lignin β -O-4 models and authentic ligninZhen Fang,^{a,b} Michael G. Flynn,^a James E. Jackson^{*c} and Eric L. Hegg^{*a,b}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Avoiding the use of expensive catalysts and harsh conditions such as elevated temperatures and high pressures is a critical goal in lignin depolymerization and valorization. In this study, we present a thio-assisted electrocatalytic reductive approach using inexpensive reticulated vitreous carbon (RVC) as the working cathode to cleave the β -O-4-type linkages in keto aryl ethers. In the presence of a pre-electrolyzed disulfide (2,2'-dithiodiethanol) and a radical inhibitor (BHT) at room temperature at a current density of 2.5 mA/cm², cathodic reduction of nonphenolic β -O-4 dimers afforded over 90% of the corresponding monomeric C-O cleavage products in only 1.5 h. Extended to DDQ-oxidized poplar lignin, this combination of electric current and disulfide, applied over 6 h, released 36 wt% of ethyl acetate soluble fragments and 26 wt% of aqueous soluble fragments, leaving only 38 wt% of insoluble residue. These findings represent a significant improvement over the current alone values (24 wt% ethyl acetate soluble; 22 wt% aqueous soluble; 54 wt% insoluble residue) and represent an important next step in our efforts to develop a mild electrochemical method for reductive lignin deconstruction.

Introduction

Lignin is the most abundant renewable aromatic biopolymer on earth, and it has been recognized as a promising feedstock for fuels and chemicals.¹⁻⁵ The complex three-dimensional structure and diverse functionalities of lignin, however, complicate the deconstruction of this macromolecule. The alkyl and aryl units of lignin's heterogeneous polymer skeleton are connected via ether or carbon-carbon bonds,¹ with the β -O-4 ether linkage being the most abundant, accounting for approximately 50% and 60% of the linkages in softwoods and hardwoods, respectively.¹ Efficient cleavage of the β -O-4 linkage is therefore critical for lignin depolymerization, and indeed, aerobic oxidation,⁶⁻¹¹ reduction,¹²⁻¹⁴ and hydrolysis¹⁵⁻¹⁷ of this bond have been extensively studied as depolymerization strategies.^{1, 2, 4, 5}

An ideal lignin depolymerization process would i) avoid costly catalysts,^{18, 19} severe conditions such as high pressures/temperatures,^{12, 14, 20} and the use of hazardous gaseous reagents such as hydrogen; ii) produce a minimum amount of chemical waste; and iii) retain the maximum amount of the feedstock carbon and energy content.²¹ Compared to conventional reductants and oxidants, electric current is an inherently clean and inexpensive reagent, capable of cleaving lignin model compounds²²⁻²⁴ and depolymerizing real lignin.²⁵ Because only electrons and protons are removed or added

during electrolysis, generation of additional pollutants or reagent wastes is largely avoided.²⁶ Moreover, the counter-electrode cell offers the opportunity to perform additional desired organic transformations, or simply to split water. For instance, in a reductive process, the resulting pure O₂ byproduct from water-splitting is a "free" non-polluting oxidant which may be discharged or used in other non-electrolytic reactions.^{27, 28} Most existing reports on electrolytic lignin decomposition describe anodic oxidation, beginning with the pioneering studies by Utley *et al.* where mechanisms of electro-oxidative cleavage of lignin models were explored using nickel anodes in various solvent systems.²⁹ Most related oxidative strategies^{30, 31} have generally required expensive electrode materials such as Au³² or IrO₂,³³ or the assistance of photo-catalysts.^{34, 35} Some reductive approaches have also used electrodes made of catalytic metals such as Raney nickel or Pd.^{22, 23, 36} Electrocatalytic hydrogenation/hydrogenolysis (ECH) with such metal electrodes typically leads to saturation of potentially desirable aromatic products, though we have recently identified promising strategies to modulate selectivities.³⁷ These further reactions add to the complexity of the cleavage product streams, complicating isolation, purification, and further direct utilization of targeted products. Thus, the development of a simple, low-cost approach to selectively and efficiently cleave the ether bonds in β -O-4 linkages remains a critical need for effective lignin depolymerization and upgrading to useful products.

We previously reported the successful reductive cleavage of lignin-relevant α -keto aryl ether dimers by various small organic thiols.^{38, 39} This bio-inspired approach formed phenol and acetophenone quantitatively from 2-phenoxyacetophenone after 24 h treatment with 100 eq β -mercaptoethanol (BME) in refluxing acetonitrile. The resulting disulfide byproduct could


^a Department of Biochemistry & Molecular Biology, Michigan State University, 603 Wilson Rd, East Lansing, Michigan 48824, USA. Email: erichegg@msu.edu

^b Great Lakes Bioenergy Research Center, Michigan State University, 164 Food Safety and Toxicology Building, East Lansing, Michigan 48824, USA

^c Department of Chemistry, Michigan State University, 578 S Shaw Ln, East Lansing, Michigan 48824, USA. Email: jackson@chemistry.msu.edu

Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x





Scheme 1. Schematic illustration of disulfide reduction via successive one-electron transfer steps.




Figure 1. Proposed potential schematic mechanism of lignin deconstruction in a thiol-assisted electrolytic system.

theoretically be reduced back to the active thiol by various reductants. Of greatest interest here would be electrochemical reduction, regenerating the thiolate with no reagent byproducts (Scheme 1).⁴⁰ This approach could potentially lead to a sustainable cycle connecting electrochemical reduction to lignin cleavage via thiols as small diffusible redox carriers (Fig. 1). We now report a thio-assisted electro-reductive approach using inexpensive reticulated vitreous carbon (RVC) as the working cathode to cleave α -oxidized β -O-4-type linkages in model compounds and in lignin samples. This study, intended to extend our biomimetic thiol-mediated reductive cleavage method to electrochemical reduction, reveals additional cleavage processes that are not completely interpretable within the above mechanistic framework, but promise significant value nonetheless.

Results and Discussion

Thio-assisted electrolysis of 2-phenoxyacetophenone

Treatment of 2-phenoxyacetophenone (4.7 mM in 20 mL) with current alone was first performed as a control at room temperature in an H-type electrochemical cell. DMF, the polar aprotic solvent that most effectively promoted keto aryl ether bond cleavage in the previous S_N2 mechanism,³⁸ was utilized as the solvent on the cathode side. For most experiments, a constant current of 5 mA was applied, passing through a 2 cm² reticulated vitreous carbon (RVC) cathode (Table 1), providing an area-based current density of 2.5 mA/cm²; herein these conditions are simply described as “5 mA.” Interestingly, current alone cleaved 2-phenoxyacetophenone (compound I in Table 1) to phenol and acetophenone in 2.5 h (Entry 1), although the substrate barely decayed in the first 45 min (Fig. 2a). Pinacol (compound IV in Table 1), known to form from the reductive dimerization of acetophenone,⁴¹ was also produced as the reaction progressed. As expected, successful cleavage




Figure 2. Electrolysis of 2-phenoxyacetophenone (compound I) under 5 mA at room temperature with (a) current alone; and (b) current with direct (*i.e.*, applied simultaneous with) addition of RSSR.

required the α -keto (oxidized) form of this β -O-4 linkage model; no reaction occurred with the parent alcohol analog under 5 mA electric current alone (Fig. S1).

Stoichiometrically, only two equivalents of thiolate are needed to reductively cleave 2-phenoxyacetophenone via a reaction mechanism consisting of an S_N2 displacement of phenoxide by thiolate, followed by nucleophilic attack of thiolate on the thioether intermediate (compound V in table 1) to replace 1-phenylethen-1-olate (the enolate of acetophenone).³⁸ To probe the susceptibility of disulfide to electrolytic cleavage and its effect on the deconstruction of lignin dimer model compounds, 1 equivalent of 2,2'-dithiodiethanol (the disulfide derived from β -mercaptoethanol oxidation) was included in the DMF solution in the cathode cell (Entry 2). The 2-phenoxyacetophenone cleavage displayed a reaction profile similar to that in Fig. 2a, with virtually no formation of phenol and acetophenone in the first 1 h (Fig. 2b) at room temperature. In addition to 2,2'-dithiodiethanol, other disulfides including cystine (Fig. S2) exhibited comparable rates of substrate consumption. The critical nucleophile that cleaved the ether bond in the previous studies,^{38, 39} 2-hydroxyethanethiolate (here referred to as RS^-), was initially presumed to be the active species in the electrochemically driven reaction. At the 5 mA current used in these experiments, two equivalents of electrons were calculated to pass through the cathode in 1 h, and the RSSR should ideally then be reduced to 2 equivalents of RS^- . Therefore, RSSR was pre-electrolyzed in DMF at 5 mA for 1 h to reduce the disulfide, the current was then removed, and the substrate 2-phenoxyacetophenone was added (Entry 3). The substrate decayed only very slowly (Fig. 3a) under these conditions, consistent with our previous finding that heat was required to get reasonable reaction rates in the thiolate-enabled dimer cleavage.³⁸ In a parallel control, the combination of β -mercaptoethanol (hereafter referred to as RSH) with



ARTICLE

Table 1. Cathodic electrolysis of 2-phenoxyacetophenone (**I**, 4.7 mM) in the presence of 2,2'-dithiodiethanol (hereafter referred to as RSSR, 4.7 mM) with 5 mA electric current (2.5 mA/cm²); Anode: Pt wire in pH 8 phosphate buffer.

Entry	Current (5 mA) ^a	RSSR	Pre-electrolysis of RSSR	BHT	N ₂ atmosphere	Conv. (%)	Yields (%)			Current efficiency (%) ^c	
							II	III	IV		
1	yes	-	-	-	-	95	81	46	36	40	
2	yes	yes	-	-	-	> 99	97	24	68	53	
3	Removed after pre-electrolysis of RSSR	yes	1 hr	-	-	25	16	-	-	16	
4	yes	yes	1 hr	-	-	> 99	> 99	0.3	73	43	
5	yes	yes	1 hr	1 eq	-	> 99	96	15	88	45	
6	yes	yes	30 min (with BHT together)	1 eq	-	> 99	96	79	17	35	
7	yes	yes	-	-	yes	> 99	84	0.8	86	51	
8	yes	yes	-	1 eq	yes	> 99	82	60	8	25	
9	yes	yes	1 hr	-	yes	> 99 ^b	94	-	65	51	
10	yes	yes	1 hr	1 eq	yes	> 99	86	2	60	37	

Note: ^aPassage of two equivalents of electrons required approximately 1 h under these conditions; ^b1-phenylethanol (6%) was formed as well. ^cPre-electrolysis time was included in the calculation.





Figure 3. Effect of flowing current (5 mA) on cleavage of 2-phenoxyacetophenone (**I**) with 1 h pre-electrolyzed RSSR: (a) current was removed after **I** was added (Entry 3); (b) current was maintained throughout the entire reaction (Entry 4).



either an insoluble (K₂CO₃) or a soluble (Et₃N) base in DMF also showed little or no cleavage of 2-phenoxyacetophenone at room temperature (Fig. S3 and S4).

When the current was maintained following pre-electrolysis of RSSR, 2-phenoxyacetophenone decayed significantly faster (Fig. 3b). As expected, cleavage was faster after an hour of pre-electrolysis (*i.e.*, two equivalents of electrons, the quantity expected to fully reduce the disulfide), than with only 30 min of pre-electrolysis (Fig. S5). Notably, the thioether (compound **V** in






Figure 4. Effect of BHT on the electrolysis of 2-phenoxyacetophenone (I) using 5 mA of current. Condition: Prior to the addition of I, (a) BHT was added after RSSR was pre-electrolyzed for 1 h (Entry 5); (b) BHT and RSSR were pre-electrolyzed together for 30 min (Entry 6).

table 1), the critical diagnostic intermediate in the S_N2 mechanism previously studied, was not observed in the cathode cell. One potential explanation is that some species other than simple RS^- was involved in attacking the substrate during electrolysis.

Thiyl radical (referred to as RS^{\bullet}) formed during the cathodic disulfide reduction might contribute to the substrate decomposition. To probe this possibility, the current was turned off once the RSSR was pre-electrolyzed for 1 h, and then the

radical trap butylated hydroxytoluene (BHT) was added together with the substrate (current was still absent). Although 2-phenoxyacetophenone was cleaved to a small extent, the reaction ceased after 45 min (Fig. S6), showing a similar profile to that in Fig. 3a, where BHT was not employed. In addition, when RSSR was pre-electrolyzed for only 30 min, the current removed, BHT added, and the resulting solution stirred for an additional 30 min, no cleavage of substrate was observed (Fig. S7); however, if no BHT was added, a slow reaction did occur (Fig. S8). In a parallel control, a thiyl radical generated from RSH and AIBN (azobisisobutyronitrile) yielded essentially no substrate cleavage in the absence of current (Fig. S9). Thus, as in the previous thiol-based cleavage studies, we infer that dimer cleavage does not involve free thiyl radical processes,³⁸ suggesting instead that the electric current forms diffusible electron-carrier species, which activate ether bond cleavage.

In the absence of the radical trap BHT, when RSSR was pre-electrolyzed for 1 h before addition of 2-phenoxyacetophenone substrate, pinacol began forming almost immediately and only small quantities of acetophenone were observed as current flow continued and the starting material was cleaved (Fig. 3b). In contrast, addition of BHT to 1 h pre-electrolyzed RSSR (Entry 5) allowed acetophenone to accumulate to 50% yield before it was further reduced and dimerized to pinacol (Fig. 4a).

With these results in mind, BHT and RSSR (1:1, molar ratio) were pre-electrolyzed together for 1 h prior to addition of 2-phenoxyacetophenone and 5 mA of current was maintained throughout the entire reaction. Compared to pre-electrolyzing RSSR alone (for 30 min) in the absence of BHT (Fig. S5), in which acetophenone only accumulated to 14% yield and up to 70% of pinacol was formed, with the BHT present in the pre-electrolysis mixture (Entry 6), 2-phenoxyacetophenone was completely cleaved in 1.5 h with a quantitative yield of phenol and up to




Figure 5. Thio-assisted cathodic electrolysis of 2-phenoxyacetophenone (I) under N_2 atmosphere. (a) Direct addition (*i.e.*, applied simultaneous with) of RSSR (Entry 7) with I; (b) Direct addition of RSSR and BHT with I (Entry 8); (c) RSSR alone was pre-electrolyzed for 1 h prior to addition of I (Entry 9); (d) RSSR and BHT were together pre-electrolyzed for 1 h prior to addition of I (Entry 10).



80% of acetophenone (Fig. 4b). In addition, no pinacol was observed until all substrate had been cleaved (Fig. 4b) at 1.5 h. Doubling the quantity of BHT used did not lead to a statistically significant change in the reaction profile (Fig. S10).

To study further the potential role of a disulfide-derived radical species during substrate cleavage in the cathode cell, the effect of O₂, a known radical trap,⁴² was evaluated. Even when RSSR was not pre-electrolyzed, the substrate decayed much faster under anoxic conditions; under a N₂ atmosphere (Entry 7), the reaction was nearly complete after 2 h with pinacol and phenol being the major products (Fig. 5a), while 40% of the substrate remained unreacted after 2 h when the same reaction was performed under fully aerobic conditions (Fig. 2b). Moreover, addition of BHT to reactions performed under N₂ (Entry 8) again inhibited the formation of pinacol (Fig. 5b), but still showed faster cleavage of 2-phenoxyacetophenone than in the open air (Fig. S11).

One pathway considered for the C-O ether bond cleavage in the keto aryl ether substrate was one-electron reduction by an electron transfer agent. In that scenario, the disulfide radical anion (RSSR^{•-}) might act as an electron carrier. Oxygen could (a) consume an electron from RSSR^{•-},⁴³ slowing the rate of 2-phenoxyacetophenone cleavage relative to that under N₂ (Scheme 2a), or (b) react with RS[•] or RS⁻ to form the RSO₂[•] radical or anion (Scheme 2b).⁴⁴⁻⁴⁶ To test this possibility, RSSR was pre-electrolyzed for 1 h under N₂; again, after substrate addition, pinacol and phenol were the major products (Entry 9, Fig. 5c). Here, the substrate was completely consumed in only 1 h while previous controls had required at least 1.5 h. Including BHT during RSSR pre-electrolysis under N₂ (Entry 10) did not slow the 2-phenoxyacetophenone deconstruction (Fig. 5d), indicating again that the cleavage did not solely depend on the thiyl radical and that oxygen competed to abstract the electron from the electron transfer agent, thereby diverting the reaction.

Importantly, we found that the electrolyte used in this study, LiBF₄, was critical in promoting the cleavage of 2-phenoxyacetophenone. Li⁺, a known ion pairing agent and Lewis

acid,⁴⁷ favored the formation of pinacol (Scheme S1) and presumably shifted the equilibrium of the cleavage reaction. When NaBF₄ was employed as the electrolyte, approximately 50% of the substrate remained unreacted and pinacol was not observed even after 2.5 h (Fig. S12). Under these same conditions using LiBF₄ as the electrolyte, however, the substrate 2-phenoxyacetophenone was nearly completely cleaved in 1.5 h and pinacol was formed (Fig. 3b).

Current efficiencies (CE%) of all reactions were calculated as follows⁴⁸:




$$CE\% = \left(\frac{\text{mol}_{\text{prod}} \times F \times n}{C_{\text{total}}} \right) \times 100\%$$

where mol_{prod} is the moles of reduction products (phenol and pinacol); *F* is the Faraday constant (96485 C/mol); *n* is the number of electrons per reaction and *C*_{total} is the total charge passed. Overall, entry 2 shows the highest faradaic efficiency (~53%).

Thio-assisted electrolysis of other β-O-4 model compounds

In our previous study, 2-phenoxyacetophenone was reacted with 100 eq of β-mercaptoethanol (BME) and a large excess of insoluble base (K₂CO₃) in refluxing acetonitrile (~85 °C) for 24 h to achieve complete cleavage of substrate.³⁸ In the present work, however, 2-phenoxyacetophenone was completely decomposed in only 1.5 h at room temperature in the presence of only 1 eq of 2,2'-dithiodiethanol (the disulfide form of BME) and 5 mA electric current, producing a quantitative yield of phenol and up to 80% of acetophenone (Fig. 4b). In addition, the formation of pinacol was inhibited by BHT until after 1.5 h. Encouraged by the success in thio-assisted electrolytic cleavage of 2-phenoxyacetophenone, we explored this system with other α-keto β-O-4 dimers (**1a-c**) more closely related to real lignin. We posited that the presence on the aryl ketone of the electron-donating 4-methoxy or -hydroxy groups commonly found in lignin might inhibit the pinacol-forming reductive dimerization.

Table 2. Thio-assisted electrolysis of other β-O-4 dimers bearing methoxy groups

Dimer	Current alone ^a			Current + disulfide and BHT ^b		
	I (%)	II (%)	Conv. (%)	I (%)	II (%)	Conv. (%)
 1a	37	58	54	91	90	97
 1b	12	22	20	22	28	30
 1c	27	36	33	90	> 99	96

Note: ^aThe reaction was performed in DMF with 5 mA only (no disulfide) at room temperature for 2h; ^bThe disulfide (2,2'-dithiodiethanol) and BHT (1:1 molar ratio) were pre-electrolyzed together for 40 min in DMF at 5 mA, and the dimer was then added and the reaction stirred for 1.5 h.



ARTICLE

The results of these experiments are summarized in Table 2 and Figs. S13–S18. Electric current (5 mA) alone still cleaved the β -O-4 dimers bearing methoxy substituents. As expected, with pre-electrolyzed 2,2'-dithiodiethanol (1 eq), the cleavage time was shortened, except in the case of dimer **1b** where the substituent *para* to the ketone moiety is -OH. We speculate that this phenolic -OH in **1b** may be deprotonated, forming a quinone methide intermediate which would resist further reduction and cleavage of the ether bond (Scheme S2). Without addition of the disulfide, an "induction period" (around 45 min) was found in the electrolysis of both dimers **1a** and **1c** where very limited dimer cleavages occurred (Fig. S13 and S17), and acetoveratrone was generated only in moderate yields. Pre-electrolysis of the disulfide shortened the induction time by half (Fig. S14 and S18) and improved the yields of acetoveratrone to 72% (dimer **1a**) and 45% (dimer **1c**), with quantitative yields of guaiacol (dimer **1a**) and 53% syringol (dimer **1c**), respectively. Termination of the reaction prior to acetoveratrone decay, which usually occurred after the dimers were consumed (*i.e.*, at 1.5 h for **1a** and at 2 h for **1c**), would help maximize the monomer yields (Fig. S14 and S18). As expected, more promising results were observed when BHT and RSSR were pre-electrolyzed together. All the monomeric products were produced in 1.5 h with over 90% yields (Fig. S19 and S20). Notably, dimer **1a**, which required 24 h to produce a quantitative yield of acetoveratrone in the previous study (employing refluxing acetonitrile with 10 eq β -mercaptoethanol and a large excess of K_2CO_3),³⁸ decomposed to acetoveratrone (91%) and guaiacol (90%) using 1 eq of 2,2'-dithiodiethanol at room temperature in only 1.5 h (Fig. S19) in the current study. Together, the shortened reaction time, lower reaction temperature, and significantly reduced thio-compound loading suggest that electrochemical depolymerization might succeed under practical conditions with authentic lignin.

Thio-assisted electrolysis of lignin^{ox}

The successful cleavage of the model β -O-4 lignin dimers described above encouraged us to apply the optimized conditions to authentic lignin samples. The lignin used in this study was extracted from hybrid poplar following the previous described copper-catalyzed alkaline peroxide (Cu-AHP) strategy.⁴⁹ Cu-AHP lignin was oxidized (referred to as lignin^{ox}) with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ),¹⁰ which was confirmed by 2D HSQC spectra (Fig. S21). To improve the conductivity of the reaction mixture and the solubility of lignin^{ox}, a solvent mixture of pH 8 phosphate buffer/isopropanol (2:1, v/v) was employed and the reaction was performed under a higher current density (10 mA/cm²) for 6 h.




Figure 6. Composition of each fraction from electrolysis of lignin; wt% of each fraction was calculated based on the weight of lignin^{ox}. Note: wt% of aqueous-soluble fraction was determined by 1 - (wt % of EA soluble + wt % of insoluble)




Figure 7. Phenolic monomers from electrolysis of lignin^{ox}; Products were determined by GC-MS.

Electric current was found to be critical in this aqueous reaction system as lignin^{ox} showed essentially no change in the absence of flowing current. This is consistent with our previous finding that water inhibited the thiol-assisted keto aryl ether cleavage. In the previous study, increased thiol loading, elevated temperature and N₂ protection were required to break the ether bond.³⁸ In contrast, at room temperature, the combination of electric current and disulfide successfully solubilized 62% of lignin^{ox} in only 6 h (Fig. 6) while only 46% of the lignin^{ox} was solubilized under analogous conditions in the absence of the disulfide, again demonstrating that the thiol assists in the lignin cleavage process. In the control reaction where no disulfide was added, electric current (20 mA) alone afforded 24 wt% of ethyl acetate soluble (EA soluble) products from lignin^{ox} (Fig. 6), including various functionalized S/G/H type phenolic monomers (Fig. 7) along with a series of dimers, trimers, tetramers and other oligomers (Fig. S22). The addition of RSSR (2,2'-dithiodiethanol) and BHT to the reaction mixture improved the yield of EA soluble products to 36 wt%, with a concomitant decrease in the insoluble material from 54 wt% to 38 wt% (Fig. 6). Although slight differences were observed via LCMS-QTOF in the distribution of products in the EA soluble fractions (Fig. S23), the profiles were relatively similar, indicating



that the addition of the disulfide did not significantly complicate the product streams. Addition of disulfide and BHT increased the amount of aqueous soluble products from 22 wt% to 26 wt%, and analysis by gel permeation chromatography (GPC) indicated that the molecular weight of the major products ranged from approximately 600 - 1000 Da (Fig. S24). Overall, at ambient temperature and atmosphere, electro reductive degradation of lignin^{ox} solubilized a significant fraction of the lignin, leaving only 38 - 54 wt% of insoluble residual. Compared to previous work on depolymerization of different types of lignin, *i.e.* via catalytic electrolysis (77 - 82 wt% residual),⁵⁰ aerobic oxidation-hydrolysis (47 - 78 wt% residual)⁹ or a two-step oxidative approach (50 - 91 wt% residual),¹⁰ the present study represents an encouraging first step in developing a simple new strategy for lignin depolymerization and valorization.

Conclusion

Building on our previously reported thiolate-based reductive cleavage of β -O-4 linkages via an S_N2 pathway,^{38, 39} this study aimed to electrochemically reduce the disulfide byproduct to achieve an electrocatalytic cycle (Fig. 1). Electrolytic reductive treatment of α -keto β -O-4 model dimers together with disulfides does indeed result in ether cleavage. With pre-electrolyzed disulfide and BHT, this approach cleaved the β -O-4 models and produced the corresponding phenolic and keto monomers with over 90% yields in only 1.5 h. Using only 1 eq of disulfide at room temperature, this surprisingly mild and rapid cleavage represented an unexpectedly large improvement over our previous mechanistically motivated work in which 10 eq. of thiol in refluxing acetonitrile required 24 h to completely cleave the oxidized β -O-4 dimers. Applied to pre-oxidized authentic lignin, the electrochemical method showed promising levels of cleavage and solubilization. In the electrochemical context, the use of thiols, originally envisioned to serve as small diffusible redox carriers via the thiol/disulfide couple, enables a significantly milder approach to deconstruction of lignin to small-molecule fragments. Though the mechanistic details of these improvements remain incompletely understood, we summarize our findings as follows: 1) pre-reduction of the disulfide forms active species that promote cleavage of 2-phenoxyacetophenone; 2) for complete substrate conversion, electric current must be supplied throughout the reaction time; 3) dimer cleavage is not likely to involve free thiyl radicals; and 4) substrate is cleaved faster under a N_2 atmosphere than under air. These results represent a substantial step forward in linking the reducing power of electrical current to the deconstruction and potential valorization of lignin as an organic feedstock.

Experimental Section

General Materials

Acetone (ACS grade), 3,4-dimethoxyacetophenone (98%), 4-hydroxy-3-methoxyacetophenone (acetovanillone, 98%), 2,6-di-tert-butyl-4-methylphenol (BHT, >99% GC), 1-phenylethanol

(98%), acetonitrile (99.5+%), dichloromethane (ACS grade), *N*-bromosuccinimide (NBS, 99%), lithium tetrafluoroborate ($LiBF_4$, 98%), *N,N*-dimethylformamide (DMF, 99.8%, anhydrous), syringol, 2,3-diphenyl-butane-2,3-diol (pinacol), hexane (>98.5%), and plastic silica gel 60 F-254 plates (for thin layer chromatography, TLC) were purchased from Sigma-Aldrich (St. Louis, MO). Acetophenone (98.8%) and triethyl amine were purchased from J.T. Baker (Phillipsburg, NJ) while benzyl chloride and 2-phenoxyacetophenone (>98%) were purchased from TCI, America, Ltd (Portland, OR). Guaiacol (99+%) was purchased from Acros Organics (New Jersey). Tetrahydrofuran (THF, 99+% with 250 ppm BHT) and phenol (99+%) were purchased from Alfa Aesar (Haverhill, MA). Sodium sulfate (anhydrous) and ethyl acetate (99.9%) were purchased from Fisher Chemicals (Pittsburgh, PA), and *p*-toluenesulfonic acid (TsOH) was purchased from Spectrum (New Brunswick, NJ). Finally, Chloroform-*d* ($CDCl_3$, 99.8 atom% D) was purchased from Cambridge Isotope Laboratories (Andover, MA).

Characterization of Products

Column chromatography was performed on Silicycle (Quebec City, Canada) SiliaFlash P60 silica gel (40-63 μ m). 1H and ^{13}C NMR spectra were acquired on Agilent 500/54 premium shielded instruments with tetramethylsilane or residual solvent as the internal reference. High performance liquid chromatography (HPLC) was conducted on an Agilent 1260 Infinity equipped with an Agilent G1315D1260 diode array detector VL, monitoring at 250 nm (for experiments related to dimer **1a-c**) or 280 nm (for experiments related to 2-phenoxyacetophenone) and recording 190-400 nm. A 10 L sample injection using a Supelco Ascentis Express C18 column (15 cm x 4.6 cm, 2.7 μ m, St. Louis, MO) at 0.5 mL/min, with a mobile phase of 35/65 and 50/50 acetonitrile/water (with 0.5 M H_2SO_4) was applied at 30 °C on analyses of **1a-c** and 2-phenoxyacetophenone related experiments, respectively. Yields of products were calibrated using authentic compounds as external standards. Gel permeation chromatography (GPC) was performed using a Waters Ultrahydrogel 250 7.8 x 300 mm column equipped with a Waters Ultrahydrogel 6 x 40 mm guard column at 40 °C with a mobile phase of 5 mM NaOH in 80/20 0.1 M aqueous sodium nitrate/acetonitrile at a flow rate of 0.7 mL/min. Polystyrene sulfonic acid and poly(ethylene glycol) were employed as external standards. $^1H^{13}C$ -gradient heteronuclear single quantum coherence (HSQC) was performed at ambient temperature on a 500 MHz Bruker NMR spectrometer equipped with a 5 mm iProbe (BBO probe). Spectra were collected by utilizing the Bruker pulse sequence "hsqcetdpsisp2.3" with an acquisition time of 63.9 ms (F2, 512 complex points for 1H) and 63.9 ms (F1, 1024 increments for the ^{13}C dimension), using a delay of 1.5 s with 48 scans per increment with spectral widths of 8013 Hz (1H) and 20 kHz (^{13}C).

General Procedure for Thio-assisted Cathodic Electrolysis of Dimers

Reactions were conducted at room temperature up to 2.5 h in an H-type electrochemical cell (separated by a Nafion 117



membrane) wherein a platinum wire in 20 mL of pH 8 phosphate buffer was placed in the anode half-cell and the reticulated vitreous carbon (RVC, 2 cm² immersed) was employed as the cathode. The start of the reaction (t=0) was defined as the time point when substrate (2-phenoxyacetophenone or dimers **1a-c**) was added. (Note: reactions were run in open air unless otherwise noted).

For reactions run without pre-electrolysis, 20 mg of the dimer (2-phenoxyacetophenone or **1a-c**), 2,2'-dithiodiethanol (none or 1 eq.) and BHT (none or 1 eq.) were directly added to 20 mL DMF in the cathode half-cell and a 5 mA electric current was applied for the duration of the reaction (the voltage of the system ranged from 5.6 - 6.0 V in DMF under our conditions). Sample aliquots (250 μ L) were taken from the cathode cell at timed intervals and diluted with another 250 μ L of acetonitrile before HPLC analysis.

For reactions involving pre-electrolysis of the disulfide (2,2'-dithiodiethanol), 2,2'-dithiodiethanol alone was added to 20 mL DMF and the 5 mA electric current was applied for 30 min or 1 h. Then 20 mg of the dimer (2-phenoxyacetophenone or **1a-c**) and BHT (none or 1 eq.) was added in solid form. Sampling for HPLC analysis was performed as above.

For reactions run under a N₂ atmosphere: 20 mL of DMF alone was placed in the sealed cathode half-cell and the cell was purged with N₂ for 30 min with stirring. Subsequent steps were as described above except that a N₂ balloon was connected to the cathode half-cell throughout the reaction. At different time intervals, 500 μ L samples were taken from the cathode cell for HPLC analysis.

Synthesis of β -O-4 Dimers (**1a-c**)

Dimers **1a-c** were synthesized based on previous reports with only minor modifications.³⁸

1-(3,4-dimethoxyphenyl)-2-(2-methoxyphenoxy)ethan-1-one (**1a**).

A solution of N-bromosuccinimide (NBS, 1.96 g, 11.1 mmol) dissolved in anhydrous acetonitrile (20 mL) was added dropwise with stirring to a solution of 3,4-dimethoxyacetophenone (2 g, 11.1 mmol) and toluenesulfonic acid (TsOH, 3.2 g, 16.6 mmol) in anhydrous acetonitrile (30 mL). The mixture was heated to 100 °C and stirred for 2 h under a N₂ atmosphere. After cooling to room temperature, the solvent was evaporated *in vacuo*. The crude product was re-dissolved in dichloromethane (DCM, 30 mL) and washed three times with 30 mL aliquots of deionized water. The organic layer was separated and dried with anhydrous Na₂SO₄ and concentrated under vacuum. The crude product was purified by column chromatography on silica gel (hexane/ethyl acetate = 2/1) to produce 2-bromo-1-(3,4-dimethoxyphenyl)ethanone (2.54 g, 9.8 mmol) as a white solid in 88% yield.

A mixture of 2-bromo-1-(3,4-dimethoxyphenyl)ethanone (2.54 g, 9.8 mmol), guaiacol (1.21 g, 9.8 mmol), and K₂CO₃ (1.35 g, 9.8 mmol) in acetone (50 mL) was stirred at room temperature overnight. The solid was removed by vacuum filtration and the solution was extracted with three 20 mL aliquots of DCM. The organic layer was separated and dried with anhydrous Na₂SO₄ and concentrated under vacuum. The crude product was purified by column chromatography on silica

gel (hexane/ethyl acetate = 2/1) to produce **1a** (2.65 g, 8.8 mmol) as a white solid in 89% yield. DOI: 10.1039/D0GC03597A

¹H NMR (500 MHz, CDCl₃): 7.69 (dd, *J* = 8.4, 2.0 Hz, 1H), 7.61 (d, *J* = 2.0 Hz, 1H), 7.01 - 6.94 (m, 1H), 6.94 - 6.88 (m, 2H), 6.88 - 6.82 (m, 2H), 5.31 (s, 2H), 3.96 (s, 3H), 3.94 (s, 3H), 3.90 (s, 3H); ¹³C NMR (126 MHz, CDCl₃): 193.2, 153.7, 149.6, 149.1, 147.5, 127.8, 122.7, 122.3, 120.8, 114.6, 112.0, 110.4, 110.1, 71.9, 56.1, 56.0, 55.9. Spectral data are in accordance with those previously reported.³⁸

2-(2,6-dimethoxyphenoxy)-1-(4-hydroxy-3-methoxyphenyl)ethan-1-one (**1b**).

Benzyl chloride (9.1 g, 72 mmol) was added dropwise to a solution of DMF containing acetovanillone (10 g, 60 mmol), K₂CO₃ (12.5 g, 90 mmol) and KI (0.2 g, 2 wt%) and the resulting solution was stirred at 40 °C overnight. The mixture was poured into ice water and the solid precipitate was filtered and washed with DI water. Drying in the vacuum oven (50 °C) gave 14.3 g of the benzyl protected acetovanillone in a yield of 93%.

A solution of N-bromosuccinimide (NBS, 6.96 g, 39 mmol) dissolved in anhydrous acetonitrile (30 mL) was added dropwise with stirring to a solution of the above benzyl protected acetovanillone (10 g, 39 mmol) and toluenesulfonic acid (TsOH, 10 g, 58.6 mmol) in anhydrous acetonitrile (70 mL). The mixture was then heated and stirred at 100 °C for 2 h under a N₂ atmosphere. After cooling to room temperature, the solvent was evaporated *in vacuo*. The crude product was re-dissolved in dichloromethane (DCM, 70 mL) and washed three times with 30 mL aliquots of with deionized water. The organic layer was separated and dried with anhydrous Na₂SO₄ and concentrated under vacuum. Without further purification, the resulting crude protected bromoacetovanillone (5 g, 14.9 mmol) was combined with syringol (2.3 g, 14.9 mmol) and K₂CO₃ (2.06 g, 14.9 mmol) in acetone (100 mL) and stirred at room temperature overnight. The solid KBr byproduct was removed by vacuum filtration and the solution was extracted with three 30 mL aliquots of DCM. The organic layers were combined and dried with anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude protected dimer (2 g) was dissolved in methanol (50 mL), 10 % Pd/C (0.2 g) was added, and the mixture was stirred for 3 h at room temperature. The solid catalyst was separated by vacuum filtration and washed with methanol. Then the filtrate was concentrated under vacuum and the crude product was purified by column chromatography on silica gel (hexane/ethyl acetate = 2/1) to give a yellow solid.

¹H NMR (500 MHz, CDCl₃): 7.69 - 7.64 (m, 2H), 7.02 (t, *J* = 8.4 Hz, 1H), 6.95 (d, *J* = 8.1 Hz, 1H), 6.59 (d, *J* = 8.4 Hz, 2H), 5.16 (s, 2H), 3.96 (s, 3H), 3.82 (s, 6H); ¹³C NMR (126 MHz, CDCl₃): 193.5, 153.2, 150.5, 146.6, 136.6, 128.0, 124.0, 123.5, 113.8, 110.4, 105.3, 75.2, 56.1, 56.1. Spectral data are in accordance with those previously reported.³⁸

2-(2,6-dimethoxyphenoxy)-1-(3,4-dimethoxyphenyl)ethan-1-one (**1c**).

A mixture of 2-bromo-1-(3,4-dimethoxyphenyl)ethanone (5 g, 19.3 mmol), syringol (3 g, 19.3 mmol) and K₂CO₃ (2.66 g, 19.3 mmol) in acetone (100 mL) was stirred at room temperature overnight. The solid was removed by vacuum filtration and the solution was extracted with three 30 mL aliquots of DCM. The



organic layer was separated and dried with anhydrous Na₂SO₄ and concentrated under vacuum. The crude product was purified by column chromatography on silica gel (hexane/ethyl acetate = 2/1) to produce **1c** (4.8 g, 14.4 mmol) as a white solid in 75% yield.

¹H NMR (500 MHz, CDCl₃): 7.73 (dd, *J* = 8.3, 2.0 Hz, 1H), 7.66 (d, *J* = 2.0 Hz, 1H), 7.02 (t, *J* = 8.4 Hz, 1H), 6.90 (d, *J* = 8.4 Hz, 1H), 6.59 (d, *J* = 8.5 Hz, 2H), 5.16 (s, 2H), 3.95 (d, *J* = 1.8 Hz, 6H), 3.82 (s, 6H); ¹³C NMR (126 MHz, CDCl₃): 193.7, 153.4, 153.3, 153.2, 148.9, 136.6, 128.3, 124.0, 123.0, 110.6, 110.0, 105.2, 105.2, 75.2, 56.0, 56.0, 55.9. Spectral data are in accordance with those previously reported.³⁸

Oxidation of Cu-AHP lignin

Lignin from alkaline pre-extracted hybrid poplar was isolated via the Cu-AHP pretreatment process as previously described using 1 mM of copper and 2 mM of bipyridine.⁴⁹ Hydrogen peroxide (H₂O₂, 100 mg per g of biomass) was added in 10 equal aliquots at 1 h intervals while shaking at 30 °C. DDQ-catalyzed oxidation of Cu-AHP lignin was performed following a previous literature report.¹⁰ Briefly, 3 g of lignin was solubilized in 1,2-dimethoxyethane/2-ethoxyethanol (v/v = 3:2, 42 mL) and then DDQ (10 wt%) and t-BuONO (10 wt%) were added. The solution was heated at 80 °C overnight under an oxygen atmosphere. The oxidized lignin product (lignin^{ox}) was precipitated by addition of 500 mL of diethyl ether, isolated by filtration, washed with additional diethyl ether (200 mL) and dried *in vacuo*, yielding 2.7 g of light-brown lignin^{ox} was obtained.

General procedure for thio-assisted electrolysis of lignin^{ox}

Reactions were conducted at room temperature in an H-type electrochemical cell (separated by a Nafion 117 membrane) with 20 mL of pH 8 phosphate buffer and 20 mL of pH phosphate 8 buffer/isopropanol in the anodic and cathodic cell, in which a platinum wire was used as the anode and the reticulated vitreous carbon (RVC) was employed as the cathode, respectively. A mixture of LiBF₄ (100 mg) and BHT (100 mg) was added into the cathodic cell. After 1 h of pre-electrolysis at 20 mA (current density = 10 mA/cm²), lignin^{ox} (100 mg) and 2,2'-dithiodiethanol (RSSR, 100 mg) were added to the catholyte and the reaction continued at 20 mA for an additional 6 h. The solvent was removed under a N₂ flow, and the crude product was solubilized in 20 mL EtOAc/water (1:1, v/v). The solution was acidified with 60% H₂SO₄ to pH 2, precipitating insoluble material. The precipitate was filtered, washed with water, dried *in vacuo* and the remaining filtrate was extracted with 30 mL EtOAc (3 x). The organic layers were combined, dried with anhydrous Na₂SO₄ and concentrated *in vacuo*. In the parallel control reaction, all set-up and work-up protocols were the same except that the RSSR and BHT components were omitted.

Conflicts of interest

A patent application has been submitted by E. L. Hegg, J. E. Jackson, and G. E. Klinger (Methods for Lignin Depolymerization Using Thiols - WO/2018/195000). As holders of this patent

application, we could potentially benefit financially from the technology discussed in this manuscript. DOI: 10.1039/D0GC03597A

Acknowledgements

This material is based upon work supported by the Great Lakes Bioenergy Research Center, U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under Award Number DE-SC0018409. The authors gratefully acknowledge Casey Johnny and Tony Schillmiller in the Research Technology Support Facility at Michigan State University for the GC-MS and LCMS-TOF analyses. ZF thanks Nicholas Schlecht for help in performing selected reactions and Dr. Asmaul Hoque at UW-Madison for running the cyclic voltammetry of 2,2'-dithiodiethanol (Fig. S31).

Notes and references

1. P. C. A. B. Joseph Zakzeski, Anna L. Jongerius, and Bert M. Weckhuysen, *Chem. Rev.*, 2010, **110**, 3552-3599.
2. Z. Sun; B. Fridrich; A. de Santi; S. Elangovan; K. Barta, *Chem. Rev.*, 2018, **118** (2), 614-678.
3. W. Schutyser; T. Renders; S. Van den Bosch; S. F. Koelewijn; G. T. Beckham; B. F. Sels, *Chem. Soc. Rev.*, 2018, **47** (3), 852-908.
4. B. M. Upton; A. M. Kasko, *Chem. Rev.*, 2016, **116** (4), 2275-2306.
5. C. Li; X. Zhao; A. Wang; G. W. Huber; T. Zhang, *Chem. Rev.*, 2015, **115** (21), 11559-11624.
6. I. Hasegawa; Y. Inoue; Y. Muranaka; T. Yasukawa; K. Mae, *Energy Fuels*, 2011, **25** (2), 791-796.
7. A. Rahimi; A. Azarpira; H. Kim; J. Ralph; S. S. Stahl, *J. Am. Chem. Soc.*, 2013, **135** (17), 6415-6418.
8. A. Rahimi; A. Ulbrich; J. J. Coon; S. S. Stahl, *Nature*, 2014, **515** (7526), 249-252.
9. A. Das; A. Rahimi; A. Ulbrich; M. Alherech; A. H. Motagamwala; A. Bhalla; L. da Costa Sousa; V. Balan; J. A. Dumesic; E. L. Hegg; B. E. Dale; J. Ralph; J. J. Coon; S. S. Stahl, *ACS Sustain. Chem. Eng.*, 2018, **6** (3), 3367-3374.
10. Y. Song; A. H. Motagamwala; S. D. Karlen; J. A. Dumesic; J. Ralph; J. K. Mobley; M. Crocker, *Green Chem.*, 2019, **21** (14), 3940-3947.
11. O. Musil; M. Holzlechner; S. Winkler; G. Gübitz; A. Potthast; T. Rosenau; S. Böhmendorfer, *ACS Sustain. Chem. Eng.*, 2019, **7** (18), 15163-15172.
12. Y. Li; B. Demir; L. M. Vázquez Ramos; M. Chen; J. A. Dumesic; J. Ralph, *Green Chem.*, 2019, **21** (13), 3561-3572.
13. E. Feghali; G. Carrot; P. Thuéry; C. Genre; T. Cantat, *Energy Environ. Sci.*, 2015, **8** (9), 2734-2743.
14. E. M. Anderson; R. Katahira; M. Reed; M. G. Resch; E. M. Karp; G. T. Beckham; Y. Román-Leshkov, *ACS Sustain. Chem. Eng.*, 2016, **4** (12), 6940-6950.
15. L. Fan; Y. Zhang; S. Liu; N. Zhou; P. Chen; Y. Cheng; M. Addy; Q. Lu; M. M. Omar; Y. Liu; Y. Wang; L. Dai; E. Anderson; P. Peng; H. Lei; R. Ruan, *Bioresour. Technol.*, 2017, **241**, 1118-1126.
16. G. Jiang; D. J. Nowakowski; A. V. Bridgwater, *Energy Fuels*, 2010, **24** (8), 4470-4475.
17. P. R. Patwardhan; R. C. Brown; B. H. Shanks, *ChemSusChem*, 2011, **4** (11), 1629-1636.
18. P. J. Deuss; K. Barta, *Coord. Chem. Rev.*, 2016, **306**, 510-532.
19. S. K. Hanson; R. T. Baker, *Acc. Chem. Res.*, 2015, **48** (7), 2037-2048.



20. S. Van den Bosch; W. Schutyser; S. F. Koelewijn; T. Renders; C. M. Courtin; B. F. Sels, *ChemComm*, 2015, **51** (67), 13158-13161.
21. C. H. Lam; S. Das; N. C. Erickson; C. D. Hyzer; M. Garedew; J. E. Anderson; T. J. Wallington; M. A. Tamor; J. E. Jackson; C. M. Saffron, *Sustain. Energy Fuels*, 2017, **1** (2), 258-266.
22. B. Mahdavi, A. Lafrance, A. Martel, J. Lessard, H. Me, and L. Brossard, *J. Appl. Electrochem.*, 1997, **27**, 605-611.
23. A. Cyr; F. Chiltz; P. Jeanson; A. Martel; L. Brossard; J. Lessard; H. Ménard, *Can. J. Chem.*, 2000, **78** (3), 307-315.
24. X. Du; H. Zhang; K. P. Sullivan; P. Gogoi; Y. Deng, *ChemSusChem*, 2020, **13** (17), 4318-4343.
25. S. Stiefel; J. Lölsberg; L. Kipshagen; R. Möller-Gulland; M. Wessling, *Electrochem. Commun.*, 2015, **61**, 49-52.
26. B. A. Frontana-Urbe; R. D. Little; J. G. Ibanez; A. Palma; R. Vasquez-Medrano, *Green Chem.*, 2010, **12** (12), 2099-2119.
27. B. H. Nguyen; R. J. Perkins; J. A. Smith; K. D. Moeller, *J. Org. Chem.*, 2015, **80** (24), 11953-11962.
28. T. Wu; B. H. Nguyen; M. C. Daugherty; K. D. Moeller, *Angew. Chem.*, 2019, **58** (11), 3562-3565.
29. V. L. Pardini, Smith, C.Z., Utley, J.H., Vargas, R.R. and Viertler, H., *J. Org. Chem.*, 1991, **56** (26), 7305-7313.
30. H. Zhu; L. Wang; Y. Chen; G. Li; H. Li; Y. Tang; P. Wan, *RSC Advances*, 2014, **4** (56), 29917-29924.
31. Y.-s. Wang; F. Yang; Z.-h. Liu; L. Yuan; G. Li, *Catal. Commun.*, 2015, **67**, 49-53.
32. P. Parpot; A. P. Bettencourt; A. M. Carvalho; E. M. Belgsir, *J. Appl. Electrochem.*, 2000, **30** (6), 727-731.
33. R. Tolba; M. Tian; J. Wen; Z.-H. Jiang; A. Chen, *J. Electroanal. Chem.*, 2010, **649** (1-2), 9-15.
34. I. Bosque; G. Magallanes; M. Rigoulet; M. D. Karkas; C. R. J. Stephenson, *ACS Cent. Sci.*, 2017, **3** (6), 621-628.
35. M. Tian; J. Wen; D. MacDonald; R. M. Asmussen; A. Chen, *Electrochem. Commun.*, 2010, **12** (4), 527-530.
36. P. C. Dabo, Andre; Lessard, Jean; Brossard, Louis; Menard, Hugues, *Can. J. Chem.*, 1999, **77**, 1225-1229.
37. Y. Zhou; G. E. Klinger; E. L. Hegg; C. M. Saffron; J. E. Jackson, *J. Am. Chem. Soc.*, 2020, **142** (8), 4037-4050.
38. G. E. Klinger; Y. Zhou; P. Hao; J. Robbins; J. M. Aquilina; J. E. Jackson; E. L. Hegg, *ChemSusChem*, 2019, **12** (21), 4775-4779.
39. G. E. Klinger; Y. Zhou; J. A. Foote; A. M. Wester; Y. Cui; M. Alherrech; S. S. Stahl; J. E. Jackson; E. L. J. C. Hegg, *ChemSusChem*, 2020, DOI: org/10.1002/cssc.202001238.
40. S. Antonello, Benassi, R., Gavioli, G., Taddei, F. and Maran, F., *J. Am. Chem. Soc.*, 2002, **124** (25), 7529-7538.
41. K. Nakahara; K. Naba; T. Saitoh; T. Sugai; R. Obata; S. Nishiyama; Y. Einaga; T. Yamamoto, *ChemElectroChem*, 2019, **6** (16), 4153-4157.
42. A. Fava; G. Reichenbach; U. J. J. o. t. A. C. S. Peron, *J. Am. Chem. Soc.*, 1967, **89** (25), 6696-6700.
43. S. a. A. Mezyk, D., *J. Chem. Soc., Perkin Trans.*, 1999, **2** (7), 1411-1420.
44. H. Wariishi, Khadar Valli, V. Renganathan, and Michael H. Gold, *J. Biol. Chem.*, 1989, **264** (24), 14185-14191.
45. X. Zhang, Zhang, N., Schuchmann, H.P. and von Sonntag, C., *J. Phys. Chem.*, 1994, **98**, 6541-6547.
46. M. P. Bertrand, *Org. Prep. Proced. Int.*, 1994, **26** (3), 257-290.
47. C. P. Andrieux, Robert, M. and Saveant, J.M., *J. Am. Chem. Soc.*, 1995, **117**, 9340-9346.
48. C. H. Lam; C. B. Lowe; Z. Li; K. N. Longe; J. T. Rayburn; M. A. Caldwell; C. E. Houdek; J. B. Maguire; C. M. Saffron; D. J. Miller; J. E. Jackson, *Green Chemistry*, 2015, **17** (1), 601-609.
49. A. Bhalla; N. Bansal; R. J. Stoklosa; M. Fountain; J. Ralph; D. B. Hodge; E. L. Hegg, *Biotechnol. Biofuels*, 2016, **9**, DOI: org/10.1186/s13068-016-0442-0
50. X. Du; W. Liu; Z. Zhang; A. Mulyadi; A. Brittain; J. Gong; Y. Deng, *ChemSusChem*, 2017, **10** (5), 847-854.

