Tetrahedron Letters 50 (2009) 1236-1239

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet

Stereoselective construction of an *anti*-β-alkoxy ether by Ireland–Claisen rearrangement for medium-ring ether synthesis

Kenshu Fujiwara^{a,*}, Natsumi Kawamura^a, Hidetoshi Kawai^{a,b}, Takanori Suzuki^a

^a Division of Chemistry, Graduate School of Science, Hokkaido University, Sapporo 060-0810, Japan
 ^b PRESTO, Japan Science and Technology Agency (JST), 4-1-8 Honcho Kawaguchi, Saitama 332-0012, Japan

ARTICLE INFO

Article history: Received 1 December 2008 Revised 5 January 2009 Accepted 8 January 2009 Available online 10 January 2009

Keywords: Stereoselective synthesis Cyclic ethers Ireland–Claisen rearrangement Ring-closing olefin metathesis

ABSTRACT

The asymmetric synthesis of an acyclic *anti*- β -alkoxy ether was achieved by the Ireland–Claisen rearrangement of Z-3-alkoxy-2-propenyl glycolate ester, prepared from Garner's aldehyde, a glycolic acid derivative, and ethynyl *N*,*N*-diisopropylcarbamate. The resulting acyclic ether was facilely converted to seven- and eight-membered cyclic ethers via processes involving ring-closing olefin metatheses. © 2009 Elsevier Ltd. All rights reserved.

Natural medium ring ether compounds, which are often bioactive, have attracted synthetic interest due to their particular structural features, such as medium-sized ring systems or stereochemical complexity around the ether linkage.¹ While a large number of synthetic methods have been developed by many research groups,² ring-closing olefin metathesis (RCM)³ based approaches have been extensively explored,^{4,5} because RCM realizes efficient ring-closure under mild catalytic conditions and tolerates a wide variety of functional groups in its substrates. However, the construction of the stereochemically complex ether linkage in the diene substrates of RCM for medium ring ethers is still difficult.⁶ Therefore, approaches toward these systems have focused on the establishment of an efficient method for the stereoselective synthesis of acyclic ethers.

In our recent studies on the construction of medium ring ethers using RCM,⁷ we found that the Ireland–Claisen rearrangement of an acyclic *Z*-3-alkoxy-2-propenyl glycolate ester stereoselectively produced an acyclic *anti*- β -alkoxy ether, corresponding to an ether moiety of a natural *trans*-fused polycyclic ether.^{7a} Here, we describe the development of an improved variant of this rearrangement, which accomplishes the asymmetric synthesis of an acyclic *anti*- β -alkoxy ether from easily available starting materials for efficient medium ring ether synthesis.

The goal of this project was to devise a facile method to construct a terminal medium ring in a *trans*-fused polycyclic ether system (1) (Scheme 1). Toward this goal, a synthetic route was



Scheme 1.





^{*} Corresponding author. Tel.: +81 11 706 2701; fax: +81 11 706 4924. E-mail address: fjwkn@sci.hokudai.ac.jp (K. Fujiwara).

^{0040-4039/\$ -} see front matter \odot 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2009.01.011

designed that relied on RCM for the ring closure of the terminal medium ring of 1 and chirality-transferring Ireland-Claisen rearrangement for setting up the *anti*- β -alkoxy ether substrate (2) stereoselectively. Based on our previous study that showed exclusive anti selectivity in the rearrangement of a simple Z-3-alkoxy-2-propenyl glycolate ester,^{7a} 1-substituted Z-3-alkoxy-2-propenyl ester 4 was employed as a precursor of 2. We also expected that the chirality at C1 of the propenyl group of 4 would be efficiently transferred to the newly formed stereocenters of 2 via putative transition state **3**.⁸ Ester **4** would be prepared from polycyclic ether 5, having a glycolic acid moiety, and chiral Z-3-alkoxy-2-propenol 6. We chose to synthesize 6 by stereoselective addition of protected vnol **9** to aldehvde **8** with the assistance of the R^2 group as a chiral auxiliary, followed by Lindlar hydrogenation. Thus, the proposed synthetic route was demonstrated by the model synthesis of oxane-fused medium ring ethers 13 and 14 from oxane-connected glycolic acid 10⁹ corresponding to 5, Garner's aldehyde (11),¹⁰ reported to be efficient for stereoselective addition reactions,¹¹ and ethynyl N,N-diisopropylcarbamate (**12**),¹² a stable ynol derivative.

First, chiral Z-3-carbamoyloxy-2-propenol **17**, corresponding to **6**, was selectively synthesized (Scheme 2). The addition reaction of **11** with the acetylide derived from **12** in the presence of LiBr¹³ produced alcohol **15**¹⁴ (70%) with almost complete diastereoselectivity. The configuration of the newly generated stereocenter of **15** was determined by the presence of an NOE between Ha and Hb of **16**, derived from **15** by hydrogenation and subsequent basic treatment. Lindlar hydrogenation of **15**, followed by simple filtration, quantitatively gave **17**¹⁴ in an almost pure form.

The chirality-transferring Ireland–Claisen rearrangement of ester **18**,¹⁴ prepared by the condensation of **10** and **17** with EDCI-HCl (74%), was found to proceed successfully under the following conditions. After ester **18** was deprotonated with KHMDS (5 equiv)¹⁵ in THF at –78 °C for 30 min, the resulting enolate was treated successively with TMSCl (5 equiv) and diethyl malonate (5 equiv). Subsequent warming of the reaction mixture to ambient tempera-



Scheme 2. Reagents and conditions: (a) *i*-Pr₂NH, MeLi (including LiBr), **12**, THF, $-78 \degree$ C, 30 min, then **11**, $-78 \degree$ C, 12 h, 70%; (b) H₂, 10% Pd/C (cat.), EtOH, 24 °C, 1.5 h; (c) NaH, DMF, 24 °C, 12.5 h, 59% from **15**; (d) H₂, Lindlar's cat., PhH, 25 °C, 7.5 h, \sim 100%; (e) **10**, EDCI-HCI, DMAP, CH₂Cl₂, 1 h, 74% from **17**; (f) KHMDS (5 equiv), $-78 \degree$ C, 30 min, then TMSCI (5 equiv), 30 min, then diethyl malonate (5 equiv), 30 min, then 24 °C, 1 h; (g) TMSCHN₂, MeOH, 24 °C, 10 min, 70% (**19+20**; **19:20** = 3.4:1) from **18**.

ture produced rearrangement products, which were methylated with TMSCHN2¹⁶ to afford a separable mixture of anti- and synethers having an E-olefin (19 and 20, respectively) in a combined yield of 70% with 3.4:1 anti-selectivity. Diethyl malonate was added as a scavenger of the excess base to avoid elimination of the carbamoyloxy group from the rearrangement products. The moderate anti-selectivity was attributed to the moderate Z-preference in the ketene silyl acetal formation step from 18 under either kinetic or thermodynamic conditions. Since neither the extended time (1 h) for the deprotonation nor the addition of HMPA as a co-solvent affected the diastereoselectivity, the equilibrium thermodynamic ratio of Z/E ketene silvl acetals in this reaction is suggested to be 3.4:1. On the other hand, the kinetic formation of the ketene silvl acetal of 18 by treatment with TMSCl prior to deprotonation with KHMDS showed a somewhat decreased anti-selectivity (2.2:1). Thus, thanks to the thermodynamic equilibration, antiether 19 was obtained in 55% isolated yield and was available for the medium ring formation.

Next, oxane-fused eight-membered ring ethers 13 and 23 were synthesized from 19 and 20, respectively (Scheme 3). The conversion of the 3-Boc-2,2-dimethyl-1,3-oxazoline-4-yl group of 19 to a hydroxymethyl group was first performed. A three-step process, including acidic methanolysis of 19, oxidative cleavage of the resulting 1,2-aminoalcohol, and Luche reduction¹⁷ produced **21** in good yield (85% from 19). Allyl alcohol 21 was transformed to terminal alkene 22 (89%) by a modified Movassaghi's procedure,¹⁸ in which an excess amount of 1-hexene was used as a scavenger of free diimide generated during the reaction. Treatment of 22 with a catalytic amount of second-generation Grubbs' catalyst¹⁹ in refluxing CH_2Cl_2 promoted RCM to furnish **13**²⁰ in high yield (80%). The same five-step process from 20 also provided bicyclic ether 23²⁰ facilely. Thus, a process for the construction of an eight-membered ether ring in five steps after the Ireland-Claisen rearrangement step was established. The *N*,*N*-diisopropylcarbamoyl (Cb) group of **13** could also be removed by stepwise treatment with LiAlH₄ and MeLi to produce **24** in an acceptable vield (57%).

The stereochemistries of **13** and **23** were determined by NMR analysis, in which an NOE correlation H4/H10 and a large J_{H9-H10} (9.9 Hz) were detected in **13**, and NOE correlations H5/H10 and H9/H10 as well as a small J_{H9-H10} (~0 Hz) were observed in **23**,



Scheme 3. Reagents and conditions: (a) 1 M HCl in MeOH, 23 °C, 3 h; (b) NalO₄, 1,4-dioxane–pH 7 buffer (1:1), 25 °C, 1 h; (c) NaBH₄, CeCl₃·7H₂O, MeOH, -78 °C, 10 min, 85% from **19**. (d) IPNBSH, Ph₃P, DEAD, THF–1-hexene (5:1), 0 °C–25 °C, 2 h, then CF₃CH₂OH–H₂O (1:1), 25 °C, 9 h, 89%; (e) (H₂IMes)(PCy₃)Cl₂RuCHPh (cat.), CH₂Cl₂, reflux, 3.5 h, 80%; (f) LiAlH₄, THF, 0 °C–23 °C, 1.5 h; (g) MeLi, THF, 0 °C, 1.5 h, 57% from **13**; (h) TFA–CH₂Cl₂ (1:2), 25 °C, 10 min; (i) NalO₄, 1,4-dioxane–pH 7 buffer (1:1), 25 °C, 2 h; (j) NaBH₄, CeCl₃·7H₂O, MeOH, -78 °C, 10 min, 44% from **20**; (k) IPNBSH, Ph₃P, DEAD, THF–1-hexene (5:1), 0 °C–25 °C, 2 h, then CF₃CH₂OH–H₂O (1:1), 25 °C, 9 h, 45%; (l) (H₂IMes)(PCy₃)Cl₂RuCHPh (cat.), CH₂Cl₂, reflux, 3.5 h, 92%.



Scheme 4. Reagents and conditions: (a) LiAlH₄, THF, -20 °C, 1 h; (b) NaH, BnBr, Bu₄NI, THF, 0 °C→25 °C, 14 h, 87% from **19**; (c) TFA-CH₂Cl₂ (1:2), 25 °C, 10 min; (d) NalO₄, 1,4-dioxane-pH 7 buffer (1:1), 25 °C, 2 h; (e) NaBH₄, CeCl₃ 7H₂O, MeOH, -78 °C, 10 min, 65% from 26; (f) NaH, allyl bromide, Bu₄NI, THF, 0 °C→25 °C, 18 h, 87%; (g) (H₂IMes)(PCy₃)Cl₂RuCHPh (cat.), CH₂Cl₂, reflux, 12 h, 96%.

thereby confirming the configurations of rearrangement products 19 and 20.

Construction of an oxane-fused seven-membered ring ether from 19 was also carried out (Scheme 4). The immediate treatment of 19 with second-generation Grubbs' catalyst¹⁹ did not afford RCM product 25 but gave only recovered starting material. This suggested that steric congestion of the CbO- and the 3-Boc-2,2-dimethyl-1,3-oxazoline-4-yl groups would inhibit the access of any active ruthenium species to the internal olefin. Therefore, a stepwise route from **19** including the removal of the oxazolinyl group and a relay-RCM process,²¹ an effective method for the RCM of sterically hindered substrates, was undertaken for the assembly of seven-membered ring ether 14. After methyl ester 19 was converted to benzyl ether 26 by reduction and subsequent benzylation (87%), the 3-Boc-2,2-dimethyl-1,3-oxazoline-4-yl group of 26 was transformed to an allyl alcohol group by the same three-step process as described above (65%). The resulting alcohol 27 was allylated under Williamson conditions to provide 28 (87%), which was cyclized with second-generation Grubbs' catalyst¹⁹ to furnish 14²⁰ in high yield (96%). Thus, a simple procedure was set up for the seven-membered ring ether synthesis subsequent to the Ireland-Claisen step.

In conclusion, the asymmetric synthesis of an acyclic anti-β-alkoxy ether was achieved by the Ireland-Claisen rearrangement of Z-3-alkoxy-2-propenyl glycolate ester, prepared from Garner's aldehyde, a glycolic acid derivative, and ethynyl N,N-diisopropylcarbamate. The resulting acyclic ether was facilely converted to seven- and eight-membered cyclic ethers via processes involving ring-closing olefin metatheses. Further improvement of the diastereoselectivity of the rearrangement step and applications to the synthesis of natural medium ring ethers are now in progress.

Acknowledgments

We thank Mr. Kenji Watanabe and Dr. Eri Fukushi (GC-MS and NMR Laboratory, Graduate School of Agriculture, Hokkaido University) for the measurements of mass spectra. This work was supported by a Global COE Program (B01: Catalysis as the Basis for Innovation in Materials Science) and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japanese Government.

References and notes

Reviews for natural cyclic ethers, see: (a) Yasumoto, T. Chem. Rec. 2001, 3, 228; (b) Yasumoto, T.; Murata, M. Nat. Prod. Rep. 2000, 17, 293; (c) Scheuer, P. J. Tetrahedron 1994, 50, 3; (d) Shimizu, Y. Chem. Rev. 1993, 93, 1658; (e) Yasumoto, T.; Murata, M. Chem. Rev. 1993, 93, 1895; (f) Erickson, K. L. In Marine Natural Products, Chemical and Biological Perspectives; Scheuer, P. J., Ed.; Academic Press: New York, 1983; Vol. 5, p 131; See also: (g) Blunt, J. W.; Copp, B. R.; Munro, M. H. G.; Northcore, P. T.; Prinsep, M. R. Nat. Prod. Rep. 2001, 18. 1. and Ref. 3a.

- 2 Recent reviews, see: (a) Fujiwara, K.. In Topics in Heterocyclic Chemistry; Gupta, R. R., Kiyota, H., Eds.; Springer: Berlin, 2006; Vol. 5, p 97; (b) Nakamura, I.; Yamamoto, Y. Chem. Rev. 2004, 104, 2127; (c) Elliot, M. C. J. Chem. Soc., Perkin Trans. 1 2002, 2301; (d) Elliot, M. C.; Williams, E. J. Chem. Soc., Perkin Trans. 1 2001, 2303; (e) Yet, L. Chem. Rev. 2000, 100, 2963; (f) Hoberg, J. O. Tetrahedron 1998, 54, 12631; (g) Alverz, E.; Candenas, M.-L.; Perez, R.; Ravelo, J. L.; Martin, J. D. Chem. Rev. 1995, 95, 1953.
- Handbook of Metathesis; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, 2003. 3.
- 4. For early applications to medium-ring ethers, see: (a) Linderman, R. J.; Siedlecki, J.; O'Neil, S. A.; Sum, H. J. Am. Chem. Soc. 1997, 119, 6919; (b) Crimmins, M. T.; Choy, A. L. J. Org. Chem. 1997, 62, 7548.
- For recent applications to medium-ring ethers, see: (a) Marvin, C. C.; Voight, E. A.; Suh, J. M.; Paradise, C. L.; Burke, S. D. J. Org. Chem. 2008, 73, 8452; (b) Oishi, T.; Hasegawa, F.; Torikai, K.; Konoki, K.; Matsumori, N.; Murata, M. Org. Lett. 2008, 10, 3599; (c) Crimmins, M. T.; Ellis, J. M. J. Org. Chem. 2008, 73, 1649; (d) Sato, K.; Sasaki, M. Angew. Chem., Int. Ed. 2007, 46, 2518; (e) Roberts, S. W.; Rainier, J. D. Org. Lett. 2007, 9, 2227; (f) Clark, J. S.; Conroy, J.; Blake, A. J. Org. Lett. 2007, 9, 2091; (g) Inoue, M.; Miyazaki, K.; Ishihara, Y.; Tatami, A.; Ohnuma, Y.; Kawada, Y.; Komano, K.; Yamashita, S.; Lee, N.; Hirama, M. J. Am. Chem. Soc. 2006, 128, 9352; (h) Kadota, I.; Nishii, H.; Ishioka, H.; Takamura, H.; Yamamoto, Y. J. Org. Chem. 2006, 71, 4183; (i) Crimmins, M. T.; McDougall, P. J.; Ellis, J. M. Org. Lett. 2006, 8, 4079; (j) Ortega, N.; Martín, T.; Martín, V. S. Org. Lett. 2006, 8, 871.
- 6. For related stereoselective construction of β-hydroxy ethers for cyclic ether synthesis: (a) Crimmins, M. T.; McDougall, P. J. Org. Lett. 2003, 5, 591; (b) Crimmins, M. T.; She, J. Synlett 2004, 1371; (c) Crimmins, M. T.; McDougall, P. J.; Emmitte, K. A. Org. Lett. 2005, 7, 4033.
- (a) Fujiwara, K.; Goto, A.; Sato, D.; Kawai, H.; Suzuki, T. Tetrahedron Lett. 2005, 46, 3465; (b) Goto, A.; Fujiwara, K.; Kawai, A.; Kawai, H.; Suzuki, T. Org. Lett. 2007, 9, 5373; (c) Sato, D.; Fujiwara, K.; Kawai, H.; Suzuki, T. Tetrahedron Lett. 2008. 49. 1514.
- McFarland, C. M.; McIntosh, M. C. In The Claisen Rearrangement; Hiersemann, M., Nubbemeyer, U., Eds.; Wiley-VCH: Weinheim, 2007; p 117.
- Acid **10** was prepared from known (2R,3S)-2-vinyltetrahydropyran-3-ol by a two-step process [(i) tert-butyl bromoacetate, Bu4NHSO4, 50% NaOH-CH2Cl2 (2:1), 22 °C, 1.5 h, 54%; (ii) TFA-CH₂Cl₂ (1:2), 24 °C, 1.5 h, ~100%]. The synthesis of (2R,3S)-2-vinyltetrahydropyran-3-ol, see: (a) Nicolaou, K. C.; Hwang, C. K.; Marron, B. E.; DeFrees, S. A.; Couladouros, E. A.; Abe, Y.; Carrol, P. J.; Snyder, J. P. J. Am. Chem. Soc. **1990**, 112, 3040–3054; (b) Alvarez, E.; Delgado, M.; Diaz, M. T.; Hanxing, L.; Perez, R.; Martin, J. D. Tetrahedron Lett. 1996, 37, 2865.
- 10. Garner, P.; Park, J. M. Org. Synth. 1991, 70, 18.
- 11. Liang, X.; Andersch, J.; Bols, M. J. Chem. Soc., Perkin Trans. 1 2001, 2136.
- 12. Gralla, G.; Wibbeling, B.; Hoppe, D. Org. Lett. 2002, 4, 2193.
- The presence of LiBr enhanced the yield of **12**. In order to include LiBr in the 13. reaction solution conveniently, acetylene 11 was deprotonated with LDA prepared from *i*-Pr₂NH and a commercially available ether solution of MeLi containing LiBr (Kanto Chemical Co. Ltd).
- The compounds **15**, **17**, and **18** were stable enough to be purified by silica gel 14. column chromatography with 5% Et₃ N containing eluent.
- An excess amount of KHMDS was required to obtain reproducible yields of the 15. products. When 18 was treated with LDA, the terminal vinvlic proton adjacent to the carbamoyloxy group was selectively deprotonated and substituted by a TMS group. LHMDS was unreactive to 18.
- 16. Hashimoto, N.; Aoyama, T.; Shioiri, T. Chem. Pharm. Bull. 1981, 29, 1457.
- 17. Gemal, A. L.; Luche, J.-L. J Am. Chem. Soc. 1981, 103, 5454.
- (a) Movassaghi, M.; Ahmad, O. K. J. Org. Chem. 2007, 72, 1838; (b) Movassaghi, 18. M.; Piizzi, G.; Siegel, D. S.; Piersanti, G. Angew. Chem., Int. Ed. 2006, 45, 5859; (c) Myers, A. G.; Zheng, B.; Movassaghi, M. J. Org. Chem. 1997, 62, 7507.
- Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. Org. Lett. **1999**, *1*, 953.
 Selected spectral data: **13**: [α]_D²⁵ +46 (c 0.035, CHCl₃); IR (neat) ν_{max} 3030, 2965, 2933, 2873, 2853, 1754, 1693, 1439, 1369, 1344, 1286, 1218, 1118, 1094, 1062, (2014) 1025, 769 cm⁻¹; ¹H NMR (400 MHz, C_6D_6) δ 1.07 (6H, d, J = 6.8 Hz), 1.08 (6H, d, J = 6.8 Hz), 1.19–1.38 (3H, m), 1.94–2.01 (1H, m), 2.52 (1H, ddd, J = 2.8, 6.7, 14.1 Hz), 2.67 (1H, dddd, J = 1.1, 3.5, 9.9, 14.1 Hz), 2.82–2.90 (1H, m), 3.12–3.20 (1H, m), 3.27-3.65 (1H, m), 3.40 (3H, s), 3.56-3.62 (1H, m), 3.71 (1H, br ddd, J = 1.7, 5.0, 8.6 Hz), 3.80-4.20 (1H, m), 4.27 (1H, d, J = 9.9 Hz), 5.60-5.69 (1H, m), 5.74 (1H, br td, J = 3.1, 9.9 Hz), 5.92 (1H, br dd, J = 5.0, 11.3 Hz); ¹³C NMR (CH), 81.8 (CH), 125.1 (CH), 136.5 (CH), 154.2 (C), 171.2 (C); HR-EIMS calcd for $C_{19}H_{31}NO_6$ [M⁺]: 369.2151, found: 369.2169. Compound **23**: $[\alpha]_D^{25}$ +139 (*c* 0.050, CHCl₃); IR (neat) ν_{max} 3030, 2957, 2927, 2850, 1760, 1736, 1686, 1440, 1369, 1290, 1210, 1147, 1121, 1075, 1030, 949, 769 cm⁻¹; ¹H NMR (400 MHz, C_6D_6) δ 1.00–1.42 (15H, m), 1.78–1.86 (1H, m), 1.92 (1H, tdd, J = 1.7, 7.3, 14.8 Hz), 2.81 (1H, ddd, J = 5.9, 8.2, 14.8 Hz), 2.92 (1H, br dt, J = 2.3, 12.0 Hz), 3.40-3.65 (4H, m), 3.45 (3H, s), 3.87 (1H, br s), 4.05-4.35 (1H, m), 5.67 (1H, br d, J = 5.7 Hz), 5.69–5.77 (1H, m), 5.89 (1H, ddd, J = 1.7, 7.2, 10.9 Hz); ¹³C NMR (75 MHz, CDCl₃) δ 20.6 (CH₃ × 2), 21.3 (CH₃ × 2), 26.0 (CH₂), 27.3 (CH₂), 31.0 (CH2), 45.3 (CH), 46.6 (CH), 52.4 (CH3), 67.9 (CH2), 74.6 (CH), 74.7 (CH), 76.4

(CH), 78.1 (CH), 126.5 (CH), 132.0 (CH), 154.7 (C), 171.0 (C); HR-EIMS calcd for C₁₉H₃₁NO₆ [M⁺]: 369.2151, found: 369.2163. Compound **14**: $[\alpha]_{D}^{D^5}$ +15.8 (c 0.120, CHCl₃); IR (neat) ν_{max} 3089, 3064, 3031, 2970, 2929, 2867, 1693, 1439, 1369, 1335, 1287, 1263, 1215, 1133, 1097, 1048, 1031, 959, 769, 737, 699 cm⁻¹; ¹H NMR (400 MHz, C₆D₆) δ 0.86–1.26 (13H, m), 1.26–1.43 (2H, m), 1.93–2.00 (1H, m), 2.87–2.95 (1H, m), 3.21–3.29 (1H, m), 3.54–3.90 (4H, m), a3.64 (1H, dd, J = 6.5, 10.5 Hz), 3.73 (1H, dd, J = 2.3, 10.5 Hz), 3.94 (1H, ddd, J = 2.3, 6.5, 9.5 Hz), 4.41 (1H, d, J = 12.3 Hz), 4.50 (1H, d, J = 12.3 Hz), 5.72 (1H, br td, J = 2.5, 5.72 (1H, br td, J = 2.5).

12.8 Hz), 5.84 (1H, br td, J = 2.1, 12.8 Hz), 5.87 (1H, br qd, J = 2.4, 9.5 Hz), 7.05– 7.11 (1H, m), 7.14–7.21 (2H, m), 7.31–7.36 (2H, m); ¹³C NMR (100 MHz, CDCl₃) δ 20.5 (CH₃ × 2), 21.6 (CH₃ × 2), 25.5 (CH₂), 31.0 (CH₂), 45.4 (CH), 46.5 (CH), 67.6 (CH₂), 71.0 (CH₂), 72.2 (CH₂), 73.5 (CH), 80.4 (CH), 81.0 (CH), 82.7 (CH), 127.5 (CH), 127.6 (CH × 2), 128.3 (CH × 2), 130.5 (CH), 132.2 (CH), 138.4 (C), 154.2 (C); HR-EIMS calcd for C24H35N05 [M⁺]: 417.2515, found: 417.2502.

 Hoye, T. R.; Jeffrey, C. S.; Tennakoon, M. A.; Wang, J.; Zhao, H. J. Am. Chem. Soc. 2004, 126, 10210.