

Interlocking the Catalyst: Thread versus Rotaxane-Mediated Enantiodivergent Michael Addition of Ketones to β -Nitrostyrene

Alberto Martinez-Cuezva,^{*,†} Marta Marin-Luna,[§] Diego A. Alonso,[‡] Diego Ros-Ñíguez,[‡] Mateo Alajarin,[†] and Jose Berna^{*,†}


[†]Departamento de Química Orgánica, Facultad de Química, Regional Campus of International Excellence “Campus Mare Nostrum”, Universidad de Murcia, E-30100 Murcia, Spain

[§]Departamento de Química Orgánica, Universidade de Vigo, Campus Lagoas-Marcosende, E-36310 Vigo, Spain

[‡]Departamento Química Orgánica e Instituto de Síntesis Orgánica, Facultad de Ciencias, Universidad de Alicante, E-03080 Alicante, Spain

S Supporting Information

ABSTRACT: Fumaramide threads bearing one L-prolinamide fragment have been designed as templates for promoting the efficient formation of novel Leigh’s [2]rotaxanes. Both threads and rotaxanes are shown to catalyze the asymmetric addition of ketones to β -nitrostyrene in an enantio- and diastereoselective manner. Interestingly, the enantioselective course of these processes is reversed simply by changing from thread to rotaxane as catalyst. DFT computations have allowed to rationalize the stereodivergence shown by the interlocked and noninterlocked catalysts.



One of the most challenging issues in asymmetric catalysis is to obtain the two possible enantiomers of the desired chiral products in a selected transformation.¹ Most of the chiral catalysts or ligands employed to access enantioenriched systems are generally obtained from natural sources as a unique enantiomer.² The synthesis of the alternative enantiomer of the catalyst is not an easy task, usually requiring several steps. To tackle this significant limitation, many researchers have approached to methods of enantio- and/or stereodivergent catalysis.³ Enantiodivergent approaches involve a single enantiomer of the catalyst, which is able to catalyze a process and access both enantiomers of the desired final product just simply changing by the reaction conditions,⁴ adding additives,⁵ or chemically tuning the catalyst structure.⁶ Of particular interest is the recent design of switchable molecular machines enabled to catalyze a chemical transformation in stereodivergent fashions.⁷

In the last decades the synthesis and applications of mechanically interlocked molecules has received increased attention.⁸ Owing to the particular properties that the mechanical bond endows to such systems,⁹ they have become ideal candidates to be applied in different fields, remarking their use as catalysts¹⁰ or ligands in metal-catalyzed processes.¹¹ Switchable rotaxane-based catalysts, able to change their catalytic activity (rate, chemoselective control, activation mode), are available.¹² Also specially selective, rigid and confined¹³ interlocked systems have been designed, by

which the chemo- or the enantioselectivity of a selected process is controlled and even enhanced due to the presence of the mechanical bond.¹⁴

Herein we describe the design and synthesis of a series of chiral prolinamide-based threads incorporating a fumaramide-ester template to allow their assembly into hydrogen-bonded [2]rotaxanes. The threads and the corresponding rotaxanes were employed as organocatalysts in the well-known asymmetric Michael addition of ketones to β -nitrostyrene,¹⁵ aiming to detect remarkable alterations in their reactivity and/or selectivity due to the presence of the mechanical bond. Interestingly, a marked effect in the course of the process was found, enabling to obtain both possible enantiomers in an enantiodivergent approach just by picking either an interlocked or noninterlocked system as catalyst.

Having in mind the elegant reported works describing the utilization of interlocked species as enhanced catalysts or ligands, especially those in which the mechanical bond participates in the stabilization of the transition states,¹⁶ we designed a pair of interlocked prolinamides (Figure 1). These systems incorporate bulky groups at the end of their threads which avoid the disassembly of the mechanically interlocked compound. Both bear a fumaric fragment as a suitable template able to ring around the hydrogen-bonded polyamide macro-

Received: May 22, 2019





Figure 1. Catalyst design: (a) noninterlocked prolinamide as thread; (b) interlocked prolinamide with a polyamide ring.

cycle. The relative position of the macrocycle and the active site of the prolinamide backbone is crucial in this design. The ring should be spatially close to the catalyst active site in order to better control and stabilize the transition states of putative processes by shielding one of the two faces where the substrates can react and creating a more restricted pocket. Importantly, this space should be large enough to allow the substrates interact with the prolinamide, without inhibiting the reaction or the catalyst turnover.

To approach this goal, we synthesized two prolinamide-based rotaxanes differing in the degree of substitution at the amide N atom (secondary or tertiary amide) (Scheme 1).

Scheme 1. Synthesis of Prolinamide-Based Threads 4a,b and Rotaxanes 6a,b^a



^aReaction conditions: (i) EDCI, HOBT, amine (2,2-diphenylethylamine or dibenzylamine), DIPEA, CH₂Cl₂, 25 °C, overnight; (ii) (E)-4-((2,2-diphenylethyl)amino)-4-oxobut-2-enoic acid (**S1**), EDCI, DMAP, CH₂Cl₂, 25 °C, overnight; (iii) TFA, CHCl₃, overnight; (iv) *p*-xylylenediamine, isophthaloyl chloride, Et₃N, CHCl₃, 25 °C, 4 h.

Starting from the commercially available protected *trans*-4-hydroxy-L-proline **1**, the synthesis of the corresponding *N*-Boc-protected threads **3a,b** was accomplished. First, an amidation reaction was carried out employing the suitable amine (2,2-diphenylethylamine for **2a** or dibenzylamine for **2b**) in the presence of the coupling system EDCI/HOBT. Then, an esterification reaction between (E)-4-((2,2-diphenylethylamino)-4-oxobut-2-enoic acid (**S1**) and fragments **2a/2b** provided the corresponding protected threads **3a,b**, which can be deprotected in the presence of TFA to afford the catalytically active threads **4a,b**.

Both threads **3a,b** own an amido-ester function which has been previously utilized as hydrogen-bond acceptor in the synthesis of polyamide-based rotaxanes although with a moderate templating ability.¹⁷ The [2]rotaxanes **5a,b** were obtained in reasonable yields (24–28%) by a five-component reaction with *p*-xylylenediamine and isophthaloyl chloride in the presence of Et₃N (see the Supporting Information).

Finally, Boc deprotection of compounds **5a,b** in the presence of TFA yielded the prolinamide-based rotaxanes **6a,b**, quantitatively. An exhaustive ¹H NMR analysis of thread **4a** and rotaxane **6a** allowed us to acquire a precise idea about the relative position of the macrocycle along the thread (Figure S1). By comparison with reported examples,¹⁸ the displacement of the proton signals of the thread after rotaxanation indicates that the ring is located over the fumaramide binding unit. In this scenario, the active site of the pyrrolidine core remains available to participate in asymmetric reactions catalyzed by the cyclic secondary amine. The photoisomerization of **6a** afforded the maleamide-based system *Z*-**6a** enabling its use as catalyst (see the SI for further details).

At this point, we were intrigued by the behavior of our systems¹⁹ as chiral catalysts in the asymmetric Michael addition of different ketones **7** to β -nitrostyrene **8**. Initially, we assayed dichloromethane as a noncompetitive solvent in which the hydrogen-bonding network between the thread and the macrocycle in the interlocked systems should remain intact. The presence of an acid as additive,²⁰ commonly employed in this enamine-mediated transformations, other solvents, and temperature were also tested (see Tables S1–S4). Thus, we assayed the reaction between acetone **7a** and β -nitrostyrene **8** in the presence of catalytic amounts (10 mol %) of the corresponding threads **4** and rotaxanes **6** (Table 1).

Table 1. Asymmetric Michael Reaction between Acetone 7a and β -Nitrostyrene 8^a

entry	catalyst	conversion ^b (%)	er ^c	config
1	4a	59	69:31	S
2	6a	57	26:74	R
3	4b	100	75:25	S
4	6b	100	29:71	R
5 ^d	4b	100	72:28	S
6 ^d	6b	100	27:73	R
7	5b	0	-	-


^aReaction conditions: nitrostyrene **8** (0.025 mmol), acetone **7a** (0.25 mmol), CH₂Cl₂ (100 μ L), and catalyst (10 mol %) were stirred at 25 °C for 48 h. ^bDetermined by ¹H NMR. ^cDetermined by chiral HPLC. ^dReaction conducted in the presence of 10 mol % of *p*-nitrobenzoic acid.

Thread **4b** and rotaxane **6b** having a tertiary amide attached to the pyrrolidine core notably showed higher activities when compared with **4a** and **6a** (with a secondary amide attached to the pyrrolidine core), obtaining 100% conversion to the Michael adduct **9a** in 2 days (Table 1, compare entries 1, 2 and 3, 4).

The lower conversions obtained with the systems **4a** and **6a** could be due to the formation of a stable imidazolidinone intermediate by reaction of the enamine with the nitrogen of the pendant amide group, dramatically slowing the reaction rates.²¹ The use of *p*-nitrobenzoic acid as additive slightly increases the er when **6b** is employed (Table 1, entry 6). As we expected, the reaction with the Boc-protected **5b** was unproductive, further proving the active participation of the secondary amine as catalytic site in the deprotected systems (Table 1, entry 7).

The results obtained for acetone **7a** were extended to the use of cyclic ketones **7**, which are prochiral and thus lead to the formation of a second stereocenter in the process. The conjugate additions of cyclohexanone **7b** and cyclopentanone **7c** afforded the corresponding *syn* diastereoisomers as main products regardless of the catalyst employed, although in different diastereomeric ratios (Scheme 2). In the case of

Scheme 2. Enantiodivergent Michael Reaction of Ketones 7a–c with β -Nitrostyrene 8^{a–c}



^aReaction conditions: nitrostyrene **8** (0.025 mmol), ketone **7a–c** (0.25 mmol), CH₂Cl₂ (100 μ L), *p*-nitrobenzoic acid (10 mol %), and catalyst **4b** or **6b** (10 mol %) were stirred at 25 $^{\circ}$ C for 48 h (complete conversion). ^bDiastereomeric ratio (*syn/anti*) determined by ¹H NMR. ^cEnantiomeric ratio of the *syn* diastereoisomer (major) determined by chiral HPLC.

adduct **9b**, the diastereoselectivity increased when the interlocked **6b** was the catalyst if compared with the unthreaded **4b** (Scheme 2, **6b**, 10:1 vs **4b**, 4:1 dr). In contrast, the dr was almost not affected by the presence of the mechanical bond when cyclopentanone **7c** was used. More importantly, as initially occurred for adduct **9a**, we were able to obtain the two enantiomerically enriched *syn* diastereoisomers **9b,c** as major byproducts just by rotaxanation of the catalyst. To the best of our knowledge, this is the first time that the presence or not of the mechanical bond at a catalyst backbone is able to control the enantioselective course of a process by accessing both enantiomers in an enantiodivergent fashion.

In order to rationalize the stereodivergence shown by the interlocked and non-interlocked catalysts and its enantioselective outcomes, we performed a DFT study of the reactions using acetone (**7a**) as the carbonyl reactant. For computational efficiency, we used as simplified models the enamine intermediates **4c'** and **6c'** in which methyl groups replace the stoppers of the experimental systems (Figure 2). The transition structures for the *Si*- and *Re*-attack of both thread-enamine **4c'** and rotaxane-enamine **6c'** over the two prochiral faces of (*E*)- β -nitrostyrene (**8**) were computed at the PCM(dichloromethane)/wB97XD/cc-pVDZ theoretical level (Supporting Information). The TS_{4c'-Si8} transition state for the addition of **4c'** to the *Si*-face of the nitrostyrene, leading to adduct (*S*)-**9a** was computed to be 7.0 kJ/mol lower in energy than TS_{4c'-Re8} leading to (*R*)-**9a** (*Re*-attack) (Figure 2a). The




Figure 2. Nucleophilic attack of the enamine of (a) thread **4c'** and (b) rotaxane **6c'** to the (*E*)- β -nitrostyrene (**8**). Energy differences, $\Delta\Delta G^{\ddagger}$ (Si–Re), between the two respective TSs are shown in kJ/mol. Distances and angles are shown in angstroms and degrees, respectively. (c) Definition of d , θ , and Φ .

establishment of a hydrogen bond between the amide NH group and the nitro group (1.99 Å) stabilizes both TS structures, bringing near the electrostatically complementary enamine and nitro N atoms ($d_{N-N} = 3.84$ and 3.87 Å for TS_{4c'-Si8} and TS_{4c'-Re8}, respectively). The gauche orientation between the phenyl group of the nitrostyrene and the double bond of the enamine moiety ($\Phi = -57.3^{\circ}$) in TS_{4c'-Si8} favors an optimal trajectory of the reactants²² with a newly forming C–C bond distance $d = 2.2$ Å and following a Bürgi–Dunitz trajectory of $\theta = 107.5^{\circ}$ very close to the ideal one.²³ In contrast, the antiperiplanar conformation ($\Phi = +2.0^{\circ}$) of TS_{4c'-Re8} showing a new forming C–C bond distance $d = 2.00$ Å counts with an approaching trajectory of the reactants $\theta = 114.4^{\circ}$ far away from the optimal one.

With rotaxane **6c'** as catalyst, the addition over the *Re*-face of nitrostyrene is preferred by 7.3 kJ/mol with respect to the *Si*-attack (Figure 2b). This preference is mainly attributed to the NH \cdots ONO hydrogen bond established in the gauche conformation ($\Phi = +79.4^{\circ}$) of TS_{6c'-Re8}, involving now one NH group of the ring, which preorganizes the nucleophile–electrophile pairing ($d_{N-N} = 3.49$ Å). In this structure, the length of newly forming C–C bond $d = 2.15$ Å and the approaching angle $\theta = 112.1^{\circ}$ are close to the ideal ones. In stark contrast, no hydrogen bonds linking both reactants are present in the computed TS for the *Si*-attack TS_{6c'-Si8}, showing an antiperiplanar orientation between the phenyl and enamine motifs ($\Phi = +175.9^{\circ}$), a $d = 2.17$ Å newly forming C–C bond distance and a $\theta = 115.1^{\circ}$ approaching trajectory.

In conclusion, we have synthesized a series of interlocked and noninterlocked prolinamides and evaluated their catalytic activity in the Michael reaction between different ketones and

β -nitrostyrene. Interestingly, the mechanical bond deeply affects the enantiocontrol of the process allowing an enantiodivergent synthesis of γ -nitroketones. Starting from the same chiral prolinamide backbone, both enantiomers of the final products were accessed just by selecting the interlocked or noninterlocked catalysts. A DFT study shows that with the rotaxanes as catalysts the macrocycle plays a crucial role by establishing a hydrogen bond with the electrophile in the TS, thus favoring the attack of the nucleophile to the Si face of the nitrostyrene. On the contrary, when the nude threads were employed the enantioselective courses of these processes were reversed as result of a different hydrogen bond between both reactants now involving the NH group of the thread. These examples could hopefully help to the future design of new interlocked catalysts by placing anchoring units for reactants in thread and ring, by means of which the mechanical bond could be further exploited in the field of asymmetric synthesis.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.9b01791](https://doi.org/10.1021/acs.orglett.9b01791).

Experimental procedures, NMR spectra, computational details, and HPLC traces (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: amcuezva@um.es.

*E-mail: ppberna@um.es.

ORCID

Alberto Martinez-Cuezva: 0000-0001-8093-7888

Diego A. Alonso: 0000-0002-9069-1188

Mateo Alajarin: 0000-0002-7112-5578

Jose Berna: 0000-0001-7775-3703

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the MINECO (CTQ2017-87231-P and CTQ2015-66624-P) with joint financing by FEDER Funds and Fundación Seneca-CARM (Project 20811/PI/18). A.M.-C. thanks MICINN for a Ramon y Cajal contract and funding (RYC-2017-22700). M.M.L. thanks Xunta de Galicia for her postdoctoral contract (ED418B 2016/166-0).

■ REFERENCES

- (1) Bartók, M. Unexpected inversions in asymmetric reactions: reactions with chiral metal complexes, chiral organocatalysts, and heterogeneous chiral catalysts. *Chem. Rev.* **2010**, *110*, 1663–1705.
- (2) Blaser, H. U. The chiral pool as a source of enantioselective catalysts and auxiliaries. *Chem. Rev.* **1992**, *92*, 935–952.
- (3) (a) Sibi, M. P.; Liu, M. Reversal of stereochemistry in enantioselective transformations. Can they be planned or are they just accidental? *Curr. Org. Chem.* **2001**, *5*, 719–755. (b) Kim, Y. H. Dual enantioselective control in asymmetric synthesis. *Acc. Chem. Res.* **2001**, *34*, 955–962. (c) Escorihuela, J.; Burguete, M. I.; Luis, S. V. New advances in dual stereocontrol for asymmetric reactions. *Chem. Soc. Rev.* **2013**, *42*, 5595–5617. (d) Beletskaya, I. P.; Najera, C.; Yus, M. Stereodivergent catalysis. *Chem. Rev.* **2018**, *118*, 5080–5200.
- (4) Solvent variation: (a) Sohtome, Y.; Tanaka, S.; Takada, K.; Yamaguchi, T.; Nagasawa, K. Solvent-dependent enantiodivergent Mannich-type reaction: utilizing a conformationally flexible guanidine/bisthiourea organocatalyst. *Angew. Chem., Int. Ed.* **2010**, *49*, 9254–9257. Reactant variation: (b) Stymiest, J. L.; Bagutski, V.; French, R. M.; Aggarwal, V. K. Enantiodivergent conversion of chiral secondary alcohols into tertiary alcohols. *Nature* **2008**, *456*, 778–782.
- (5) (a) Blackmond, D. G.; Moran, A.; Hughes, M.; Armstrong, A. Unusual reversal of enantioselectivity in the proline-mediated α -amination of aldehydes induced by tertiary amine additives. *J. Am. Chem. Soc.* **2010**, *132*, 7598–7599. (b) Macharia, J.; Wambua, V.; Hong, Y.; Harris, L.; Hirschi, J. S.; Evans, G. B.; Vetticatt, M. J. A designed approach to enantiodivergent enamine catalysis. *Angew. Chem., Int. Ed.* **2017**, *56*, 8756–8760.
- (6) Soai, K.; Ookawa, A.; Kaba, T.; Ogawa, K. Catalytic asymmetric induction. Highly enantioselective addition of dialkylzincs to aldehydes using chiral pyrrolidinyldimethanols and their metal salts. *J. Am. Chem. Soc.* **1987**, *109*, 7111–7115.
- (7) (a) Wang, J.; Feringa, B. L. Dynamic control of chiral space in a catalytic asymmetric reaction using a molecular motor. *Science* **2011**, *331*, 1429–1432. (b) Vlatković, M.; Bernardi, L.; Otten, E.; Feringa, B. L. Dual stereocontrol over the Henry reaction using a light- and heat-triggered organocatalyst. *Chem. Commun.* **2014**, *50*, 7773–7775. (c) Zhao, D.; Neubauer, T. M.; Feringa, B. L. Dynamic control of chirality in phosphine ligands for enantioselective catalysis. *Nat. Commun.* **2015**, *6*, 6652. (d) Kassem, S.; Lee, A. T. L.; Leigh, D. A.; Marcos, V.; Palmer, L. I.; Pisano, S. Stereodivergent synthesis with a programmable molecular machine. *Nature* **2017**, *549*, 374–378.
- (8) Bruns, C. J.; Stoddart, J. F. *The Nature of the Mechanical Bond: From Molecules to Machines*; Wiley: New York, 2016.
- (9) (a) Neal, E. A.; Goldup, S. M. Chemical consequences of mechanical bonding in catenanes and rotaxanes: isomerism, modification, catalysis and molecular machines for synthesis. *Chem. Commun.* **2014**, *50*, 5128–5142. (b) Lewis, J. E. M.; Galli, M.; Goldup, S. M. Properties and emerging applications of mechanically interlocked ligands. *Chem. Commun.* **2017**, *53*, 298–312. (c) Martinez-Cuezva, A.; Bautista, D.; Alajarin, M.; Berna, J. Enantioselective formation of 2-azetidinones by ring-assisted cyclization of interlocked N-(α -methyl)benzyl fumaramides. *Angew. Chem., Int. Ed.* **2018**, *57*, 6563–6567. (d) Martinez-Cuezva, A.; Lopez-Leonardo, C.; Bautista, D.; Alajarin, M.; Berna, J. Stereocontrolled synthesis of β -lactams within [2]rotaxanes: showcasing the chemical consequences of the mechanical bond. *J. Am. Chem. Soc.* **2016**, *138*, 8726–8729.
- (10) (a) Thordarson, P.; Bijsterveld, E. J. A.; Rowan, A. E.; Nolte, R. J. M. Epoxidation of polybutadiene by a topologically linked catalyst. *Nature* **2003**, *424*, 915–918. (b) Berna, J.; Alajarin, M.; Orenes, R.-A. Azodicarboxamides as template binding motifs for the building of hydrogen-bonded molecular shuttles. *J. Am. Chem. Soc.* **2010**, *132*, 10741–10747. (c) Lewandowski, B.; De Bo, G.; Ward, J. W.; Pappmeyer, M.; Kuschel, S.; Aldegunde, M. J.; Gramlich, P. M. E.; Heckmann, D.; Goldup, S. M.; D'Souza, D. M.; Fernandes, A. E.; Leigh, D. A. Sequence-specific peptide synthesis by an artificial small-molecule machine. *Science* **2013**, *339*, 189–193. (d) Blanco, V.; Leigh, D. A.; Marcos, V. Artificial switchable catalysts. *Chem. Soc. Rev.* **2015**, *44*, 5341–5370.
- (11) (a) Tachibana, Y.; Kihara, N.; Takata, T. Asymmetric benzoin condensation catalyzed by chiral rotaxanes tethering a thiazolium salt moiety via the cooperation of the component: Can rotaxane be an effective reaction field? *J. Am. Chem. Soc.* **2004**, *126*, 3438–3439. (b) Hattori, G.; Hori, T.; Miyake, Y.; Nishibayashi, Y. Design and preparation of a chiral ligand based on a pseudorotaxane skeleton: application to rhodium-catalyzed enantioselective hydrogenation of enamides. *J. Am. Chem. Soc.* **2007**, *129*, 12930–12931. (c) Suzuki, Y.; Shimada, K.; Chihara, E.; Saito, T.; Tsuchido, Y.; Osakada, K. [3]Rotaxane-based dinuclear palladium catalysts for ring-closure Mizoroki–Heck reaction. *Org. Lett.* **2011**, *13*, 3774–3777. (d) Galli, M.; Lewis, J. E. M.; Goldup, S. M. A stimuli-responsive–gold catalyst: regulation of activity and diastereoselectivity. *Angew. Chem., Int. Ed.* **2015**, *54*, 13545–13549. (e) Marcos, V.; Stephens, A.

- J.; Jaramillo-Garcia, J.; Nussbaumer, A. L.; Woltering, S. L.; Valero, A.; Lemonnier, J.-F.; Vitorica-Yrezabal, I. J.; Leigh, D. A. Allosteric initiation and regulation of catalysis with a molecular knot. *Science* **2016**, *352*, 1555–1559.
- (12) (a) Blanco, V.; Carlone, A.; Hänni, K. D.; Leigh, D. A.; Lewandowski, B. A rotaxane-based switchable organocatalyst. *Angew. Chem., Int. Ed.* **2012**, *51*, 5166–5169. (b) Blanco, V.; Leigh, D. A.; Lewandowska, U.; Lewandowski, B.; Marcos, V. Exploring the activation modes of a rotaxane-based switchable organocatalyst. *J. Am. Chem. Soc.* **2014**, *136*, 15775–15780. (c) Martinez-Cuezva, A.; Saura-Sanmartin, A.; Nicolas-Garcia, T.; Navarro, C.; Orenes, R.-A.; Alajarin, M.; Berna, J. Photoswitchable interlocked thiodiglycolamide as a cocatalyst of a chalcogeno-Baylis–Hillman reaction. *Chem. Sci.* **2017**, *8*, 3775–3780. (d) Eichstaedt, K.; Jaramillo-Garcia, J.; Leigh, D. A.; Marcos, V.; Pisano, S.; Singleton, T. A. Switching between anion-binding catalysis and aminocatalysis with a rotaxane dual-function catalyst. *J. Am. Chem. Soc.* **2017**, *139*, 9376–9381. (e) Lee, Y.-J.; Liu, K.-S.; Lai, C.-C.; Liu, Y.-H.; Peng, S.-M.; Cheng, R. P.; Chiu, S.-H. Na⁺ ions induce the pirouetting motion and catalytic activity of [2]rotaxanes. *Chem. - Eur. J.* **2017**, *23*, 9756–9760.
- (13) Examples of highly confined catalysts: (a) Shenoy, S. R.; Crisóstomo, F. R. P.; Iwasawa, T.; Rebek, J. Organocatalysis in a synthetic receptor with an inwardly directed carboxylic acid. *J. Am. Chem. Soc.* **2008**, *130*, 5658–5659. (b) Ćorić, I.; List, B. Asymmetric spiroacetalization catalysed by confined Brønsted acids. *Nature* **2012**, *483*, 315–319.
- (14) (a) Hoekman, S.; Kitching, M. O.; Leigh, D. A.; Pappmeyer, M.; Roke, D. Goldberg active template synthesis of a [2]rotaxane ligand for asymmetric transition-metal catalysis. *J. Am. Chem. Soc.* **2015**, *137*, 7656–7659. (b) Cakmak, Y.; Erbas-Cakmak, S.; Leigh, D. A. Asymmetric catalysis with a mechanically point-chiral rotaxane. *J. Am. Chem. Soc.* **2016**, *138*, 1749–1751. (c) Xu, K.; Nakazono, K.; Takata, T. Design of rotaxane catalyst for O-acylative asymmetric desymmetrization of *meso*-1,2-diol utilizing the cooperative effect of the components. *Chem. Lett.* **2016**, *45*, 1274–1276. (d) Mitra, R.; Zhu, H.; Grimme, S.; Niemeyer, J. Functional mechanically interlocked molecules: asymmetric organocatalysis with a catenated bifunctional Brønsted acid. *Angew. Chem., Int. Ed.* **2017**, *56*, 11456–11459.
- (15) (a) List, B.; Pojarliev, P.; Martin, H. J. Efficient proline-catalyzed Michael additions of unmodified ketones to nitro olefins. *Org. Lett.* **2001**, *3*, 2423–2425. (b) Mase, N.; Watanabe, K.; Yoda, H.; Takabe, K.; Tanaka, F.; Barbas, C. F. Organocatalytic direct Michael reaction of ketones and aldehydes with β -nitrostyrene in brine. *J. Am. Chem. Soc.* **2006**, *128*, 4966–4967. (c) Cao, C.-L.; Ye, M.-C.; Sun, X.-L.; Tang, Y. Pyrrolidine–thiourea as a bifunctional organocatalyst: highly enantioselective Michael addition of cyclohexanone to nitroolefins. *Org. Lett.* **2006**, *8*, 2901–2904. (d) Alonso, D. A.; Baeza, A.; Chinchilla, R.; Gómez, C.; Guillena, G.; Pastor, I. M.; Ramón, D. J. Recent advances in asymmetric organocatalyzed conjugate additions to nitroalkenes. *Molecules* **2017**, *22*, 895–946.
- (16) (a) De Bo, G.; Dolphijn, G.; McTernan, C. T.; Leigh, D. A. [2]Rotaxane formation by transition state stabilization. *J. Am. Chem. Soc.* **2017**, *139*, 8455–8457. (b) Fielden, S. D. P.; Leigh, D. A.; McTernan, C. T.; Pérez-Saavedra, B.; Vitorica-Yrezabal, I. J. Spontaneous assembly of rotaxanes from a primary amine, crown ether and electrophile. *J. Am. Chem. Soc.* **2018**, *140*, 6049–6052.
- (17) (a) Saura-Sanmartin, A.; Martinez-Cuezva, A.; Pastor, A.; Berna, J.; Bautista, D. Light-driven exchange between extended and contracted lasso-like isomers of a bistable [1]rotaxane. *Org. Biomol. Chem.* **2018**, *16*, 6980–6987. (b) Naranjo, T.; Lemishko, K. M.; de Lorenzo, S.; Somoza, A.; Ritort, F.; Pérez, E. M.; Ibarra, B. Dynamics of individual molecular shuttles under mechanical force. *Nat. Commun.* **2018**, *9*, 4512.
- (18) Gatti, F. G.; Leigh, D. A.; Nepogodiev, S. A.; Slawin, A. M. Z.; Teat, S. J.; Wong, J. K. Y. Stiff, and sticky in the right places: the dramatic influence of preorganizing guest binding sites on the hydrogen bond-directed assembly of rotaxanes. *J. Am. Chem. Soc.* **2001**, *123*, 5983–5989.
- (19) Z-6a is unstable when used as catalyst. During the reaction, the formation of the former 6a was observed.
- (20) Hong, L.; Sun, W.; Yang, D.; Li, G.; Wang, R. Additive effects on asymmetric catalysis. *Chem. Rev.* **2016**, *116*, 4006–4123.
- (21) Gryko, D.; Lipiński, R. L-prolinethioamides—efficient organocatalysts for the direct asymmetric aldol reaction. *Adv. Synth. Catal.* **2005**, *347*, 1948–1952.
- (22) Ruiz-Olalla, A.; Retamosa, M. de G.; Cossío, F. P. Densely substituted L-proline esters as catalysts for asymmetric Michael additions of ketones to nitroalkenes. *J. Org. Chem.* **2015**, *80*, 5588–5599.
- (23) Bürgi, H. B.; Dunitz, J. D. From crystal statics to chemical dynamics. *Acc. Chem. Res.* **1983**, *16*, 153–161.