COMMUNICATION

www.rsc.org/obc

B

A pincer auxiliary to force difficult lactamisations †

Hans Bieräugel,^a Hans E. Schoemaker,^b Henk Hiemstra^a and Jan H. van Maarseveen **

^a Institute of Molecular Chemistry, University of Amsterdam, Nieuwe Achtergracht 129,

1018 WS Amsterdam, The Netherlands. E-mail: jvm@science.uva.nl

^b DSM Research, Life Science Products, PO Box 18, 6160 MD Geleen, The Netherlands

Received 7th April 2003, Accepted 7th May 2003 First published as an Advance Article on the web 13th May 2003

A new method to cyclise peptides is reported based on insertion of a salicylaldehyde derived pincer auxiliary in the linear precursor sequence. H- β Ala-Phe-OH and H-Phe- β Ala-OH were chosen as representative model peptides.

Cyclic peptides constitute a versatile compound class.¹ Especially small cyclic peptides are promising in drug,² materials³ and catalysis applications.⁴ However, in general head-to-tail ring-closure of small cyclopeptides (strategy A, Scheme 1) has proven to be difficult.⁵ In addition and in vast contrast to the cyclodipeptides (diketopiperazines), the majority of the (7-membered) monocyclic homodiketopiperazines, made up of an α - and a linear β -amino acid, are difficult to cyclise. Consequently, only a limited number of cyclotetrapeptides⁶ and homodiketopiperazines⁷ have been described. These rings have been closed successfully only due to the presence of turn-inducing residues and the application of highdilution conditions. Also, the correct choice of the coupling site has shown to be crucial. To say it concisely, to date, no generally applicable and thus sequence independent methods exist to effect small peptide cyclisations. The main reason for the cyclisation failure is the predominant trans arrangement of the amide bond(s) in the linear precursor, preventing a correct spatial positioning of the terminal amine and the activated carboxylic group for cyclisation.^{5,8} Unexpectedly, the inclusion of secondary amino acids (e.g. proline or sarcosine) in the linear precursor sequence, resulting in a conformationally unbiased tertiary amide bond, does not necessarily lead to efficient cyclisations.9

Recently, we have developed a novel auxiliary mediated

† Electronic supplementary information (ESI) available: experimental details. See http://www.rsc.org/suppdata/ob/b3/b303836j/

combined tethered-templated strategy to cyclise medium-sized lactams.¹⁰ In our search for new sequence independent methods for small peptide cyclisations we report an improved auxiliary, giving access to lactams containing multiple endocyclic amide bonds. Our approach is based on the use of a pincer auxiliary that facilitates *two* subsequent lactamisations by playing a dual tethering and templating role (strategy **B**, Scheme 1).

In strategy **B**, in the first macrolactamisation step (*i.e.* $3\rightarrow 4$) the auxiliary serves as an internal tether (or hinge) allowing an unconstrained approach of the mutually reactive amino and activated ester groups, followed by the final templated lactamisation by a transannular ring-contracting $O\rightarrow N$ acyl transfer reaction (*i.e.* $4\rightarrow 1$).

Another strategy (strategy C, Scheme 1), which has been applied successfully in the cyclisation of pentapeptides in a pioneering study by Meutermans and coworkers,¹¹ relies on functionalisation of the *N*-terminus of the peptide with a similar auxiliary now serving as an external tether. Activation of the carboxylic acid leads to a macrolactonisation reaction (*i.e.* $5 \rightarrow 4$), still hampered by the predominant *trans* amide bonds in the linear precursor, followed by the same templated transannular ring-contraction to give the final lactam.

In the present communication the power of our novel internal auxiliary based method (strategy **B**, Scheme 1) will be demonstrated as compared to the other strategies (strategies **A** and **C**, Scheme 1). In our study the monocyclic homodiketopiperazine 3-benzyl[1,4]diazepane-2,5-dione (or cyclo(-Phe- β Ala-)) **1a**, composed of phenylalanine and β -alanine, was used as a difficult to lactamise model cyclopeptide (Scheme 2).

Strategy A: That indeed ring-closure to give cyclo(-Phe- β Ala-) **1a** by direct head-to-tail lactamisation of H- β Ala-Phe-OH **2a** or H-Phe- β Ala-OH **2b** (Scheme 3) is difficult is underscored by the finding that no cyclisation could be



Scheme 1 Cyclisation strategies towards bislactams.

DOI: 10.1039/b303836



Scheme 3 Strategy A; direct cyclisation. *Reagents and conditions:* EDC (5 equiv), HOBt (4 equiv), DMF–CH₂Cl₂ (1:4), RT.

accomplished using the most powerful coupling reagents (*i.e.* EDC-HOBt, BOP, TBTU and PfPyU) at high dilution. Lactamisation of the amide *N*-benzylated precursors H- β Ala-BnPhe-OH **2c** and H-Phe-Bn β Ala-OH **2d**, employing a tertiary amide bond under high-dilution conditions, indeed gave the corresponding homodiketopiperazines cyclo(- β Ala-BnPhe-) **1c** and cyclo(-Phe-Bn β Ala-) **1d**, albeit in yields of only 11% and 30%, respectively.

Strategy B: The recent results published by Meutermans and coworkers¹¹ and our preliminary study,¹⁰ prompted us to use salicylaldehyde-derived auxiliaries for the combined tethered–templated approaches towards cyclo(-Phe- β Ala-) **1a**. Salicylaldehyde derived auxiliaries combine the high reactivity of an aryl ester towards aminolysis (enthalpic activation) with a correct spatial positioning of the secondary amine and carbonyl functional groups for the transannular ring-contraction reaction (entropic activation).

Synthesis of the linear cyclisation precursors 3a and 3b to test our internal tethered strategy was accomplished by a reductive amination reaction of H- β Ala-OBn and H-Phe-OBn with 3-*tert*-butoxy-2-hydroxybenzaldehyde 6,¹² followed by Bocprotection of the resulting amine and esterification with Cbz-Phe-OH or Cbz- β Ala-OH, respectively (Scheme 4). After liberation of the carboxylic acid and amine functional groups by catalytic hydrogenolysis under neutral conditions, the tethered lactamisation precursors **3a** and **3b** were isolated as stable compounds.

The introduction of the sterically demanding *tert*-butoxy group at the *ortho* position with respect to the ester group has shown to be crucial because preliminary experiments, starting from salicylaldehyde, suffered from extensive side-reactions due to premature intermolecular aminolysis.¹³

Activation of the carboxylic acids of **3a** and **3b** by treatment with EDC–HOBt effected the tethered cyclisation reactions to give the macrocyclic lactams **4a** and **4b** in yields of 84% and 71%, respectively. Removal of the Boc and *tert*-butyl protective groups by stirring in TFA–dichloromethane was followed by the addition of solid NaHCO₃ to induce the final templated transannular $O \rightarrow N$ acyl transfer reaction furnishing the final homodiketopiperazines **1g** and **1h**, both in yields of 79%.

These results show that our approach, featuring an auxiliary which is inserted in the peptide bond, and serves as a tethering hinge or molecular pincer facilitating the mutually reactive terminal functional groups to approach one and another, is sequence independent. Also, due to the connection of the auxiliary onto the C-terminal amino acid as a tertiary amide (*i.e.* compounds **3**), this residue is shielded from racemisation, which often occurs with difficult peptide cyclisations.⁵

Removal of the remaining 1-(2,3-dihydroxybenzyl) group was accomplished by reduction with sodium in liquid ammonia after methylation of the phenolic-OH groups to give cyclo-(-Phe- β Ala-) 1a in overall yields of 57% (from 1g) and 65% (from 1h) (Scheme 4).

Strategy C: For the synthesis of **1a** via strategy C, the N-termini of H- β Ala-Phe-OBn and H-Phe- β Ala-OBn were reductively alkylated with salicylaldehyde **7**, followed by liberation of the carboxyl group by catalytic hydrogenolysis to furnish **5a** and **5b**, respectively.

The consecutive macrolactonisation and ring-contractive lactamisation reactions were conducted using the optimised conditions as developed by Meutermans and coworkers.¹¹ Activation of **5a** with BOP (1.1 equiv) using DiPEA (2 equiv) as the base in DMF under high dilution (1 mM) only gave complex reaction mixtures. On the other hand, **5b** cyclised smoothly to give homodiketopiperazine **1f** in a yield of 65% (Scheme 5). Although the reason for the cyclisation failure of **5a** is unclear to us,¹⁴ it is obvious that strategy B, featuring an external tethering auxiliary, is strongly sequence dependent.

In conclusion, we have developed a new and powerful auxiliary mediated lactamisation strategy towards small cyclopeptides. This was demonstrated by the successful



Scheme 4 Strategy B; internal tether-template. *Reagents and conditions:* i, H- β Ala-OBn or H-Phe-OBn, Na(OAc)₃BH, THF; ii, (Boc)₂O, CH₂Cl₂; iii, Cbz-Phe-OH or Cbz- β Ala-OH, DCC, DMAP, CH₂Cl₂; iv, H₂, Pd/C, EtOAc-iPrOH = 4 : 1; v, EDC-HOBt (5 equiv); vi, TFA-CH₂Cl₂ = 1 : 1; vii, NaHCO₃, EtOAc; viii, MeI, K₂CO₃, followed by Na, NH₃(I).



Scheme 5 Strategy C; external tether-template. Reagents and conditions: i, H- β Ala-Phe-OBn or H-Phe- β Ala-OBn, Na(OAc)₃BH, CH₂Cl₂; ii, H₂, Pd/C, EtOAc-iPrOH = 4 : 1; iii, BOP (1.1 equiv), DiPEA (2 equiv), DMF (1 mM).

sequence independent ring-closure to a homodiketopiperazine, which could not be prepared efficiently using the currently known lactamisation methods. Currently, our novel internal auxiliary mediated combined tethered-templated approach is expanded towards the synthesis, of to date, inaccessible homodetic and heterodetic cyclotetrapeptides.

Acknowledgements

Dr. Olivier David is acknowledged for the helpful discussions. We thank Mr. A. van den Hoogenband and Mr. K. Stegman (Solvay Pharmaceuticals, Weesp, The Netherlands) for measuring the ESI-MS spectra.

Notes and references

- (a) J. N. Lambert, J. P. Mitchell and K. D. Roberts, J. Chem. Soc., Perkin Trans. 1, 2001, 471–484; (b) H. Kessler, Angew. Chem., Int. Ed. Engl., 1982, 21, 512–523; (c) P. Li, P. P. Roller and J. Xu, Curr. Org. Chem., 2002, 6, 411–440.
- 2 (a) J. Taunton, J. L. Collins and S. L. Schreiber, J. Am. Chem. Soc., 1996, 118, 10412–10422; (b) P. J. Murray, M. Kranz, M. Ladlow,

S. Taylor, F. Berst, A. B. Holmes, K. N. Keavey, A. Jaxa-Chamiec, P. W. Seale, P. Stead, R. J. Upton, S. L. Croft, W. Clegg and M. R. J. Elsegood, *Bioorg. Med. Chem. Lett.*, 2001, **11**, 773–776.

- 3 (a) J. Sanchez-Quesada, M. R. Ghadiri, H. Bayley and O. Braha, J. Am. Chem. Soc., 2000, **122**, 11757–11766; (b) D. T. Bong, T. D. Clark, J. R. Granja and M. R. Ghadiri, Angew. Chem., Int. Ed., 2001, **40**, 988–1011.
- 4 E. R. Jarvo and S. J. Miller, Tetrahedron, 2002, 58, 2481-2495.
- 5 (a) U. Schmidt and J. Langner, J. Peptide Res., 1997, 49, 67–73; (b)
 F. Cavelier-Frontin, G. Pèpe, J. Verducci, D. Siri and R. J. Jacquier, J. Am. Chem. Soc., 1992, 114, 8885–8890.
- 6 M. P. Glenn, M. J. Kelso, J. D. A. Tyndall and D. P. Fairlie, J. Am. Chem. Soc., 2002, **125**, 640–641 and references cited therein.
- 7 (a) O. El Mahdi, J.-P. Lavergne, J. Martinez, P. Viallefont, E. M. Essassi and C. Riche, *Eur. J. Org. Chem.*, 2000, 251–255. In this paper only closure of a configurationally favourable 6,7-trans-difunctionalised homodiketopiperazine succeeded. For solid phase approaches towards monocyclic diketopiperazines see; (b) R. Sarfati, M. Païs and F. X. Jarreau, *Bull. Soc. Chim. Fr.*, 1971, **38**, 255–263; (c) A. Nefzi, J. M. Ostresh and R. A. Houghten, *Tetrahedron Lett.*, 1997, **38**, 4943–4946; (d) V. Krchnák and A. S. Weichsel, *Tetrahedron Lett.*, 1997, **38**, 7299–7302; (e) J. Giovannoni, G. Subra, M. Amblard and J. Martinez, *Tetrahedron Lett.*, 2001, **42**, 5389–5392; (f) J. C. D. Müller-Hartwieg, K. G. Akyel and J. Zimmerman, *J. Peptide Sci.*, 2003, **9**, 187–199.
- 8 B. Calas, R. Michelot, J. P. Leacer, A. Cave, J. Parello and P. Potier, Int. J. Pept. Protein Res., 1987, **29**, 170–183.
- 9 F. Cavelier-Frontin, S. Achmad, J. Verducci, R. Jacquier and G. Pèpe, *THEOCHEM*, 1993, 286, 125–130.
- 10 H. Bieräugel, T. P. Jansen, H. E. Schoemaker, H. Hiemstra and J. H. Van Maarseveen, Org. Lett., 2002, 4, 2673–2674.
- 11 W. D. F. Meutermans, S. W. Golding, G. T. Bourne, L. P. Miranda, M. J. Dooley, P. F. Alewood and M. L. Smythe, *J. Am. Chem. Soc.*, 1999, **121**, 9790–9796.
- 12 The synthesis of the novel auxiliary **6** was easily accomplished in one step from commercially available 2,3-dihydroxybenzaldehyde (see electronic supplementary material).[†]
- 13 The shielding effect of *ortho*-alkoxy groups for aminolysis reactions of aryl esters has been described before. After removal of the *tert*-butyl group, the reactivity of the phenolic ester is enhanced due to the formation of a hydrogen bond between the *ortho*-phenolic OH and the ester carbonyl group. See: (a) G. T. Bourne, S. W. Golding, R. P. McGeary, W. D. F. Meutermans, A. Jones, G. R. Marshall, P. F. Alewood and M. L. Smythe, *J. Org. Chem.*, 2001, **66**, 7706–7713; (b) C. L. Beech, J. F. Coope, G. Fairley, P. S. Gilbert, B. G. Main and K. Plé, *J. Org. Chem.*, 2001, **66**, 2240–2245.
- 14 In pentapeptide cyclisation studies it has been shown that steric effects at the C-terminus play an important role, although not so pronounced as found by us. See: X.-M. Gao, Y.-H. Ye, M. Bernd and B. Kutscher, J. Peptide Sci., 2002, 8, 418–430.