Accepted Manuscript

Utilization of flow chemistry in catalysis: New avenues for the selective synthesis of Bis(indolyl)methanes

Swapna S. Mohapatra, Zoe E. Wilson, Sujit Roy, Steven V. Ley

PII: S0040-4020(17)30158-8

DOI: 10.1016/j.tet.2017.02.026

Reference: TET 28467

To appear in: *Tetrahedron*

- Received Date: 19 December 2016
- Revised Date: 1 February 2017

Accepted Date: 13 February 2017

Please cite this article as: Mohapatra SS, Wilson ZE, Roy S, Ley SV, Utilization of flow chemistry in catalysis: New avenues for the selective synthesis of Bis(indolyl)methanes, *Tetrahedron* (2017), doi: 10.1016/j.tet.2017.02.026.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Graphical Abstract

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.





Tetrahedron journal homepage: www.elsevier.com



Tetrahedror

Utilization of flow chemistry in catalysis: new avenues for the selective synthesis of Bis(indolyl)methanes.

Swapna S. Mohapatra^{*a,b*}, Zoe E. Wilson^{*a*}, Sujit Roy^{*b*} and Steven V. Ley^{*a*}*

^a Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK

^b Organometallics & Catalysis Laboratory, School of Basic Sciences, Indian Institute of Technology, Bhubaneswar 751013, India

ARTICLE INFO

ABSTRACT

Article history: Received Received in revised form Accepted Available online

Keywords: Alkylation Scandium Triflate Bis(indolyl)methanes Flow Catalysis

1. Introduction

Flow chemistry has emerged as an excellent technique for reaction profiling and optimization, and has the benefit of additionally aiding in the scale up of reactions, with its characteristic rapid and controlled mixing and excellent temperature control often leading to higher yields and or selectivity over batch processes.^{1,2} In this work, we report a scandium triflate (Sc(OTf)₃) catalyzed flow chemistry preparation of bis(indolyl) alkanes, which are structural motifs frequently found in natural products, pharmaceuticals, and other functional materials³. Particularly, bis(indolyl) methane (BIM) derivatives possess promising biological effects such as antipyretic, anti-fungal, anti-inflammatory, anti-convulsant. cardiovascular, and selective COX-2 inhibitory activities.^{4,5} In addition, oxidized forms of BIMs are utilized as dyes as well as colorimetric chemosensors.⁵

Owing to their wide spread occurrence, there has been significant interest in the synthesis of BIMs. To date, most methods have employed conventional Lewis acids as well as protic acids as catalysts to promote the electrophilic substitution reaction of indoles with various aldehydes or carbonyl compounds.^{5–8} A variety of organo-^{9–13}, transition metal^{14–20}, ionic liquid^{21–26} and nanomaterial^{27–32} catalysts have also been reported. However, many of these reported methods suffer disadvantages such as requiring high catalyst loading, long reaction times and/or monotonous work-up procedures.^{33–36} The development of a more sustainable alternative protocol for the preparation of BIMs would therefore be desirable. Scandium triflate is an attractive Lewis acid to employ for this purpose as it

Flow chemistry enables the preparation of bis(indolyl)methanes from various indoles and structurally divergent aldehydes using $Sc(OTf)_3$ catalysis. The reaction is regioselective for C-3 functionalization of the indoles, occurring over short reaction times allowing for rapid investigation of scope with straightforward work up facilitating product isolation.

2009 Elsevier Ltd. All rights reserved.

is highly soluble in a range of organic solvents while being stable in the presence of water, unlike the majority of Lewis acids, meaning that it can be recovered and reused if desired.³⁷

Prior to this work there was a single report of the use of $Sc(OTf)_3$ proceeding by an acid catalyzed mechanism^{5,39} to prepare **3a** (Scheme 1).³⁸ It was envisaged that by using flow methodology it would be possible to rapidly optimise and explore the scope of this process leading to decorated tri-substituted methanes with an indole core using Sc(OTf)₃ catalysis.



Scheme 1. Synthesis of BIM 3a in batch by Sato and Sato.³⁸

2. Results and Discussion

Initially, the assembly of BIM 3a from benzaldehyde 1a (0.66 mmol) and indole 2a (1.32 mmol) was repeated in batch by Sato's method to afford the desired product in a comparable 91% yield with 3.5 h reaction time, confirming the robustness of this process.

Next, we transferred this reaction to flow. Initially the flow equipment was set up according to the scheme of Table 1, using

Syrris Asia syringe pumps. Indole 2a and benzaldehyde 1a were taken up in acetonitrile and filled into loop 1. Sc(OTf)₃ (5 mol%) was taken up in acetonitrile and filled into loop 2. The two loops were simultaneously injected into streams of acetonitrile and the plugs met at a T- piece before passing through a 5 mL reactor coil, for a reaction time of 30 min. The reaction stream was pressurised by a 100 psi back pressure regulator (BPR) and the outflow collected. The solvent was removed in vacuo and the crude directly purified by flash chromatography to afford the desired product in 54% yield (entry 1). When the concentration of reactants and catalyst was doubled, the yield was found to improve to 81% (entry 2) but further increasing the concentration led to a drop in yield to 75% (entry 3). Increasing the flow rate (halving the reaction time) at the optimized concentration resulted in a decrease in yield to 72% (entry 4) whereas increasing reaction time to 45 min resulted in a slight increase to 82% (entry 5). At this point the reaction coil was changed to a 20 mL coil to allow a reaction time of 60 min with faster flow rates and resulted in an unchanged yield of 82% (entry 6). Next the solvent was changed to tetrahydrofuran (THF), resulting in the yield of the desired product 3a increasing to 92% (entry 7). As before, when the concentration was lowered the yield of desired product 3a was reduced to 89% (entry 8). When the reaction was performed at 40 °C at the optimal concentration, the yield of the desired product 3a was decreased to 82% (entry 9). Therefore, it was decided to carry out all reactions at room temperature (entry 7), using THF as a solvent.

With optimized flow conditions established, the scope of the reaction was evaluated with a variety of aldehydes (1) and indoles (2) employed as detailed in Table 2. It was seen that 5-bromo indole **2b** (entry 2), and 5-methoxy indole **2c** (entry 3)

reacted with benzaldehyde 1a to form the corresponding products 3b (89%) and 3c (95%) respectively. Similarly, benzaldehyde 1a reacted with 5-nitro indole 2d to form 3d as the product with 83% yield (entry 4). It was found that the reaction of 1a with 4methoxy indole 2e (entry 5), resulted in a modest 50% of 3e. Pleasingly, it was found that N-methyl indole 2f reacted to form the product 3f (88%) with 4-chloro benzaldehyde 1b (entry 6). Additionally, it was found that a wide range of substituted benzaldehydes such as 2,3,4,5,6-pentafluoro benzaldehyde 1c (entry 7), 3-nitro benzaldehyde 1d (entry 8), and 4-chloro benzaldehyde 1e (entry 9) reacted with indole 2a to form the corresponding products 3g, 3h, and 3i in yields ranging from 80-97%. It was even seen that one equivalent of terephthalaldehyde 1f reacted with four equivalents of 2a to form 3j in 65% yield (entry 10). Due to poor solubility, when reacting 10-chloro-9anthraldehyde 1g and 4-methoxy indole 2e in THF the concentration of the solution was lowered to 0.08 M (entry 11), with the corresponding novel BIM 3k obtained in a modest 20% When 5-methoxy indole **2c** reacted yield. with **1h**, 3-chlorobenzaldehyde 4-nitrobenzaldehyde 1i. and mesitaldehyde 1j (entries 12-14), the products formed in 80% (31), 97% (3m), 83% (3n) yield respectively. Finally, the scope was further extended by exchanging the benzaldehyde for a heterocyclic aldehyde. It was found that using furfural 1k (entry 15) and 2-thiophene carboxaldehyde 11 (entry 16) as the aldehyde coupling partners resulted in 78% and 93% of the desired products 30 and 3p respectively. However, when 2-pyridine carboxaldehyde 1m and 3-pyridine carboxaldehyde 1n were used, the reactions did not go to completion in the time frame of the flow reaction, with yields reduced to 30% (3q) and 10% (3r) (entries 17 and 18).

Table 1. Optimization of conditions for the synthesis of bis(indolyl)methane 3a.



entry	Solvent	Flow rate	Reaction	Conc.	Vol.	Temp.	Yield 3a
		(µL/min/pump)	time (min)	(M)	(mL)		(%)
1	CH ₃ CN	83	30	0.33	5	RT	54
2	CH ₃ CN	83	30	0.66	5	RT	81
3	CH ₃ CN	83	30	0.99	5	RT	75
4	CH ₃ CN	166	15	0.66	5	RT	72
5	CH ₃ CN	55	45	0.66	5	RT	82
6	CH ₃ CN	166	60	0.66	20	RT	82
7	THF	166	60	0.66	20	RT	92
8	THF	166	60	0.33	20	RT	89
9	THF	83	60	0.66	10	40 °C	82

ACCEPTED MANUSCRIPT





ACCEPTED MANUSCRIPT



¹ Four equivalents of the indole were used.

4



Scheme 3. Continuous flow synthesis of 3b.

Additionally, as proof of scalability the synthesis of **3b** was run continuously for 5 h 46 min on 28.6 mmol scale (determined by our laboratories supply of indole **2b**) to afford 12.4 g of product (91% yield). For the purposes of this reaction the flow equipment used was changed to acid-resistant Vapourtec R-series pumps. The reaction coil volume was increased to 60 mL to allow the pumps to be run with flow rates of 0.5 mL.min⁻¹ per pump without changing the reaction time, and a 40 psi BPR was used to pressurize the reaction stream (Scheme 3).

3. Conclusions

Flow chemistry has been found to be an effective method for the synthesis of bis(indolyl)methanes from a range of indoles with structurally diverse aldehydes using $Sc(OTf)_3$ as catalyst in both plug and continuous flow modes. It was found that shorter reaction times were needed and only a straightforward work up required. The successful synthesis of a library of 18 BIMs, including novel structures **3e**, **3k**, **3l** and **3n**, provides the opportunity to further explore the bioactivity of this family of compounds.

4. Experimental Details

4.1. General Experimental Details

Reactions were performed using oven-dried glassware (200 °C) under an atmosphere of argon unless otherwise stated. All plug flow reactions were performed using a Syrris Asia flow system⁴⁰ and a Polar Bear Plus cooling/heating unit (Cambridge reactor design)⁴¹. The continuous flow reaction was carried out using a Vapourtec R-series flow machine with acid resistant pumps⁴². Solvents were freshly distilled over calcium hydride and lithium aluminium hydride (tetrahydrofuran) or calcium hydride (acetonitrile, ethyl acetate and 40-60 petroleum ether). All additional reagents and solvents were obtained from commercial sources and used without further purification. Flash column chromatography was performed using high-purity grade silica gel (grade 9385) with 60-120 mesh particle size under air pressure. Analytical thin layer chromatography (TLC) was performed using silica gel 60 F254 pre-coated glass backed plates and visualized by ultraviolet radiation (254 nm).

All NMR spectra except **3d** were recorded on a 600 MHz Avance 600 BBI Spectrometer. ¹H and ¹³C NMR spectra of **3d** were recorded on a Bruker-AC 400 MHz Spectrometer. Unless otherwise stated, all samples were run at room temperature in deuterated solvent, with chemical shift (δ) reported to the nearest 0.01 (¹H)/0.1 (¹³C) ppm, relative to the residual protic solvent; δ (CDCl₃) = 7.26 (¹H)/77.16 (¹³C) ppm, δ (CD₃COCD₃) = 2.05 (¹H)/ 29.84, 206.26 (¹³C) or δ (CD₃SOCD₃) = 2.50 (¹H)/39.52 (¹³C). All ¹³C NMR spectra were run with broadband proton decoupling. Multiplicity of a signal in ¹H NMR is indicated by: s = singlet, d = doublet, t = triplet, m = multiplet, or a combination thereof. Multiplets are reported as the range of ppm values covered by the signals, otherwise the center of the signal is given. Coupling constants, *J*, are quoted in Hz and recorded to the nearest 0.1 Hz. Assignments (detailed in the supporting information) were confirmed using Distortionless Enhanced Polarisation Transfer NMR (DEPT 135) and two dimensional NMR (1H–1H Correlation Spectroscopy (COSY), Heteronuclear Single Quantum Coherence (HSQC) and Heteronuclear Multiple Bond Correlation (HMBC)) experiments gave information used to assign both the ¹H NMR and ¹³C NMR spectra.

Infrared spectra were recorded neat as thin films on a Perkin-Elmer Spectrum One FTIR spectrometer. Absorbances were recorded in the range $4000 \square 650 \text{ cm}^{\square 1}$.

High resolution mass spectrometry (HRMS) was performed using positive/negative electrospray ionisation (ESI+/ESI-), on either a Waters Micromass LCT Premier spectrometer or using a Bruker Bioapex 47e FTICR spectrometer. All m/z values are reported to 4 decimal places and are within \pm 5 ppm of theoretical values.

Melting points were measured on a Stuart Scientific SMP3 melting point apparatus using a gradient of 0.5 °C.min⁻¹. Samples were recrystallized from ethanol and water prior to measuring.

Additional data related to this publication is available at the University of Cambridge Institutional Data Repository (https://doi.org/10.17863/CAM.6695).

4.2. Batch Procedure for the synthesis of 3,3'- (phenylmethylene)bis(1H-indole) (3a)

Benzaldehyde **1a** (70 mg, 0.66 mmol), indole **2a** (154 mg, 1.32 mmol) and ScOTf₃ (8 mg, 0.02 mmol) were taken up in acetonitrile (4 mL) and was stirred at room temperature (20 °C) for 3.5 h. The solvent was removed *in vacuo* and the mixture was purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (4:1) as eluent to afford the *title compound* **3a** (193 mg, 0.60 mmol, 91%) as an off white solid. See below for characterization.

4.3. General Procedure for the synthesis of BIMs in Flow (plug flow)

The flow equipment was set up according to Table 2. Indole 2 (1.65 mmol) and aldehyde 1 (0.82 mmol) were taken up in THF (5 mL) and filled into loop 1. $Sc(OTf)_3$ (5 mol%, 0.041 mmol) was taken up in THF (5 mL) and filled into loop 2. The two loops were simultaneously injected and the plugs met at a T-piece

before passing through a 20 mL reactor coil pressurised by a M 100 psi back pressure regulator and the outflow collected. The solvent was removed *in vacuo* and the mixture purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate as eluent to afford the *title compound* **3**.

4.4. Continuous flow synthesis of 3,3'-(phenylmethylene)bis(5bromo-1H-indole) (3b)

The flow equipment was set up according to Scheme 3. 5-bromoindole **2b** (11.2 g, 57.1 mmol) and benzaldehyde **1a** (2.9 mL, 28.6 mmol) were taken up in THF (173 mL). $Sc(OTf)_3$ (0.81 g, 1.6 mmol) was taken up in THF (200 mL, 0.034 M). The pump for the catalyst solution was started 30 s before the other pump to ensure overlap of streams. The reaction streams met at a T-piece before passing through a 60 mL reactor coil, pressurised by a 40 psi back pressure regulator and the outflow collected, with the lines being flushed out with THF once the solution has all been passed through the system. The solvent was removed *in vacuo* and the mixture purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (4:1, $R_f = 0.32$) as eluent to afford the *title compound* **3b** (12.4 g, 25.9 mmol, 91%) as an off white solid. See below for characterization.

4.5. Characterization of BIMs synthesized in flow.

4.5.1. 3,3'-(phenylmethylene)bis(1H-indole) (3a)

According to the flow general method indole **2a** was coupled to benzaldehyde **1a** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (4:1, $R_f = 0.4$) as eluent to afford the title compound **3a** (243 mg, 0.75 mmol, 92%) as a pink solid, mp = 91.6 – 92.7 °C, lit. mp = 93 – 95 °C.⁴³ **¹H NMR (600 MHz, CDCl_3):** δ 7.63 (s, 2H), 7.47 (d, *J* = 8.0 Hz, 2H), 7.41 (d, *J* = 7.3 Hz, 2H), 7.38 – 7.19 (m, 7H), 7.09 (t, *J* = 8.0 Hz, 2H), 6.55 (s, 2H), 5.95 (s, 1H); ¹³C NMR (**150 MHz, CDCl_3):** δ 144.1, 136.6, 128.7, 128.3, 127.1, 126.2, 123.7, 121.9, 119.9, 119.5, 119.2, 111.2, 40.2; **FTIR (v_{max}, cm⁻¹):** 3405, 3055, 3024, 2833, 1723, 1599, 1492, 1455, 1417, 1336, 1216, 1092, 1040, 1009, 762, 739, 700; **HRMS (ESI):** found 321.1399, [M-H]⁻ C₂₃H₁₇N₂ requires 321.1397. All spectroscopic data was in agreement with that published in the literature.⁴⁴

4.5.2. 3,3'-(phenylmethylene)bis(5-bromo-1Hindole) (3b)

According to the flow general method indole 2b was coupled to benzaldehyde 1a and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (4:1, $R_f = 0.32$) as eluent to afford the title compound **3b** (350 mg, 0.73 mmol, 89%) as an off white solid, mp = 253.4 - 255.0 °C, lit. mp = $246 - 248 \text{ °C}.^{45} \text{ }^{1}\text{H} \text{ NMR}$ (600 MHz, CD₃COCD₃): $\delta 10.27$ (s, 2H), 7.54 (s, 2H), 7.45 – 7.37 (m, 4H), 7.31 (t, J = 7.7 Hz, 2H), 7.22 – 7.19 (m, 3H), 6.89 (s, 2H), 5.95 (s, 1H); ¹³C NMR (150 **MHz**, **CD**₃**COCD**₃): δ 145.0, 136.7, 129.7, 129.3, 129.0, 127.0, 126.2, 124.8, 122.5, 119.2, 114.1, 112.3, 40.5; **FTIR** (v_{max}, cm⁻¹): 3418, 1720, 1597, 1564, 1492, 1443, 1417, 1373, 1241, 1225, 1095, 1040, 977, 881, 793, 745, 705; HRMS (ESI): found 476.9609, $[M-H]^-$ C₂₃H₁₅Br₂N₂ requires 476.9596. All spectroscopic data was in agreement with that published in the literature.

4.5.3. 3,3'-(phenylmethylene)bis(5-methoxy-1Hindole) (3c)

According to the flow general method indole **2c** was coupled to benzaldehyde **1a** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (4:1, $R_f = 0.3$) as eluent to afford the title compound **3c** (297 mg, 0.78 mmol, 95%) as an off white solid, mp = 218.0 – 220.0 °C, lit. mp = 217 – 218 °C.⁴⁶ ¹H NMR (600 MHz, CD₃COCD₃): δ 9.83 (s, 2H), 7.42 (dd, J = 7.3, 1.4 Hz, 2H), 7.29 – 7.17 (m, 4H), 7.18 (d, J = 7.4 Hz, 1H), 6.86 – 6.79 (m, 4H), 6.74 (dd, J = 8.8, 2.4 Hz, 2H), 5.84 (s, 1H), 3.62 (s, 6H); ¹³C NMR (150 MHz, CD₃COCD₃): δ 154.4, 146.0, 133.2, 129.5, 128.8, 128.5, 126.6, 125.3, 119.5, 112.7, 112.0, 102.5, 55.7, 41.2; FTIR (v_{max} , cm⁻¹): 3392, 3317, 3008, 2964, 2936, 1623, 1583, 1484, 1453, 1441, 1292, 1207, 1171, 1126, 1038, 1021, 923, 832, 808, 719; HRMS (ESI): found 383.1737, [M+H]⁺ C₂₅H₂₃N₂O₂ requires 383.1754. All spectroscopic data was in agreement with that published in the literature.⁴⁴

4.5.4. 3,3'-(phenylmethylene)bis(5-nitro-1H-indole) (3d)

According to the flow general method indole 2d was coupled to benzaldehyde **1a** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.46$) as eluent to afford the title compound 3d (280 mg, 0.68) mmol, 83%) as a yellow solid, mp = >300 °C, lit. mp = >300°C⁴⁷. ¹H NMR (400 MHz, CD₃SOCD₃): δ 11.66 (s, 2H), 8.30 (d, J = 2.2 Hz, 2H), 7.97 (dd, J = 9.0, 2.3 Hz, 2H), 7.54 (d, J = 9.0Hz, 2H), 7.39 (d, J = 7.2 Hz, 2H), 7.32 (t, J = 7.6 Hz, 2H), 7.22 (t, J = 7.3 Hz, 1H), 7.13 (d, J = 1.9 Hz, 2H), 6.19 (s, 1H); ¹³C NMR (101 MHz, CD₃SOCD₃): δ 144.1, 140.6, 140.2, 128.9, 128.6, 128.0, 126.8, 126.2, 120.9, 117.0, 116.6, 112.5, 38.9; **FTIR** (**v**_{max}, **cm**⁻¹): 3368, 1623, 1580, 1514, 1470, 1424, 1377, 1324, 1239, 1090, 1040, 893, 817, 739, 701; HRMS (ESI): found 413.1232, $[M+H]^+$ $C_{23}H_{17}N_4O_4$ requires 413.1244. All spectroscopic data was in agreement with that published in the literature.4

4.5.5. 3,3'-(phenylmethylene)bis(4-methoxy-1H-indole) (3e)

According to the flow general method indole **2e** was coupled to benzaldehyde **1a** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.8$) as eluent to afford the title compound **3e** (156 mg, 41 mmol, 50%) as a white solid, mp = 268 – 270 °C (decomposes). **¹H NMR (600 MHz, CD₃COCD₃):** δ 9.82 (s, 2H), 7.32 (d, J =7.5 Hz, 2H), 7.22 (t, J = 7.5 Hz, 2H), 7.10 (t, J = 7.5 Hz, 1H), 6.98 – 6.92 (m, 4H), 6.79 (s, 1H), 6.54 (d, J = 1.7 Hz, 2H), 6.43 – 6.34 (m, 2H), 3.61 (s, 6H); ¹³C NMR (151 MHz, CD₃COCD₃): δ 155.9, 148.3, 139.5, 129.6, 128.2, 125.7, 123.1, 122.7, 122.1, 118.3, 105.5, 100.1, 55.4, 42.1; FTIR (v_{max}, cm⁻¹): 3438, 3398, 3020, 2837, 1611, 1581, 1504, 1462, 1453, 1358, 1259, 1118, 1085, 1038, 766, 728; HRMS (ESI): found 382.1676, [M]⁺ C₂₅H₂₂N₂O₂ requires 382.1673.

4.5.6. 3,3'-((4-chlorophenyl)methylene)bis(1methyl-1H-indole) (3f)

According to the flow general method indole **2f** was coupled to benzaldehyde **1b** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.91$) as eluent to afford the title compound **3f** (277 mg, 0.72 mmol, 88%) as a light pink solid, mp = 208.2 – 209.3 °C, lit. mp = 208 – 209 °C⁴³. ¹H NMR (600 MHz, CD₃COCD₃): δ 7.42 (d, J = 7.7 Hz, 2H), 7.39 – 7.23 (m, 8H), 7.07 (t, J = 7.7 Hz, 2H), 6.58 (s, 2H), 5.92 (s, 1H), 3.73 (s, 6H); ¹³C NMR (150 MHz, CD₃COCD₃): δ 143.1, 137.5, 131.7, 130.1, 128.4, 128.3, 127.3, 121.6, 112.0, 118.8, 117.8, 109.2, 39.6, 32.8; FTIR (v_{max}, cm⁻¹): 3024, 3050, 2998, 2865, 2817, 1731, 1616, 1548, 1472, 1484, 1422, 1370, 1329, 1223, 1199, 1152, 1128, 1085, 1056, 1010, 920, 862, 800, 734; HRMS (ESI): found 383.1311, [M-H]⁻ C₂₅H₂₀N₂CI requires 383.1310. All spectroscopic data was in agreement with that published in the literature.¹⁸

4.5.7. 3,3'-((perfluorophenyl)methylene)bis(1Hindole)(3g)

According to the flow general method indole 2a was coupled to benzaldehyde 1c and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.8$) as eluent to afford the title compound 3g (324 mg, 0.78 mmol, 96%) as an off white solid, mp = 125.4 - 127.0 °C, lit. mp = 129 - 130 °C⁴⁹. ¹H NMR (600 MHz, CD₃COCD₃): δ 10.25 (s, 2H), 7.45 (d, J = 8.2 Hz, 2H), 7.42 (d, J = 8.0 Hz, 2H), 7.18 (d, J = 1.6 Hz, 2H), 7.13 (m, 2H) 7.05 – 6.97 (m, 2H), 6.41 (s, 1H); $^{13}\mathrm{C}$ NMR (150 **MHz**, **CD**₃**COCD**₃): δ 147.0, 145.3, 139.5, 137.8, 127.6, 124.8, 122.4, 119.9, 119.6, 119.3, 115.1, 112.4, 39.6, 32.8; **FTIR** (v_{max}, cm⁻¹): 3407, 3048, 1718, 1651, 1621, 1519, 1497, 1457, 1418, 1338, 1114, 1045, 986, 950, 774, 740; HRMS (ESI): found $C_{23}H_{12}N_2F_5$ requires 411.0915. 411.0911, [M-H]⁻ A11 spectroscopic data was in agreement with that published in the literature.^{49,50}

4.5.8. 3,3'-((3-nitrophenyl)methylene)bis(1H-indole) (3h)

According to the flow general method indole 2a was coupled benzaldehyde 1d and purified by flash column to chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.76$) as eluent to afford the title compound **3h** (289 mg, 0.78 mmol, 96%) as an off white solid, mp = 255.1 - 258.3 °C, lit. mp = $261 - 263 \degree C.^{51}$ ¹H NMR (600 MHz, CDCl₃): δ 8.23 (s, 1H), 8.16 - 8.02 (m, 1H), 7.90 (s, 2H), 7.69 (d, J = 7.7 Hz, 1H), 7.45 -7.33 (m, 5H), 7.23 (t, J = 7.6 Hz, 2H), 7.06 (t, J = 7.6 Hz, 2H), 6.59 (d, J = 2.2 Hz, 2H), 6.00 (s, 1H); ¹³C NMR (150 MHz, **CDCl₃**): δ 148.4, 146.4, 136.7, 135.0, 129.2, 126.6, 123.8, 123.6, 122.3, 121.5, 119.5, 118.2, 111.4, 40.0; **FTIR** (**v**_{max}, **cm**⁻¹): 3406, 3057, 2976, 1723, 1618, 1522, 1488, 1456, 1417, 1345, 1243, 1216, 1124, 1093, 1040, 1010, 927, 832, 793, 739, 728; HRMS (ESI): found 366.1236, [M-H]⁻ C₂₃H₁₆N₃O₂ requires 366.1237. All spectroscopic data was in agreement with that published in the literature.⁵²

4.5.9. 3,3'-((4-chlorophenyl)methylene)bis(1H-indole)(3i)

According to the flow general method indole **1a** was coupled to benzaldehyde **2a** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.78$) as eluent to afford the title compound **3i** (281 mg, 0.78 mmol, 96%) as an off white solid, mp = 102.7 – 104.2 °C, lit. mp = 104 – 106 °C.⁵³ ¹H NMR (600 MHz, CDCl₃): δ 7.73 (s, 2H), 7.44 (d, *J* = 7.5 Hz, 2H), 7.35 (d, *J* = 7.5 Hz, 2H), 7.33 – 7.27 (m, 4H), 7.25 (t, *J* = 7.5 Hz, 2H), 7.09 (t, *J* = 7.5 Hz, 2H), 6.57 (s, 2H), 5.90 (s, 1H); ¹³C NMR (150 MHz, CDCl₃): δ 142.6, 136.7, 131.8, 130.1, 128.4, 126.9, 123.7, 122.1, 119.8, 119.4, 119.1, 111.2, 39.6; FTIR (v_{max}, cm⁻¹): 3408, 3054, 2837, 1722, 1617, 1550, 1487, 1455, 1416, 1336, 1243, 1215, 1088, 1038, 1012, 856, 785, 738; HRMS (ESI): found 355.0998, [M-H]⁻ C₂₃H₁₆N₂Cl requires 355.0997. All spectroscopic data was in agreement with that published in the literature.⁵⁴

4.5.10. 1, 4-bis(di(1H-indol-3-yl)methyl)benzene (3j)

According to the flow general method **4 equivalents** of indole **2a** were coupled to benzaldehyde **1f** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.65$) as eluent to afford the title compound **3j** (302 mg, 0.53 mmol, 65%) as an off white solid, mp = 246.5 – 248.0 °C, lit. mp = 194 – 195 °C⁵⁵. ¹H NMR (600 MHz, CDCl₃): δ 7.40 (d, J = 7.6 Hz, 4H), 7.18 (s, 4H), 7.15 (t, J = 7.6 Hz, 4H), 7.08 – 7.05 (m, 8H), 6.67 (s, 4H), 5.79 (s, 4H), 5.69 (s, 2H); ¹³C NMR (150 MHz, CDCl₃): δ 141.5, 136.5, 128.6, 127.0, 123.8, 121.7, 119.9, 119.2, 118.9, 111.4, 39.9; FTIR (v_{max} cm⁻¹): 3408, 3051, 1615, 1455, 1417, 1337, 1215, 1092, 1010, 738; HRMS (ESI): found 565.2378, [M-H]⁻ C₄₀H₂₉N₄ requires 565.2387. All spectroscopic data was in agreement with that published in the literature.⁵⁶

4.5.11. 3,3'-{(10-chloroanthracen-9-

yl)methylene)bis(4-methoxy-1H-indole) (3k)

According to the flow general method indole **2e** was coupled to benzaldehyde **1g** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.8$) as eluent to afford the title compound **3k** (85 mg, 0.16 mmol, 20%) as an off white solid, mp = 202.8 – 204.3 °C. ¹H **NMR (600 MHz, CDCl₃)**: δ 8.84 (d, J = 8.6 Hz, 2H), 8.56 (d, J = 8.7 Hz, 2H), 8.00 (s, 1H), 7.81 (s, 2H), 7.48 (s, 2H), 7.27 (s, 2H), 7.02 (s, 2H), 6.91 (d, J = 7.9 Hz, 2H), 6.69 (s, 2H), 6.31 (s, 2H), 3.13 (s, 6H); ¹³C **NMR (150 MHz, CDCl₃)**: δ 154.9, 139.1, 138.2, 131.0, 129.0, 127.2, 126.9, 125.7, 124.9, 124.4, 122.5, 117.5, 104.1, 99.7, 54.6, 37.4; **FTIR (v_{max}, cm⁻¹)**: 3409, 3385, 2925, 2833, 1614, 1582, 1504, 1434, 1360, 1323, 1253, 1118, 1083, 928, 770, 740; **HRMS (ESI)**: found 515.1516, [M-H]⁻ C₃₃H₂₄N₂O₂Cl requires 515.1521.

4.5.12. 3,3'-((3-chlorophenyl)methylene)bis(5methoxy-1H-indole)(31)

According to the flow general method indole 2c was coupled benzaldehyde 1h and purified by flash column to chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.7$) as eluent to afford the title compound **31** (273 mg, 0.65 mmol, 80%) as an off white solid, mp = 242.6 - 243 °C. ¹H NMR (600 MHz, CD₃COCD₃): δ 9.90 (s, 2H), 7.43 (s, 1H), 7.38 (d, J = 7.7 Hz, 1H), 7.31 (d, J = 8.8 Hz, 2H), 7.29 (t, J = 17.7 Hz, 10.1 Hz)2H), 7.24 - 7.20 (m, 1H), 6.90 - 6.85 (m, 4H), 6.77 (dd, J = 8.8, 2.4 Hz, 2H), 5.90 (s, 1H), 3.64 (s, 6H); ¹³C NMR (150 MHz, **CD₃COCD₃**): δ 154.4, 148.6, 134.2, 133.2, 130.4, 129.3, 128.2, 128.0, 126.7, 125.3, 118.6, 112.8, 112.1, 102.3, 55.7, 40.8; FTIR (**v**_{max}, **cm**⁻¹): 3393, 2992, 2948, 2825, 1624, 1584, 1484, 1453, 1440, 1287, 1207, 1172, 1127, 1091, 1041, 1025, 926, 832, 812, 767, 737, 704.52; HRMS (ESI): found 415.1203, [M-H]⁻ C₂₅H₂₀N₂O₂Cl requires 415.1208.

4.5.13. Bis(5-methoxyindol-3-yl)(4'nitrophenyl)methane (3m)

According to the flow general method indole 2c was coupled to benzaldehyde 1i and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.67$) as eluent to afford the title compound 3m (340 mg, 0.79 mmol, 97%) as off white solid, mp =138.5 - 140.0 °C, lit. mp = 142 °C⁵⁷. ¹H NMR (600 MHz, CDCl₃): δ 8.12 (d, J = 8.6 Hz, 2H), 8.02 (s, 2H), 7.49 (d, J = 8.6 Hz, 2H), 7.26 (d, J = 8.8 Hz, 2H), 6.89 (dd, J = 8.8, 2.0 Hz, 2H), 6.81 (d, J = 2.0 Hz, 2H), 6.65 (d, J = 1.3 Hz, 2H), 5.90 (s, 1H), 3.74 (s, 6H); ¹³C NMR (150 MHz, CDCl₃): δ 153.9, 151.9, 146.4, 131.9, 129.5, 127.1, 124.6, 123.6, 117.5, 112.12, 112.13, 101.8, 55.9, 40.2; **FTIR** (**v**_{max}, **cm**⁻¹): 3411, 2938, 2830, 1624, 1583, 1513, 1482, 1453, 1438, 1341, 1292, 1206, 1170, 1106, 1040, 925, 834, 795, 749, 721; HRMS (ESI): found 428.1602, $[M+H]^+ C_{25}H_{22}N_3O_4$ requires 428.1605. All spectroscopic data was in agreement with that published in the literature.5

4.5.14. 3,3'-(mesitylmethylene)bis(5-methoxy-1H-indole) (3n)

According to the flow general method indole **2c** was coupled to benzaldehyde **1j** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $\mathbf{R}_{\rm f} = 0.92$) as eluent to afford the title compound **3n** (289 mg, 0.68 mmol, 83%) as an off white solid, mp = 204.0 – 205.0 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.83 (s, 2H), 7.26 (d, J = 8.8 Hz, 2H), 6.88 (dd, J = 8.8, 2.4 Hz, 2H), 6.75 (d, J = 2.4 Hz, 2H), 6.67 (d, J = 1.0 Hz, 2H), 6.16 (s, 1H), 3.72 (s, 6H), 2.32 (s, 3H), 2.18 (br. s, 6H); ¹³C NMR (150 MHz, CDCl₃): δ 153.6, 137.2, 136.6, 135.4, 131.9, 130.1, 127.9, 124.5, 117.4, 111.83, 111.75, 102.0, 55.9, 35.8, 21.4, 20.9; FTIR (v_{max} , cm⁻¹): 3412, 2942, 1722, 1622, 1581,

1482, 1439, 1207, 1171, 1041, 925, 794, 707; **HRMS (ESI):** $M \neq 150$ °C⁶¹, ¹**H NMR (600 MHz, CD₃COCD₃):** δ 10.13 (s, 2H), found 423.2060, [M-H] C₂₈H₂₇N₂O₂ requires 423.2067. 8.71 (s, 1H), 8.43 (s, 1H), 7.72 (d, J = 7.7 Hz, 1H), 7.41 (d, J = 7.7 Hz, 1H), 7.41

4.5.15. 3,3'-(furan-2-ylmethylene)bis(1H-indole) (30)

According to the flow general method indole 2a was coupled benzaldehyde 1k and purified by flash column to chromatography using 40-60 petroleum ether:ethyl acetate (4:1, $R_f = 0.38$) as eluent to afford the title compound **30** (199 mg, 0.63) mmol, 78%) as a brown solid, mp = >300 °C, lit. mp = 322 - 324°C. ¹H NMR (600 MHz, CDCl₃): δ 7.64 (s, 2H), 7.57 (d, J = 8.0Hz, 2H), 7.41 (d, J = 0.9 Hz, 1H), 7.30 - 7.21 (m, 4H), 7.17 -7.11 (m, 2H), 6.73 (d, J = 2.1 Hz, 2H), 6.37 (dd, J = 3.1, 1.9 Hz, 1H), 6.13 (d, J = 3.1 Hz, 1H), 6.00 (s, 1H); ¹³C NMR (150 MHz, **CDCl₃**): δ 157.1, 141.2, 136.4, 126.7, 123.2, 121.9, 119.7, 119.4, 116.9, 111.3, 110.2, 106.7, 34.1; **FTIR** (**v**_{max}, **cm**⁻¹): 3405, 3119, 3053, 1719, 1615, 1589, 1548, 1504, 1455, 1418, 1336, 1240, 1216, 1147, 1092, 1038, 1092, 1008, 927, 781, 738; HRMS (ESI): found 311.1186, [M-H]⁻C₂₁H₁₅N₂O requires 311.1179. All spectroscopic data was in agreement with that published in the literature.

4.5.16. Bis(5-methoxyindol-3-yl)(2-thienyl)methane (3p)

According to the flow general method indole 1a was coupled benzaldehyde 2a and purified by flash column to chromatography using 40-60 petroleum ether:ethyl acetate (4:1, $R_f = 0.26$) as eluent to afford the title compound **3p** (296 mg, 0.76 mmol, 93%) as an off-white solid, mp = 197.8 - 199.3 °C, lit. mp = $180 - 182 \circ C^{58}$. ¹H NMR (600 MHz, CDCl₃): δ 7.86 (s, 2H), 7.27 (d, J = 5.6 Hz, 2H), 7.17 (dd, J = 4.7, 1.6 Hz, 1H), 6.96 -6.92 (m, 2H), 6.90 (d, J = 2.3 Hz, 2H), 6.88 – 6.83 (m, 4H), 6.07 (s, 1H), 3.74 (s, 6H); ¹³C NMR (150 MHz, CDCl₃): δ 153.9, 148.7, 131.8, 127.3, 126.5, 125.2, 124.0, 123.7, 119.4, 112.1, 111.9, 101.9, 56.0, 35.5; **FTIR** (**v**_{max}, **cm**⁻¹): 3389, 3326, 3107, 3059, 3008, 2936, 2829, 1621, 1585, 1484, 1439, 1330, 1298, 1280, 1208, 1171, 1151, 1127, 1100, 1021, 923, 830, 808, 764, 7371; **HRMS (ESI):** found 389.1311, $[M+H]^+ C_{23}H_{21}N_2O_2S$ requires 389.1318. All spectroscopic data was in agreement with that published in the literature.59

4.5.17. 3,3'-(pyridin-2-ylmethylene)bis(1H-indole) (3q)

According to the flow general method indole 2a was coupled to benzaldehyde 1m and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.28$) as eluent to afford the title compound **3q** (79 mg, 0.24) mmol, 30%) as a brown solid, mp = 232.5 - 235.1 °C, lit. mp = 238.0 – 240.0 °C⁴⁵. ¹H NMR (600 MHz, CDCl₃): δ 8.59 (d, J =4.3 Hz, 1H), 8.08 (s, 2H), 7.60 (dd, J = 10.9, 4.4 Hz, 1H), 7.39 (d, J = 7.8 Hz, 2H), 7.35 - 7.34 (m, 3H), 7.18 - 7.14 (m, 3H),7.00 (t, J = 7.8 Hz, 2H), 6.78 (s, 2H), 6.08 (s, 1H); ¹³C NMR (150 MHz, CDCl₃+ CD₃SOCD₃): δ 162.9, 147.7, 135.7, 135.2, 125.7, 122.6, 121.6, 120.0, 119.9, 118.1, 117.3, 116.0, 110.3, 42.0; FTIR (v_{max}, cm⁻¹): 3449, 3144, 2922, 1587, 1567, 1471, 1456, 1435, 1339, 1214, 1090, 1002, 864, 797, 766, 737; HRMS (ESI): found 324.1503, $[M+H]^+ C_{22}H_{18}N_3$ requires 324.1495. All spectroscopic data was in agreement with that published in the literature. 45,60

4.5.18. 3,3'-(pyridin-3-ylmethylene)bis(1H-indole) (3r)

According to the flow general method indole **2a** was coupled to benzaldehyde **1n** and purified by flash column chromatography using 40-60 petroleum ether:ethyl acetate (1:1, $R_f = 0.17$) as eluent to afford the title compound **3r** (26 mg, 0.08 mmol, 10%) as an off white solid, mp = 145.2 - 147.0 °C, lit. mp

Acknowledgements

We gratefully acknowledge funding from the British Council, UK and DST-Inspire, India (Newton-Bhabha Ph.D Placement Programme, SSM) and the Engineering and Physical Sciences Research Council (grants EP/K009494/1 and EP/K039520/1).

References

- 1. Pastre, J. C.; Browne, D. L.; Ley, S. V. *Chem. Soc. Rev.* **2013**, *42* (23), 8849.
- 2. Ley, S. V. Catal. Sci. Technol. 2016, 6 (13), 4676.
- 3. Bandini, M.; Eichholzer, A. Angew. Chem., Int. Ed. 2009, 48 (51), 9608.
- 4. Sujatha, K.; Perumal, P. T.; Muralidharan, D.; Rajendran, M. *Indian J. Chem. B* **2009**, *48*, 267.
- Shiri, M.; Zolfigol, M. A.; Kruger, H. G.; Tanbakouchian, Z. Chem. Rev, 2009, 110 (4), 2250.
- Ekbote, S. S.; Deshmukh, K. M.; Qureshi, Z. S.; Bhanage, B. M. Green Chem. Lett. Rev. 2011, 4 (2), 177.
- 7. Qu, H. E.; Xiao, C.; Wang, N.; Yu, K. H.; Hu, Q. S.; Liu, L. X.
- Molecules 2011, 16 (5), 3855.
- 8. Shirini, F.; Khaligh, N. G.; Jolodar, O. G. *Dyes and Pigments* **2013**, *98* (2), 290.
- Ghorbani-Vaghei, R.; Malaekehpoor, S. M. Org. Prep. Proced. Int. 2010, 42 (2), 175.
- Yang, Q.; Yin, Z. L.; Ouyang, B. L.; Peng, Y. Y. Chinese Chem. Lett. 2011, 22 (5), 515.
- Kumar, A.; Gupta, M. K.; Kumar, M. *Green Chem.* 2012, *14* (2), 290.
 Huo, C.; Sun, C.; Wang, C.; Jia, X.; Chang, W. ACS Sustain. Chem.
- *Eng.* **2013**, *I* (5), 549. 13. Chen, L.; Wang, C.; Zhou, L.; Sun, J. Adv. Synth. Catal. **2014**, 356
- (10), 2224.
- 14. Shaikh, A. C.; Chen, C. J. Chinese Chem. Soc. 2011, 58 (7), 899.
- Yang, Y. L.; Wan, N. N.; Wang, W. P.; Xie, Z. F.; De Wang, J. Chinese Chem. Lett. 2011, 22 (9), 1071.
- Jafarpour, M.; Rezaeifard, A.; Gazkar, S.; Danehchin, M. *Transit. Met. Chem.* 2011, *36* (7), 685.
- Chu, X.-Q.; Zi, Y.; Lu, X.-M.; Wang, S.-Y.; Ji, S.-J. *Tetrahedron* 2014, 70 (2), 232.
- Mohapatra, S. S.; Mukhi, P.; Mohanty, A.; Pal, S.; Sahoo, A. O.; Das, D.; Roy, S. *Tetrahedron Lett.* **2015**, *56* (42), 5709.
- Ebrahimipour, S. Y.; Khabazadeh, H.; Castro, J.; Sheikhshoaie, I.; Crochet, A.; Fromm, K. M. *Inorganica Chim. Acta* 2015, 427, 52.
- Baig, N.; Shelke, G. M.; Kumar, A.; Sah, A. K. Catal. Lett. 2016, 146 (2), 333.
- 21. Zolfigol, M. A.; Khazaei, A.; Moosavi-Zare, A. R.; Zare, A. *Org. Prep. Proced. Int.* **2010**, *42* (1), 95.
- 22. Mulla, S. A. R.; Sudalai, A.; Pathan, M. Y.; Siddique, S. A.; Inamdar, S. M.; Chavan, S. S.; Reddy, R. S. *RSC Adv.* **2012**, *2* (8), 3525.
- 23. Das, P. J.; Das, J. Tetrahedron Lett. 2012, 53 (35), 4718.
- Zare, A.; Bahrami, F.; Merajoddin, M.; Bandari, M.; Moosavi-Zare, A. R.; Zolfigol, M. A.; Hasaninejad, A.; Shekouhy, M.; Beyzavi, M. H.; Khakyzadeh, V. Org. Prep. Proced. Int. 2013, 45 (3), 211.
- 25. Song, Y.; Cheng, C.; Jing, H. Chem. Eur. J. 2014, 20 (40), 12894.
- 26. Goli-Jolodar, O.; Shirini, F. J. Iran. Chem. Soc. 2016, 13 (6), 1077.
- 27. Bhardwaj, M.; Sahi, S.; Mahajan, H.; Paul, S.; Clark, J. H. J. Mol. Catal. A Chem. 2015, 408, 48.
- 28. Rafiee, E.; Khodayari, M. Res. Chem. Intermed. 2016, 42 (4), 3523.
- Sadeghi, B.; Tavasoli, F. A.; Hassanabadi, A. Synth. React. Inorganic, Met. Nano-Metal Chem. 2015, 45 (9), 1396.
- 30. Saffar-Teluri, A. Res. Chem. Intermed. 2014, 40 (3), 1061.

- Shirini, F.; Akbari-Dadamahaleh, S.; Mohammad-Khah, A. Comptes Guo, X.; Pan, S.; Liu, J.; Li, Z. J. Org. Chem. 2009, 74 (22), 8848. 31. Rendus Chim. 2013, 16 (10), 945. 10
- Wang, A.; Liu, X.; Su, Z.; Jing, H. Catal. Sci. Technol. 2014, 4 (1), 71. 32.
- 33. Babu, G.; Sridhar, N.; Perumal, P. T. Synth. Commun. 2000, 30 (9), 1609.
- 34. Chakrabarty, M.; Ghosh, N.; Basak, R.; Harigaya, Y. Tetrahedron Lett. 2002, 43 (22), 4075.
- 35. Kamal, A.; Qureshi, A. A. Tetrahedron 1963, 19 (4), 513.
- 36. Chatterji, A.; Manna, S.; Banerji, J. Perkin Trans. 1980, 1, 533.
- Kobayashi, S. Eur. J. Org. Chem. 1999, 1999 (1), 15. 37.
- 38. Sato, S.; Sato, T. Carbohydr. Res. 2005, 340 (14), 2251.
- Burr, G. O.; Gortner, R. A. J. Am. Chem. Soc. 1924, 46 (5), 1224. 39. 40. Syrris webpage: http://syrris.com/flow products/asia flow
- chemistry (accessed Nov 17, 2016). 41. Cambridge Reactor Design webpage:
- http://www.cambridgereactordesign.com/polarbearplus/polar?bear?plu s.html. (accessed Nov 17, 2016).
- 42. Vapourtec webpage: http://www.vapourtec.com (accessed Nov 17, 2016).
- 43. Li, J.-T.; Sun, M.-X.; He, G.-Y.; Xu, X.-Y. Ultrasonic Sonochemistry 2011, 18 (1), 412.
- 44. Karam, A.; Alonso, J. C.; Gerganova, T. I.; Ferreira, P.; Bion, N.; Barrault, J.; Jérôme, F. Chem. Commun. 2009, No. 45, 7000.
- Liu, S.-T.; Liao, B.-S.; Chen, J.-T.; Liao, Bei-Sih; Chen, Jwu-Ting; 45. Liu, S.-T. Synthesis 2007, 20 (20), 3125.
- 46. Whitney, S.; Grigg, R.; Derrick, A.; Keep, A. Org. Lett. 2007, 9 (17), 3299.
- 47. Wang, L.; Wei, W.; Guo, Y.; Xu, J.; Shao, S. Spectrochim. Acta, Part A 2011, 78 (2), 726.

- Fujii, S.; Kimoto, H.; Nishida, M.; Cohen, L. A. J. Fluorine Chem. 1990, 46 (3), 479.
- 50. Gadekar, S. C.; Reddy, B. K.; Panchal, S. P.; Anand, V. G. Chem Commun. 2016, 52 (24), 4565.
- 51. Heravi, M. M.; Bakhtiari, K.; Fatehi, A.; Bamoharram, F. F. Catal. Commun. 2008, 9 (2), 289.
- 52. Qu, H.-E.; Xiao, C.; Wang, N.; Yu, K.-H.; Hu, Q.-S.; Liu, L.-X. Molecules 2011, 16 (5), 3855.
- 53. Ghorbani-Vaghei, R.; Veisi, H.; Keypour, H.; Dehghani-Firouzabadi, A. A. Mol. Divers. 2010, 14 (1), 87.
- 54. Lin, X.; Cui, S.; Wang, Y. Synth. Commun. 2006, 36 (21), 3153.
- 55 Veisi, H.; Gholbedaghi, R.; Malakootikhah, J.; Sedrpoushan, A.; Maleki, B.; Kordestani, D. J. Heterocycl. Chem. 2010, 47 (6), 1398.
- 56. Nair, V.; Abhilash, K. G.; Vidya, N. Org. Lett. 2005, 7 (26), 5857.
- 57. Chakrabarty, M.; Mukherji, A.; Karmakar, S.; Arima, S.; Harigaya, Y.; Chakrabarty, Manas; Mukherji, Ajanta; Karmakar, Sulakshana; Arima, Shiho; Harigaya, Y. Heterocycles 2006, 68 (2), 331.
- Bharate, S. B.; Bharate, J. B.; Khan, S. I.; Tekwani, B. L.; Jacob, M. 58. R.; Mudududdla, R.; Yadav, R. R.; Singh, B.; Sharma, P. R.; Maity, S.; Singh, B.; Khan, I. A.; Vishwakarma, R. A. Eur. J. Med. Chem. 2013, 63, 435.
- 59. Hikawa, H.; Yokoyama, Y. RSC Adv. 2013, 3 (4), 1061.
- 60. Dhumaskar, K. L.; Tilve, S. G. Green Chem. Lett. Rev. 2012, 5 (3), 353
- 61. Neshat, A. RSC Adv. 2016, 6 (39), 32839.
- 62. Naidu, K. R. M.; Khalivulla, S. I.; Kumar, P. C. R.; Lasekan, O. Org. Commun. 2012, 5 (3), 150.