Journal of Medicinal Chemistry

Article

Subscriber access provided by Purdue University Libraries

Discovery and kinetic profiling of 7-aryl-1,2,4-triazolo[4,3-a]pyridines: positive allosteric modulators of the metabotropic glutamate receptor 2.

Maarten L. J. Doornbos, José María Cid, Jordi Haubrich, Alexandro Nunes, Jasper W. van de Sande, Sophie C. Vermond, Thea Mulder-Krieger, Andrés A. Trabanco, Abdellah Ahnaou, Wilhelmus H. Drinkenburg, Hilde Lavreysen, Laura H. Heitman, Adriaan P. IJzerman, and Gary Tresadern

J. Med. Chem., Just Accepted Manuscript • DOI: 10.1021/acs.jmedchem.7b00669 • Publication Date (Web): 13 Jul 2017 Downloaded from http://pubs.acs.org on July 13, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of Medicinal Chemistry is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Discovery and kinetic profiling of 7-aryl-1,2,4triazolo[4,3-a]pyridines: positive allosteric modulators of the metabotropic glutamate receptor 2.

Maarten L. J. Doornbos¹, José María Cid², Jordi Haubrich¹, Alexandro Nunes¹, Jasper W. van de Sande¹, Sophie C. Vermond¹, Thea Mulder-Krieger¹, Andrés A. Trabanco², Abdellah Ahnaou³, Wilhelmus H Drinkenburg³, Hilde Lavreysen³, Laura H. Heitman¹, Adriaan P. IJzerman^{1*}, Gary Tresadern^{2*}.

¹ Division of Medicinal Chemistry, Leiden Academic Centre for Drug Research (LACDR), Leiden University, P.O. Box 9502, 2300RA Leiden, The Netherlands; ² Janssen Research and Development, Calle Jarama 75A, 45007, Toledo, Spain; ³ Janssen Research and Development, Turnhoutseweg 30, 2340 Beerse, Belgium.

ABSTRACT.

We report the synthesis and biological evaluation of a series of 7-aryl-1,2,4-triazolo[4,3a]pyridines with mGlu₂ positive allosteric modulator (PAM) activity and affinity. Besides traditional *in vitro* parameters of potency and affinity, kinetic parameters k_{on} , k_{off} and residence time (RT) were determined. The PAMs showed various kinetic profiles; k_{on} values ranged over two orders of magnitude, whereas RT values were within a 10-fold range. Association rate constant k_{on} was linearly correlated to affinity. Evaluation of a short, medium and long RT compound in a label-free assay indicated a correlation between RT and functional effect. The effects of long RT compound **9** on sleep-wake states indicated long RT was translated into sustained inhibition of rapid eye movement (REM) *in vivo*. These results show that affinity-only driven selection would have resulted in mGlu₂ PAMs with high values for k_{on} , but not necessarily optimized RT, which is key to predict optimal efficacy *in vivo*.

INTRODUCTION

Glutamate is an important neurotransmitter in the human central nervous system (CNS), where it modulates synaptic responses by activating ionotropic glutamate receptors (iGlu) and metabotropic glutamate receptors (mGlu).¹ The eight known mGlu receptors, which belong to the class C G protein-coupled receptors (GPCRs), have been divided into three subclasses based on sequence homology, signaling pathway and pharmacological profile: group I (mGlu_{1, 5}), group II (mGlu_{2,3}) and group III (mGlu_{4,6,7,8}).² Structurally, class C GPCRs consist of a large extracellular orthosteric binding domain, the so-called venus flytrap (VFT), a cysteine-rich domain connecting the VFT to the seven-transmembrane (7TM) domain.^{2,3} Group II mGlu receptors are coupled to the $G\alpha_{i/0}$ protein and are generally expressed presynaptically in the periphery of the synapse where they inhibit the release of glutamate upon activation.⁴ The activation of mGlu₂ receptors has been shown to be a potential strategy for the treatment of psychiatric diseases involving abnormal glutamatergic neurotransmission, such as schizophrenia and anxiety disorder.^{5,6} Drug discovery efforts include the development of positive allosteric modulators (PAMs) for which subtype-selectivity can be more easily obtained as they bind within the less conserved 7TM domain and have little or no intrinsic activity on their own.^{7,8} The N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3mGlu₂ PAMs vlmethylamine (1, LY487379)⁹, Biphenyl-indanone A (2, BINA)¹⁰, N-(4-((2-(Trifluoromethyl)--Hydroxy-4-(Isobutyryl)phenoxy)methyl)benzyl)-1-Methyl-1*H*-Imidazole-4-Carboxamide (3, THIIC, also known as LY2607540)¹¹ and 3-cyano-1-cyclopropylmethyl-4-(4-phenyl-piperidin-1vl)-pyridine-2(1*H*)-one (4, JNJ-40068782)¹² have been characterized both *in vitro* and *in vivo*. To date, two mGlu₂ PAMs have reached clinical trials. The development of 7-methyl-5-[3-(piperazin-1-ylmethyl)-1,2,4-oxadiazol-5-yl]-2-[[4-(trifluoromethoxy)phenyl]methyl]-3H-

isoindol-1-one (**5**, AZD8529) by AstraZeneca was discontinued after a phase 2 schizophrenia trial in which it did not show clinical efficacy when tested as monotherapy.^{13,14} Compound **5** recently finished a phase II clinical trial to study smoking cessation in female smokers.¹⁵

Meanwhile, the Janssen/Addex mGlu₂ PAM 1-butyl-3-chloro-4-(4-phenyl-1-piperidinyl)-(1*H*)pyridone (**6**, JNJ-40411813/ADX71149) met the primary objectives of safety and tolerability in an exploratory phase IIa study in schizophrenia. Moreover, patients treated with antipsychotics who experience residual negative symptoms were identified as the subgroup that may potentially benefit from add-on treatment with **6**, although this is yet to be established in a formal proof-ofconcept study. In addition, in a second phase IIa study with **6** as adjunctive therapy in patients having major depressive disorder (MDD) with significant anxiety symptoms, **6** did not meet the criterion for efficacy vs placebo. Although **6** showed efficacy signals on several anxiety and depression measures, the primary end point was not met and overall the results did not suggest efficacy for **6** as an adjunctive treatment for patients with MDD with significant anxious symptoms.^{16–22}

To understand more about the key *in vitro* parameters that could drive *in vivo* efficacy, we have focused on the ligand binding kinetics in addition to more classical *in vitro* parameters. It is now becoming increasingly apparent that *in vivo* efficacy can depend on optimized kinetic behavior.²³ Indeed, in the last decade the concept of ligand binding kinetics has received an increasing amount of interest.^{24–27} Both the association and dissociation rate constants k_{on} and k_{off} have been studied for various drug targets including GPCRs.²⁸ The relevance of residence time (RT), defined as the reciprocal of the dissociation rate constant (RT = $1/k_{off}$), for *in vivo* efficacy was retrospectively shown for multiple marketed GPCR drugs, such as the long acting M₃ receptor antagonist tiotropium and fast dissociating D₂ antagonists with lower side-effect

burden.^{29–31} Whereas dissociation rates have been the focus of most kinetic studies, association rates have also been described to be important for fast drug action and high receptor occupancy.^{32–34}

Here, we report the synthesis and biological evaluation of a novel series of 7-aryl-1,2,4triazolo[4,3-a]pyridines, which evolved from a pyridone series (**4** and **6**) through a scaffold hopping drug-design strategy previously described.^{35,36} These compounds were characterized as mGlu₂ PAMs in a functional assay and additionally, radioligand binding experiments were performed to determine their affinity and kinetic binding parameters. We demonstrate that the differentiated binding kinetics result in a varied response in a label-free assay on whole cells. In this way we have been able for the first time to obtain both structure-activity/affinity relationships and structure-kinetic relationships for a class C GPCR.

CHEMISTRY

The target compounds **8–48** were prepared via Suzuki cross coupling of the corresponding 7halotriazolopyridines (**49a,b**; **50a-c** and **51a,b**) with an array of boronic esters (**52a-v**) as shown in Scheme 1. The chemical structures of compounds **8–48** are shown in Tables 3-6).

The key 7-halotriazolopyridines (**49a,b**; **50a-c** and **51a,b**) and non-commercially available boronic esters (**52a-v**) were prepared following the procedures previously described.^{37,38}

BIOLOGY

The potency of the compounds was determined by quantifying the increase in response of a fixed EC₂₀ glutamate concentration (4 μ M) observed in a functional [³⁵S]GTP γ S binding assay. The affinity of all compounds was determined in a full curve radioligand displacement assay in the presence of 6 nM of the mGlu₂ PAM 3-(Cyclopropylmethyl)-7-[(4-phenyl-1piperidinyl)methyl]-8-(trifluoromethyl)-1,2,4-triazolo[4,3-a]pyridine (7, JNJ-46281222), which was radioactively labelled with tritium.³⁹ In order to determine the kinetic parameters k_{on} and k_{off} , we performed a radioligand competition association assay characterized by the equations of Motulsky and Mahan that describe the binding between two ligands competing for the same receptor binding site.⁴⁰ For this purpose, first the association rate constant k_{on} and dissociation rate constant k_{off} of 7 were determined in radioligand association and dissociation experiments using a scintillation proximity assay (SPA).⁴¹ Kinetic parameters were determined at 28°C, presumably these will increase at 37°C. All kinetic assays were performed in the presence of 1 mM glutamate; this condition induced a monophasic association process of the PAM, which is required for the determination of the kinetic parameters of unlabeled PAMs.³⁹ The presence of glutamate was shown to not affect affinity and kinetic parameters of 7.³⁹ Ultimately, three

compounds with representative RT values were selected for comparison in a label-free, impedance based assay (xCELLigence) which measures ligand-induced responses on whole cells. This assay provided further understanding of the influence of RT on the duration of PAM-induced receptor activation.

RESULTS AND DISCUSSION

Kinetic rate constants k_{on} and k_{off} for 7 were similar when obtained in the association and dissociation assays compared to the competition association assay (Table 1), which validated the use of the competition association assay for all other compounds. Kinetic K_D values were comparable to the values reported before.³⁹

Reference mGlu₂ PAMs 2-5 (Table 2) showed moderate to good potency and affinity at the mGlu₂ receptor, in agreement with previous reports.^{39,42,43} Kinetic parameters k_{on} , k_{off} and RT were also determined. Reference compound 2 was shown to have a RT of 11.6 minutes. The other reference PAMs had RTs in the same range, from 7.8 minutes for 5 to 17.3 minutes for 7 (Table 1).

The analysis of the triazolopyridines containing a substituted pyridine (Table 3) began with elaboration of the R¹ substitution. Compound **8** was taken as the starting point. Substitution of chlorine at the R¹ position by trifluoromethyl (**9**) resulted in an increase in potency, while affinity remained the same. Introduction of methyl (**10**) or iso-propyl (**11**) led to an approx. 10-fold decrease in affinity and potency. Interestingly both k_{on} and RT values were similar for **8**, **10** and **11** but 2-fold (k_{on}) and approx. 3-fold (RT) higher for **9**. Substitution of fluorine by chlorine at the R² position resulted in increased potency and affinity for **12**, **13** and **14**. In both cases, the PAMs with CF₃ at R¹ (**9** and **13**) showed modestly longer RTs of 30 and 23 minutes, respectively, along with highest potency and affinity.

Exploration of the pyridine R^3 revealed the importance of the position of the nitrogen. Compound 15 showed decreased potency and affinity compared to 8. A similar trend was seen for 16 that showed values comparable to those reported previously which revealed decreased

 potency and affinity compared to 9.⁴² Values for k_{on} followed this trend whereas RTs remained the same for 15, but decreased from 30 to 11 minutes for 16. Subsequently the alkyl substituents on the pyridine ring were evaluated. Substitution of methyl in 15 by ethyl in 17, led to an increased potency and affinity. Addition of a methyl group on the 5-position of 16, did not change affinity or potency in 18, nor were large changes seen in kinetic parameters. Introduction of the bulky and lipophilic iso-propyl in 19 and 20 increased potency, affinity and k_{on} slightly. Interestingly, the RT of 20 was reduced to only 5 minutes.

Substitution of the cyclopropylmethyl substituent at the R⁴ position by 2,2,2-trifluoroethyl (**21**) resulted in a slight decrease in affinity, similar values for potency and RT, and a 5-fold increase in k_{on} compared to **8**. Next, the ethoxymethyl substituent was evaluated at R⁴ (**22**), which yielded a slight decrease in potency, a 10-fold decrease in k_{on} and similar values for affinity and RT compared to **21**. Substitution of F in **22** by CF₃ at R¹ did not yield a drastic increase in affinity and potency (**23**). Together, **21-23** showed that a cyclopropyl methyl substituent at R⁴ is preferred, as in all three cases the analogue bearing cyclopropyl methyl at R⁴ showed a higher affinity (**8** or **9**).

The structure of substituted pyridine of **8** was changed to the short aliphatic iso- or cyclopropyl substituent, yielding **24-30** (Table 4). Potency and affinity of **24** were relatively low (pEC₅₀ 6.65; pK_i 7.21). Changing the R¹ substituent to CF₃ (**25**) increased affinity and potency again (pEC₅₀ 7.03; pK_i 8.05). Substitution of the fluorine on R² to chlorine (**26**) resulted in a -3-fold increase in affinity and potency.

Substitution of i-Pr at the R^3 position by a c-Pr induced an increase in affinity and potency in all cases (27-29). As observed in Table 3, replacing F at R^2 in 27 by Cl in 30 led to an increased

affinity and potency and a concomitant increase in k_{on} . The data in Table 4 show that k_{on} values followed affinity, whereas RT values were all within a narrow range (8.2 – 11.9 min).

In addition to the results obtained with the first sub-series of triazolopyridines 8-23, a 4methoxy-3-pyridine moiety connected by a methylene spacer was introduced, leading to compounds 31-35 (Table 5). These molecules share a moderate to good potency and affinity (pEC₅₀ 6.8 – 7.8; pK_i 7.9 – 8.7). Removal of the fluorine at the R² position (32) did not significantly change affinity, potency or kinetic parameters when compared to 31. Changing the ether linker in 31 and 32 to an amine in 33 and 34 resulted in an approx. 3-fold increase in affinity and potency. A moderate increase in k_{on} was observed and interestingly a 4- and 2-fold increase in RT for 33 and 34, respectively. This may be explained by the increased flexibility given by the methylene bridge between the triazolopyridine core and the pyridine substituent. As seen in many examples, substitution of CF₃ in 34 to Cl in 35 led to a decrease in affinity, potency and kinetic parameter values.

Reducing aromaticity and introducing more aliphatic and sp3 character can sometimes be beneficial for physicochemical properties and drug-likeness.⁴⁴ In order to evaluate the effects of aromaticity, the 4-methoxy-3-pyridyl substituent was replaced with a 4-tetrahydropyranyl ring (**36**, Table 6). This change resulted in a (pK_i) and potency (pEC₅₀) of 6.65 and 7.64, respectively, which were increased in **37** by the introduction of CF₃ in the R¹ position. Compound **37** has a high affinity (pK_i 8.28) and increased k_{on} (2.0 x 10⁷ M⁻¹min⁻¹), however, contrary to the high affinity, its RT (4.1 min) was the shortest found throughout this study. As seen in the first series of ligands (Table 2), affinity and potency did not change by altering the linker atom from ether to

Journal of Medicinal Chemistry

amine (**38**, **39**). Increasing the linker length by a single methylene unit resulted in an approx. 10fold loss in affinity, (**40**; pK_i 7.21). Values for k_{on} followed affinity, but RT values were all in the same range.

The tetrahydropyran moiety was then replaced by a cyclohexanol ring, leading to 41-48. Again, compounds bearing CF_3 at R^1 (42, 46) showed better affinity and potency than their chlorine analogues (41, 45). Molecule 42 not only showed a high potency (pEC_{50} 8.03), affinity (pK_i 8.66) and k_{on} (1.1 x 10⁷ M⁻¹min⁻¹), but also a longer RT of 31 minutes. Its cis-diastereomer 43 showed a 3-fold reduction in potency (pEC₅₀ 7.38) and affinity (pK₁ 8.03), a 2-fold reduction in k_{on} (5.7 x 10⁶ M⁻¹min⁻¹) and a small reduction in RT (17 min). Introduction of a c-Pr substituent at R^3 (44) led to an increase in potency (pEC₅₀ 8.95), affinity (pK_i 9.05) and k_{on} (1.4 x 10⁷ M⁻¹min⁻¹), and also to a prolonged RT of 50 minutes, which was the highest value observed in this study. It may be that the cyclopropyl at R^3 directs the hydrophilic hydroxyl into a better position for interaction with the receptor. Changing R^4 from c-Pr to CF₃ (45) did not change any parameter compared to 41. Changing the linker group from amine in 43 to ether in 46 did not affect affinity and potency. However, RT seemed to be somewhat increased. Differences between diastereomers were seen again in 47 and 48. Whereas the potencies were similar, affinity was decreased 3-fold and k_{on} remained similar, while RT was prolonged by 2-fold when moving from a *trans* to *cis* configuration.

In general, CF_3 was shown to be the preferred substituent at R^1 . Compounds bearing this substituent typically had a higher potency and affinity. Interestingly, all PAMs with a RT longer than 20 minutes bear this moiety, including **44** with the longest RT seen in this study (50 min). On the other hand, molecule **37** had the shortest RT (4.1 min) and also contains the CF₃ group,

indicating that CF_3 in itself is not the driver for long RT. At the R² position chlorine was shown to be the best substituent for high potency and affinity. PAMs bearing a substituted pyridine revealed the importance of its substitutions in combination with the position of the nitrogen in the ring for potency, affinity and kinetic parameters.

The rather small differences in RT induced by substitutions at the R¹ and R² position at the central triazolopyridine core indicated that this region does not induce large differences in RT. On the other hand, small differences in the distal tail of the compounds induced large differences in RT. As we have shown previously, this triazolopyridine series binds with the scaffold and R⁴ substituent in the deepest part of the receptor, whilst the distal tail points to the extracellular side.³⁹ These differences in RT may be explained by the tight receptor-ligand interactions of the triazolopyridine core compared to looser interactions on the extracellular side. Certain molecules may trap the more flexible ECL2 (or the tops of TMs 3 and 5) in conformations leading to longer RT. Such induced fit modulation of RT can involve ordering of loops to partially block the binding pocket and prevent ligand escape. This mechanism has been seen for multiple targets from different classes, amongst which the 5-HT_{2B} receptor as reported in the recently published LSD-bound crystal structure.^{24,45} Selectivity of this series of mGlu₂ PAMs for other members of the mGlu family was good. Representatives (**12, 27, 37** and **45**) showed no activity at mGlu1, 3, 4, 5, 6, 7 or 8, see Supporting Information table S1.

In order to compare the kinetic parameters and affinity, a kinetic map was generated (Fig. 2). In this map k_{on} (x-axis), k_{off} (y-axis) and K_D (diagonal lines) were plotted. Whereas k_{off} values were all within a small range of about one order of magnitude between 0.020 min⁻¹ (44) and 0.241 min⁻¹ (37), k_{on} values were more diverse and were spread over almost two orders of

magnitude between 1.7 x 10^6 M⁻¹min⁻¹ (40) to 7.7 x 10^7 M⁻¹min⁻¹ (28 and 29). The molecules were plotted per subgroup, as presented in the different tables. This revealed that 8-23 and 24-30 were mostly diverse in on-rates (8-23: 0.17 – 6.2 x 10^7 M⁻¹min⁻¹; 24-30: 0.48 – 7.7 x 10^7 M⁻¹min⁻¹), whereas 31-35 and 36-48 were more diverse in off-rates (31-35: 0.044 – 0.17 min⁻¹; 36-48: 0.020 – 0.24 min⁻¹), indicating that an increased affinity between close analogues may be achieved by optimizing one kinetic parameter at a time. Interestingly, the five reference mGlu₂ PAMs 2-5 and 7, which are very diverse in structure and affinity, differed mostly in terms of onrate (0.21 – 4.6 x 10^6 M⁻¹min⁻¹), whereas off-rates were very close (0.086 – 0.13 min⁻¹). As expected, the compounds with the highest affinity either have a long RT (44) or a fast k_{on} (28-29). Ideally, these optimized parameters may be combined leading to an mGlu₂ PAM with even more increased affinity, which would be represented in the top right corner of the kinetic map. For further comparison, the compounds were arbitrarily divided in short RT (RT < 10 min), medium RT (10 – 20 min) and long RT (> 20 min) (Fig. 2).

To gain more insight in the relationships between the different parameters studied, correlation plots were made (Fig. 3). First, the affinity (pK_i) and potency (pEC₅₀) were shown to be linearly correlated (Fig. 3A). Secondly, the affinity obtained from the classical radioligand displacement assay (pK_i) and affinity obtained from kinetic parameters (pK_D), were shown to be correlated (Fig. 3B). This further supported the use of the kinetic SPA radioligand binding assay. Long RT PAMs were shown to share a high affinity and potency, but by itself a high potency or affinity does not predict a long RT (Fig. 3A, 3B, S1). Then, the dissociation rate constants (pk_{off}) were plotted against the kinetic affinity data (pK_D) (Fig. 3C), which did not show correlation. Interestingly, the association rate constants (log k_{on}) were strongly correlated to affinity (pK_D) (Fig 3D). These parameters were both within a similar range of roughly two orders of magnitude for the compounds of our series. Interestingly, although having different structures and in some cases lower affinities and k_{on} values, the reference mGlu₂ PAMs (**2-5**) also fitted well in this correlation (Fig. 3D). This indicated that this correlation has likely revealed a general trend for PAMs and how they bind at the mGlu₂ receptor.

Correlations between kinetic parameters and affinity have been made before. Most of these studies found a correlation between k_{off} values and affinity or efficacy, for instance on the M₃ and A_{2A} receptors.^{46–48} However, a correlation between association rate constants and affinity, as was shown here for mGlu₂ PAMs, has also been observed before. Exemplary GPCRs are the OX₂ receptor and β_2 -adrenoreceptors,^{49,50} and other targets such as the hERG channel.^{51,52} These correlations are receptor-specific and may therefore be caused by differences in receptor structure, dynamics, or local environment of the receptor.

Recent simulation studies have stipulated the importance of fast association rates for receptor occupancy and drug dosing.^{33,53} One of the mechanisms which may be responsible for these effects is receptor rebinding, which is described as the binding of freshly dissociated ligands from the local environment of the receptor.⁵⁴ These high local concentrations may lead to a prolonged activity of the drug, even when concentrations within the effect compartment have fallen below therapeutic levels.⁵⁵ For the mGlu₂ receptor, rebinding may play a role. The physiological synaptic environment of the receptor may permit more rebinding as its interstitial location will result in less diffusion and therefore higher local ligand concentrations.^{56,57} Another reason for receptor rebinding would be binding of compound to or in the cell membrane close to the receptor.⁵⁶ This would, however, require a correlation between lipophilicity and association

Journal of Medicinal Chemistry

rate constants, as lipophilic compounds are more likely to bind to the cell membrane. Since the plot in figure 3E shows no such correlation for the studied mGlu₂ PAMs, membrane binding is not likely to be involved for these compounds.

xCELLigence

We used a label-free, impedance-based technology (xCELLigence) to evaluate whether a long receptor binding RT leads to a prolonged functional activity. This method is based on the measurement of cellular impedance, expressed as Cell Index, and enables real-time recordings.⁵⁸ Responses induced by compound addition can thus be measured and quantified. Earlier studies have shown that results obtained were comparable to more classical endpoint assays.⁵⁹ As the assay is performed on whole cells at 37°C it is considered a valuable translational step towards in vivo experiments.⁶⁰ Endogenous glutamate levels at time of experiments were around 90 µM. Compound-induced responses therefore represent the PAM effects. As a control, reference compound 7 was used to check the response on CHO-K1 WT cells compared to CHO-K1 hmGlu₂ cells (Fig. S2). No response was found on the WT cells, indicating that the responses were mGlu₂ receptor-mediated. For further experiments we selected compounds with diversity in RT (measured at 28°C): 37 (4.1 min), its close analogue 39 with 3-4 -fold longer RT (15 min) and 44 with again a 3-fold longer RT compared to 39 (50 min; Fig. 4A-D). We determined their potency in the label-free assay using CHO-K1 hmGlu₂ cells and found pEC_{50} values of 7.05 ± 0.04 , 7.73 ± 0.23 and 7.95 ± 0.12 , respectively. Then we performed a wash-out assay, in order to evaluate differences in compound-induced responses after washing. This enabled us to trace the remaining response exerted by only receptor-bound compound. To this end we first determined the EC₈₀ concentrations of short RT **37**, medium RT **39** and long RT **44**,

which were 600, 200 and 60 nM, respectively. Then, cells were stimulated with this EC₈₀ concentration and just before maximal response was reached, after 5 minutes, cells were washed and fresh medium was added (Fig. 4E-G). By comparing these traces with control traces of unwashed wells, we evaluated the differences in responses between **37**, **39** and **44** as the ratio between the AUC of washed and unwashed wells (Fig. 4H). The functional effect of **37** and **39** were partially lost in this experimental set-up, while the effect of **44** remained almost unchanged. The wash-out assay showed a significant difference between short RT **37** and long RT **44**. The value for **39** was in between those of **37** and **44**, in agreement with its RT. Together, these data showed that a longer RT also leads to an increased functional effect under non-equilibrium conditions and that these parameters are positively correlated. Since equilibrium conditions are not always present in physiological conditions, these findings may be considered a translational step from *in vitro* towards *in vivo* experiments.

In vivo sleep-wake effects

Glutamate neurotransmission plays a key role in sleep-wake mechanisms and these processes have translational value for central activity and target engagement. Previous *in vivo* studies have repeatedly shown that positive allosteric modulation of the mGlu₂ receptor results in distinct changes in rodent sleep-wake organization, more specifically in a dose-dependent reduction in so-called Rapid Eye Movement Sleep or REM sleep.^{61,62} Changes in sleep-wake states in the rat are of a dynamic nature, i.e. with more frequent transitions between sleep-wake states as compared to humans: therefore the effects of mGlu₂ PAM compounds on REM sleep are expressed as averages over for example 2 hour or 4 hours periods after compound administration. In order to investigate whether the different RT values are reflected in the sleep-

Journal of Medicinal Chemistry

wake related pharmacodynamic read-out, sleep-wake analysis of **9** and **20** (selected as examples with long versus short *in vitro* RT, respectively, yet having appropriate and similar pharmacokinetic behavior) was even done on a 5 minute basis to optimally capture sleep changes in function of time. All methods, (circadian) timings and analyses otherwise were similar to those reported earlier by Ahnaou et al. (2016).⁶²

Figure 5 shows the effects of **9** and **20** on REM sleep reduction in rats for 24 consecutive 5min periods after a single oral administration of 1, 3, and10 mg/kg and 20% CD vehicle control. The first 6 consecutive 5-minute periods can be discarded due to the confounding arousing effect of the oral administration procedure per se, the animal returning to undisturbed REM sleep baseline values after about 30 minutes. Molecule **9** showed a longer *in vitro* RT of 30 minutes whilst for **20** it was 5 minutes. Correspondingly, at the higher dose of 10 mg/kg, the effect of **9** on REM reduction seemed more immediate and constant compared to **20**. For both compounds no effects on total sleep were seen (Fig. S3).

Compound **9** and **20** showed similar *in vitro* binding affinities with pK_i values of 8.59 and 8.44, respectively. At 10 mg/kg, estimated free levels in brain 1 hour post-administration are 5 and 3.5 ng/g for **9** and **20**, respectively. Hence the two molecules show similar affinity and roughly similar range of free concentrations in brain. While the underlying mechanism causing this different *in vivo* effect is still to be elucidated, this finding provides a first hint that different RT's may impact the *in vivo* effect. The interplay of RT with elimination/clearance from brain is important for the resulting *in vivo* effect, but with similar levels of **9** and **20** after 1 hour, it seems that RT may play a role in this case. To strengthen these observations, future experiments could

include the synthesis of PET ligands with different residence times and further *in vivo* studies with a different endpoint such as cognition.

While potent GPCR antagonists can act via a long RT with expected improved clinical efficacy there is limited published data assessing the relationship between agonist/PAM RT and *in vivo* efficacy. Agonist responses are usually regulated by receptor desensitization and internalization, which can act to limit the effect and duration of receptor signaling. PAMs do not activate receptors in the absence of agonist, and hence allosteric modulators with prolonged action may not be negated by internalization and loss of function due to longer RT of the PAM-receptor complex.

CONCLUSIONS

In this study we described a series of 7-aryl-1,2,4-triazolo[4,3-a]pyridines with potent mGlu₂ PAM activity and a high affinity. An SPA radioligand binding assay was developed to study the kinetic parameters of these compounds. We showed that association rate constants range within three orders of magnitude, whereas RT values are more confined, from 4.1 minutes (**37**) up to 50 minutes (**44**). Correlation plots between affinity and kinetic parameters revealed a strong correlation between affinity and association rate constants, whereas no such correlation was found between RT and affinity.

Structure-kinetics relationships were explored in addition to the more traditional structureactivity/affinity relationships analysis. We learned that a CF₃ substituent at R^1 increased potency and affinity and was also important for RT, but did not lead to a long RT *per se* as shown by its presence in **37** ('short' RT) and **44** ('long' RT). Chlorine was the preferred substituent for R^2 , it increased potency and affinity but not RT. Differences in the distal tail of the molecules, such as small changes in the pyridine moiety or single substitutions to the cyclohexanol, induced the biggest differences in RT. This was exemplified by the introduction of c-Pr in **44**, leading to a 3fold increase in RT compared to **43**.

To evaluate whether a long RT also leads to a prolonged functional effect, **37**, **39** and **44** were evaluated in a wash-out assay using the label-free impedance based xCELLigence technique. The response of **44** on whole cells at physiological temperature remained more sustained after washing than that of **37** and **39**. These data suggested that an increased lifetime of the receptor-ligand complex (RT) is correlated to an increased functional effect under non-equilibrium conditions.

Together, these results show for the first time for a class C GPCR that optimization of ligand binding kinetics in addition to potency and affinity is possible. Given that pure PAMs do not exert an effect in the absence of glutamate and are thus less likely to induce on-target toxicity, a long RT seems a promising strategy for these ligands. As rebinding may be involved in mGlu₂ receptor occupancy, a RT for **44** in our system of 50 minutes may be prolonged in the interstitial synaptic environment. Without any on-target toxicity occurring, the design of mGlu₂ PAMs with even longer RTs may be a logical next step. Lastly, a first attempt was made to relate the compound's RT to its *in vivo* pharmacodynamic effect. Ultimately, this study may contribute to the development of compounds with a high affinity and efficacy *in vitro* and *in vivo*, not only for the mGlu₂ receptor but also for other GPCRs.

EXPERIMENTAL SECTION

Chemistry

Unless otherwise noted, all reagents and solvents were obtained from commercial suppliers and used without further purification. Thin layer chromatography (TLC) was carried out on silica gel 60 F254 plates (Merck). Flash column chromatography was performed on silica gel, particle size 60 Å, mesh of 230-400 (Merck) under standard techniques. Microwave assisted reactions were performed in a single-mode reactor, Biotage Initiator Sixty microwave reactor (Biotage), or in a multimode reactor, MicroSYNTH Labstation (Milestone, Inc.). Nuclear magnetic resonance (NMR) spectra were recorded with either a Bruker DPX-400 or a Bruker AV-500 spectrometer (Bruker AG) with standard pulse sequences, operating at 400 and 500 MHz, respectively, using CDCl₃ and DMSO-d₆ as solvents. Chemical shifts (δ) are reported in parts per million (ppm) downfield from tetramethylsilane ($\delta = 0$). Coupling constants are reported in hertz. Splitting patterns are defined by s (singlet), d (doublet), dd (double doublet), t (triplet), q (quartet), quin (quintet), sex (sextet), sep (septet), or m (multiplet). Liquid chromatography combined with mass spectrometry (LCMS) was performed on either a HP 1100 HPLC system (Agilent Technologies) or Advanced Chromatography Technologies system composed of a quaternary or binary pump with degasser, an autosampler, a column oven, a diode array detector (DAD), and a column as specified in the respective methods below. Flow from the column was split to a MS spectrometer. The MS detector was configured with either an electrospray ionization source or an ESCI dual ionization source (electrospray combined with atmospheric pressure chemical ionization). Nitrogen was used as the nebulizer gas. Data acquisition was performed with MassLynx-Openlynx software or with Chemsation-Agilent data browser software. Melting point values are peak values and were obtained with experimental uncertainties that are commonly associated with this analytical method. Melting points were determined in open capillary tubes on a Mettler FP62 apparatus with a temperature gradient of 10 °C/min. Maximum temperature was 300 °C.

Purities of all new compounds were determined by analytical RP HPLC using the area percentage method on the UV trace recorded at a wavelength of 254 nm, and compounds were found to have \geq 95% purity unless otherwise specified.

8-Chloro-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (8). To a stirred suspension of **50a** (1.7 g, 5.09 mmol) and **52a** (2.1 g, 6.12 mmol) in a saturated aqueous solution of NaHCO₃ (18 mL) and 1,4-dioxane (36 mL) was added Pd(PPh₃)₄ (0.589 g, 0.51 mmol). The mixture was heated at 150 °C for 10 min under microwave irradiation. The mixture was cooled to room temperature and filtered through a Celite pad. The filtrate was diluted with water (20 mL) and extracted with EtOAc (2 × 15 mL). The organic layer was washed with brine (15 mL), dried over anhydrous Na₂SO₄, and concentrated in vacuo. The crude was purified by flash column chromatography (silica gel, EtOAc in DCM, 0/100 to 20/80) to give the desired product **8** as a white solid (1.3 g, 60%). Mp 207.2 °C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.32-0.42 (m, 2H), 0.61-0.69 (m, 2H), 1.17-1.28 (m, 1H), 2.54 (s, 3H), 2.55 (s, 3H), 3.13 (d, *J* = 6.9 Hz, 2H), 6.87 (d, *J* = 6.9 Hz, 1H), 6.92 (t, *J* = 8.4 Hz, 1H), 7.02 (d, *J* = 8.4 Hz, 1H), 7.16 (d, *J* = 8.4 Hz, 1H), 7.25 (d, *J* = 9.2 Hz, 1H), 7.41 (dd, *J* = 11.3, 1.7 Hz, 1H), 7.98 (d, *J* = 6.9 Hz, 1H). LC–MS *m/z* 423 [M + H]⁺, t_R = 2.86 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl [1,2,4] triazolo [4,3-a]-pyridine (9). Starting from 49a (0.150 g, 0.54 mmol) and 52a (0.242 g, 0.707 mmol) and following the procedure described for 8, compound 9 was

obtained as a white solid (0.139 g, 56%). ¹H NMR (400 MHz, CDCl₃) δ ppm 0.31-0.43 (m, 2H), 0.61-0.70 (m, 2H), 1.16-1.30 (m, 1H), 2.53 (s, 3H), 2.55 (s, 3H), 3.15 (d, *J* = 6.7 Hz, 2H), 6.79 (d, *J* = 7.2 Hz, 1H), 6.89 (t, *J* = 8.3 Hz, 1H), 7.01 (d, *J* = 8.3 Hz, 1H), 7.05 (br d, J = 8.6 Hz, 1H), 7.14 (d, *J* = 8.3 Hz, 1H), 7.22 (dd, *J* = 10.9, 2.1 Hz, 1H), 8.11 (d, *J* = 7.2 Hz, 1H). LC–MS *m/z* 457 [M + H]⁺, t_R = 3.03 min.

8-Methyl-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl[**1,2,4**]**triazolo**[**4,3-a**]**-pyridine (10).** Starting from **51a** (0.250 g, 1.127 mmol) and **52a** (0.464 g, 1.35 mmol) and following the procedure described for **8**, compound **10** was obtained as a white solid (0.103 g, 23%). Mp 147.4 °C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.32 - 0.39 (m, 2H), 0.60 - 0.67 (m, 2H), 1.16 - 1.29 (m, 1H), 2.55 (s, 6H), 2.65 (s, 3H) 3.11 (d, *J* = 6.65 Hz, 2H), 6.78 (d, *J* = 7.22 Hz, 1H), 6.93 (t, *J* = 8.38 Hz, 1H), 7.00 (d, *J* = 8.38 Hz, 1H), 7.07 (d, *J* = 8.38 Hz, 1H), 7.13 (d, *J* = 8.09 Hz, 1H), 7.23 (dd, *J* = 11.27, 2.02 Hz, 1H), 7.89 (d, *J* = 6.94 Hz, 1H). LC–MS *m/z* 403 [M + H]⁺, t_R = 2.07 min.

8-Cyclopropyl-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl[1,2,4]triazolo[4,3-a]-pyridine (11). Starting from **51b** (0.230 g, 0.93 mmol) and **52a** (0.382 g, 1.11 mmol) and following the procedure described for **8**, compound **11** was obtained as a white solid (0.269 g, 68%). Mp 166.4 °C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.29 - 0.38 (m, 2H), 0.58 - 0.67 (m, 2H), 0.90 - 1.00 (m, 2H), 1.14 - 1.31 (m, 1H), 1.55 - 1.70 (m, 2H) 2.10 (tt, *J*=8.64, 5.46 Hz, 1H), 2.54 (s, 3H), 2.55 (s, 3H), 3.06 (d, *J*=6.70 Hz, 2H), 6.73 (d, *J*=7.17 Hz, 1H), 6.93 (t, *J*=8.44 Hz, 1H), 7.00 (d, *J*=8.32 Hz, 1H), 7.13 (d, *J*=8.32 Hz, 1H), 7.15 - 7.20 (m, 1H), 7.32 (dd, *J* = 11.21, 1.97 Hz, 1H), 7.80 (d, *J* = 7.17 Hz, 1H). LC–MS *m/z* 429 [M + H]⁺, t_R = 2.24 min.

chlorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (12). Starting from 50a (0.250 g, 0.75 mmol) and 52b (0.385 g, 0.75 mmol) and following the procedure described for 8, compound 12 was obtained as a white solid (0.100 g, 70%). Mp 182.7 °C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.31 - 0.41 (m, 2H), 0.57 - 0.70 (m, 2H), 1.21 - 1.33 (m, 1H), 2.51 (s, 3H), 2.56 (s, 3H), 3.13 (d, J =6.94 Hz, 2H), 6.82 (d, J = 8.55 Hz, 1H), 6.87 (d, J = 6.94 Hz, 1H), 7.03 (d, J = 8.32 Hz, 1H), 7.16 (d, J = 8.09 Hz, 1H), 7.37 (dd, J = 8.55, 2.31 Hz, 1H), 7.66 (d, J = 2.31 Hz, 1H), 7.97 (d, J =7.17 Hz, 1H). LC–MS m/z 439 [M + H]⁺, t_R = 3.08 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

chlorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (13). Starting from 49a (0.2 g, 0.73 mmol) and 52b (0.26 g, 0.72 mmol) and following the procedure described for 8, compound 13 was obtained as a white solid (0.217 g, 61%). Mp > 300°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.34 - 0.41 (m, 2H), 0.62 - 0.70 (m, 2H), 1.16 - 1.30 (m, 1H), 2.50 (s, 3H), 2.56 (s, 3H), 3.15 (d, *J* = 6.70 Hz, 2H), 6.79 (d, *J* = 8.09 Hz, 2H), 7.03 (d, *J* = 8.09 Hz, 1H), 7.11 - 7.19 (m, 2H), 7.49 (d, *J* = 2.31 Hz, 1H), 8.10 (d, *J* = 7.17 Hz, 1H). LC–MS *m/z* 473 [M + H]⁺, t_R = 3.22 min.

8-Methyl-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

chlorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (14). Starting from 60a (0.250 g, 0.127 mmol) and 52b (0.464 g, 1.35 mmol) and following the procedure described for 8, compound 14 was obtained as a white solid (0.107 g, 22%). Mp 121 °C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.36 (q, J = 5.01 Hz, 2H), 0.58 - 0.67 (m, 2H), 1.17 - 1.30 (m, 1H), 2.52 (s, 3H), 2.56 (s, 3H), 2.64 (s, 3H), 3.11 (d, J = 6.65 Hz, 2H), 6.78 (d, J = 6.94 Hz, 1H), 6.83 (d, J = 8.67 Hz, 1H), 7.02 (d, J = 8.38 Hz, 1H), 7.13 (d, J = 8.09 Hz, 1H), 7.18 (dd, J = 8.38, 2.02 Hz, 1H), 7.49 (d, J = 2.31 Hz, 1H), 7.89 (d, J = 6.94 Hz, 1H). LC–MS m/z 419 [M + H]⁺, t_R = 2.23 min.

8-Chloro-3-cyclopropylmethyl-7-{4-[(2-methylpyridin-4-yl)oxy]-3-

fluororophenyl[**1,2,4**]**triazolo**[**4,3-a**]**-pyridine (15)**. Starting from **50a** (1.71 g, 5.13 mmol) and **52c** (1.4 g, 5.67 mmol) and following the procedure described for **8**, compound **15** was obtained as a white solid (0.100 g, 70%). Mp > 300°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.34 - 0.41 (m, 2H), 0.63 - 0.70 (m, 2H), 1.18 - 1.28 (m, 1H), 2.55 (s, 3H), 3.14 (d, *J* = 6.65 Hz, 2H), 6.73 (dd, *J* = 5.78, 2.31 Hz, 1H), 6.78 (d, *J* = 2.31 Hz, 1H), 6.91 (d, *J* = 7.22 Hz, 1H), 7.27 - 7.34 (m, 1H), 7.36 - 7.41 (m, 1H), 7.45 (dd, *J* = 10.69, 2.02 Hz, 1H), 8.01 (d, *J* = 6.94 Hz, 1H), 8.41 (d, *J* = 5.78 Hz, 1H). LC–MS *m/z* 409 [M + H]⁺, t_R = 2.57 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2-methylpyridin-4-yl)oxy]-3-

fluororophenyl [1,2,4]triazolo [4,3-a]-pyridine (16). Starting from 49a (1.4 g, 5.08 mmol) and 52c (1.84 g, 5.59 mmol) and following the procedure described for 8, compound 16 was obtained as a white solid (0.233 g, 10%). Mp 194.6°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.34 - 0.44 (m, 2 H), 0.61 - 0.73 (m, 2H), 1.18 - 1.29 (m, 1H), 2.55 (s, 3H), 3.17 (d, *J* = 6.66 Hz, 2H), 6.70 (dd, *J* = 5.8, 2.6 Hz, 1H), 6.76 (d, *J* = 2.3 Hz, 1H), 6.83 (d, *J* = 7.2 Hz, 1H), 8.20 (br d, J = 8.4 Hz, 1H), 7.23 - 7.31 (m, 2H), 8.14 (d, *J* = 7.2 Hz, 1H), 8.41 (d, *J* = 5.8 Hz, 1H). LC–MS *m/z* 443 [M + H]⁺, t_R = 2.70 min.

8-Chloro-3-cyclopropylmethyl-7-{4-[(2-ethylpyridin-4-yl)oxy]-3-

fluororophenyl}[1,2,4]triazolo[4,3-a]-pyridine (17). Starting from 50a (0.26 g, 0.78 mmol) and 52d (0.293 g, 0.857 mmol) and following the procedure described for 8, compound 17 was obtained as a white solid (0.316 g, 96%). Mp > 300°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.31 - 0.43 (m, 2H), 0.60 - 0.72 (m, 2H), 1.15 - 1.29 (m, 1H), 1.31 (t, J = 7.7 Hz, 3H), 2.82 (q, J = 7.6 Hz, 2H), 3.14 (d, J = 6.6 Hz, 2H), 6.72 (dd, J = 5.8, 2.3 Hz, 1H), 6.81 (d, J = 2.3 Hz, 1H), 6.91

(d, J = 6.9 Hz, 1H), 7.31 (t, J = 8.2 Hz, 1H), 7.35 - 7.42 (m, 1H), 7.45 (dd, J = 10.7, 2.0 Hz, 1H), 8.01 (d, J = 6.9 Hz, 1H), 8.44 (d, J = 5.8 Hz, 1H). LC–MS m/z 423 [M + H]⁺, t_R = 2.04 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2,6-dimethylpyridin-4-yl)oxy]-3-

fluororophenyl[1,2,4]triazolo[4,3-a]-pyridine (18). Starting from **49a** (0.185 g, 0.67 mmol) and **52e** (0.3 g, 0.874 mmol) and following the procedure described for **8**, compound **18** was obtained as a white solid (0.100 g, 33%). Mp 232.5°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.24 - 0.50 (m, 2H) 0.55 - 0.79 (m, 2H) 1.05 - 1.39 (m, 1H) 2.50 (s, 6H) 3.17 (d, *J*=6.70 Hz, 2H) 6.56 (s, 2H) 6.84 (d, *J*=7.17 Hz, 1H) 7.16 - 7.21 (m, 1H) 7.22 - 7.29 (m, 2H) 8.14 (d, *J*=7.17 Hz, 1 H). LC–MS *m/z* 457 [M + H]⁺, t_R = 2.94 min.

8-Chloro-3-cyclopropylmethyl-7-{4-[(2-cyclopropylpyridin-4-yl)oxy]-3-

fluororophenyl[**1,2,4**]**triazolo**[**4,3-a**]**-pyridine (19)**. Starting from **50a** (0.103 g, 0.310 mmol) and **52f** (0.22 g, 0.310 mmol) and following the procedure described for **8**, compound **19** was obtained as a white solid (0.06 g, 45%). Mp > 300°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.38 (q, *J* = 5.11 Hz, 2H), 0.62 - 0.71 (m, 2H), 0.95 - 1.09 (m, 4H), 1.16 - 1.29 (m, 1H), 1.94 - 2.03 (m, 1H), 3.14 (d, *J* = 6.65 Hz, 2H), 6.66 (dd, *J* = 5.64, 2.46 Hz, 1H), 6.78 (d, *J* = 2.31 Hz, 1H), 6.90 (d, *J* = 6.94 Hz, 1H), 7.28 - 7.33 (m, 1H), 7.34 - 7.40 (m, 1H), 7.44 (dd, *J* = 10.69, 2.02 Hz, 1H), 8.00 (d, *J* = 6.94 Hz, 1H), 8.35 (d, *J* = 5.78 Hz, 1H). LC–MS *m/z* 435 [M + H]⁺, t_R = 4.51 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2-cyclopropylpyridin-4-yl)oxy]-3-

fluororophenyl}[1,2,4]triazolo[4,3-a]-pyridine (20). Starting from 49a (0.2 g, 0.725 mmol) and 52f (0.283 g, 0.798 mmol) and following the procedure described for 8, compound 20 was obtained as a white solid (0.058 g, 17%). Mp 211.1°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.32-0.45 (m, 2H), 0.53 - 0.75 (m, 2H), 0.96 - 1.03 (m, 2H), 1.02 - 1.08 (m, 2H), 1.16 - 1.30 (m, 1H),

1.91 - 2.03 (m, 1H), 3.16 (d, J = 6.7 Hz, 2H), 6.63 (dd, J = 5.8, 2.3 Hz, 1H), 6.75 (d, J = 2.3 Hz, 1H), 6.83 (d, J = 7.2 Hz, 1H), 7.15-7.22 (m, 1H), 7.22 - 7.31 (m, 2H), 8.15 (d, J = 6.9 Hz, 1H), 8.35 (d, J = 5.5 Hz, 1H). LC-MS m/z 469 [M + H]⁺, t_R = 3.31 min.

8-Chloro-3-(2,2,2-trifluoroethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl[1,2,4]triazolo[4,3-a]-pyridine (21). Starting from **56c** (0.2 g, 0.55 mmol) and **52a** (0.228 g, 0.664 mmol) and following the procedure described for **8**, compound **21** was obtained as a white solid (0.032 g, 13%). Mp 164.5°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 2.53 (s, 3H), 2.55 (s, 3H), 4.11 (q, J = 9.9 Hz, 2H), 6.93 (t, J = 8.3 Hz, 1H), 6.98 (d, J = 7.2 Hz, 1H), 7.02 (d, J = 8.3 Hz, 1H), 7.16 (d, J = 8.3 Hz, 1H), 7.23 - 7.28 (m, 1H), 7.42 (dd, J = 11.1, 2.1 Hz, 1H), 8.01 (d, J = 7.2 Hz, 1H). LC-MS *m/z* 451 [M + H]⁺, t_R = 3.24 min.

8-Chloro-3-(ethoxymethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (22). Starting from 50b (0.4 g, 1.19 mmol) and 52a (0.488 g, 1.42 mmol) and following the procedure described for 8, compound 22 was obtained as a white solid (0.310 g, 61%). Mp 180.3°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.23 (t, *J* = 6.94 Hz, 3H), 2.54 (s, 3 H), 2.55 (s, 3H), 3.58 (q, *J* = 6.94 Hz, 2H), 5.10 (s, 2H), 6.87 - 6.95 (m, 2H), 7.02 (d, *J* = 8.09 Hz, 1H), 7.16 (d, *J* = 8.09 Hz, 1H), 7.22 - 7.26 (m, 1H), 7.42 (dd, *J* = 11.27, 2.31 Hz, 1H), 8.25 (d, *J*=7.22 Hz, 1H). LC–MS *m/z* 427 [M + H]⁺, t_R = 2.77 min.

8-Trifluoromethyl-3-(ethoxymethyl-7-{4-[(2,6-dimethylpyridin-3-yl)oxy]-3-

fluorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (23). Starting from 49b (0.150 g, 0.536 mmol) and 52a (0.239 g, 0.697 mmol) and following the procedure described for 8, compound 23 was obtained as a white solid (0.150 g, 61%). Mp 142.2°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.23 (t, *J*=7.08 Hz, 3H), 2.53 (s, 3H), 2.54 (s, 3H), 3.59 (q, *J* = 7.13 Hz, 2H), 5.12 (s, 2H), 6.82 (d, *J* = 7.22 Hz, 1H), 6.90 (t, *J* = 8.24 Hz, 1H), 7.01 (d, *J* = 8.38 Hz, 1H), 7.06 (d, *J* = 8.09 Hz, 1H), 7.14

(d, J = 8.38 Hz, 1H), 7.23 (dd, J = 10.69, 2.02 Hz, 1H), 8.40 (d, J = 6.94 Hz, 1H). LC–MS m/z461 [M + H]⁺, t_R = 2.89 min.

8-Chloro-3-cyclopropylmethyl-7-[3-fluoro-4-(isopropylamino)-phenyl][1,2,4]triazolo[4,3a]-pyridine (24). Starting from 50a (0.350 g, 1.049 mmol) and 52g (0.381 g, 1.36 mmol) and following the procedure described for 8, compound 24 was obtained as a white solid (0.097 g, 26%). Mp 196.4°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.35 (q, J = 5.20 Hz, 2H), 0.60 - 0.67 (m, 2H), 1.15 - 1.26 (m, 1H), 1.29 (d, J = 6.36 Hz, 6H), 3.10 (d, J = 6.65 Hz, H), 3.72 (dq, J =13.19, 6.49 Hz, 1H), 3.98 (br d, J = 5.20 Hz, 1H), 6.70 - 6.81 (m, 1H), 6.87 (d, J = 6.94 Hz, 1H), 7.17 - 7.31 (m, 2H) 7.91 (d, J = 7.22 Hz, 1H). LC–MS *m/z* 359 [M + H]⁺, t_R = 3.11 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-fluoro-4-(isopropylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (25). Starting from 49a (0.350 g, 1.27 mmol) and 52g (0.461 g, 1.65 mmol) and following the procedure described for 8, compound 25 was obtained as a white solid (0.250 g, 50%). Mp 196.9°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.28-0.42 (m, 2H), 0.57 - 0.71 (m, 2H), 1.12 - 1.26 (m, 1H), 1.29 (d, *J* = 6.4 Hz, 6H), 3.12 (d, *J* = 6.6 Hz, 2H), 3.64-3.77 (m, 1H), 3.96 (d, *J* = 4.9 Hz, 1H), 6.74 (t, *J* = 8.4 Hz, 1H), 6.80 (d, *J* = 7.2 Hz, 1H), 7.02 (d, *J* = 10.1 Hz, 2H), 8.04 (d, *J* = 6.9 Hz, 1H). LC–MS *m/z* 393 [M + H]⁺, t_R = 3.33 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(isopropylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (26). Starting from 49a (0.350 g, 1.27 mmol) and 52h (0.488 g, 1.65 mmol) and following the procedure described for 8, compound 26 was obtained as a white solid (0.250 g, 50%). Mp 230°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.29 - 0.42 (m, 2H), 0.55 - 0.71 (m, 2H), 1.11 - 1.27 (m, 1H), 1.31 (d, *J* = 6.36 Hz, 6H), 3.13 (d, *J* = 6.65 Hz, 2H), 3.73 (dq, *J* = 13.11, 6.51 Hz, 1H), 4.38 (br d, *J* = 7.51 Hz, 1H), 6.71 (d, *J* = 8.67 Hz, 1H), 6.79

 (d, J = 7.22 Hz, 1H), 7.16 (dd, J = 8.38, 1.44 Hz, 1H), 7.30 (d, J = 2.02 Hz, 1H), 8.03 (d, J = 6.94 Hz, 1H). LC–MS m/z 409 [M + H]⁺, t_R = 3.71 min.

8-Chloro-3-cyclopropylmethyl-7-[3-fluoro-4-(cyclopropylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (27). Starting from 50a (0.30 g, 0.9 mmol) and 52i (0.3 g, 1.08 mmol) and following the procedure described for 8, compound 27 was obtained as a white solid (0.147 g, 46%). ¹H NMR (500 MHz, CDCl₃) δ ppm 0.28 - 0.40 (m, 2H), 0.57 - 0.69 (m, 4H), 0.78 - 0.87 (m, 2H), 1.15 - 1.28 (m, 1H), 2.45 - 2.56 (m, 1H), 3.11 (d, *J* = 6.65 Hz, 2H), 4.60 (br s, 1H), 6.88 (d, *J* = 7.22 Hz, 1H), 7.13 - 7.20 (m, 1H), 7.21 - 7.31 (m, 2H), 7.92 (d, *J* = 6.94 Hz, 1H). LC–MS *m/z* 357 [M + H]⁺, t_R = 3.30 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-fluoro-4-(cyclopropylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (28). Starting from 49a (0.20 g, 0.73 mmol) and 52i (0.221 g, 0.8 mmol) and following the procedure described for 8, compound 28 was obtained as a white solid (0.169 g, 60%). Mp. > 300° C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.28 - 0.43 (m, 2H), 0.51 - 0.72 (m, 4H), 0.75 - 0.91 (m, 2H), 1.12 - 1.31 (m, 1H), 2.50 (tt, *J* = 6.56, 3.27 Hz, 1H), 3.13 (d, *J* = 6.70 Hz, 2H), 4.58 (br s, 1H), 6.80 (d, *J* = 6.94 Hz, 1H), 6.98 - 7.09 (m, 2H), 7.10 - 7.18 (m, 1H), 8.05 (d, *J* = 7.17 Hz, 1H). LC–MS *m/z* 391 [M + H]⁺, t_R = 3.18 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(cyclopropylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (29). Starting from 49a (0.20 g, 0.73 mmol) and 52j (0.34 g, 0.8 mmol) and following the procedure described for 8, compound 29 was obtained as a white solid (0.109 g, 37%). ¹H NMR (400 MHz, CDCl₃) δ ppm 0.31 - 0.40 (m, 2H), 0.59 - 0.69 (m, 4H), 0.81 - 0.89 (m, 2H), 1.15 - 1.27 (m, 1H), 2.46 - 2.56 (m, 1H), 3.13 (d, *J* = 6.70 Hz, 2H), 4.93 (s, 1H), 6.80 (d, *J* = 7.17 Hz, 1H), 7.11 - 7.16 (m, 1H), 7.17 - 7.23 (m, 1H), 7.28 (d, *J* = 1.85 Hz, 1H), 8.05 (d, *J* = 7.17 Hz, 1H). LC–MS *m/z* 407 [M + H]⁺, t_R = 3.48 min.

phenyl][1,2,4]triazolo[4,3-a]-pyridine (30). Starting from 50a (0.30 g, 0.9 mmol) and 52j (0.32 g, 1.08 mmol) and following the procedure described for 8, compound 30 was obtained as a white solid (0.147 g, 46%). Mp > 300°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.29 - 0.42 (m, 2H), 0.57 - 0.70 (m, 4H), 0.78 - 0.92 (m, 2H), 1.15 - 1.27 (m, 1H), 2.49 - 2.56 (m, 1H), 3.11 (d, J = 6.7 Hz, 2H), 4.60 (br s, 1H), 6.87 (d, J = 6.9 Hz, 1H), 7.18 (d, J = 8.3 Hz, 1H), 7.42 dd, J = 8.3, 2.1 Hz, 1H), 7.47 (d, J = 1.8 Hz, 1H), 7.92 (d, J = 7.2 Hz, 1H). LC–MS *m/z* 373 [M + H]⁺, t_R = 3.83 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2-methoxypyridin-5-yl)methoxy]-3-

fluorophenyl} **[1,2,4]triazolo[4,3-a]-pyridine (31).** Starting from **49a** (0.3 g, 1.088 mmol) and **52k** (0.721 g, 1.306 mmol) and following the procedure described for **8**, compound **31** was obtained as a white solid (0.198 g, 38%). Mp 148.5°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.37 (q, J = 5.09 Hz, 2H), 0.56 - 0.74 (m, 2H), 1.09 - 1.32 (m, 1H), 3.14 (d, J = 6.70 Hz, 2H), 3.96 (s, 3H), 5.12 (s, 2H), 6.79 (dd, J = 13.87, 7.63 Hz, 2H), 7.00 - 7.19 (m, 3H), 7.71 (dd, J = 8.55, 2.54 Hz, 1H) 8.09 (d, J = 7.17 Hz, 1H), 8.24 (d, J = 2.31 Hz, 1H). LC–MS m/z 473 [M + H]⁺, t_R = 2.79 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2-methoxypyridin-5-yl)methoxy]- phenyl }[1,2,4]triazolo[4,3-a]-pyridine (32). Starting from 49a (0.315 g, 1.15 mmol) and 52l (0.430 g, 1.26 mmol) and following the procedure described for 8, compound 32 was obtained as a white solid (0.286 g, 60%). Mp 159.1°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.29 - 0.45 (m, 2H), 0.55 - 0.72 (m, 2H), 1.10 - 1.33 (m, 1H), 3.14 (d, *J* = 6.70 Hz, 2H), 3.96 (s, 3H), 5.04 (s, 2H), 6.67 - 6.87 (m, 2H), 7.00 - 7.12 (m, 2H), 7.32 (d, *J* = 8.55 Hz, 2H), 7.70 (dd, *J* = 8.55, 2.31 Hz, 1H), 8.08 (d, *J* = 7.17 Hz, 1H), 8.25 (d, *J* = 2.31 Hz, 1H). LC–MS *m/z* 455 [M + H]⁺, t_R = 3.73 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2-methoxypyridin-5-yl)methylamino]-3fluorophenyl}[1,2,4]triazolo[4,3-a]-pyridine (33). Starting from 49a (0.530 g, 1.92 mmol) and 52m (0.758 g, 2.11 mmol) and following the procedure described for 8, compound 33 was obtained as a white solid (0.460 g, 51%). Mp 160.4°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.30 -0.42 (m, 2H), 0.59 - 0.71 (m, 2H), 1.15 - 1.27 (m, 1H), 3.13 (d, *J* = 6.70 Hz, 2H), 3.95 (s, 3H), 4.36 (d, *J* = 4.86 Hz, 2H), 4.44 (br s, 1H), 6.69 - 6.83 (m, 3H), 6.95 - 7.12 (m, 2H), 7.62 (dd, *J* = 8.44, 2.43 Hz, 1H), 8.05 (d, *J* = 7.17 Hz, 1H), 8.19 (d, *J* = 2.54 Hz, 1H). LC–MS *m/z* 472 [M + H]⁺, t_R = 2.67 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-{4-[(2-methoxypyridin-5-yl)methylamino]-3phenyl}[1,2,4]triazolo[4,3-a]-pyridine (34). Starting from 49a (0.150 g, 0.544 mmol) and 52n (0.204 g, 0.6 mmol) and following the procedure described for 8, compound 34 was obtained as a white solid (0.115 g, 46%). Mp 127.5°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.29 - 0.42 (m, 2H), 0.60 - 0.68 (m, 2H), 1.17 - 1.24 (m, 1H), 3.12 (d, *J* = 6.94 Hz, 2H), 3.95 (s, 3H), 4.20 (br s, 1H), 4.32 (d, *J* = 4.91 Hz, 2H), 6.70 (d, *J* = 8.38 Hz, 2H), 6.76 (d, *J* = 8.38 Hz, 1H), 6.80 (d, *J* = 6.94 Hz, 1H), 7.21 (d, *J* = 8.67 Hz, 2H), 7.61 (dd, *J* = 8.53, 2.46 Hz, 1H), 8.03 (d, *J* = 7.22 Hz, 1H), 8.18 (d, *J* = 2.31 Hz, 1H). LC–MS *m/z* 454 [M + H]⁺, t_R = 2.05 min.

8-Chloro-3-cyclopropylmethyl-7-{4-[(2-methoxypyridin-5-yl)methylamino]-3-phenyl}

[1,2,4]triazolo[4,3-a]-pyridine (35). Starting from 50a (0.150 g, 0.45 mmol) and 52n (0.168 g, 0.494 mmol) and following the procedure described for 8, compound 35 was obtained as a white solid (0.098 g, 51%). Mp 158.1°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.30 - 0.40 (m, 2H), 0.57 - 0.70 (m, 2H), 1.13 - 1.39 (m, 1H), 3.10 (d, *J* = 6.70 Hz, 2H), 3.95 (s, 3H), 4.33 (s, 3H), 6.69 - 6.79 (m, 3H), 6.89 (d, *J* = 7.17 Hz, 1H), 7.38 - 7.46 (m, 2H), 7.62 (dd, *J* = 8.55, 2.54 Hz, 1H), 7.91 (d, *J* = 7.17 Hz, 1H), 8.19 (d, *J* = 2.08 Hz, 1H). LC–MS *m/z* 420 [M + H]⁺, t_R = 1.93 min.

8-Chloro-3-cyclopropylmethyl-7-[3-chloro-4-(tetrahydro-pyran-4-yloxy)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (36). Starting from 50a (0.20 g, 0.6 mmol) and 52o (0.243 g, 0.72 mmol) and following the procedure described for 8, compound 36 was obtained as a white solid (0.160 g, 64%). Mp. 189.1°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.30 - 0.43 (m, 2H), 0.56 - 0.73 (m, 2H), 1.15 - 1.30 (m, 1H), 1.92 (ddt, J = 17.05, 7.28, 3.55, 3.55 Hz, 2H), 2.02 - 2.17 (m, 2H), 3.12 (d, J = 6.70 Hz, 2H), 3.65 (ddd, J = 11.44, 7.40, 3.58 Hz, 2H), 4.05 (ddd, J = 11.44, 7.40, 3.58 Hz, 2H), 4.66 (tt, J = 7.11, 3.64 Hz, 1H), 6.86 (d, J = 6.94 Hz, 1H), 7.06 (d, J = 8.55 Hz, 1H), 7.43 (dd, J = 8.55, 2.31 Hz, 1H), 7.58 (d, J = 2.31 Hz, 1H), 7.96 (d, J = 7.17 Hz, 1H). LC–MS m/z 418 [M + H]⁺, t_R = 3.25 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(tetrahydro-pyran-4-yloxy)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (37). Starting from 49a (0.09 g, 0.33 mmol) and 52o (0.138 g, 0.41 mmol) and following the procedure described for 8, compound 37 was obtained as a white solid (0.083 g, 56%). Mp. 176.9°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.30 - 0.43 (m, 2H), 0.58 - 0.73 (m, 2H), 1.16 - 1.28 (m, 1H), 1.86-1.97 (m, 2H), 2.02 - 2.12 (m, 2H), 3.14 (d, *J* = 6.70 Hz, H), 3.59-3.69 (m, 2H), 4-4.09 (m, 2H), 4.61-4.68 (m, 2H), 6.78 (d, *J* = 7.2 Hz, 1H), 7.02 (d, *J* = 8.6 Hz, 1H), 7.20 (dd, *J* = 8.6, 2.1 Hz, 1H), 7.41 (d, *J* = 2.1 Hz, 1H), 8.09 (d, *J* = 7.2 Hz, 1H). LC–MS *m/z* 452 [M + H]⁺, t_R = 3.33 min.

8-Chloro-3-cyclopropylmethyl-7-[3-chloro-4-(tetrahydro-pyran-4-ylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (38). Starting from 50a (0.20 g, 0.6 mmol) and 52p (0.242 g, 0.72 mmol) and following the procedure described for 8, compound 38 was obtained as a white solid (0.160 g, 64%). Mp. 221.3°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.28 - 0.41 (m, 2H), 0.56 - 0.71 (m, 2H), 1.13 - 1.29 (m, 1H), 1.45 - 1.71 (m, 2H), 2.10 (br d, J = 13.29 Hz, 2H), 3.11 (d, J = 6.65 Hz, 2H), 3.52 - 3.60 (m, 2H), 3.62 (br dd, J = 6.94, 3.18 Hz, 1H), 4.05 (dt, J = 13.29 Hz, 2H), 4.05

Journal of Medicinal Chemistry

11.78, 3.65 Hz, 2H), 4.49 (br d, J = 7.80 Hz, 1H), 6.78 (d, J = 8.38 Hz, 1H), 6.86 (d, J = 7.22 Hz, 1H), 7.39 (dd, J = 8.53, 2.17 Hz, 1H), 7.51 (d, J = 2.31 Hz, 1H), 7.92 (d, J = 7.22 Hz, 1H). LC-MS m/z 417 [M + H]⁺, t_R = 4.08 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(tetrahydro-pyran-4-ylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (39). Starting from **49a** (0.07 g, 0.25 mmol) and **52p** (0.107 g, 0.32 mmol) and following the procedure described for **8**, compound **39** was obtained as a white solid (0.045 g, 39%). Mp. 198.4°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.31 - 0.42 (m, 2H), 0.58 - 0.71 (m, 2H), 1.16 - 1.27 (m, 1H), 1.55 - 1.68 (m, 2H), 2.09 (br d, J = 12.7 Hz, 2H), 3.13 (d, J = 6.7 Hz, 2H), 3.56 (td, J = 11.8, 2.3 Hz, 1H), 3.56-3.67 (m, 1H), 4.05 (dt, J = 11.7, 3.7 Hz, 2H), 4.47 (d, J = 7.6 Hz, 1H), 6.74 (d, J = 8.6 Hz, 1H), 6.78 (d, J = 7.2 Hz, 1H), 7.16 (dd, J = 8.3, 1.8 Hz, 1H), 7.32 (d, J = 2.1 Hz, 1H), 8.05 (d, J = 7.2 Hz, 1H). LC–MS *m/z* 451 [M + H]⁺, t_R = 3.29 min.

8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(tetrahydro-pyran-4-

ylaminomethyl)-phenyl][1,2,4]triazolo[4,3-a]-pyridine (40). Starting from 49a (0.2 g, 0.73 mmol) and 52q (0.306 g, 0.87 mmol) and following the procedure described for 8, compound 40 was obtained as a white solid (0.060 g, 18%). ¹H NMR (400 MHz, CDCl₃) δ ppm 0.37 (q, J = 5.09 Hz, 2H), 0.59 - 0.74 (m, 2H), 1.10 - 1.32 (m, 2H), 1.41 - 1.73 (m, 2H), 1.92 (br dd, J = 12.60, 1.73 Hz, 2H), 2.69 - 2.86 (m, 1H), 3.15 (d, J = 6.70 Hz, 2H), 3.34 - 3.49 (m, 2H), 3.97 - 4.01 (m, 3H), 4.01 - 4.04 (m, 1H), 6.76 (d, J = 7.17 Hz, 1H), 7.22 - 7.28 (m, 1H), 7.37 (d, J = 1.62 Hz, 1H), 7.56 (d, J = 7.86 Hz, 1H), 8.10 (d, J = 7.17 Hz, 1H). LC–MS *m/z* 465 [M + H]⁺, t_R = 2.09 min.

Trans-8-Chloro-3-cyclopropylmethyl-7-[3-chloro-4-(4-hydroxy-cyclohexylamino)phenyl][1,2,4]triazolo[4,3-a]-pyridine (41). Starting from 50a (0.13 g, 0.388 mmol) and 52r (0.15 g, 0.43 mmol) and following the procedure described for **8**, compound **41** was obtained as a white solid (0.087 g, 52%). Mp 270.9°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.29 - 0.42 (m, 2H), 0.56 - 0.71 (m, 2H), 1.17 - 1.25 (m, 1H), 1.47 (br s, 1H), 1.73 - 1.8 (m, 4H), 1.80-1.91 (m, 4H), 3.11 (d, J=6.7 Hz, 2H), 3.46 - 3.57 (m, 1H), 3.98 (br s, 1H), 4.60 (br d, J=7.6 Hz, 1H), 6.76 (d, J=8.8 Hz, 1H), 6.87 (d, J=7.2 Hz, 1H), 7.39 (dd, J=8.3, 2.3 Hz, 1H), 7.49 (d, J=2.1 Hz, 1H), 7.91 (d, J=7.2 Hz, 1H), LC-MS *m/z* 431 [M + H]⁺, t_R = 3.49 min.

Trans-8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(4-hydroxy-

cyclohexylamino)-phenyl][1,2,4]triazolo[4,3-a]-pyridine (42). Starting from 49a (0.3 g, 1.088 mmol) and 52r (0.459 g, 1.306 mmol) and following the procedure described for 8, compound 42 was obtained as a white solid (0.209 g, 41%). Mp. 290.7°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.3 - 0.43 (m, 2H), 0.58 - 0.71 (m, 2H), 1.16 - 1.25 (m, 1H), 1.29-1.42 (m, 2H), 1.42 - 1.53 (m, 3H), 2.03-2.12 (m, 2H), 2.20 (br d, *J* = 12.0 Hz, 2H), 3.13 (d, *J*=6.7 Hz, 2H), 3.32 - 3.43 (m, 1H), 3.70-3.80 (m, 1H), 4.39 (d, *J* = 7.6 Hz, 1H), 6.72 (d, *J* = 8.6 Hz, 1H), 6.79 (d, *J* = 7.2 Hz, 1H), 7.16 (dd, *J* = 8.6, 2.1 Hz, 1H), 7.30 (d, *J* = 2.1 Hz, 1H), 8.04 (d, *J* = 7.2 Hz, 1H). LC–MS m/z 465 [M + H]⁺, t_R = 2.50 min.

Cis-8-Chloro-3-cyclopropylmethyl-7-[3-chloro-4-(4-hydroxy-cyclohexylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (43). Starting from 50a (0.13 g, 0.388 mmol) and 52s (0.15 g, 0.43 mmol) and following the procedure described for 8, compound 43 was obtained as a white solid (0.060 g, 36%). Mp. 273.7°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.27 - 0.43 (m, 2H) 0.57 - 0.70 (m, 2H), 1.02 - 1.32 (m, 2H), 1.68 - 1.98 (m, 8H), 3.11 (d, *J* = 6.70 Hz, 2H), 3.44 - 3.62 (m, 1H), 3.98 (br s, 1H), 4.60 (br d, *J* = 7.63 Hz, 1H), 6.76 (d, *J* = 8.79 Hz, 1H), 6.87 (d, *J* = 7.17 Hz, 1H), 7.39 (dd, *J* = 8.44, 2.20 Hz, 1H), 7.49 (d, *J* = 2.08 Hz, 1H), 7.91 (d, *J* = 7.17 Hz, 1H). LC-MS m/z 431 [M + H]⁺, t_R = 3.08 min.

Cis-8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(4-hydroxy-4-cyclopropylcyclohexylamino)-phenyl][1,2,4]triazolo[4,3-a]-pyridine (44). Starting from 49a (0.143 g, 0.52 mmol) and 52t (0.235 g, 0.6 mmol) and following the procedure described for 8, compound 44 was obtained as a white solid (0.047 g, 18%). Mp > 300°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 0.04 - 0.16 (m, 2H), 0.18 - 0.33 (m, 5H), 0.34 - 0.43 (m, 2H), 0.60 (tt, *J* = 8.31, 5.56 Hz, 1H), 0.80 - 1.00 (m, 1H), 1.11 - 1.25 (m, 2H), 1.31 - 1.56 (m, 4H), 1.63 - 1.79 (m, 2H), 2.49 (d, *J* = 6.65 Hz, 2H), 2.97 (tdt, *J* = 11.13, 11.13, 7.37, 3.97, 3.97 Hz, 1H), 4.45 (d, *J* = 7.51 Hz, 1H), 5.95 (d, *J* = 7.22 Hz, 1H), 6.53 (d, *J* = 8.67 Hz, 1H), 6.73 (d, *J* = 7.22 Hz, 1H), 7.16 (s, 1H) 7.28 (d, *J* = 2.02 Hz, 1H). LC–MS *m/z* 465 [M + H]⁺, t_R = 3.49 min.

Trans-8-Chloro-3-(2,2,2-trifluoroethyl)-7-[3-chloro-4-(4-hydroxy-cyclohexylamino)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (45). Starting from 56c (0.11 g, 0.304 mmol) and 52r (0.128 g, 0.365 mmol) and following the procedure described for 8, compound 45 was obtained as a white solid (0.054 g, 39%). Mp > 300°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.28 - 1.52 (m, 5H), 2.08 (br d, J = 10.63 Hz, 2H), 2.20 (br d, J = 12.02 Hz, 2H), 3.26 - 3.46 (m, 1H), 3.62 - 3.85 (m, 1H), 4.08 (q, J = 9.71 Hz, 2H), 4.44 (d, J = 7.63 Hz, 1H), 6.77 (d, J = 8.55 Hz, 1H), 6.97 (d, J = 7.17 Hz, 1H), 7.39 (dd, J = 8.55, 2.08 Hz, 1H), 7.50 (d, J = 2.08 Hz, 1H), 7.94 (d, J = 7.17 Hz, 1H). LC–MS m/z 459 [M + H]⁺, t_R = 3.23 min.

Trans-8-Trifluoromethyl-3-cyclopropylmethyl-7-[3-chloro-4-(4-hydroxy-cyclohexyloxy)phenyl][1,2,4]triazolo[4,3-a]-pyridine (46). Starting from 49a (0.119 g, 0.432 mmol) and 52u (0.183g, 0.518 mmol) and following the procedure described for 8, compound 46 was obtained as a white solid (0.022 g, 27%). ¹H NMR (400 MHz, CDCl₃) δ ppm 0.31 - 0.41 (m, 2H), 0.61 - 0.71 (m, 2H), 1.09 - 1.32 (m, 2H), 1.42 - 1.56 (m, 2H), 1.67 - 1.80 (m, 2H), 2.05 - 2.23 (m, 4H), 3.14 (d, *J* = 6.70 Hz, 2H), 3.83 - 3.96 (m, 1H), 4.36 - 4.49 (m, 1H), 6.78 (d, *J* = 7.17 Hz, 1H), 7.02 (d, J = 8.79 Hz, 1H), 7.20 (dd, J = 8.44, 2.20 Hz, 1H), 7.39 (d, J = 2.31 Hz, 1H), 8.09 (d, J = 7.17 Hz, 1H). LC-MS m/z 466 [M + H]⁺, t_R = 2.96 min.

Trans-8-Chloro-3-(2,2,2-trifluoroethyl)-7-[3-chloro-4-(4-hydroxy-cyclohexyloxy)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (47). Starting from 56c (0.190 g, 0.526 mmol) and 52u (0.222 g, 0.631 mmol) and following the procedure described for **8**, compound 47 was obtained as a white solid (0.065 g, 27%). Mp >300°C. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.42 (d, J = 3.76 Hz, 1H), 1.46 - 1.55 (m, 2H), 1.68 - 1.79 (m, 2H), 2.10 (td, J = 7.73, 3.61 Hz, 2H), 2.18 (td, J = 7.66, 3.47 Hz, 2H), 3.91 (td, J = 8.09, 4.05 Hz, 1H), 4.10 (q, J = 9.73 Hz, 2H), 4.46 (tt, J = 8.06, 3.79 Hz, 1H), 6.97 (d, J = 7.22 Hz, 1H), 7.08 (d, J = 8.38 Hz, 1H), 7.42 (dd, J = 8.53, 2.17 Hz, 1H), 7.57 (d, J = 2.31 Hz, 1H), 7.98 (d, J = 7.22 Hz, 1H). LC–MS *m/z* 460 [M + H]⁺, t_R = 3.23 min.

Cis-8-Chloro-3-(2,2,2-trifluoroethyl)-7-[3-chloro-4-(4-hydroxy-cyclohexyloxy)-

phenyl][1,2,4]triazolo[4,3-a]-pyridine (48). Starting from 56c (0.150 g, 0.415 mmol) and 52v (0.176g, 0.498 mmol) and following the procedure described for 8, compound 48 was obtained as a white solid (0.044 g, 23%). Mp 196.7°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.39 - 1.48 (m, 1H), 1.66 - 1.93 (m, 6H), 2.06 - 2.21 (m, 2H), 3.81 (tq, J = 8.44, 4.24 Hz, 1H), 4.10 (q, J = 9.71 Hz, 2H), 4.56 (tt, J = 5.20, 2.77 Hz, 1H), 6.98 (d, J = 7.17 Hz, 1H), 7.06 (d, J = 8.55 Hz, 1H), 7.43 (dd, J = 8.55, 2.31 Hz, 1H), 7.57 (d, J = 2.31 Hz, 1H), 7.99 (d, J = 7.17 Hz, 1H). LC–MS m/z 460 [M + H]⁺, t_R = 3.14 min.

8-Trifluoromethyl-3-(ethoxymethyl)-7-[3-chloro-1,2,4-triazolo[4,3-a]-pyridine (49b). Mp 104°C. ¹H NMR (400 MHz, CDCl₃) δ ppm ¹H NMR (500 MHz, CHLOROFORM-d) δ ppm 1.22 (t, J=6.94 Hz, 3 H) 3.57 (q, J=6.94 Hz, 2 H) 5.08 (s, 2 H) 6.95 (d, J=7.22 Hz, 1 H) 8.35 (d, J=7.51 Hz, 1 H). LC-MS *m/z* 280 [M + H]⁺, t_R = 1.32 min.

Biology

Membrane preparation

CHO-K1 cells stably expressing the wildtype hmGlu₂ receptor (CHO-K1_hmGlu₂) were grown in DMEM medium supplemented with 10% (v/v) fetal calf serum, 200 IU/mL penicillin, 200 µg/mL streptomycin, 30.5 µg/mL L proline and 400 µg/mL G418 at 37°C and 5% CO₂. Sodium butyrate (final concentration 5 mM) was added to the plates when cells growth reached 70% confluency.⁶³ 24 hours later, cells were detached from the plates by scraping into 5 ml of PBS and centrifuged for 5 min at 1500 rpm. Pellets were resuspended into ice-cold Tris buffer (50 mM Tris-HCl pH 7.4) and homogenized using an Ultra Turrax homogenizer (IKA Werke GmbH & Co.KG, Staufen, Germany). An Optima LE 80 K ultracentrifuge (Beckman Coulter, Fullerton, CA) at 31,000 rpm was used for separation of membranes and the cytosolic fraction at 4°C for 20 min. Pellets were resuspended in 10 ml Tris buffer and the centrifugation and homogenization steps were repeated. Remaining pellets were suspended into assay buffer (50 mM Tris-HCl pH 7.4, 2 mM CaCl₂, 10 mM MgCl₂) and the homogenization step was repeated. Aliquots were stored at -80°C. BCA protein determination was used to determine the membrane protein concentrations.

[³⁵S]GTP_yS Binding Assay

[³⁵S]GTPγS binding experiments were performed as previously described.¹²

Radioligand Binding Assays

After thawing, membranes were homogenized by an Ultra Turrax homogenizer. **4** (10 μ M) was used to determine nonspecific binding. DMSO concentrations were $\leq 0.25\%$. For all experiments, radioligand concentrations were such that <10% of the amount added was receptor-bound to avoid ligand depletion.

For displacement assays, membrane protein aliquots (30 μ g) were incubated with 6 nM [³H]-7 and 10 concentrations of competing ligand diluted by an HP D300 digital dispenser (Tecan, Giessen, The Netherlands) in assay buffer to a total volume of 100 μ l. After 60 minutes at 15°C, incubation was terminated by rapid filtration over GF/C filterplates (PerkinElmer, Groningen, The Netherlands) on a PerkinElmer filtermate harvester. Filterplates were washed 10 times with ice-cold wash buffer (50 mM Tris-HCl pH 7.4) and filter-bound radioactivity was determined in a Microbeta 2450² microplate counter (PerkinElmer).

For association, dissociation and competition association experiments a scintillation proximity assay (SPA) was developed and used. Membrane protein (20 μ g) and pre-wetted wheat-germ agglutinin coated SPA beads (0.2 mg; RPNQ0001, PerkinElmer) were pre-coupled in assay buffer while gently shaking at room temperature for 30 minutes. Then, this membrane bead mixture was added to an Isoplate-96 (PerkinElmer) together with 6 nM [³H]-7 and competing ligand in case of competition association. After addition, the plate was rapidly placed in a Microbeta 2450² microplate counter. Plates were recorded for 120 minutes measuring every 30 seconds at ambient temperature of 28°C. Binding values were determined in corrected counts per minute (CCPM). For dissociation assays, radioligand dissociation was initiated after 60 minutes of incubation by addition of 5 μ l 4 (final concentration: 10 μ M). Subsequently, plates were recorded for 120 minutes being measuring every 30 seconds.

Label-free whole-cell assays

Label-free whole-cell assays were performed using the xCELLigence real-time cell analyzer (RTCA DP) system as described previously ^{58,64}. Baseline impedance was measured using 45 μ l culture medium (as described under membrane preparation) per well in 16 well E-plates (Westburg, Leusden, The Netherlands). 40,000 cells per well were added in a volume of 50 μ l.

Journal of Medicinal Chemistry

After resting for 30 minutes at room temperature, E-plates were placed in the recording station within a humidified 37°C, 5% CO₂ incubator. Impedance (represented in the arbitrary unit Cell Index) was measured every 15 minutes overnight. After 18 hours, cells were stimulated with the indicated concentration of compound. Impedance was measured every 15 seconds for 20 minutes, then 5 minute intervals were used for the next 40 minutes, followed by 15 minute intervals thereafter. For washing experiments, cells were stimulated with an EC₈₀ equivalent concentration of compound in a volume of 5 μ l. Wells were washed after 5 minutes by aspiration of the medium, addition of 100 μ l fresh medium, which was subsequently aspirated and replaced by another 100 μ l fresh medium. This washing step was chosen such that it was just before the maximum Cell Index level. In control wells, medium was pipetted up and down in order to induce similar mechanical cell stress.

Data Analysis

Data analyses were performed using Prism 6.04 (GraphPad software, San Diego, CA, USA). pIC₅₀ values were acquired using non-linear regression curve fitting into a sigmoidal concentration-response curve using the equation: Y=Bottom + (Top-Bottom)/(1+10^(X-LogIC₅₀)). (p)K_i values were acquired from (p)IC₅₀ values using the Cheng-Prusoff equation.⁶⁵ Dissociation rate constants k_{off} were determined using an exponential decay analysis of radioligand binding. Association rate constants k_{on} were determined using the equation $k_{on} = (k_{obs} - k_{off})/[L]$, in which L is the concentration of radioligand used for association experiments and k_{obs} was determined using exponential association analysis.

Journal of Medicinal Chemistry

Association and dissociation rate constants for unlabelled mGlu₂ PAMs were determined by nonlinear regression analysis of competition association data as described by Motulsky and Mahan.⁴⁰

$$K_{A} = k_{1}[L] \cdot 10^{-9} + k_{2}$$

$$K_{B} = k_{3}[I] \cdot 10^{-9} + k_{4}$$

$$S = \sqrt{(K_{A} - K_{B})^{2} + 4 \cdot k_{1} \cdot k_{3} \cdot L \cdot I \cdot 10^{-18}}$$

$$K_{F} = 0.5(K_{A} + K_{B} + S)$$

$$K_{S} = 0.5(K_{A} + K_{B} - S)$$

$$Q = \frac{B_{\max} \cdot k_{1} \cdot L \cdot 10^{-9}}{K_{F} - K_{S}}$$

$$Y = Q \cdot (\frac{k_{4} \cdot (K_{F} - K_{S})}{K_{F} \cdot K_{S}} + \frac{k_{4} - K_{F}}{K_{F}} e^{(-K_{F} \cdot X)} - \frac{k_{4} - K_{S}}{K_{S}} e^{(-K_{S} \cdot X)})$$

Experimental data are reported as k_{on} and k_{off} and the corresponding conversion of k_{off} to residence time (RT) is performed to be consistent with current state of the art.²⁴

Data shown are the mean \pm SEM of at least three individual experiments performed in duplicate unless stated otherwise. Statistical analysis was performed if indicated, using a two-tailed unpaired Student's t-test. Observed differences with p-values <0.05 were considered statistically significant. clogP values were calculated using ChemDraw Professional v15.0.0.106 (PerkinElmer).

ASSOCIATED CONTENT

*S Supporting Information

The Supporting Information is available free of charge on the ACS Publications website and includes: Molecule SMILES strings and bioactivity, additional correlation plots, mGlu₂ PAM response on CHO-K1 WT cells, effect of PAM treatment on total sleep and mGlu receptor selectivity data.

AUTHOR INFORMATION

Corresponding Authors

Adriaan P. IJzerman

Leiden Academic Centre for Drug Research, Leiden University, P.O. Box 9502, 2300 RA

Leiden, The Netherlands

Phone: +31 (0)71 527 4651

E-mail: ijzerman@lacdr.leidenuniv.nl

Gary Tresadern

Janssen Research and Development, Calle Jarama 75A, 45007, Toledo, Spain

Phone: +34 925 24 5777

E-mail: gtresade@its.jnj.com

Notes

The authors declare no competing financial interest.

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENTS

The authors thank Lieve Heylen for technical assistance and Heidi Huysmans for her support of the in vivo studies and analyses.

Funding Sources

This project was financially supported by Vlaams Agentschap Innoveren & Ondernemen project number: 120491.

ABBREVIATIONS

AUC, area under the curve; BCA, bicinchoninic acid; CHO, Chinese hamster ovary; DMEM, Dulbecco's modified Eagle's medium; DMSO, dimethyl sulfoxide; GPCR, G protein-coupled receptor; [³⁵S]GTPγS, guanosine 5'-O-[gamma-thio]triphosphate; mGlu, metabotropic glutamate; PAM, positive allosteric modulator; PBS, phosphate-buffered saline; RT, residence time; SEM, standard error of the mean; SPA, scintillation proximity assay; VFT, venus flytrap domain; 7TM, seven transmembrane domain; (1) Kew, J. N. C.; Kemp, J. A. Ionotropic and Metabotropic Glutamate Receptor Structure and Pharmacology. *Psychopharmacology (Berl)*. **2005**, *179*, 4–29.

(2) Niswender, C. M.; Conn, P. J. Metabotropic Glutamate Receptors: Physiology, Pharmacology, and Disease. *Annu. Rev. Pharmacol. Toxicol.* **2010**, *50*, 295–322.

(3) Muto, T.; Tsuchiya, D.; Morikawa, K.; Jingami, H. Structures of the Extracellular Regions of the Group II/III Metabotropic Glutamate Receptors. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 3759–3764.

(4) Nicoletti, F.; Bockaert, J.; Collingridge, G. L.; Conn, P. J.; Ferraguti, F.; Schoepp, D. D.; Wroblewski, J. T.; Pin, J. P. Metabotropic Glutamate Receptors: From the Workbench to the Bedside. *Neuropharmacology* **2011**, *60*, 1017–1041.

(5) Dunayevich, E.; Erickson, J.; Levine, L.; Landbloom, R.; Schoepp, D. D.; Tollefson, G.
D. Efficacy and Tolerability of an mGlu2/3 Agonist in the Treatment of Generalized Anxiety
Disorder. *Neuropsychopharmacology* 2008, *33*, 1603–1610.

(6) Patil, S. T.; Zhang, L.; Martenyi, F.; Lowe, S. L.; Jackson, K. A.; Andreev, B. V; Avedisova, A. S.; Bardenstein, L. M.; Gurovich, I. Y.; Morozova, M. A.; Mosolov, S. N.; Neznanov, N. G.; Reznik, A. M.; Smulevich, A. B.; Tochilov, V. A.; Johnson, B. G.; Monn, J. A.; Schoepp, D. D. Activation of mGlu2/3 Receptors as a New Approach to Treat Schizophrenia: A Randomized Phase 2 Clinical Trial. *Nat. Med.* **2007**, *13*, 1102–1107.

(7) Keov, P.; Sexton, P. M.; Christopoulos, A. Allosteric Modulation of G Protein-Coupled Receptors: A Pharmacological Perspective. *Neuropharmacology* **2011**, *60*, 24–35.

(8) Trabanco, A. A.; Cid, J. M. mGluR2 Positive Allosteric Modulators: A Patent Review (2009 - Present). *Expert Opin. Ther. Pat.* **2013**, *23*, 629–647.

Journal of Medicinal Chemistry

(9) Johnson, M. P.; Baez, M.; Jagdmann, G. E.; Britton, T. C.; Large, T. H.; Callagaro, D. O.; Tizzano, J. P.; Monn, J. A.; Schoepp, D. D. Discovery of Allosteric Potentiators for the Metabotropic Glutamate 2 Receptor: Synthesis and Subtype Selectivity of N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2- Trifluoroethylsulfonyl)pyrid-3-Ylmethylamine. *J. Med. Chem.* **2003**, *46*, 3189–3192.

(10) Galici, R.; Jones, C. K.; Hemstapat, K.; Nong, Y.; Echemendia, N. G.; Williams, L. C.; de Paulis, T.; Conn, P. J. Biphenyl-Indanone A, a Positive Allosteric Modulator of the Metabotropic Glutamate Receptor Subtype 2, Has Antipsychotic- and Anxiolytic-like Effects in Mice. *J. Pharmacol. Exp. Ther.* **2006**, *318*, 173–185.

(11) Fell, M. J.; Witkin, J. M.; Falcone, J. F.; Katner, J. S.; Perry, K. W.; Hart, J.; Rorick-Kehn, L.; Overshiner, C. D.; Rasmussen, K.; Chaney, S. F.; Benvenga, M. J.; Li, X.; Marlow, D. L.; Thompson, L. K.; Luecke, S. K.; Wafford, K. A.; Seidel, W. F.; Edgar, D. M.; Quets, A. T.; Felder, C. C.; Wang, X.; Heinz, B. A.; Nikolayev, A.; Kuo, M.; Mayhugh, D.; Khilevich, A.; Zhang, D.; Ebert, P. J.; Eckstein, J. A.; Ackermann, B. L.; Swanson, S. P.; Catlow, J. T.; Dean, R. A.; Jackson, K.; Tauscher-Wisniewski, S.; Marek, G. J.; Schkeryantz, J. M.; Svensson, K. A. N-(4-((2-(Trifluoromethyl)-3-Hydroxy-4-(Isobutyryl)phenoxy)methyl)benzyl)-1-Methyl-1H-Imidazole-4-Carboxamide (THIIC), a Novel Metabotropic Glutamate 2 Potentiator with Potential Anxiolytic/antidepressant Properties: In Vivo Profiling Suggests a Link between Behavioral and Central Nervous System Neurochemical Changes. *J. Pharmacol. Exp. Ther.* 2011, *336*, 165–177.

(12) Lavreysen, H.; Langlois, X.; Ahnaou, A.; Drinkenburg, W.; te Riele, P.; Biesmans, I.; Van der Linden, I.; Peeters, L.; Megens, A.; Wintmolders, C.; Cid, J. M.; Trabanco, A. A.; Andrés, J. I.; Dautzenberg, F. M.; Lütjens, R.; Macdonald, G.; Atack, J. R. Pharmacological Characterization of JNJ-40068782, a New Potent, Selective, and Systemically Active Positive Allosteric Modulator of the mGlu2 Receptor and Its Radioligand [3H]JNJ-40068782. *J. Pharmacol. Exp. Ther.* **2013**, *346*, 514–527.

(13) Cook, D.; Brown, D.; Alexander, R.; March, R.; Morgan, P.; Satterthwaite, G.; Pangalos,
M. N. Lessons Learned from the Fate of AstraZeneca's Drug Pipeline: A Five-Dimensional
Framework. *Nat. Rev. Drug Discovery* 2014, *13*, 419–431.

(14) Homepage: http://clinicaltrials.gov. The Effects AZD8529 on Cognition and Negative Symptoms in Schizophrenics: https://clinicaltrials.gov/show/NCT00986531 (accessed December 22, 2016).

(15) Homepage: http://clinicaltrials.gov. The Study of AZD8529 for Smoking Cessation inFemaleSmokers:

https://www.clinicaltrials.gov/ct2/show/NCT02401022?term=AZD8529&rank=1 (accessed April 3, 2017).

(16) Cid, J. M.; Tresadern, G.; Duvey, G.; Lütjens, R.; Finn, T.; Rocher, J.; Poli, S.; Vega, J. A.; de Lucas, A. I.; Matesanz, E.; Linares, M. L.; Andrés, J. I.; Alcazar, J.; Alonso, J. M.; Macdonald, G. J.; Oehlrich, D.; Lavreysen, H.; Ahnaou, A.; Drinkenburg, W.; Mackie, C.; Pype, S.; Gallacher, D.; Trabanco, A. A. Discovery of 1-Butyl-3-Chloro-4-(4-Phenyl-1-Piperidinyl)-(1H)-Pyridone (JNJ-40411813): A Novel Positive Allosteric Modulator of the Metabotropic Glutamate 2 Receptor. *J. Med. Chem.* 2014, *57*, 6495–6512.

(17) Lavreysen, H.; Ahnaou, A.; Drinkenburg, W.; Langlois, X.; Mackie, C.; Pype, S.; Lütjens, R.; Le Poul, E.; Trabanco, A. A.; Cid Nuñez, J. M. Pharmacological and Pharmacokinetic Properties of JNJ-40411813, a Positive Allosteric Modulator of the mGlu2 Receptor. *Pharmacol. Res. Perspect.* **2015**, *3*, e00096.

Journal of Medicinal Chemistry

(18) Lavreysen, H.; Langlois, X.; Ver Donck, L.; Cid Nuñez, J. M.; Pype, S.; Lütjens, R.; Megens, A. Preclinical Evaluation of the Antipsychotic Potential of the mGlu2-Positive Allosteric Modulator JNJ-40411813. *Pharmacol. Res. Perspect.* **2015**, *3*, e00097.

(19) Salih, H.; Anghelescu, I.; Kezic, I.; Sinha, V.; Hoeben, E.; Van Nueten, L.; De Smedt,
H.; De Boer, P. Pharmacokinetic and Pharmacodynamic Characterisation of JNJ-40411813, a
Positive Allosteric Modulator of mGluR2, in Two Randomised, Double-Blind Phase-I Studies. *J. Psychopharmacol.* 2015, *29*, 414–425.

(20) Kent, J. M.; Daly, E.; Kezic, I.; Lane, R.; Lim, P.; De Smedt, H.; De Boer, P.; Van Nueten, L.; Drevets, W. C.; Ceusters, M. Efficacy and Safety of an Adjunctive mGlu2 Receptor Positive Allosteric Modulator to a SSRI/SNRI in Anxious Depression. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **2016**, *67*, 66–73.

(21) http://www.addextherapeutics.com/ investors/press-releases/news-details/article/addex-reports-top-linedata-from-a-successful-phase-2a-clinical-study-with-adx71149-inschizophrenia/ (accessed July 28, 2016).

(22) http:// clinicaltrials.gov/show/NCT01582815 (accessed July 28, 2016).

(23) Swinney, D. C. The Role of Binding Kinetics in Therapeutically Useful Drug Action. *Curr. Opin. Drug Discovery Dev.* **2009**, *12*, 31–39.

(24) Copeland, R. A. The Drug–target Residence Time Model: A 10-Year Retrospective. *Nat. Rev. Drug Discovery* **2016**, *15*, 87–95.

(25) Guo, D.; Heitman, L. H.; IJzerman, A. P. The Role of Target Binding Kinetics in Drug Discovery. *ChemMedChem* **2015**, *10*, 1793–1796.

(26) Swinney, D. C.; Haubrich, B. A.; Liefde, I. Van; Vauquelin, G. The Role of Binding Kinetics in GPCR Drug Discovery. *Curr. Top. Med. Chem.* **2015**, *15*, 2504–2522.

(27) Guo, D.; Heitman, L. H.; IJzerman, A. P. Kinetic Aspects of the Interaction between Ligand and G Protein-Coupled Receptor: The Case of the Adenosine Receptors. *Chem. Rev.* **2017**, *117*, 38–66.

(28) Guo, D.; Hillger, J. M.; IJzerman, A. P.; Heitman, L. H. Drug-Target Residence Time-a Case for G Protein-Coupled Receptors. *Med. Res. Rev.* **2014**, *34*, 856–892.

(29) Dowling, M. R.; Charlton, S. J. Quantifying the Association and Dissociation Rates of Unlabelled Antagonists at the Muscarinic M3 Receptor. *Br. J. Pharmacol.* **2006**, *148*, 927–937.

(30) Tresadern, G.; Bartolome, J. M.; Macdonald, G. J.; Langlois, X. Molecular Properties Affecting Fast Dissociation from the D2 Receptor. *Bioorg. Med. Chem.* **2011**, *19*, 2231–2241.

(31) Kapur, S.; Seeman, P. Does Fast Dissociation From the Dopamine D2 Receptor Explain the Action of Atypical Antipsychotics?: A New Hypothesis. *Am. J. Psychiatry* **2001**, *158*, 360–369.

(32) Yin, N.; Pei, J.; Lai, L. A Comprehensive Analysis of the Influence of Drug Binding Kinetics on Drug Action at Molecular and Systems Levels. *Mol. Biosyst.* **2013**, *9*, 1381–1389.

(33) Vauquelin, G. Effects of Target Binding Kinetics on in Vivo Drug Efficacy: K off, K on and Rebinding. *Br. J. Pharmacol.* **2016**, *173*, 2319–2334.

(34) Schoop, A.; Dey, F. On-Rate Based Optimization of Structure–kinetic Relationship – Surfing the Kinetic Map. *Drug Discovery Today: Technol.* **2015**, *17*, 9–15.

(35) Cid, J. M.; Tresadern, G.; Vega, J. A.; de Lucas, A. I.; Matesanz, E.; Iturrino, L.; Linares, M. L.; Garcia, A.; Andrés, J. I.; Macdonald, G. J.; Oehlrich, D.; Lavreysen, H.; Megens, A.; Ahnaou, A.; Drinkenburg, W.; Mackie, C.; Pype, S.; Gallacher, D.; Trabanco, A. A. Discovery of 3-Cyclopropylmethyl-7-(4-Phenylpiperidin-1-Yl)-8-trifluoromethyl[1,2,4]triazolo[4,3-

Journal of Medicinal Chemistry

A]pyridine (JNJ-42153605): A Positive Allosteric Modulator of the Metabotropic Glutamate 2
Receptor. J. Med. Chem. 2012, 55, 8770–8789.
(36) Tresadern, G.; Cid, J.-M. M.; Trabanco, A. A. QSAR Design of Triazolopyridine mGlu2
Receptor Positive Allosteric Modulators. J. Mol. Graph. Model. 2014, 53, 82–91.
(37) Cid-Nunez, J. M.; De Lucas Olivares, A. I.; Trabanco-Suarez, A. A.; MacDonald, G. J. 7Aryl-1,2,4-triazolo[4,3-A]pyridine Derivatives and Their Use as Positive Allosteric Modulators of mGluR2 Receptors. WO2010130423 A1, 2010.
(38) Cid, J. M.; Duvey, G.; Tresadern, G.; Nhem, V.; Furnari, R.; Cluzeau, P.; Vega, J. A.; de
Lucas, A. I.; Matesanz, E.; Alonso, J. M.; Linares, M. L.; Andrés, J. I.; Poli, S. M.; Lutjens, R.;
Himogai, H.; Rocher, J.; Macdonald, G. J.; Oehlrich, D.; Lavreysen, H.; Ahnaou, A.;
Drinkenburg, W.; Mackie, C.; Trabanco, A. A. Discovery of 1,4-Disubstituted 3-Cyano-2Pyridones: A New Class of Positive Allosteric Modulators of the Metabotropic Glutamate 2
Receptor. J. Med. Chem. 2012, 55, 2388–2405.

(39) Doornbos, M. L. J.; Pérez-Benito, L.; Tresadern, G.; Mulder-Krieger, T.; Biesmans, I.; Trabanco, A. A.; Cid, J. M.; Lavreysen, H.; IJzerman, A. P.; Heitman, L. H. Molecular Mechanism of Positive Allosteric Modulation of the Metabotropic Glutamate Receptor 2 by JNJ-46281222. *Br. J. Pharmacol.* **2016**, *173*, 588–600.

(40) Motulsky, H. J.; Mahan, L. C. The Kinetics of Competitive Radioligand Binding Predicted Mass Action by the Law of Mass Action. *Mol. Pharmacol.* **1984**, *25*, 1–9.

(41) Xia, L.; de Vries, H.; IJzerman, A. P.; Heitman, L. H. Scintillation Proximity Assay (SPA) as a New Approach to Determine a Ligand's Kinetic Profile. A Case in Point for the Adenosine A1 Receptor. *Purinergic Signalling* **2016**, *12*, 115–126.

(42) Farinha, A.; Lavreysen, H.; Peeters, L.; Russo, B.; Masure, S.; Trabanco, A. A.; Cid, J.; Tresadern, G. Molecular Determinants of Positive Allosteric Modulation of the Human Metabotropic Glutamate Receptor 2. *Br. J. Pharmacol.* **2015**, *172*, 2383–2396.

(43) Pérez-Benito, L.; Doornbos, M. L. J.; Cordomí, A.; Peeters, L.; Lavreysen, H.; Pardo, L.;
Tresadern, G. Molecular Switches of Allosteric Modulation of the Metabotropic Glutamate 2
Receptor. *Structure* 2017, *25*, 1–10.

(44) Lovering, F.; Bikker, J.; Humblet, C. Escape from Flatland: Increasing Saturation as an Approach to Improving Clinical Success. *J. Med. Chem.* **2009**, *52*, 6752–6756.

(45) Wacker, D.; Wang, S.; McCorvy, J. D.; Betz, R. M.; Venkatakrishnan, A. J.; Levit, A.; Lansu, K.; Schools, Z. L.; Che, T.; Nichols, D. E.; Shoichet, B. K.; Dror, R. O.; Roth, B. L. Crystal Structure of an LSD-Bound Human Serotonin Receptor. *Cell* **2017**, *168*, 377–389.e12.

(46) Sykes, D. A.; Dowling, M. R.; Charlton, S. J. Exploring the Mechanism of Agonist Efficacy: A Relationship between Efficacy and Agonist Dissociation Rate at the Muscarinic M3 Receptor. *Mol. Pharmacol.* **2009**, *76*, 543–551.

(47) Guo, D.; Mulder-Krieger, T.; IJzerman, A. P.; Heitman, L. H. Functional Efficacy of Adenosine A2A Receptor Agonists Is Positively Correlated to Their Receptor Residence Time. *Br. J. Pharmacol.* **2012**, *166*, 1846–1859.

(48) Deyrup, M. D.; Nowicki, S. T.; Richards, N. G.; Otero, D. H.; Harrison, J. K.; Baker, S.

P. Structure-Affinity Profile of 8-Hydroxycarbostyril-Based Agonists That Dissociate Slowly from the beta2-Adrenoceptor. *Naunyn-Schmiedebergs's Arch. Pharmacol.* **1999**, *359*, 168–177.

(49) Mould, R.; Brown, J.; Marshall, F. H.; Langmead, C. J. Binding Kinetics Differentiates Functional Antagonism of Orexin-2 Receptor Ligands. *Br. J. Pharmacol.* **2014**, *171*, 351–363.

Journal of Medicinal Chemistry

(50) Sykes, D. a; Charlton, S. J. Slow Receptor Dissociation Is Not a Key Factor in the Duration of Action of Inhaled Long-Acting β 2-Adrenoceptor Agonists. *Br. J. Pharmacol.* **2012**, *165*, 2672–2683.

(51) Yu, Z.; Van Veldhoven, J. P. D.; Louvel, J.; 'T Hart, I. M. E.; Rook, M. B.; Van Der Heyden, M. A. G.; Heitman, L. H.; IJzerman, A. P. Structure-Affinity Relationships (SARs) and Structure-Kinetics Relationships (SKRs) of Kv11.1 Blockers. *J. Med. Chem.* **2015**, *58*, 5916–5929.

(52) Yu, Z.; IJzerman, A. P.; Heitman, L. H. K v 11.1 (hERG)-Induced Cardiotoxicity: A Molecular Insight from a Binding Kinetics Study of Prototypical K v 11.1 (hERG) Inhibitors. *Br. J. Pharmacol.* 2015, *172*, 940–955.

(53) de Witte, W. E. A.; Danhof, M.; van der Graaf, P. H.; de Lange, E. C. M. In Vivo Target Residence Time and Kinetic Selectivity: The Association Rate Constant as Determinant. *Trends Pharmacol. Sci.* **2016**, *37*, 831–842.

(54) Vauquelin, G.; Charlton, S. J. Long-Lasting Target Binding and Rebinding as Mechanisms to Prolong in Vivo Drug Action. *Br. J. Pharmacol.* **2010**, *161*, 488–508.

(55) Vauquelin, G. On the "micro"-Pharmacodynamic and Pharmacokinetic Mechanisms That Contribute to Long-Lasting Drug Action. *Expert Opin. Drug Discovery* **2015**, *10*, 1085–1098.

(56) Vauquelin, G. Cell Membranes... and How Long Drugs May Exert Beneficial Pharmacological Activity in Vivo. *Br. J. Clin. Pharmacol.* **2016**, *82*, 673–682.

(57) Perry, D. C.; Mullis, K. B.; Øie, S.; Sadée, W. Opiate Antagonist Receptor Binding in Vivo: Evidence for a New Receptor Binding Model. *Brain Res.* **1980**, *199*, 49–61.

(58) Yu, N.; Atienza, J. M.; Bernard, J.; Blanc, S.; Zhu, J.; Wang, X.; Xu, X.; Abassi, Y. Real-Time Monitoring of Morphological Changes in Living Cells by Electronic Cell Sensors Arrays: An Approach to Study G Protein-Coupled Receptors. *Anal. Chem.* **2006**, *78*, 35–43.

(59) Hillger, J. M.; Diehl, C.; van Spronsen, E.; Boomsma, D. I.; Slagboom, P. E.; Heitman,
L. H.; IJzerman, A. P. Getting Personal: Endogenous Adenosine Receptor Signaling in
Lymphoblastoid Cell Lines. *Biochem. Pharmacol.* 2016, *115*, 114–122.

(60) Xi, B.; Yu, N.; Wang, X.; Xu, X.; Abassi, Y. A. The Application of Cell-Based Label-Free Technology in Drug Discovery. *Biotechnol. J.* **2008**, *3*, 484–495.

(61) Ahnaou, A.; Dautzenberg, F. M.; Geys, H.; Imogai, H.; Gibelin, A.; Moechars, D.; Steckler, T.; Drinkenburg, W. H. I. M. Modulation of Group II Metabotropic Glutamate Receptor (mGlu2) Elicits Common Changes in Rat and Mice Sleep-Wake Architecture. *Eur. J. Pharmacol.* **2009**, *603*, 62–72.

(62) Ahnaou, A.; De Boer, P.; Lavreysen, H.; Huysmans, H.; Sinha, V.; Raeymaekers, L.; Van De Casteele, T.; Cid, J. M.; Van Nueten, L.; MacDonald, G. J.; Kemp, J. A.; Drinkenburg, W. H.
I. M. Translational Neurophysiological Markers for Activity of the Metabotropic Glutamate Receptor (mGluR2) Modulator JNJ-40411813: Sleep EEG Correlates in Rodents and Healthy Men. *Neuropharmacology* 2016, *103*, 290–305.

(63) Cuisset, L.; Tichonicky, L.; Jaffray, P.; Delpech, M. The Effects of Sodium Butyrate on Transcription Are Mediated through Activation of a Protein Phosphatase. *J. Biol. Chem.* **1997**, *272*, 24148–24153.

(64) Hillger, J. M.; Schoop, J.; Boomsma, D. I.; Eline Slagboom, P.; IJzerman, A. P.; Heitman, L. H. Whole-Cell Biosensor for Label-Free Detection of GPCR-Mediated Drug Responses in Personal Cell Lines. *Biosens. Bioelectron.* **2015**, *74*, 233–242.

1
2
3
4
5
0
6
1
8
9
10
11
12
13
1/
14
10
16
17
18
19
20
21
22
22
23
24
25
26
27
28
29
20
30
31
32
33
34
35
36
37
20
30
39
40
41
42
43
44
45
46
47
41 40
48
49
50
51
52
53
54
55
55
30
57
58

(65) Cheng, Y.; Prusoff, W. H. Relationship between the Inhibition Constant (KI) and the Concentration of Inhibitor Which Causes 50 per Cent Inhibition (I50) of an Enzymatic Reaction. *Biochem. Pharmacol.* **1973**, *22*, 3099–3108.

FIGURES



Figure 1. Structures of reference mGlu₂ PAMs. The position of the tritium label of $[^{3}H]$ -7 is denoted by *.



Figure 2. Kinetic map of all tested compounds. The dissociation rate $(pk_{off}, k_{off}: min^{-1})$ is plotted on the *y*-axis; the association rate $(\log k_{on}, k_{on}: M^{-1}min^{-1})$ on the *x*-axis. Identical affinity (pK_D) values may result from a different combination of k_{on} and k_{off} ($K_D = k_{off} / k_{on}$, diagonal blue lines). Horizontal lines indicating a RT of 10 and 20 minutes divide the compounds in short, medium and long RT.



Figure 3. Correlation between affinity (pK_i) and potency (pEC₅₀) (A); affinity determined from $[^{3}H]$ -7 displacement assays (pK_i) and affinity determined based on kinetic parameters k_{on} and k_{off} obtained from $[^{3}H]$ -7 competition association experiments (pK_D) (B); affinity (pK_D) and dissociation rate (p k_{off}) (C); affinity (pK_D) and association rate (log k_{on}) (D); association rate (log k_{on}) and partition coefficient (clogP) (E).





Figure 4. Concentration-dependent effects of 37 (A), 39 (B) and 44 (C) on CHO-K1 cells stably expressing the human mGlu₂ receptor. A representative example is shown of a baseline-corrected response, which was repeated three times in duplicate. (D) Concentration-effect curve of 37, 39 and 44, derived from the AUC up to 150 minutes. Data are from three individual experiments performed in duplicate and are expressed as percentage of maximum AUC. Responses in a label-free, impedance-based assay induced by an EC₈₀ equivalent concentration of 37 (600 nM) (E), 39 (200 nM) (F) and 44 (60 nM) (G) with and without washing-step induced 5 minutes after stimulation. A representative example is shown of a baseline-corrected response, which was repeated three times in duplicate. (H) Ligand-induced response that is left after washing step. Data are expressed as the ratio between the AUC of the trace of the washed and unwashed wells.

Data are from three individual experiments performed in duplicate and are expressed as the mean

± SEM. * p < 0.05.



Figure 5. Average amount of REM sleep in minutes for 24 consecutive 5-min period during the first 2 hours of the recording session after oral administration of **9** (top panel) and **20** (bottom panel), each at 1, 3, 10 mg/kg, p.o. to Sprague-Dawley rats (n=8 for each condition). Black lines indicate control vehicle condition (20% CD), while the green, blue, and red lines indicate doses 1, 3, 10 mg/kg, respectively. Error flags indicate SEM values: please note that the higher time resolution of 5 minute bins also causes REM sleep changes to be expressed more dynamically and less 'consistent' as compared to reduction score averaged over the whole 2 hour post-administration period. * indicate p < 0.05 vehicle compared to 10 mg/kg, Mixed Model ANOVA.

SCHEMES





^a Reagents and conditions: (a) Pd(PPh₃)₄, NaHCO₃, H₂O/1,4-dioxane, 150°C, 5-30 min, microwave.

Table 1.	Comparison	of kinetic para	meters of 7	determined	by two	different methods ^a
	1	1			2	

Assay	$K_D(\mathrm{nM})^{\mathrm{b}}$	$k_{on}(\mathrm{M}^{-1}\mathrm{min}^{-1})$	$k_{off}(\min^{-1})$	RT (min)
Association and dissociation ^c	1.1 ± 0.22	$(7.1 \pm 0.43) \ge 10^7$	0.080 ± 0.015	12.5 ± 2.3
Competition association ^d	2.5 ± 0.71	$(2.3 \pm 0.60) \ge 10^7$	0.058 ± 0.005	17.3 ± 1.6

^aValues represent the mean \pm SEM of at least three individual experiments performed in duplicate. ^bKinetic K_D values, defined by $K_D = k_{off} / k_{on}$. ^cAssociation and dissociation rate constants of [³H]-7 in a classical radioligand kinetic binding assay at 28°C using SPA. ^dAssociation and dissociation rate constants of 7 measured in a competition association assay at 28°C using SPA.

10						
11 12	pEC ₅₀	pK_i^{b}	$k_{on} \left(\mathrm{M}^{-1} \min^{-1} \right)^{\mathrm{c}}$	$k_{off} (\min^{-1})^{d}$	RT (min) ^e	$K_D \left(\mathrm{nM} \right)^{\mathrm{f}}$
13 2	7.03 ± 0.14^a	7.22 ± 0.15	$(4.6 \pm 2.3) \ge 10^6$	0.086 ± 0.022	11.6 ± 2.9	190 ± 100
14 15 3	6.97 ± 0.08^a	7.10 ± 0.04	$(1.2 \pm 0.38) \ge 10^6$	0.120 ± 0.020	8.4 ± 1.4	100 ± 36
¹⁶ 4	7.1 ± 0.1^{a}	7.58 ± 0.08	$(1.7 \pm 0.70) \ge 10^6$	0.090 ± 0.015	11.1 ± 1.9	51 ± 22
18 5	6.25 ± 0.06	6.43 ± 0.03	$(2.1 \pm 0.65) \ge 10^5$	0.128 ± 0.040	7.8 ± 2.4	600 ± 260
19 <u>20</u>	As previo	ously described b	y a) Farinha et al. (20	$(15)^{42}$, b) Doornbos	<i>et al.</i> $(2016)^{39}$.	^{,f} Values

As previously described by a) Farinha *et al.* (2015)⁴², b) Doornbos *et al.* (2016)³⁹, ^{c,d,e,t} Values represent the mean \pm SEM of at least three individual experiments. ^eRT = $1/k_{off}$. ^f Kinetic K_D values, defined by $K_D = k_{off} / k_{on}$.

Table 3. Functional activity (pEC₅₀), affinity (pK_i) and kinetic parameters (k_{on} , k_{off} , RT) for

 mGlu₂ PAMs 8-23



18 19 20	R^1	R ²	R ³	R^4	pEC ₅₀ ^a	pK_i^b	$k_{on}(\mathrm{M}^{-1}\mathrm{min}^{-1})^{\mathrm{c}}$	$k_{off}(\min^{-1})^{d}$	RT (min) ^e	$K_D \left(\mathrm{nM} \right)^{\mathrm{f}}$
21 22 ₈ 23	Cl	F	*	CH ₂ -(c-Pr)	7.39 (7.35;7.43)	8.71 ± 0.12	$(6.3 \pm 1.5) \ge 10^6$	0.097 ± 0.024	10.3 ± 2.5	15.3 ± 5.3
24 25 ₉ 26	CF ₃	F	*	CH ₂ -(c-Pr)	8.06 ± 0.14	8.59 ± 0.10	$(1.3 \pm 0.49) \ge 10^7$	0.033 ± 0.009	30.1 ± 8.2	2.5 ± 1.2
28 29 ¹⁰	Me	F	*	CH ₂ -(c-Pr)	7.11 (7.02; 7.19)	7.66 ± 0.33	$(8.6 \pm 3.3) \ge 10^6$	0.086 ± 0.013	11.6 ± 1.8	10.0 ± 4.1
30 31 32 ¹¹	i-Pr	F	*	CH ₂ -(c-Pr)	7.55 (7.55; 7.54)	7.60 ± 0.24	$(6.5 \pm 1.4) \ge 10^6$	0.070 ± 0.017	14.2 ± 3.5	10.7 ± 3.5
34 35 12 36	Cl	Cl	*	CH ₂ -(c-Pr)	7.91 (7.71; 8.11)	8.22 ± 0.13	$(6.2 \pm 2.4) \ge 10^7$	0.061 ± 0.009	16.3 ± 2.4	1.0 ± 0.40
37 38 13 39	CF ₃	Cl	*	CH ₂ -(c-Pr)	8.54 (8.31; 8.78)	8.82 ± 0.22	$(6.1 \pm 2.4) \ge 10^6$	0.044 ± 0.006	22.7 ± 2.9	7.3 ± 3.0
40 41 14 42	Me	Cl	*	CH ₂ -(c-Pr)	7.76 ±0.26	7.83 ± 0.25	$(5.5 \pm 1.7) \ge 10^6$	0.096 ± 0.021	10.5 ± 2.3	17.4 ± 6.5
43 44 15 45	Cl	F	* N	CH ₂ -(c-Pr)	6.60 (6.64; 6.55)	7.20 ± 0.14	$(2.6 \pm 0.72) \ge 10^6$	0.096 ± 0.031	10.4 ± 3.4	37.6 ± 16
46 47 16 48	CF ₃	F	*	CH ₂ -(c-Pr)	7.25 ± 0.32	8.15 ± 0.05	$(9.8 \pm 5.2) \ge 10^6$	0.090 ± 0.015	11.1 ± 1.8	9.2 ± 5.1
49 ⁵⁰ 17 51	Cl	F	* N	CH ₂ -(c-Pr)	7.00 ± 0.10	7.73 ± 0.08	$(3.0 \pm 0.97) \ge 10^6$	0.069 ± 0.011	14.4 ± 2.3	23.3 ± 8.4
52 53 54 18 55 56	CF ₃	F	* N	CH ₂ -(c-Pr)	7.19 ± 0.10	8.00 ± 0.16	$(8.3 \pm 0.60) \ge 10^6$	0.095 ± 0.020	10.5 ± 2.2	11.5 ± 2.5
57										

ACS Paragon Plus Environment

2 3			. ^							
4 5 19 6 7	Cl	F		CH ₂ -(c-Pr)	7.06 ± 0.07	7.27 ± 0.20	$(1.7 \pm 0.82) \ge 10^6$	0.073 ± 0.004	13.7 ± 0.8	42.7 ± 21
8 9 20 10	CF ₃	F	*N	CH ₂ -(c-Pr)	7.73 ± 0.14	8.44 ± 0.07	$(2.2 \pm 0.42) \ge 10^7$	0.189 ± 0.082	5.3 ± 2.3	8.6±4.1
11 12 ₂₁ 13	Cl	F	*	CH ₂ -CF ₃	7.53 ± 0.11	8.10 ± 0.24	$(2.9 \pm 0.49) \ge 10^7$	0.096 ± 0.013	10.5 ± 1.5	3.3 ± 0.7
14 15 ₂₂ 16 17	Cl	F	*	CH ₂ -OEt	7.13 (7.27; 6.99)	8.19 ± 0.10	$(3.0 \pm 0.11) \ge 10^6$	0.076 ± 0.004	13.2 ± 0.7	25.2 ± 1.6
18 ₂₃ 19 ²	CF ₃	F	*	CH ₂ -OEt	7.53 ± 0.19	7.93 ± 0.26	$(3.6 \pm 2.1) \ge 10^7$	0.075 ± 0.006	13.4 ± 1.0	2.1 ± 1.2

^aValues represent the mean followed by individual values in parentheses when of two experiments and the mean \pm SEM when of three or more experiments. ^{b,c,d,e,f} Values represent the mean \pm SEM of at least three individual experiments, all performed in duplicate. ^eRT = $1/k_{off}$. ^f Kinetic K_D values, defined by $K_D = k_{off}/k_{on}$.

Table 4. Functional activity (pEC₅₀), affinity (pK_i) and kinetic parameters (k_{on} , k_{off} , RT) for

mGlu₂ PAMs 24-30

 R^{2}

	\mathbb{R}^1	\mathbb{R}^2	R ³	pEC ₅₀ ^a	$pK_i^{\ b}$	$k_{on} (\mathrm{M}^{-1}\mathrm{min}^{-1})^{\mathrm{c}}$	$k_{off}(\min^{-1})^{d}$	RT (min) ^e	$K_D \left(\mathrm{nM} \right)^{\mathrm{f}}$
24	Cl	F	i-Pr	6.65 (6.43; 6.88)	7.21 ± 0.07	$(5.0 \pm 1.2) \ge 10^6$	0.101 ± 0.009	9.9 ± 0.9	20.3 ± 5.4
25	CF_3	F	i-Pr	7.03 (7.44; 6.62)	8.05 ± 0.10	$(4.6 \pm 1.1) \ge 10^7$	0.100 ± 0.029	10.0 ± 2.9	2.2 ± 0.8
26	CF_3	Cl	i-Pr	7.76 (7.55; 7.96)	8.41 ± 0.07	$(3.6 \pm 1.3) \ge 10^7$	0.121 ± 0.035	8.2 ± 2.4	3.4 ± 1.6
27	Cl	F	c-Pr	7.20 (7.19; 7.21)	7.99 ± 0.05	$(4.8 \pm 1.5) \ge 10^6$	0.061 ± 0.006	16.4 ± 1.7	12.7 ± 4.1
28	CF_3	F	c-Pr	7.52 (7.60; 7.43)	8.44 ± 0.07	$(7.7 \pm 4.8) \ge 10^7$	0.086 ± 0.018	11.9 ± 2.3	1.1 ± 0.7
29	CF ₃	Cl	c-Pr	8.11 (8.04; 8.18)	8.65 ± 0.11	$(7.7 \pm 3.8) \ge 10^7$	0.088 ± 0.007	11.3 ± 0.9	1.2 ± 0.6
30	Cl	Cl	c-Pr	7.53 (7.54; 7.51)	8.17 ± 0.03	$(1.4 \pm 0.35) \ge 10^7$	0.084 ± 0.016	11.9 ± 2.3	6.1 ± 1.9

^aValues represent the mean followed by individual values in parentheses. ^{b,c,d,e,f} Values represent the mean \pm SEM of at least three individual experiments. ^eRT = $1/k_{off}$. ^f Kinetic K_D values, defined by $K_D = k_{off}/k_{on}$.

Journal of Medicinal Chemistry

 Table 5. Functional activity (pEC₅₀), affinity (pK_i) and kinetic parameters (k_{on} , k_{off} , RT) for mGlu₂ PAMs **31-35**



	\mathbf{R}^1	R^2	Х	pEC_{50}^{a}	pK _i ^b	$k_{on} (M^{-1} \min^{-1})^{c}$	$k_{off} (\min^{-1})^{d}$	RT (min) ^e	$K_D \left(\mathrm{nM} \right)^{\mathrm{f}}$
31	CF_3	F	0	7.39 (7.40; 7.37)	7.93 ± 0.14	$(1.1 \pm 0.40) \ge 10^7$	0.173 ± 0.007	5.8 ± 0.2	16.0 ± 5.9
32	CF_3	Н	0	7.24 ± 0.07	8.06 ± 0.16	$(1.4 \pm 0.86) \ge 10^7$	0.125 ± 0.013	8.0 ± 0.8	8.7 ± 5.4
33	CF ₃	F	NH	8.23 (8.13; 8.33)	8.69 ± 0.16	$(1.5 \pm 0.94) \ge 10^7$	0.044 ± 0.005	22.7 ± 2.8	2.9 ± 1.9
34	CF ₃	Н	NH	7.82 ± 0.05	8.50 ± 0.09	$(3.4 \pm 1.3) \ge 10^7$	0.069 ± 0.012	14.6 ± 2.6	2.0 ± 0.8
35	Cl	Н	NH	6.82 (6.94; 6.71)	8.13 ± 0.08	$(6.2 \pm 1.6) \ge 10^6$	0.097 ± 0.022	10.3 ± 2.3	15.6 ± 5.3

^aValues represent the mean followed by individual values in parentheses when of two experiments and the mean \pm SEM when of three or more experiments. ^{b,c,d,e,f} Values represent the mean \pm SEM of at least three individual experiments, all performed in duplicate. ^eRT = $1/k_{off}$. ^f Kinetic K_D values, defined by $K_D = k_{off}/k_{on}$.

Table 6. Functional activity (pEC₅₀), affinity (pK_i) and kinetic parameters (k_{on}, k_{off}, RT) for

6 mGlu ₂ PAMs 36-48 7 8 9 10 11 12 13 14 15 16 17							(R	R^1	N−N N R ⁴				
18- 19		\mathbb{R}^1	\mathbb{R}^2	R ³	\mathbb{R}^4	Х	Y	pEC ₅₀ ^a	pK_i^{b}	$k_{on} (\mathrm{M}^{-1}\min^{-1})^{\mathrm{c}}$	$k_{off} (\min^{-1})^d$	RT (min) ^e	$K_D \left(\mathrm{nM} \right)^{\mathrm{f}}$
20 21	36	Cl	-	-	CH ₂ -(c-Pr)	0	0	6.65 (6.54; 6.77)	7.64 ± 0.01	$(3.3 \pm 1.4) \ge 10^6$	0.092 ± 0.014	10.9 ± 1.7	28.1 ± 13
22 23	37	CF ₃	-	-	CH ₂ -(c-Pr)	0	0	7.42 (7.40; 7.45)	8.28 ± 0.18	$(2.0 \pm 0.61) \ge 10^7$	0.241 ± 0.063	4.1 ± 1.1	12.3 ± 5.0
24	38	Cl	-	-	CH ₂ -(c-Pr)	NH	0	6.83 (6.77; 6.89)	7.41 ± 0.21	$(7.0 \pm 2.4) \ge 10^6$	0.136 ± 0.023	7.3 ± 1.2	19.5 ± 7.5
25 26	39	CF ₃	-	-	CH ₂ -(c-Pr)	NH	0	7.73 (7.77; 7.68)	8.30 ± 0.06	$(1.8 \pm 0.61) \ge 10^7$	0.067 ± 0.008	15.0 ± 1.8	3.8 ± 1.4
27 28	40	CF ₃	-	-	CH ₂ -(c-Pr)	CH ₂ -NH	0	6.20 (6.23; 6.18)	7.21 ± 0.06	$(1.7 \pm 0.18) \ge 10^6$	0.091 ± 0.018	11.0 ± 2.1	54 ± 12
29	41	Cl	Н	OH	CH ₂ -(c-Pr)	NH	С	7.48 ± 0.22	8.52 ± 0.08	$(6.7 \pm 1.3) \ge 10^6$	0.084 ± 0.011	11.8 ± 1.6	12.7 ± 3.0
30 31	42	CF ₃	Н	ОН	CH ₂ -(c-Pr)	NH	С	8.03 ± 0.12	8.66 ± 0.01	$(1.1 \pm 0.09) \ge 10^7$	0.033 ± 0.002	30.8 ± 1.7	3.0 ± 0.3
32 33	43	CF ₃	OH	Н	CH ₂ -(c-Pr)	NH	С	7.38 ± 0.08	8.03 ± 0.17	$(5.7 \pm 1.3) \ge 10^6$	0.058 ± 0.014	17.1 ± 4.0	10.3 ± 3.4
34	44	CF ₃	ОН	c-Pr	CH ₂ -(c-Pr)	NH	С	8.95 (9.00; 8.90)	9.05 ± 0.10	$(1.4 \pm 0.16) \ge 10^7$	0.020 ± 0.005	49.5 ± 12.4	1.5 ± 0.4
36	45	Cl	Н	ОН	CH ₂ -CF ₃	NH	С	7.38 (7.41; 7.34)	8.22 ± 0.20	$(6.7 \pm 1.3) \ge 10^6$	0.074 ± 0.015	13.6 ± 2.8	10.8 ± 3.1
37 38	46	CF ₃	OH	Н	CH ₂ -(c-Pr)	0	С	7.60 (7.64; 7.57)	8.72 ± 0.15	$(1.2 \pm 0.31) \ge 10^7$	0.036 ± 0.001	28.1 ± 0.7	3.0 ± 0.8
39 40	47	Cl	Н	ОН	CH ₂ -CF ₃	0	С	6.80 (6.73; 6.88)	7.94 ± 0.04	$(8.3 \pm 3.6) \ge 10^6$	0.137 ± 0.049	7.3 ± 1.2	16.6±9.3
41	48	Cl	ОН	Н	CH ₂ -CF ₃	0	С	7.01 (7.03; 6.99)	7.49 ± 0.21	$(6.9 \pm 1.5) \ge 10^6$	0.056 ± 0.001	17.7 ± 0.2	8.1 ± 1.7
42- 43			aV	alues	represent	the me	an f	followed by	individual	values in par	entheses wh	nen of tw	0

"Values represent the mean followed by individual values in parentheses when of two experiments and the mean \pm SEM when of three or more experiments.^{b,c,d,e,f} Values represent the mean \pm SEM of at least three individual experiments, all performed in duplicate. ^eRT = $1/k_{off}$.^f Kinetic K_D values, defined by $K_D = k_{off}/k_{on}$.







42x22mm (300 x 300 DPI)