Journal of Chemical & Engineering Data

Potentiometric and Spectrophotometric Studies on the Binding Ability of a Flexible Tripodal Catecholamine Ligand toward Iron(III)

Suban K. Sahoo,^{*,†} Minati Baral,[‡] and B. K. Kanungo^{*,§}

[†]Department of Applied Chemistry, S.V. National Institute of Technology (SVNIT), Surat, Gujrat, India

[‡]Department of Chemistry, National Institute of Technology (NIT), Kurukshetra, Haryana, India

[§]Department of Chemistry, Sant Longowal Institute of Engineering & Technology (SLIET), Longowal, Punjab, India

ABSTRACT: The tripodal catecholamine ligand N^1, N^3, N^5 -tris(2-(2,3-dihydroxybenzylamino)ethyl)cyclohexane-1,3,5-tricarboxamide (CYCOENCAT, L) has been synthesized and characterized. The ligand was investigated as a chelator for Fe(III) in an aqueous medium of 0.1 M KCl at (25 ± 1) °C by potentiometric and spectrophotometric methods. Six protonation constants for the ligand were determined and were used as input data to determine the stability constants of the metal complexes. The stability constants for the FeLH₃, FeLH₂, FeLH, and FeL complexes are reported. The ligand showed the potential to form a tris(catecholate) type complex with a pFe value of 24.76 at pH = 7.4.

1. INTRODUCTION

Biomimetic approaches in research have gained immense interest in recent years due to the interface between chemistry and biology. Many biomolecules are identified in living systems, which act as chelators for the assimilation and transport of metal ions. These natural chelators have received considerable interest due to their selective and strong binding efficiency for a specific metal ion in the presence of other metal ions in living systems. Therefore, to design new chelators for a metal ions, effort has been made to mimic the molecular structure and binding sites of natural chelators through simple synthetic molecules.¹ Also, such synthetic biomimetic molecules are implemented for metal ion sequestration due to their similar selective and strong binding properties toward the metal ion. One of the most studied biomolecules for designing biomimetic synthetic chelators for iron(III) is enterobactin (Scheme 1), which is produced and excreted by bacteria in iron deficient media to bind and assimilate extracellular iron.² It contains three catechol groups appended to tripodal cyclic L-serine through an amide linkage, which has been found to be the best iron-chelating agent with the highest formation constant (log K = 52).³ The enormous iron(III) complexing ability and effective selectivity has led to the synthesis of new biomimetic analogs containing three catechol units in the tripod for: (i) their potential applications as clinical iron removal agents (iron overload is one of the most common types of poisoning), 4 (ii) the use of their iron complexes in agriculture for the prevention or treatment of chlorosis,⁵ and (iii) to mimic the essential structural features that are responsible for the different biological functions of natural compounds.⁶

In this communication, a novel biomimetic tripodal catecholamine ligand N^1 , N^3 , N^5 -tris(2-(2,3-dihydroxybenzylamino)ethyl)cyclohexane-1,3,5-tricarboxamide (CYCOENCAT, L) was investigated as a chelator for iron(III) in an aqueous medium of 0.1 M KCl at (25 \pm 1) °C by potentiometric and spectrophotometric methods. The stability constants of the various complexes and their possible structures in solution are explained.

2. EXPERIMENTAL SECTION

2.1. Materials and Measurements. The triamine CYCOEN was synthesized by following a reported method.⁷ All chemicals required for the synthesis: 1,3,5-benzenetricarboxylic acid, platinum dioxide, ethylenediamine, 2,3-dihydroxybenzaldehyde, and sodium borohydride were obtained from Sigma-Aldrich and were used directly. Anhydrous FeCl₃, absolute ethanol, methanol, KOH, HCl, and KCl were obtained from Ranbaxy Chemicals Ltd., India.

Proton NMR spectra were recorded on a Bruker DPX-300 spectrometer in DMSO- d_6 or D₂O, and chemical shifts were reported relative to Me₄Si. IR (KBr pellets, (450 to 4000) cm⁻¹) spectra were recorded on a Perkin-Elmer RX I FT-IR spectrometer. The electronic spectra were recorded on an Agilent-8453 diode array spectrometer. Elemental analyses were determined for C, H, and N using the Exeter Analytical CE-440.

2.2. Synthesis of CYCOENCAT. The Schiff base intermediate was prepared by dropwise addition of 2,3-dihydroxybenzaldehyde (0.39 g, 28.24 mmol) in 5 mL of absolute ethanol to a magnetically stirred suspended solution of CYCOEN (0.32 g, 9.33 mmol) in 10 mL of absolute ethanol. A bright yellow coloration appeared immediately. The mixture was refluxed for one hour. A dark yellow solution was obtained. On cooling in a freezer, a dark yellow precipitate was obtained, which was filtered off and washed with cold ethanol followed by ether and then dried in a vacuum. Yield = 0.49 g (75 %); IR (KBr pellet, cm⁻¹): 3372, 3215, 2918, 2717, 1623, 1581, 1512, 1445, 1354, 1310, 1235, 1155, 1115, 1021, 962, 851, 774, 652, and 636. ¹H NMR (DMSO-*d*₆, δ ppm): 1.58 (3H, d), 1.83 (3H, d), 2.21 (3H, t), 3.42 (6H, t), 3.69 (6H, t), 6.62 (3H, m), 6.76 (3H, q), 6.86 (3H, m), and 8.30 (3H, s).

To the suspension of Schiff base (0.50 g, 0.71 mmol) in 50 mL of dry methanol, sodium borohydride (0.22 g, 5.81 mmol) was

Received:	December 25, 2010
Accepted:	April 15, 2011
Published:	May 02, 2011

Scheme 1. Molecular Structures of Enterobactin and CYCOENCAT (L)



added with stirring over a period of two hours under nitrogen atmosphere. When the solid phase disappeared and the solution became colorless, 2 mL of concentrated HCl taken in 10 mL methanol was added and kept overnight at -40 °C. A white residue was removed by filtration. The solvent was evaporated under reduced pressure. The crude product obtained was dissolved in dry methanol and evaporated. This process was repeated 3-4 times, and then the residue was dissolved again in dry methanol and to it decolorizing charcoal was added. After vigorous shaking, the charcoal was filtered off. The filtrate was collected, and the solvent was evaporated under reduced pressure. A white product was obtained, which was dried under vacuum. Yield = 0.39 g (65 %); IR (KBr pellet, cm⁻¹): 3374, 3220, 2922, 2718, 1634, 1576, 1504, 1446, 1356, 1304, 1240, 1160, 1126, 1066, 1020, 960, 838, 764, 696, 640, and 616. ¹H NMR (D₂O, δ ppm): 1.35 (3H, q), 1.83 (3H, d), 2.38 (3H, t), 3.12 (6H, t), 3.46 (6H, t), 4.15 (s, 3H), 6.70-6.85 (6H, m), and 6.87 (3H, d). Elemental analysis: Anal. Calcd. (Found) for $C_{36}H_{48}N_6O_9 \cdot 3HCl \cdot 2H_2O: C, 50.58 (50.62); H,$ 6.47 (6.49); N, 9.89 (9.84).

2.3. Titration Procedure. The apparatus used, experimental details, calibration of the electrode, and titration procedures were as described before.⁸ A standard hydrochloric acid solution was titrated with a standard KOH solution, and the calculated hydrogen ion concentrations ($pK_w = 13.77$) were used to convert the pH-meter reading to hydrogen ion concentration. All solutions were prepared prior to the experiments in double-distilled deoxygenated water. KOH solution of 0.1 M was prepared and standardized against potassium hydrogen phthalate. HCl solution (0.1 M) was prepared and standardized against standard KOH. The ionic strength was maintained at 0.1 M by adding the appropriate amount of 1 M KCl. Stock solutions of 0.01 M ligand and 0.01 M iron(III) were also prepared in deoxygenated water. The final concentrations of ligand $(1 \cdot 10^{-3} \text{ M})$ and metal $(1\!\cdot\!10^{-3}~M~\text{or}~5\!\cdot\!10^{-4}~M)$ were maintained for the different potentiometric titrations. The following titrations with metal-toligand molar ratios of $C_{\rm Fe}/C_{\rm L} = 0.1, 1.1, 1.2$ were carried out, and the potentiometric data were refined using the computer program Hyperquad 2000⁹ to get the protonation constants of the ligand and the formation constants of the metal complexes. Literature data $(\log \beta)$ on the iron(III) hydroxo complexes were included in the refinement process to obtain the formation constants of the

ferric complexes.¹⁰ The method adopted for the spectrophotometric titrations are similar to the potentiometric titrations except that dilute solutions of ligand $(4.02 \cdot 10^{-5} \text{ M})$ and metal ion $(4.02 \cdot 10^{-5} \text{ M})$ were used. After each adjustment of pH, an aliquot was removed, and spectra were recorded.

3. RESULTS AND DISCUSSION

3.1. Ligand Synthesis. The synthetic procedure of CYCOEN-CAT (L) is presented in Scheme 2. The ligand was isolated as a hydrochloride salt, and care was taken to avoid contact with the atmosphere throughout the procedure. This compound is highly hygroscopic and soluble in water, methanol, and ethanol but insoluble in chloroform, ether, and acetonitrile. The ligand was characterized on the basis of elemental analyses and various spectral (electronic, IR, and ¹H NMR) data.

The infrared spectrum of the unreduced form of CYCOEN-CAT showed a weak band at 1623 cm⁻¹ for $v_{C=N}$ that merged with the broad $v_{C=O}$ band. Upon reduction, the infrared spectrum of L showed only the $v_{C=O}$ band at 1634 cm⁻¹. Usually, the normal position for the stretching vibration of non-hydrogen bonded or free hydroxyl groups absorbs strongly in the range of (3700 to 3584) cm⁻¹.¹¹ In the present case, the ligand showed a broad v_{O-H} stretching vibration at 3374 cm⁻¹ indicating the presence of intramolecular hydrogen bonding. The deformation δ_{O-H} peak was observed at 1356 cm⁻¹. The band observed at 1240 cm⁻¹ was attributed to v_{C-O} . Aromatic carbon–carbon stretching vibrations were observed at (1574 and 1504) cm⁻¹, whereas two bands at (838 and 764) cm⁻¹ were assigned to the ring substituted bending vibrations.

The experimental ¹H NMR spectrum of the unreduced form of CYCOENCAT displayed a resonance signal at 8.30 ppm due to imine hydrogens and three distinct peaks at (1.58, 1.83, and 2.21) ppm for three different types of protons present in the cyclohexane ring. Triplets at (3.42 and 3.69) ppm were characterized for the methylene groups attached to the imine and amide functional groups, respectively. Other peaks at (6.62, 6.76, and 6.86) ppm were characterized for the aromatic protons. Upon reduction, the characteristic imine band at 8.30 ppm disappeared, and the ¹H NMR spectrum of CYCOENCAT (Figure 1) showed the presence of benzylic signals at 4.15 ppm.

Scheme 2. Synthesis of CYCOENCAT. Reagents and Conditions: (a) i. EtOH, Dry HCl; ii. PtO_2/H_2 , Room Temp., 24 h; (b) Ethylenediamine (en); (c) 2,3-Dihydroxybenzaldehyde, EtOH; (d) NaBH₄, Conc. HCl, MeOH



Figure 1. ¹H NMR spectrum of CYCOENCAT (D₂O, 300 MHz).

The spectrum showed three distinct peaks for the cyclohexane ring protons. A quartet and doublet were obtained at 1.35 ppm and 1.83 ppm for the axial and equatorial protons of methylene groups, respectively, whereas a triplet at 2.38 ppm was observed for the three axial protons attached with the appended groups in the cyclohexane ring, which confirmed a *cis,cis*-equatorial configuration for the ligand. Other peaks at (3.12 and 3.46) ppm were characterized for the methylene groups attached to the amine and amide functional groups, respectively. Peaks due to aromatic protons showed multiplets between (6.70 to 6.85) ppm and a doublet at 6.87 ppm.

3.2. Ligand Protonation Constants. Potentiometric and spectrophotometric titrations of the ligand CYCOENCAT were carried out in an aqueous medium to determine the protonation constants. In the potentiometric method, the ligand was first acidified with 10-fold excess of a measured amount of standard HCl and then titrated against standard KOH at an ionic strength of $\mu = 0.1$ M KCl and (25 ± 1) °C. Equilibrium points (pH = $-\log_{10}[\text{H}^+]$) were collected by adding base into the acidified solution of ligand. The titration curve of the ligand is shown in Figure 2. The analysis of the potentiometric titration curve using the program Hyperquad 2000 gave the best fit for six protonation



Figure 2. Potentiometric titration curves of CYCOENCAT (L) in the absence and presence of Fe(III) at a 1:1 ligand-metal molar ratio.

constants (Table 1) for the ligand as proposed by the eq 1 (charges are omitted for simplicity):

$$LH_{n-1} + H \rightleftharpoons LH_n \qquad K_n = \frac{[LH_n]}{[LH_{n-1}][H]} \quad (1)$$

The ligand CYCOENCAT contained nine deprotonation sites: six catechols and three sec-amines. However, due to the very low acidity of the catechol $(pK_{a1} = 9.13 \text{ and } pK_{a2} = 13.00)^{12}$ subunits, only one proton could be deprotonated. Therefore, the neutral CYCOENCAT was considered as triprotic acid and addressed as LH_{3} , whereas the fully protonated form as LH_{6}^{+3} . The species distribution diagram as a function of pH (Figure 3) indicates that in acidic solution CYCOENCAT initially exists 100 % in the fully protonated form as LH_6^{3+} below pH < 4.0. As the pH increases, deprotonation starts with the formation of LH_5^{2+} which exists between pH 5–8 with a maximum concentration (70%) at pH \sim 7. Species LH_4^+ , LH_3 , and LH_2^- are formed successively with an increase in pH and show a maximum concentration of 55 %, 70 %, and 55 % at pH \sim 8, \sim 9, and \sim 10, respectively. At pH \sim 11, the species LH^2 shows the maximum concentration of 50 %, and thereafter the deprotonated ligand L^{3-} starts to predominate.

Spectrophotometrically, the electronic spectra [(225 to 340) nm] of the ligand were recorded in the pH range 3.2-10.2 (Figure 4a). No appreciable change in the electronic spectra was observed below the experimental pH of 3.2, and the spectra overlapped each other. With an increase in pH, the spectra showed a bathochromic shift in the absorption band for the $\pi \rightarrow \pi^*$ transition from (281 to 288) nm with a concomitant increase in absorbance. The shift continued until pH 8.12, and within this range, two isosbestic points at (266 and 286) nm were observed. The formation of isosbestic points indicate the state of equilibrium between the deprotonated and the protonated ligand and can be attributed to the formation of catecholate ions. Since the catechol group is the only significant chromophore in CYCOENCAT, the red shift in the electronic spectra is attributable to deprotonation of the hydroxy group. As the pH rises further, a hyperchromic shift was prominent, and no more isosbestic points were observed. This observation can be characterized for the deprotonation process from $-NH_2^+$ groups. The inclusion of the whole range

of data in the least-squares fitting program pHAb¹³ resulted in six protonation constants (Table 1), which accords well with the potentiometric results. The predicted electronic spectra of the species whose protonation constants have been determined are shown in Figure 4b. According to these results, the six protonation constants calculated for CYCOENCAT can now be assigned to the three amine groups and three most acidic hydroxyl groups of the catechol units.

3.3. Metal-Ligand Complex Formation. Potentiometric titrations of the ligand CYCOENCAT (L) were carried out in the presence of Fe(III) at 1:1 and 1:2 metal-to-ligand molar ratios to study the formation of metal complexes. The titrations were carried out in an aqueous medium at an ionic strength of $\mu = 0.1$ M KCl and (25 ± 1) °C. The potentiometric curve for the 1:1 metal-ligand molar ratio is shown in Figure 2. The solid symbols represent equilibrium points collected when no solid phase was present in solution, while dotted lines represent points collected when turbidity or precipitation appeared in the solution. The deviation in the metal-ligand titration curves from the free ligand curve implies the formation of metal complexes. As the pH increases, precipitation/solid phase starts to appear from pH = \sim 8.0. Considering these preliminary observations, the experimental data represented in symbols were only tested in the minimization program with many sets of possible models. The best-fit models were obtained, when the formation of the species FeLH₃, FeLH₂, FeLH, and FeL were considered. No additional species were detected from the data obtained between the 1:1 and the 1:2 metal-ligand molar ratios. The overall formation constants (log β) of the species were calculated using the Hyperquad 2000 program and are summarized in Table 2. The equilibrium reactions for the overall formation of the complexes are given below by the eqs 2 to 5 (charges are omitted for simplicity):

$$Fe + LH_6 \rightleftharpoons FeLH_3 + 3H \qquad \beta_{113} = \frac{[FeLH_3][H]^3}{[Fe][LH_6]} \quad (2)$$

$$FeLH_3 \rightleftharpoons FeLH_2 + H$$
 $\beta_{112} = \frac{[FeLH_2][H]}{[FeLH_3]}$ (3)

$$\operatorname{FeLH}_2 \rightleftharpoons \operatorname{FeLH} + H \qquad \beta_{111} = \frac{[\operatorname{FeLH}][H]}{[\operatorname{FeLH}_2]} \quad (4)$$

$$FeLH \rightleftharpoons FeL + H$$
 $\beta_{110} = \frac{[FeL][H]}{[FeLH]}$ (5)

The interaction of Fe(III) with CYCOENCAT was also studied by a spectrophotometric method to explain the different possible coordination modes of the species formed in solution. The spectrometric titration was carried out with a 1:1 metal-to-ligand ratio with the ligand and metal ion concentrations equal to $4.02 \cdot 10^{-5}$ M and varying the pH between 2.5 and 8.0. The experimental electronic spectra of L in the presence of Fe(III) is shown in Figure 5. Major spectral changes were observed after pH \sim 3.2 with the appearance of a charge-transfer band at 575 nm due to the formation of ferric complexes. A study of species distribution curves (Figure 6) for the L-Fe(III) system indicates that below pH 3, the free ligand, and the species FeLH₃ are present in appreciable amounts. Hence, the peak at 575 nm is due to the formation of FeLH₃. On addition of the Fe(III) ion, three protons from LH_6^{3+} are released upon complexation to give the species FeLH₃. These three deprotonation/coordination

Table 1. Protonation Constants (log *K*) of CYCOENCAT (L), TAMCHCAT,⁸ and TRENCAT²¹ Obtained at (25 \pm 1) °C and μ = 0.1 M (A = Potentiometry and B = Spectrophotometry)

Equilibrium	CYCOENCAT		TMACHCAT	TRENCAT
	H,N, H,N, H,N, H,N, H,N, OH		н,)з Н, М, ОН ОН	N (H ₂ N))3 H ₂ N OH OH
	Α	В	Α	Α
$L^{3-} + H^+ \leftrightarrows HL^{2-}$	11.36 ± 0.05	11.38 ± 0.07	11.26	11.20
$\mathrm{HL}^{2-} + \mathrm{H}^+ \leftrightarrows \mathrm{H}_2\mathrm{L}^-$	10.67 ± 0.04	10.65 ± 0.05	10.65	10.60
$H_2L + H^+ \leftrightarrows H_3L$	9.82 ± 0.08	9.80 ± 0.02	9.80	6.59
$H_3L + H^+ \leftrightarrows H_4L^+$	8.49 ± 0.07	8.51 ± 0.08	8.48	8.07
$H_4L^+ + H^+ \leftrightarrows H_5L^{+2}$	7.62 ± 0.06	7.59 ± 0.06	7.61	7.29
$H_5L^{+2} + H^+ \leftrightarrows H_6L^{+3}$	6.27 ± 0.10	6.29 ± 0.09	6.23	6.17



Figure 3. pH-dependent species distribution curves for CYCOENCAT (L) at $[L]_{total} = 5 \cdot 10^{-4}$ M.

processes should have resulted from the three most acidic *ortho*hydroxy groups of the catechol to form a capped-type geometry. Thus, the peak at 575 nm can now be assigned to be a charge-transfer transition from the ligand to metal (catechol \rightarrow Fe^{III}). Similar observations have been made for other tripodal catecholate ligands,¹⁴ in which on coordination of the three acidic hydroxyl groups with Fe(III) in acetonitrile, a peak at 528 nm results due to ligand-to-metal charge transfer (LMCT).

At higher pH, the other species FeLH₂, FeLH, and FeL are formed in steps with the release of three protons from FeLH₃ (Figure 6). The release of these protons from FeLH₃ can either take place from the NH₂⁺ groups for which the protonation constant values have been evaluated, or from the remaining -OH groups of catechols, for which the protonation constant values are unknown. These two probabilities lead to two distinct geometries: a nitrogenoxygen encapsulated or oxygen-oxygen encapsulated. In the first case, if the metal ion enters into the nitrogen-oxygen cavity with the release of one/two protons from NH₂⁺ upon coordination, it may encounter the repulsive force from the two or one protonated NH₂⁺ groups, whereas in the second case, the ligand coordinates through its more basic catecholic oxygens as an anionic donor to



Figure 4. (a) Experimental electronic spectral data of CYCOENCAT (L) as a function of pH (3.2–10.2) during a spectrophotometric titration: [L] = 0.041 mM; [KCI] = 0.1 M, and $T = (25 \pm 1) \,^{\circ}\text{C}$. (b) Electronic spectra of the seven species as a function of molar absorptivity and wavelength predicted from pHAb using experimental data for CYCOENCAT.

form a tris(catechol) type complex. The latter is more preferable for these hard metal ions, which can be assigned from the shifting

Table 2. Overall (log β) Formation Constants of the Metal Complexes Formed by Ligand CYCOENCAT (L) at (25 ± 1) °C and μ = 0.1 M KCl

equilibrium	$\log eta$
$[FeLH_3]/[Fe][L][H]^3$	49.82 ± 0.03
$[FeLH_2]/[Fe][L][H]^2$	46.55 ± 0.04
[FeLH]/[Fe][L][H]	42.21 ± 0.07
[FeL]/[Fe][L]	34.61 ± 0.02



Figure 5. Experimental electronic spectra at 1:1 metal-to-ligand ratio for Fe(III)-CYCOENCAT system.



Figure 6. Speciation diagram calculated for the Fe(III)-CYCOENCAT system at an equimolar ferric ion-ligand ratio ($[Fe^{3+}] = [L] = 10^{-3} \text{ M}$).

of the peak at 575 nm (pH \sim 3.2) toward \sim 500 nm characterized for the formation of the tris(catechol) type complex. Iron(III)catechol usually gives the LMCT at \sim 490 nm,¹⁵ which also supports the present proposition. This type of coordination (O,O) is expected for iron(III), because the hard metal ion Fe(III) prefers coordination of hard negatively charged oxygen as compared to the nitrogen donors. X-ray crystal structure studies of tripodal



Figure 7. Plot of pFe $(-\log [Fe^{3+}])$ vs pH. pFe was calculated for $[L] = 5 \cdot 10^{-4}$ M, $[Fe^{3+}] = 5 \cdot 10^{-5}$ M using the deprotonation constants of ligand L, and the complexation constants β_{11n} .

ligands derived from TREN possessing six oxygen functions show that in their cationic or neutral complexes, nitrogen atoms (imine or amide functions) are not involved in coordination.^{16,17} On the basis of above facts, it is suggested that the ferric ion in the tricapped geometry of FeLH₃ is not efficiently protected by the wrapped pod and remains accessible for further complexation to give the encapsulated tris(catecholate) complex.

The ligand is a weak acid, and proton competition occurs depending on their pK_a and the pH; therefore, the pFe (= $-\log[\text{Fe}^{3+}]$) value was calculated under given conditions of pH and M^{*n*+} and L concentration to quantify the complexation efficiency of CYCO-ENCAT. The calculated pFe value of CYCOENCAT at pH = 7.4 by taking $[L]_{total} = 5 \cdot 10^{-4} \text{ M}$ and $[\text{Fe}^{3+}]_{total} = 5 \cdot 10^{-5} \text{ M}$ is pFe = 24.76, which infers its ability to compete with transferrin (pFe =20.3).¹⁸ It is well-known that most of the iron circulating in blood plasma is bound to transferrin. Because of the labile nature of the iron transferrin complex, iron overload is cured by sequestering agents like desferrioxamine B (DFO), deferiprone, and deferasirox.¹⁹ Concerning the high iron(III) chelating ability of CYCOENCAT, aqueous solubility over a wide range of pH and no hydroxo species with Fe(III) make the ligand useful for iron(III) sequestration. Moreover, Santos and co-workers²⁰ suggested that sequestering ligands with free amines as in DFO may improve their usefulness as a drug, where the amine groups are used as a point of attachment to a polymeric solid matrix, which enhance the properties of removing "hard" ions from water solutions. Such free amines are present in CYCOENCAT and also in their complexes depicted in solution. Further, the pH dependence of the selectivity was examined owing to the fact that, in some particular biological compartments or circumstances, pH values exceeding 7 can be observed. The plot of pFe as a function of the pH (Figure 7) reflects that the selectivity of the ligand for Fe(III) is maintained over a pH range 4 to 8.

AUTHOR INFORMATION

Corresponding Author

*E-mail: suban sahoo@rediffmail.com and b.kanungo@vsnl.com.

REFERENCES

(1) Steed, J. W.; Atwood, J. L. Supramolecular Chemistry, 2nd ed.; John Wiley & Sons: New York, 2009.

(2) Neilands, J. B. Struct. Bonding (Berlin) 1984, 58, 1-24.

(3) Loomis, L. D.; Raymond, K. N. Solution equilibria of enterobactin and metal-enterobactin complexes. *Inorg. Chem.* **1991**, *30*, 906–911.

(4) Liu, Z. D.; Hider, R. C. Design of Clinically Useful Iron(III)-Selective Chelators. *Med. Res. Rev.* 2002, 22, 26–64.

(5) Caris, C.; Baret, P.; Beguin, C.; Serratrice, G.; Pierre, J. L.; Laulhere, J. P. Metabolization of iron by plant cells using O-Trensox, a high-affinity abiotic iron-chelating agent. *J. Biochem.* **1995**, *312*, 879– 885.

(6) Shanzer, A.; Libman, J. Iron transport and storage in microorganisms. *Plants and animals*; Sigel, A., Sigel, H., Eds.; Dekker: New York, 1998; p 239.

(7) Baral, M.; Kanungo, B. K.; Moore, P. Synthesis of cis,cis-1,3,5trisubstituted cyclohexane based chelators with polyfunctional pendant arms. J. Chem. Res. 2005, 43, 43–45.

(8) Sahoo, S. K.; Baral, M.; Kanungo, B. K. Potentiometric, spectrophotometric, theoretical studies and binding properties of a novel tripodal polycatechol-amine ligand with lanthanide(III) ions. *Polyhedron* **2006**, *25*, 722–736.

(9) Gans, P.; Sabatini, A.; Vacca, A. Investigation of equilibria in solution. Determination of equilibrium constants with the HYPER-QUAD suite of programs. *Talanta* **1996**, *43*, 1739–1753.

(10) Khoe, G. H.; Brown, P. L.; Sylva, R. N.; Robins, R. G. The hydrolysis of metal ions. part 9. iron(III) in perchlorate, nitrate and chloride media (1 mol dm⁻³). *J. Chem. Soc., Dalton Trans.* **1986**, 1901–1906.

(11) Silverstein, R. M.; Webster, F. X. Spectrometric Identification of Organic Compounds, 6th ed.; John Wiley & Sons: New York, 1996.

(12) Martell, A. E.; Smith, R. M. Critical Stability Constants, Vols. 1–6; Plenum: New York, 1974–1989.

(13) Gans, P.; Sabatini, A.; Vacca, A. Determination of equilibrium constants from spectrophotometric data obtained from solutions of known pH: the program pHab. *Ann. Chim.* **1999**, *89*, 45–49.

(14) Hayashi, M.; Hiratani, K.; Kina, S. I.; Ishii, M.; Saigo, K. Synthesis and binding property of a novel tripodal hexadentate ligand having catechol moieties. *Tetrahedron Lett.* **1998**, *39*, 6211–6214.

(15) Sever, M. J.; Wilker, J. J. Visible absorption spectra of metalcatecholate and metal-tironate complexes. *J. Chem. Soc., Dalton Trans.* **2004**, 1061–1072.

(16) Archibald, S. J.; Blake, A. J.; Schroder, M.; Winpenny, R. E. P. New ligands for complexation of lanthanoids: the synthesis and structures of a nonadentate Schiff-base ligand (L^1) and of the complexes $[ML^1(OH_2)](ClO_4)_3 \cdot 3MeNO_2$ (M = La,Pr) and $[YL^1](ClO_4)_3 \cdot 3MeCN$. J. Chem. Soc., Chem. Commun. **1994**, 1669–1670.

(17) Xu, J.; Franflin, S. J.; Whisenhunt, D. W., Jr.; Raymond, K. N. Gadolinium complex of tris[(3-hydroxy-1-methyl-2-oxo-1,2-didehydropyridine-4-carboxamido)ethyl]-amine: a new class of gadolinium magnetic resonance relaxation agents. J. Am. Chem. Soc. **1995**, 117, 7245–7246.

(18) Crisponi, G.; Remelli, M. Iron chelating agents for the treatment of iron overload. *Coord. Chem. Rev.* **2008**, 252, 1225–1240.

(19) Sharma, R. N.; Pancholi, S. S. Oral iron chelators: a new avenue for the management of thalassemia major. *J. Curr. Pharm. Res.* 2010, 01, 1–7.

(20) Gaspar, M.; Grazina, R.; Bodor, A.; Farkas, E.; Santos, M. A. Siderophore analogues. a new macrociclic tetraminetrishydroxamate ligand. synthesis and solution chemistry of the iron(III), aluminium(III) and copper(II) complexes. J. Chem. Soc., Dalton Trans. **1999**, 799–806.

(21) Bismondo, A.; Comuzzi, C.; Di Bernardo, P.; Zanonato, P. L. Complexation of thorium(IV) by tris((2,3-dihydroxybenzylamino)ethyl)amine-a new strong chelating agent. *Inorg. Chim. Acta* **1999**, 286, 103–107.