

# Highly Efficient Cyanoimidation of Aldehydes


Ping Yin,<sup>†</sup> Wen-Bo Ma,<sup>†</sup> Yue Chen,<sup>‡</sup> Wen-Cai Huang,<sup>†</sup> Yong Deng,<sup>†</sup> and Ling He<sup>\*,†</sup>

Key Laboratory of Drug-Targeting and Drug-Delivery Systems of the Ministry of Education, Department of Medicinal Chemistry, West China School of Pharmacy, Sichuan University, Chengdu, Sichuan, 610041, P. R. China, and Department of Nuclear Medicine Affiliated Hospital, Luzhou Medical College, No. 25 Taiping Street, Luzhou, Sichuan, 646000, P. R. China

lhe2001@sina.com

Received October 7, 2009

## ABSTRACT



Cyanoimidation of aldehydes using cyanamide as a nitrogen source and using NBS as an oxidant was achieved in high yields without the addition of a catalyst. The method has several advantages, including mild conditions, simple workflow, and inexpensive reagents. The reaction proceeds in a one-pot manner, giving rise to the formation of intermolecular C–N and C–O bonds. Subsequently, the substituted *N*-cyanoimide products may also undergo a cyclization reaction to give 1,2,4-triazole derivatives in high yields.

The development of efficient and convenient procedures to construct the building blocks of heterocyclic compounds has provoked interest. As such, cyanoimide derivatives have been widely used as precursors for the synthesis of heterocyclic compounds.<sup>1</sup> *N*-Cyanoimides can undergo various reactions with a variety of nucleophilic reagents in the construction of *N*-heterocyclic bioactive compounds, such as hydraziniums, imidamides(amidines), hydroxylamine,

amines, and nitriles, all of which were obtained from 3-amino-1,2,4-triazole derivatives, *s*-triazines, 1,2,4-oxadiazoles, isomeric 5-(phenylsulfonyl)pyrimidines, and substituted xanthenes by application of *N*-cyanoimides as the key intermediate.<sup>1</sup> Heterocyclic compounds generally possess important biological activities. Many kinds of natural products, pharmaceutical compounds, insecticides, dyes, and solvents are heterocyclic molecules.<sup>2</sup> It is important to synthesize heterocyclic compounds via a convenient route with high yields. Cyanoimides are attractive chemicals not only because of their unique structure and reactivity but also

<sup>†</sup> Sichuan University.

<sup>‡</sup> Luzhou Medical College.


(1) (a) He, R. J.; Ching, S. M.; Lam, Y. L. *J. Comb. Chem.* **2006**, *8*, 923. (b) Nakajima, T.; Nakajima, S. *Chem. Pharm. Bull.* **1994**, *42*, 2483. (c) M'Hamed, M. O.; M'Rabet, H.; Efrat, M. L. *C. R. Chim.* **2007**, *10*, 1147. (d) Zarguil, A.; Boukhris, S.; El Efrat, M. L. *Tetrahedron Lett.* **2008**, *49*, 5883. (e) Pérez, M. A.; Soto, J. L. *J. Chem. Soc., Perkin Trans. 1* **1985**, *1*, 87. (f) Perez, M. A.; Soto, J. L. *Synthesis* **1983**, *5*, 402. (g) McCall, J. M.; Kloosterman, D. J. *Org. Chem.* **1979**, *44*, 1562.

(2) (a) Yu, G.; Cheng, J. S. *Huaxue Gongye* **2006**, *27* (5), 24. (b) Carroll, W. A.; Perez-Medrano, A.; Peddi, S.; Florjancic, A. S. U.S. Pat. Appl. Publ., 2006025614, 02 Feb 2006. (c) Montserrat, C.; Albert, P. PCT Int. Appl., 2006010756, 02 Feb 2006. (d) Xu, D. F.; Wang, L. Q. CN1467206, 14 Jan 2004.

because they serve as versatile building blocks for heterocyclic compounds in the fields of organic synthetic chemistry, medicinal chemistry, and materials science. Although cyanoimides are important chemical reagents, there is a paucity of reliable synthetic methods for producing them.<sup>2</sup> The yields for synthesis of aryl-substituted *N*-cyanoimides are also unsatisfactory. We report an efficient and convenient synthetic route (Scheme 1) for synthesizing *N*-cyanoimides via a one-pot reaction of aldehyde derivatives in a *t*-BuONa/NBS system to give aromatic cyanoimides in good to excellent yields. The results are summarized in Tables 1 and 2.

With an interest in synthesizing potentially bioactive nitrogenous compounds, we focused on the application and prolongation of active nitrogen sources and development of the catalyzed amidation. We found that the saturated C–H bonds of cyclic ethers and tertiary amines could be activated by transition metals.<sup>3</sup> We also attended to a series of studies on metal-catalyzed amidation of aldehydes and amidation of aldehydes under metal-free conditions.<sup>4</sup> Although the methodologies mentioned above were well suited for some amides, such as carboxamide and sulfonamide, and could effectively form carbon–nitrogen bonds, the use of cyanamide as a nitrogen source has never been reported in the literature. As part of our continuing interest in catalyzed amidation, we would like to establish an effective reaction system for the oxidative amidation of aldehydes and ketones


**Scheme 1.** Cyanoimidation of Aldehydes



using cyanamide as a nitrogen source. In our first attempt to obtain amide derivatives, we chose  $PhI(OAc)_2$  as the oxidant and either  $Al_2O_3$  or  $K_2CO_3$  as the base. Unfortunately, products of amidation were not obtained; only the starting material was recovered. However, using *t*-BuOK/NBS with  $CH_3OH$  as solvent led to the formation of the nonamidated product *N*-cyanoimide **3** (Scheme 1).

A possible mechanism for the selective *N*-cyanoimidation–esterification of aldehydes is proposed in Scheme 2. According to route 1, condensation of compound A, which was determined by  $^1H$  NMR and HRMS, occurred at first. The subsequent Michael addition of  $CH_3OH$  to *N*-(benzo[d][1,3]dioxol-5-ylmethylene) cyanamide generated intermediate B.<sup>5</sup> Subse-

**Scheme 2.** Tentative Mechanism for Selective Cyanoimidation and Oxidative Esterification of Aldehyde



**Table 1.** Optimization of Selective Reaction Conditions<sup>a</sup>

$Cl-C_6H_4-CHO + CH_3OH \xrightarrow[oxidant]{H_2NCN, base} Cl-C_6H_4-C(=NCN)OCH_3 + Cl-C_6H_4-C(=O)OCH_3$			
entry	base	oxidant	% yield (conversion) <sup>b</sup> of <b>3b/4b</b>
1	<i>t</i> -BuONa	$SeO_2$	trace/–
2	<i>t</i> -BuONa	$PhI(OAc)_2$	12/–
3	<i>t</i> -BuONa	$CuCl_2^c$	trace/–
4	<i>t</i> -BuONa	NCS	24/–
5	<i>t</i> -BuONa	$I_2$	32/25
6	<i>t</i> -BuONa	NBS	64/16
7 <sup>d</sup>	<i>t</i> -BuONa	NBS	35/37 (86)
8	$K_2CO_3$	NBS	41/12
9	KOH	NBS	58/15
10 <sup>e</sup>	<i>t</i> -BuONa	NBS	78/8 (91)
11 <sup>f</sup>	<i>t</i> -BuONa	NBS	83/trace (94)
12 <sup>g</sup>	<i>t</i> -BuONa	NBS	85/trace (97)
13		NBS	–/85 (87)

<sup>a</sup> Unless otherwise noted, all reactions were carried out with molar ratio **1b**/ $H_2NCN$ /base = 1:1.2:1.5 at rt for 30 min, and then oxidant (1.5 equiv of **1**) was added at 50 °C for 12 h. <sup>b</sup> Isolated yields. <sup>c</sup> 20 mol % catalytic amounts  $CuCl_2$  of **1b** were used. <sup>d</sup> The addition of each reagent had no interval. <sup>e</sup> The molar ratio was **1b**/ $H_2NCN$ /*t*-BuONa/NBS = 1:2:2:2. <sup>f</sup> The molar ratio was **1b**/ $H_2NCN$ /*t*-BuONa/NBS = 1:3:3:3. <sup>g</sup> The molar ratio was **1b**/ $H_2NCN$ /*t*-BuONa/NBS = 1:4:4:4.

quently, the active hydrogen of intermediate B was captured by bromine to produce haloamine C in the presence of NBS, finally haloamine C which in the presence of an alkali converted to **3A** with the elimination of  $HBr$ .<sup>6</sup> Similar oxidation and elimination steps were performed via hemiacetal intermediate D and its halide E to generate carboxylic ester **4A** using NBS as an oxidant<sup>7</sup> via route 2.

(5) (a) Ishihara, M.; Togo, H. *Tetrahedron*. **2007**, 63, 1474. (b) Li, G. L.; Fronczek, F. R.; Antilla, J. C. *J. Am. Chem. Soc.* **2008**, 130, 12216.

(3) (a) He, L.; Yu, J.; Zhang, J.; Yu, X. -Q. *Org. Lett.* **2007**, 9, 2277. (b) Liu, N.; Tang, B. Y.; Chen, Y.; He, L. *Eur. J. Org. Chem.* **2009**, 26, 2059.

(4) (a) Yoo, W. J.; Li, C. J. *J. Am. Chem. Soc.* **2006**, 128, 13064. (b) Chan, J.; Baucom, K. D.; Murry, J. A. *J. Am. Chem. Soc.* **2007**, 129, 14106. (c) Chang, J. W. W.; Chan, P. W. H. *Angew. Chem., Int. Ed.* **2008**, 47, 1138. (d) Seo, S.; Marks, T. J. *Org. Lett.* **2008**, 10, 317. (e) Suto, Y.; Yamagiwa, N.; Torisawa, Y. *Tetrahedron Lett.* **2008**, 49, 5732. (f) Wang, L.; Fu, H.; Jiang, Y. Y.; Zhao, Y. F. *Chem.–Eur. J.* **2008**, 14, 10722. (g) Ekoue-Kovi, K.; Wolf, C. *Org. Lett.* **2007**, 9, 3429. (h) Gao, J.; Wang, G. W. *J. Org. Chem.* **2008**, 73, 2955.

**Table 2.** Cyanoimidation of Aldehydes<sup>a</sup>


entry <sup>a</sup>	R <sub>1</sub>	R <sub>2</sub>	3	4	% yield (conversion) <sup>b</sup> of 3/4 (conversion)
1	Ph	Me	<b>3a</b>	<b>4a</b>	87/– (96)
2	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	Me	<b>3b</b>	<b>4b</b>	85/– (97)
3 <sup>c</sup>	<i>p</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	Me	<b>3c</b>	<b>4c</b>	82/– (57)
4 <sup>d</sup>	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Me	<b>3d</b>	<b>4d</b>	67/24 (99)
5	<i>p</i> -BrC <sub>6</sub> H <sub>4</sub>	Me	<b>3e</b>	<b>4e</b>	76/16 (96)
6 <sup>c</sup>	<i>m</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	Me	<b>3f</b>	<b>4f</b>	58/– (86)
7 <sup>d</sup>	<i>m</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Me	<b>3g</b>	<b>4g</b>	72/16 (93)
8	<i>m</i> -BrC <sub>6</sub> H <sub>4</sub>	Me	<b>3h</b>	<b>4h</b>	94/– (99)
9	<i>m</i> -ClC <sub>6</sub> H <sub>4</sub>	Me	<b>3i</b>	<b>4i</b>	83/– (95)
10	1-Naph	Me	<b>3j</b>	<b>4j</b>	76/– (71)
11	6-quinolyl	Me	<b>3k</b>	<b>4k</b>	80/– (88)
12	2-thienyl	Me	<b>3l</b>	<b>4l</b>	49/– (87)
13	Ph	Et	<b>3m</b>	<b>4m</b>	74/– (98)
14	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	Et	<b>3n</b>	<b>4n</b>	79/– (90)
15	<i>p</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	Et	<b>3o</b>	<b>4o</b>	–/–
16 <sup>d</sup>	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Et	<b>3p</b>	<b>4p</b>	84/– (99)
17 <sup>e</sup>	<i>m</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	Et	<b>3q</b>	<b>4q</b>	77/– (74)
18 <sup>d</sup>	<i>m</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Et	<b>3r</b>	<b>4r</b>	81/– (96)
19	<i>m</i> -ClC <sub>6</sub> H <sub>4</sub>	Et	<b>3s</b>	<b>4s</b>	92/– (87)
20 <sup>f</sup>	Ph	<i>i</i> -Pr	<b>3t</b>	<b>4t</b>	20/– (78)
21	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	<i>n</i> -Pr	<b>3u</b>	<b>4u</b>	51/trace (87)
22	<i>o</i> -FC <sub>6</sub> H <sub>4</sub>	Me	<b>3v</b>	<b>4v</b>	90/– (99)
23	2,6-di-ClC <sub>6</sub> H <sub>3</sub>	Me	<b>3w</b>	<b>4w</b>	84/– (92)
24	<i>o</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	Me	<b>3x</b>	<b>4x</b>	85/– (98)
25	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	hydroxyethyl	<b>3y</b>	<b>4y</b>	–/67 (95)
26	Ph	hydroxyethyl	<b>3z</b>	<b>4z</b>	–/68 (92)
27 <sup>g</sup>	4-formyl C <sub>6</sub> H <sub>4</sub>	Me	<b>3I</b>	<b>4I</b>	37/40 (97)
28	4-MeC <sub>6</sub> H <sub>4</sub>	Me	<b>3II</b>	<b>4II</b>	72/– (81)
29	<i>n</i> -butyl	Me	<b>3III</b>	<b>4III</b>	–/–

<sup>a</sup> Unless otherwise noted, all reactions were carried out with molar ratio 1/H<sub>2</sub>NCN/*t*-BuONa = 1:4:4 at room temperature for 30 min. Then NBS (4 equiv of 1) was added at 50 °C for 12 h. <sup>b</sup> Yield of isolated product. <sup>c</sup> The reaction was performed at rt for 6 h. <sup>d</sup> The addition of each reagent had no interval. <sup>e</sup> The reaction temperature was room temperature, and the reaction time was 24 h. <sup>f</sup> The reaction time was 24 h. <sup>g</sup> Product **3I** was methyl-4-((cyanoimino)(methoxy)methyl)benzoate, **4I** was dimethyl *N,N*-dicyanoterephthal imide, and the molar ratio was aldehyde/H<sub>2</sub>NCN/*t*-BuONa/NBS = 1:6:6:6.

To evaluate the selectivity of esterification and cyanoimidation, the cyanoimidation reaction of 4-chlorobenzaldehyde was studied by a one-pot reaction to give *N*-cyanoimide compound **3b** and carboxylic ester compound **4b** with a ratio of 1:1 (Table 1, entry 7). When 4-chlorobenzaldehyde was used as substrate with a substrate/NH<sub>2</sub>CN/potassium *tert*-butoxide/NBS molecular ratio of 1:3:3:3, the produced cyanoimides were obtained with good yield. With the same procedure, several oxidants, such as SeO<sub>2</sub>, periodate reagents, such as PhI(OAc)<sub>2</sub> and PhI=O, and iodine were tested in

(6) (a) Liu, X. W.; Zhang, Y. M.; Wang, L.; Fu, H.; Jiang, Y. Y.; Zhao, Y. F. *J. Org. Chem.* **2008**, *73*, 6207. (b) Wang, Z.; Zhang, Y. M.; Fu, H.; Jiang, Y. Y.; Zhao, Y. F. *Org. Lett.* **2008**, *10*, 1863. (c) Fujioka, H.; Murai, K.; Kubo, O.; Ohba, Y.; Kita, Y. *Tetrahedron* **2007**, *63*, 638. (d) Fujioka, H.; Murai, K.; Ohba, Y.; Hiramatsu, A.; Kita, Y. *Tetrahedron Lett.* **2005**, *46*, 2197.

(7) (a) Travis, B. R.; Sivakumar, M.; Hollist, G. O.; Borhan, B. *Org. Lett.* **2003**, *5*, 1031. (b) Gopinath, R.; Patel, B. K. *Org. Lett.* **2000**, *2*, 577.

**Scheme 3.** Synthesis of 1,2,4-Triazoles**Table 3.** Synthesis of 1,2,4-Triazoles<sup>a</sup>

entry	substrate	product	% yield <sup>b</sup>
1			74
2			80
3			80
4			92
5			88
6			82

<sup>a</sup> Reactions were performed in CH<sub>3</sub>OH in reflux for 4 h with a substrate/PhNHNH<sub>2</sub> ratio of 1:1.2. <sup>b</sup> Isolated yields.

CH<sub>3</sub>OH in the presence of *t*-BuONa. We validated the general procedure, and the results are shown in Table 1 (entries 1, 2, and 5). It was found that these oxidants and a transition-metal salt (20 mol % of CuCl<sub>2</sub>) (Table 1, entry 3) resulted in the formation of trace products. Only NBS with the molar ratio (1b/H<sub>2</sub>NCN/*t*-BuONa/NBS = 1:3:3:3 and 1b/H<sub>2</sub>NCN/*t*-BuONa/NBS = 1:4:4:4) gave high yields of **3b** (Table 1, entries 11 and 12). When using 4-chlorobenzaldehyde as a substrate with 1b/H<sub>2</sub>NCN/*t*-BuONa/NBS in the molecular ratio of 1:1.2:0:1.5, the product methyl 4-chlorobenzoate was obtained with 85% yield, but the formation of cyanoimide was not observed (Table 1, entry 13). We also examined the effect of bases (Table 1, entries 7–9), which displayed moderate yields of **3b**. On the basis of these results, we used sodium *tert*-butoxide in a molar ratio of 1b/H<sub>2</sub>NCN/*t*-BuONa/NBS = 1:3:3:3 and 1b/H<sub>2</sub>NCN/*t*-BuONa/NBS = 1:4:4:4 for subsequent studies.

To test the universality of the reaction system and the scope of selective cyanoimide reactions, several different

aldehydes were examined. Almost all the substrates could give their corresponding product with good yields (Table 2). However, the yields were moderate or low, and the selectivity could not be ensured when using other alcohols as solvents (Table 2, entries 20, 21, 25, and 26) under identical conditions. A semiselective product **3I** was also obtained with diester **4I** using terephthalaldehyde as the substrate (Table 2, entry 27). In addition, the aliphatic aldehyde could not yield the cyanoimide product in the reaction system (Table 2, entry 29). From these results, we can conclude that this method could provide a convenient and efficient way for the preparation of substituted *N*-cyanobenimide from aromatic aldehydes with simple commercially available reagents.

Moreover, methyl *N*-cyanobenzimidate derivatives can undergo a cyclization reaction with hydrazine derivatives to give 1,2,4-triazoles,<sup>8</sup> which can play a key role in pharmacological activity.<sup>9</sup> Thus, we used the products of cyanoimide of aldehydes to construct nitrogen-containing hetero-

cyclic compounds (Scheme 3). The results demonstrated that several heterocyclic compounds could readily be obtained, all of which are shown in Table 3.

In summary, we have demonstrated an efficient one-pot procedure for the cyanoimide of aldehydes in the absence of a catalyst to form the corresponding key intermediate *N*-cyanobenimide using NBS as an oxidant. Subsequently, the substituted *N*-cyanobenimide products were found to be capable of undergoing a cyclization reaction to give 1,2,4-triazole derivatives in good yields. The method has several advantages, including mild conditions, simple workflow, high yields, and inexpensive reagents. Further investigation into the pharmacological activity of these heterocyclic compounds is underway and will be reported in due course.

**Supporting Information Available:** Detailed experimental procedures and characterization data for central compounds are available free of charge via the Internet at <http://pubs.acs.org>.

OL902207H

(8) (a) Barluenga, J.; Valdés, C.; Beltrán, G.; Escibano, M.; Aznar, F. *Angew. Chem., Int. Ed.* **2006**, *45*, 6893. (b) Lipshutz, B. H.; Taft, B. R. *Angew. Chem., Int. Ed.* **2006**, *45*, 8235. (c) M'Hamed, M. O.; M'Rabet, H.; Efrat, M. L. *C. R. Chim.* **2007**, *10*, 1147. (d) Zarguil, A.; Boukhris, S.; El Efrat, M. L.; Souizi, A.; Essassi, E. M. *Tetrahedron Lett.* **2008**, *49*, 5883.

(9) (a) Akbarzadeh, T.; Tabatabai, S. A.; Khoshnoud, M. J.; Shafaghi, B.; Shafiee, A. *Bioorg. Med. Chem.* **2003**, *11*, 769. (b) Naito, Y.; Akahoshi, F.; Takeda, S.; Okada, T.; Kajii, M.; Nishimura, H.; Sugiura, M.; Fukaya, C.; Kagitani, Y. *J. Med. Chem.* **1996**, *39*, 3019.