Copper-Catalyzed Oxidative Dimerizations of 3-*N*-Hydroxy-aminoprop-1-enes to form 1,4-Dihydroxy-2,3-diaminocyclohexanes with *C*₂ Symmetry**

Satish Ghorpade and Rai-Shung Liu*

Abstract: This work describes the one-step construction of complex and important molecular frameworks through copper-catalyzed oxidations of cheap tertiary amines. Copper-catalyzed aerobic oxidations of N-hydroxyaminopropenes to form C_2 -symmetric N- and O-functionalized cyclohexanes are described. Such catalytic oxidations proceed with remarkable stereocontrol and high efficiency. Reductive cleavage of the two N-O bonds of these products delivers 1,4-dihydroxy-2,3-diaminocyclohexanes, which are important skeletons of several bioactive molecules.

Metal-catalyzed oxidative C–H functionalizations of aminoalkane moieties have been a subject of intense interest because of their practical significance.^[1] Reported examples focus mainly on the formation of iminium ions (I) which are subsequently attacked by suitable nucleophiles [Eq. (1)].^[2-6]

These reactions are appealing surrogates for the well-known Mannich reaction^[7] because tertiary amines (1) are readily available and cheap reagents. *N*-Hydroxyaminoalkanes (1, R' = OH) are an important class of organic compounds,^[8] and their C–H oxidative functionalization generate nitrone species (**I**; $R' = O^-$) as depicted in Equation (1).^[9] Alternatively, α -carbonyl N-hydroxyamino species (3) undergo metal-free oxidations to generate reactive amidoxyl radicals (**II**), which then attack an alkene to form a new C–O bond [Eq. (2)].^[10] Despite numerous publications, the present N–CH oxidative functionalizations focus mainly on the Mannich-type reactions, rather than on the construction of complicated frameworks. Herein, we report the copper-catalyzed oxidative dimerizations of 3-*N*-hydroxy-aminoprop-1-enes (**5**) to pro-

 [*] S. Ghorpade, Prof. Dr. R.-S. Liu Department of Chemistry, National Tsing Hua University 101, Sec. 2, Kuang-Fu Rd., Hsinchu 30013 (Taiwan) E-mail: rsliu@mx.nthu.edu.tw Homepage: http://chem-en.web.nthu.edu.tw/files/13-1117-33454.php
[**] We thank the National Science Council, Taiwan for financial support of this work

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201407987.



duce highly O- and N-functionalized cyclohexanes (6) with excellent stereoselectivity (d.r. > 20:1) [Eq. (3)]. Subsequent reductive cleavage of their N–O bonds affords 1,4-dihydroxy-2,3-diaminocyclohexanes (7) efficiently. Notably, these substances do not follow current process to generate alkenylnitrone intermediates [Eq. (1)], despite the presence of a NCH₂ moiety.

This work:



These resulting O- and N-containing cyclohexane frameworks are found in bioactive molecules of various families as exemplified by the species **III–VI** (Figure 1), and their stereoselective synthesis inevitably requires long procedures.^[11–15] Some 1,2-diaminocyclitols (**III**)^[12] are potent glucocerebrosidase activators and promising pharmacological chaperones for Gaucher disease. In contrast, several conduritols (**IV**) showed biological activity as glycosidase inhibitors.^[13] The compounds (**V**) are alcohol derivatives of *trans*diaminocyclohexane dichloroplatinum (DACH-PtCl₂), which exhibits potent antitumor activity against P388 leukemia.^[14] Oseltamivir phosphate (**VI**; Tamiflu) is currently used for the treatment of influenza.^[15]

Table 1 presents optimization of the reaction conditions for the oxidative dimerizations of the 3-*N*-hydroxy-1-propene **5a** using various catalysts, oxidants, and solvents. We first



Figure 1. Representative bioactive molecules.

Angew. Chem. Int. Ed. 2014, 53, 1-5

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Weinheim Wiley Online Library These are not the final page numbers!



Table 1: Conditions for the oxidative dimerizations of 5 a.

	Ph`N OH 5a	Y ^{Ph} ⊂at s	alyst (5 mol% oxidant olvent, 25 °C	$\begin{array}{c} Ph \\ N-\\ \bullet Ph \\ O-\\ 6a \end{array}$		h
Entry	Catalyst	Oxidant	Solvent ^[a]	Additive	<i>t</i> [h]	Yield ^[b] [%]
1	CuCl	O ₂	THF	-	9	63
2	CuBr	O ₂	THF	-	9	70
3	[IPrCuCl]	O ₂	THF	-	9	84
4	$CuCl_2$	O ₂	THF	-	9	25
5	$FeCl_3$	O ₂	THF	-	9	67
6	Co(OAc) ₂	O ₂	THF	-	10	76
7	[IPrCuCl]	O ₂	toluene	-	9	69
8	[IPrCuCl]	O ₂	DCE	-	9	55
9	[IPrCuCl]	H_2O_2	THF	4 Å M.S.	9	71
10	[IPrCuCl]	<i>t</i> BuOOH	THF	4 Å M.S.	12	75
11	[IPrCuCl]	DTBP	THF	-	12	68
12	[IPrCuCl]	-	THF	-	12 ^[c]	-
13	-	O ₂	THF	-	12	-

[a] [5a] = 0.20 M. [b] Product yields are reported after purification from a silica gel column. [c] The recovery yield of 5a was 67 and 71% for entries 12 and 13, respectively. M.S. = molecular sieves, THF = tetrahydrofuran.

tested the reaction with a CuCl catalyst (5 mol %) and O_2 (1 atm) in THF under ambient conditions (25 °C, 9 h), which led to 1,4-dihydroxy-2,3-diaminocyclohexane 6a as a single diastereometic product (d.r. > 20:1) with a yield of 63% (entry 1). Among other copper catalysts tested, [IPrCuCl] (IPr = 1,3-bis(diisopropylphenyl)imidazol-2-ylidene) gave the best yield (84%) of **6a** (entry 3), whereas CuBr and CuCl₂ provided the same product in 70 and 25%, respectively (entries 2 and 4). In the presence of O₂, both FeCl₃ and $Co(OAc)_2$ were also effective catalysts, thus yielding **6a** in 67 and 76% yield, respectively (entries 5 and 6). The use of [IPrCuCl] in toluene and 1,2-dichloroethane (DCE) gave 6a in 69 and 55% yield, respectively (entries 7 and 8). With [IPrCuCl], other oxidants such as H₂O₂, tert-butyl peroxide (TBPO), and di-tert-butyl peroxide (DTBP) also give the desired 6a in 68-75% yields (entries 9-11). In the absence of an oxidant and metal catalyst, no reaction occurred over a protracted period (12 h) with 67 and 71% recovery, respectively, of the staring 5a (entries 12 and 13). The structural characterization of 6a relies on X-ray diffraction.[16] The ORTEP drawing (see the Supporting Information) indicates the presence of C_2 symmetry for the molecular framework.

We prepared additional 2-substituted 3-*N*-hydroxy-1propenes (**5b–u**) to examine the generalization of such oxidative dimerizations (Table 2). Most reactions were performed with 5 mol% [IPrCuCl]/O₂ in THF (25°C, 11–16 h) except for those involving **5j–l**, for which the reactions were run with [IPrCuCl]/tBuO₂H (3 equiv) in THF (4 Å M.S., 25°C, 11–12 h). In all cases, the resulting O- and N-functionalized cyclohexanes **6b–u** were obtained as a single diastereomer (d.r. > 20:1). We tested the oxidations with the initial substrates **5b–f**, bearing variable aniline groups (R²=4-XC₆H₄, X=F, Cl, Br, CO₂Me and Me), and the resulting products **6b–f** were obtained in 74–83% yields. We also Table 2: Scope for the oxidative dimerizations.



[а] [6 b] = 0.20 м.	Product yields	are reported	after	purification	from
a silica gel colur	nn.				

prepared their *meta*-substituted analogues 5g-i ($R^2 = 3$ - XC_6H_4 , X = Cl, Me, and OMe), which gave the desired products 6g-i in 78-89%. The dimerizations of the orthosubstituted analogues 5j-l ($R^2=2-X$ C_6H_4 , X=Cl, Br, and Me) were unsuccessful with [IPrCuCl] and O₂, but the use of [IPrCuCl] and tBuO₂H (3 equiv) enabled the production of desired compounds 6j-l in reasonable yields (51-68%). We envisage that these ortho substituents are not favorable for the coordination of copper with their hydroxyamino groups, thus rendering the oxidation difficult. We also prepared the 3-N-hydroxy-1-propenes 5m-o bearing various alkenyl substituents ($R^1 = 4-XC_6H_4$, X = Br, Cl, and Me), and their oxidative dimerizations proceeded smoothly with [IPrCuCl]/ O_2 to give the desired **6m–o** in 61-83% yields. For the substrates 5p and 5q bearing a 2- and 3-thienyl group, respectively, at the alkenyl moiety, their corresponding products 6p (82%) and 6q (69%) were obtained. To our delight, these catalytic oxidations were compatible with the substrates 5r-u, bearing various alkyl groups at their amino or alkenyl positions ($\mathbf{R}^1 = \mathbf{M}\mathbf{e}$, *i*Pr or $\mathbf{R}^2 = i$ Pr or *t*Bu), and their desired dimerization products 6r-u were obtained in 72-81% yields. These oxidative dimerizations failed to work with those substrates bearing $R^1 = CO_2Me$ and H, and thus resulted in a complex mixture of products.

We also examined the access to the O- and N-containing cyclohexanes **7** by employing reductive cleave of the N–O bonds of the products **6**. The procedures^[17] involved Pd/C and H₂ (1 atm) in MeOH/CH₂Cl₂ (1:1) near 25 °C. The results are summarized in Table 3. As shown, this N–O cleavage works

www.angewandte.org

2

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

K These are not the final page numbers!

Table 3: Reductive cleavage of the N-O bonds.



[a] [6a] = 0.04 M. Product yields are reported after purification from a silica gel column.

well for **6a–c** and **6e**, bearing various aniline groups ($R^2 = 4$ - XC_6H_4 , X = H, F, Cl, and CO_2Me), thus yielding the desired 7a-c and 7e in satisfactory yields (78-85%). This N-O cleavage is applicable to their meta- or ortho-substituted aniline analogues (6g-i and 6j-l), thus giving the highly functionalized cyclohexanes 7g-i and 7j-l in 69-83% yields. The cleavage also proceeded well with substrates 6n and 6o comprising different R^1 substituents ($R^1 = 4-XC_6H_4$, X = Cland Me), and giving the compounds 7n and 7o, respectively, in good yields. These reactions were compatible with the 2and 3-thienyl-containing species 6p and 6q, respectively, thus yielding the corresponding products 7p(72%) and 7q(63%). For the alkyl-substituted derivatives 6r-u, the reductive cleavage delivered the compounds 7r-u in 67-81% yields. Of these products, X-ray diffraction^[16] of a representative compound, 7b, was performed to confirm its structure as having all hydroxy and amino groups in axial positions.

Shown in Equations (4) and (5) (Tf = trifluoromethanesulfonyl) are control experiments to assist the understanding of the mechanism of the dimerization. Treatment of *N*hydroxyaniline (**8a**) with 2-phenyl-2-en-1-al (**8b**) with [IPr-CuCl]/O₂ in toluene (25 °C, 12 h) produced the quinoline



derivative **8c** in 54% yield through an intramolecular cyclization of the *E*-configured nitrone intermediate (*E*)-**I'**. We thus exclude the role of nitrone intermediates (**I'**) in the dimerization reactions because of their distinct chemoselectivity. We tested further the dimerization of **5a** with chiral CuOTf/**L2*** [**L2*** = 2,2-bis{(4S)-(-)-4-isopropyloxazoline}-propane], which enabled the production of (-)-**6a** in 19 and 27% *ee* in THF and toluene, respectively.^[18] The observed asymmetric induction of (-)-**6a** suggests that copper complexes participate in the formation of the first asymmetric C–C bond during the course of the dimerization.

We postulate a mechanism in Scheme 1. Under O_2 , Cu^{II} species are generated to coordinate with two molecules of the *N*-hydroxy aminopropenes **5** to form Cu-**5**, which undergoes a single-electron transfer (SET) to generate the radical cation **A**, and subsequently yields the amidoxyl radical **B**.^[10] We



Scheme 1. Postulated mechanism and stereochemical course.

envisage that **B** becomes a stable allyl radical **C** by hydrogen transfer. Concurrently, air oxidation of Cu^I to Cu^{II}, as with C, enables the same SET to form the bis(allyl) radical D, further inducing a dimerization to generate species E. Notably, E should preferably have a *trans* configuration for two bulky alkenyl groups to minimize steric hindrance. We envisage that **E** tends to release Cu^I because of its highly substituted and congested geometry, thus yielding the dimers \mathbf{F} only in *dl*forms. This process rationalizes well the asymmetric induction of the chiral copper catalyst [Eq. (5)]. As shown by the ground-state staggered conformations, F undergoes SET to generate the amidoxyl radical G to activate double radical/ alkene cyclizations,^[10] thus yielding the tertiary carbon radical H efficiently. Cu^{II} oxidation of these carbon radicals form to activate double radical/alkene cyclizations,^[10] thus yielding the stable tertiary cation I, and ultimately giving the desired compounds 6 upon ring closure. Such cascade radical cyclizations are consistent with the stereochemical course of these dimerizations.

Angew. Chem. Int. Ed. 2014, 53, 1-5

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

www.angewandte.org

Angewandte Communications

In this mechanism, we did not observe a 5-endo-trig radical cyclization for the monomeric species **B** because this process is not feasible according to the Baldwin's rule.^[19] In contrast, the 5-exo-trig cyclization of **G** is operable in accord with this rule. We postulated that its rigid staggered conformation allows the amidoxy radical close to its alkenyl group to facilitate this process.

In summary, we report the copper-catalyzed aerobic oxidation of N-hydroxyaminopropenes to form N- and Ocontaining cyclohexane derivatives. Such oxidations proceed with surprising stereocontrol and high efficiency. Reductive cleavage of the N-O bonds of these functionalized cyclohexanes affords 1,4-dihydroxy-2,3-diaminocyclohexanes which are the structural skeletons of several bioactive molecules. Our control experiments exclude the formation of nitrone intermediates, whereas chiral copper catalysts can induce asymmetric induction. We postulate a mechanism in which the key step involves the coordination of Cu^{II} with two allyl radicals to achieve a stereoselective C-C bond formation. These *dl*-configured dimeric intermediates presumably form amidoxyl radicals to undergo tandem radical/alkene cyclizations with high stereocontrol, thus giving observed products efficiently.

Received: August 5, 2014 Published online: ■■■■, ■■■■

Keywords: copper · cyclization · oxidation · radicals · synthetic methods

- For Selected reviews, see: a) S. A. Girard, T. Knauber, C.-J. Li, Angew. Chem. Int. Ed. 2014, 53, 74–100; Angew. Chem. 2014, 126, 76–103; b) A. E. Wendlandt, A. M. Suess, S. S. Stahl, Angew. Chem. Int. Ed. 2011, 50, 11062–11087; Angew. Chem. 2011, 123, 11256–11283; c) M. Klussmann, D. Sureshkumar, Synthesis 2011, 353–369; d) C. Liu, H. Zhang, W. Shi, A. Lei, Chem. Rev. 2011, 111, 1780–1824; e) C. Zhang, C. Tang, N. Jiao, Chem. Soc. Rev. 2012, 41, 3464–3484.
- [2] Selected examples, see: a) E. Boess, C. Schmits, M. Klussmann, J. Am. Chem. Soc. 2012, 134, 5317-5325; b) J. Zhang, B. Tiwari, C. Xing, X. Chen, Y. R. Chi, Angew. Chem. Int. Ed. 2012, 51, 3649-3652; Angew. Chem. 2012, 124, 3709-3712; c) M. Ghobrial, M. Schürch, M. D. Mihovilovic, J. Org. Chem. 2011, 76, 8781-8793; d) J. Xie, Z.-Z. Huang, Angew. Chem. Int. Ed. 2010, 49, 10181-10185; Angew. Chem. 2010, 122, 10379-10383; e) F. Yang, J. Li, J. Xie, Z.-Z. Huang, Org. Lett. 2010, 12, 5214-5217; f) L. Zhao, C.-J. Li, Angew. Chem. Int. Ed. 2008, 47, 7075-7078; Angew. Chem. 2008, 120, 7183-7186; g) E. Boess, D. Sureshkumar, A. Sud, C. Wirtz, C. Farès, M. Klussmann, J. Am. Chem. Soc. 2011, 133, 8106-8109; h) O. Baslé, C.-J. Li, Green Chem. 2007, 9, 1047-1050; i) Y. Shen, M. Li, S. Wang, T. Zhan, Z. Tan, C.-C. Guo, Chem. Commun. 2009, 953-955; j) O. Baslé, C.-J. Li, Chem. Commun. 2009, 4124-4126; k) J.-S. Tian, T.-P. Loh, Angew. Chem. Int. Ed. 2010, 49, 8417-8420; Angew. Chem. 2010, 122, 8595-8598.
- [3] a) W. Han, A. R. Ofial, *Chem. Commun.* 2009, 5024–5026; b) P. Liu, C.-Y. Zhou, S. Xiang, C.-M. Che, *Chem. Commun.* 2010, 46, 2739–2741; c) G. Kumaraswamy, A. N. Murthy, A. Pitchaiah, *J. Org. Chem.* 2010, 75, 3916–3919; d) E. Shirakawa, N. Uchiyama, T. Hayashi, *J. Org. Chem.* 2011, 76, 25–34; e) M. Ghobrial,

K. M. Harhammer, D. Mihovilovic, M. Schnürch, *Chem. Commun.* **2010**, *46*, 8836–8838; f) C. M. R. Volla, P. Vogel, *Org. Lett.* **2009**, *11*, 1701–1704; g) M. O. Ratnikov, X. Xu, M. P. Doyle, *J. Am. Chem. Soc.* **2013**, *135*, 9475–9479.

- [4] a) M.-Z. Wang, C.-Y. Zhou, M.-K. Wong, C.-M. Che, *Chem. Eur. J.* 2010, *16*, 5723–5735; b) S.-I. Murahashi, T. Naota, K. Yonemura, *J. Am. Chem. Soc.* 1988, *110*, 8256–8258; c) S.-I. Murahashi, T. Naota, T. Kuwabara, T. Saito, H. Kumobayashi, S. Akutagawa, *J. Am. Chem. Soc.* 1990, *112*, 7820–7822; d) S.-I. Murahashi, N. Komiya, H. Terai, T. Nakae, *J. Am. Chem. Soc.* 2003, *125*, 15312–15313; e) S.-I. Murahashi, T. Saito, T. Naota, H. Kumobayashi, S. Akutagawa, *Tetrahedron Lett.* 1991, *32*, 5991–5994; f) S.-I. Murahashi, T. Nakae, H. Terai, N. Komiya, *J. Am. Chem. Soc.* 2008, *130*, 11005–11012; g) A. Yu, Z. Gu, D. Chen, W. He, P. Tan, J. Xiang, *Catal. Commun.* 2009, *11*, 162–166.
- [5] a) A. J. Catino, J. M. Nichols, B. J. Nettles, M. P. Doyle, J. Am. Chem. Soc. 2006, 128, 5648-5649; b) M. O. Ratnikov, M. P. Doyle, J. Am. Chem. Soc. 2013, 135, 1549-1557; c) K. Alagiri, K. R. Prabhu, Org. Biomol. Chem. 2012, 10, 835-842.
- [6] a) A. Sud, D. Sureshkumar, M. Klussmann, *Chem. Commun.* 2009, 3169-3171; b) S. Singhal, S. L. Jain, B. Sain, *Chem. Commun.* 2009, 2371-2372.
- [7] a) G. Casiraghi, L. Battistini, C. Curti, G. Rassu, F. Zanardi, *Chem. Rev.* 2011, 111, 3076–3154; b) B. B. Touré, D. G. Hall, *Chem. Rev.* 2009, 109, 4439–4486.
- [8] H. C. J. Ottenheijm, J. D. M. Herscheid, Chem. Rev. 1986, 86, 697–707.
- [9] For copper-catalyzed aerobic oxidations of N-hydroxyamines to nitrone species, see: a) G. W. Gribble, T. C. Barden, J. Org. Chem. 1985, 50, 5900-5902; b) T. Sakurai, Y. Uematsu, O. Tanaka, H. Inoue, J. Chem. Soc. Perkin Trans. 2 1992, 2163-2167; c) R. K. Kawade, C. C. Tseng, R.-S. Liu, Chem. Eur. J. 2014, DOI: 10.1002/chem.201404201.
- [10] a) V. A. Schmidt, E. J. Alexanian, J. Am. Chem. Soc. 2011, 133, 11402-11405; b) B. C. Giglio, V. A. Schmidt, E. J. Alexanian, J. Am. Chem. Soc. 2011, 133, 13320-13322; c) V. A. Schmidt, E. J. Alexanian, Angew. Chem. Int. Ed. 2010, 49, 4491-4494; Angew. Chem. 2010, 122, 4593-4596; d) C. Berti, L. Grierson, J. A.-M. Grimes, M. J. Perkins, B. Terem, Angew. Chem. Int. Ed. Engl. 1990, 29, 653-655; Angew. Chem. 1990, 102, 684-685; e) V. A. Schmidt, E. J. Alexanian, Chem. Sci. 2012, 3, 1672-1674; f) R. K. Quinn, V. A. Schmidt, E. J. Alexanian, Chem. Sci. 2013, 4, 4030-4034.
- [11] For the stereoselective synthesis of the epimers of the 1,4dihydroxy-2,3-diaminocyclohexanes 7, see: C. Boga, C. Fiorelli, D. Savoia, *Synthesis* 2006, 285–292.
- [12] A. Trapero, A. Llebaria, ACS Med. Chem. Lett. 2011, 2, 614– 619.
- [13] A. Arcelli, V. Cere, F. Peri, S. Pollicino, A. Riccib, *Tetrahedron: Asymmetry* 2002, 13, 191–196.
- [14] S. Hanessian, J. Wang, Can. J. Chem. 1993, 71, 2102-2108.
- [15] J. Magano, Tetrahedron 2011, 67, 7875-7899.
- [16] CCDC 1013973 (6a) and 1013974 (7b) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [17] a) R. A. Stockman, A. Sinclair, L. G. Arini, P. Szeto, D. L. Hughes, J. Org. Chem. 2004, 69, 1598-1602; b) C. Ji, M. J. Miller, Tetrahedron Lett. 2010, 51, 3789-3791; c) T. Sasaki, S. Eguchi, T. Suzuki, J. Org. Chem. 1982, 47, 5250-5255.
- [18] A summary of asymmetric dimerizations of 5a using various chiral copper catalysts and solvents is provided in Table S1 of the Supporting Information.
- [19] J. E. Baldwin, J. Chem. Soc. Chem. Commun. 1976, 738-741.

www.angewandte.org

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

Communications



Copper-Catalyzed Oxidative Dimerizations of 3-N-Hydroxyaminoprop-1-enes to form 1,4-Dihydroxy-2,3-diaminocyclohexanes with C_2 Symmetry

Skeletons! Copper-catalyzed aerobic oxidations of *N*-hydroxyaminopropenes to form C_2 -symmetric N- and O-functionalized cyclohexanes are described. The catalytic oxidations proceed with remarkable stereocontrol and high efficiency.

 R^2

όн

Cu (5 mol %)

oxidant

THF, 25 °C

Reductive cleavage of the two N-O bonds of these products delivers 1,4dihydroxy-2,3-diaminocyclohexanes, which are important skeletons of several bioactive molecules.

 R^1

Pd/C, H₂

OH

R

einheim www.angewandte.org 5 These are not the final page numbers!