### Accepted Manuscript

One-pot, microwave-assisted synthesis of polymethylene-bridged bis(1H-1,2,4-triazol-5(3)-amines) and their tautomerism

Felicia Phei Lin Lim, Lee Ming Hu, Edward R.T. Tiekink, Anton V. Dolzhenko

PII:	S0040-4039(18)31099-2
DOI:	https://doi.org/10.1016/j.tetlet.2018.09.018
Reference:	TETL 50257
To appear in:	Tetrahedron Letters
Received Date:	18 July 2018
Revised Date:	30 August 2018
Accepted Date:	6 September 2018



Please cite this article as: Lim, F.P.L., Hu, L.M., Tiekink, E.R.T., Dolzhenko, A.V., One-pot, microwave-assisted synthesis of polymethylene-bridged bis(1*H*-1,2,4-triazol-5(3)-amines) and their tautomerism, *Tetrahedron Letters* (2018), doi: https://doi.org/10.1016/j.tetlet.2018.09.018

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### **Graphical Abstract**

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.





Tetrahedron Letters

# One-pot, microwave-assisted synthesis of polymethylene-bridged bis(1H-1,2,4-triazol-5(3)-amines) and their tautomerism

Felicia Phei Lin Lim<sup>a</sup>, Lee Ming Hu<sup>a</sup>, Edward R. T. Tiekink<sup>b</sup> and Anton V. Dolzhenko<sup>a,c</sup>\*

<sup>a</sup> School of Pharmacy, Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway, Selangor Darul Ehsan 47500, Malaysia

<sup>b</sup> Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, Bandar Sunway, Selangor Darul Ehsan 47500, Malaysia <sup>c</sup> School of Pharmacy and Biomedical Sciences, Curtin Health Innovation Research Institute, Faculty of Health Sciences, Curtin University, GPO Box U1987 Perth, Western Australia 6845, Australia

#### ARTICLE INFO

Article history: Received Received in revised form Accepted Available online

Keywords: Triazole Dicarboxylic acid Microwave-assisted synthesis One-pot synthesis Tautomerism

#### ABSTRACT

A highly selective and efficient method for the synthesis of 3,3'(5,5')-polymethylene-bis(1H-1,2,4-triazol-5(3)-amines) was developed using the reaction of dicarboxylic acids and aminoguanidine in an aqueous medium. This one-pot, microwave-promoted method was proved to be scalable affording the desired products in good yields and purity. The scope of the method was successfully explored by the preparation of a small library of polymethylene-bis(1H-1,2,4-triazol-5(3)-amines) with different alkyl chain linkers. The annular prototropic tautomerism in the prepared compounds was studied using NMR spectroscopy and X-ray crystallography.

2018 Elsevier Ltd. All rights reserved.

The twin-drugs strategy involves linking two identical pharmacophores in one molecule by a chemically and pharmacologically inert bridge. It is an efficient approach to significantly improve the effectiveness of bioactive agents. A polymethylene chain is often used as a linker between the pharmacophores in twin-drugs and the length of this chain is adjusted to achieve optimal therapeutic properties.

Many symmetrical polymethylene-bridged diamines, their derivatives and analogues, including bis(*N*-heterocycles), have been found to possess antiparasitic,<sup>1</sup> antibacterial,<sup>2</sup> and anticancer<sup>3</sup> properties. Some notable recent examples of these bioactive compounds include highly potent antimalarial diamine 1;<sup>1d</sup> bis-thiourea 2 inhibiting soluble epoxide hydrolase activity at a low nanomolar concentration range;<sup>4</sup> bis-imidazoles **3** and **4** possessing strong anti-biofilm activity against *Staphylococcus aureus* and *Pseudomonas aeruginosa*, respectively;<sup>2c</sup> compound **5** effectively bis-intercalating into the tatelomeric DNA/RNA duplex;<sup>3e</sup> and bis(1*H*-1,2,4-triazol-5-amine) **6** possessing anticancer properties<sup>3f</sup> (Fig. 1).

Bis-triazole **6** was previously synthesized using the reaction of bis-urea **8** or bis-thiourea **9**, which were prepared from sebacoyl chloride (**7**), with hydrazine (Scheme 1).<sup>3f</sup> This two-step process produced **6** in 59-61% overall yield. However, using a relatively unstable acid chloride and extended reaction times (4-8 h for each step) makes this process less practical. Moreover, the

selectivity of all steps is questionable as confirmed by the low purity of products (by reported elemental analysis data for 6, 8, and 9).<sup>3f</sup>



Figure 1. Selected bioactive symmetrical polymethylene-bridged compounds.

\* Corresponding author. Tel.: +60 3 5514 5867; fax: +60 3 5514 6364; e-mail addresses: dolzhenkoav@gmail.com, anton.dolzhenko@monash.edu (A.V. Dolzhenko).

#### ACCEPTED MANUSCRIPT Tetrahedron Letters

Scheme 1. Synthesis of octamethylene-bis(1*H*-1,2,4-triazol-5-amine) (6).<sup>3f</sup>



2

Scheme 2. Synthesis of decamethylene-bis(1H-1,2,4-triazol-5-amine) (11).<sup>5</sup>

A homologue of compound 6, decamethylene-bis(1H-1,2,4triazol-5-amine) (11) was prepared via the cyclization of bisamidoguanidine 10 (Scheme 2).<sup>5</sup> This reaction required heating at high temperature (160-180 °C) for 3 h but afforded 11 in rather modest yield (26%). Previously, we developed an effective method for the cyclization of amidoguanidines in water under irradiation microwave affording the corresponding aminotriazoles in excellent yields.6 This approach was further elaborated for the synthesis of more complex compounds in a one-pot manner.<sup>7</sup> Therefore, we decided to adopt a similar onepot, microwave-promoted methodology to prepare 3,3'(5,5')polymethylene-bis(1H-1,2,4-triazol-5(3)-amines) from readily available dicarboxylic acids.

As the model reaction, we decided to examine the condensation of succinic acid with aminoguanidine. This reaction under conventional heating was reported to suffer from a lack of selectivity resulting in the formation of a mixture of products, from which pure ethylene-bis(1*H*-1,2,4-triazol-5-amine) **12** was isolated in 8-11% yield only.<sup>8</sup> To overcome this problem, we optimized the reaction conditions under microwave irradiation.

Initially, using the Discover SP (CEM) microwave synthesizer, succinic acid was heated with aminoguanidine hydrochloride in water at 200 °C for 15 min. After cooling, an aqueous solution of potassium carbonate was added to the reaction mixture and heating was continued at 200 °C for 5 min. To our satisfaction, we observed the formation of ethylenebis(1H-1,2,4-triazol-5-amine) 12, which was isolated via filtration (Entry 1). To further improve the reaction yield, we continued the optimization process and found that increasing the amount of potassium carbonate led to a slight increase in product yield (Entry 2). However, further optimization of the reaction conditions, particularly increasing the temperature, was limited due to exessive pressure generated in the reaction vessel. Screening other bases for the reaction (Entries 3-4), we identified that under similar conditions sodium hydroxide was equally efficient as a base in terms of yield. Hence, the optimization was continued using sodium hydroxide as a base. Further variations in the reaction parameters revealed that 12 could be obtained in good yield by heating succinic acid and aminoguanidine hydrochloride in water at 220 °C for 15 min followed by the addition of an aqueous solution of sodium hydroxide and further microwave irradiation at 200 °C for 5 min (Entry 16). It should be noted that the exclusive presence of ethylene-bis(1H-1,2,4triazol-5-amine) 12 was observed in the crude product isolated from the reactions.

To explore the scope of the method, various dicarboxylic acids were used as substrates in the reaction with aminoguanidine hydrochloride to produce polymethylene-bis(1H-1,2,4-triazol-5(3)-amines). The optimized conditions<sup>9</sup> were found to be sufficiently general and suitable for the synthesis of compounds **6**, **11-17** in good yields (Table 2). In all cases, the reaction was selective and only polymethylene-bis(1H-1,2,4-triazol-5(3)-amines) were isolated in high purity *via* filtration. The reaction was also effectively performed on 3.4 mmol and 17 mmol scales with similar outcomes.

It should also be noted that the developed method was successfully tested using a Monowave 450 (Anton Paar) microwave synthesizer for the synthesis of **12** under the conditions optimized using the Discover SP (CEM). The outcome was satisfactory affording **12** in 91% yield, therefore confirming the reproducibility of the method in different microwave reactors.

Table 1. Optimization of the reaction conditions for the synthesis of ethylene-bis(1H-1,2,4-triazol-5(3)-amine) 12<sup>a</sup>



Entry	Molar ratio of succinic acid: aminoguanidine	Conditions	Isolated yield 12 (%)
1	1:2	<i>i</i> : water, 200 °C, 15 min; <i>ii</i> : K <sub>2</sub> CO <sub>3</sub> (0.5 equiv.), water, 200 °C, 5 min	33
2	1:2	<i>i</i> : water, 200 °C, 15 min; <i>ii</i> : K <sub>2</sub> CO <sub>3</sub> (1.0 equiv.), water, 200 °C, 5 min	37
3	1:2	<i>i</i> : water, 200 °C, 15 min; <i>ii</i> : NaHCO <sub>3</sub> (1.0 equiv.), water, 200 °C, 5 min	27
4	1:2	i: water, 200 °C, 15 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min	36
5	1:3	i: water, 200 °C, 15 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min	58
6	1:4	i: water, 200 °C, 15 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min,	66
7	1:5	i: water, 200 °C, 15 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min	79
8	1:6	i: water, 200 °C, 15 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min	74
9	1:7	i: water, 200 °C, 15 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min	71
10	1:5	i: water, 200 °C, 10 min; ii: NaOH (1.0 equiv.), water, 200 °C, 5 min	58
11	1:5	<i>i</i> : water, 180 °C, 30 min; <i>ii</i> : NaOH (1.0 equiv.), 200 °C, 5 min	58
12	1:5	<i>i</i> : water, 200 °C, 15 min; <i>ii</i> : NaOH (1.0 equiv.), 180 °C, 5 min	68
13	1:5	<i>i</i> : water, 180 °C, 15 min; <i>ii</i> : NaOH (1.0 equiv.), 200 °C, 5 min	46
14	1:5	i: water, 190 °C, 15 min; ii: NaOH (1.0 equiv.), 200 °C, 5 min	54
15	1:5	<i>i</i> : water, 210 °C, 15 min; <i>ii</i> : NaOH (1.0 equiv.), 200 °C, 5 min	79
16	1:5	i: water, 220 °C, 15 min; ii: NaOH (1.0 equiv.), 200 °C, 5 min	85
17	1:5	<i>i</i> : water, 230 °C, 15 min; <i>ii</i> : NaOH (1.0 equiv.), 200 °C, 5 min	77

<sup>a</sup> Reaction was performed using a Discover SP (CEM) microwave synthesizer on a 3.4 mmol scale (succinic acid) using 2 mL of water in the first step, followed by the addition of 1 mL of the specified aqueous base in the second step.

R

3

......

Tetrahedron Letters

1) H<sub>2</sub>O, MW, 220 °C, 15 min;

2) NaOH, MW, 200 °C, 5 min

Compound	Structure	Scale (mmol)	Yield (%)
12		3.4	85
	$H_2N$ $N$ $N-NH$	17	88
13		3.4	58
$H_2N^-$		17	63
14	HN-N NH2	3.4	77
	H <sub>2</sub> N N N-NH	17	66
15		3.4	52
		17	56
16	HN-N	3.4	72
		17	65
17		3.4	57
		17	54
6		3.4	63
	$H_2N \sim N$	17	63
11	HN-N A N	3.4	72
	$H_2N$	17	67

Table 2. One-pot synthesis of polymethylene-bis(1H-1,2,4-triazol-5(3)-amines).<sup>a</sup>

H<sub>2</sub>N NH o-

0 0

<sup>a</sup> Reaction was performed in water using a Discover SP CEM microwave synthesizer at 220 °C for 15 min, followed by the addition of aq. NaOH and heating at 200 °C for 5 min.

The annular prototropic tautomerism, possible in both triazole rings of the prepared compounds, might result in six different tautomers representing different combinations of the 5-amino-1H-1,2,4-triazole, 3-amino-1H-1,2,4-triazole, and 5-amino-4H-1,2,4-triazole rings (Scheme 3). These three principal forms of the 1,2,4-triazole ring can be distinguished on the basis of NMR spectroscopy data (though without allowing assigning the presence of individual tautomers). The <sup>1</sup>H NMR spectra of the

products in DMSO-*d*<sub>6</sub> gave two sets of signals for the protons of the primary amino group and the triazole ring. These signals together with two sets of signals for the triazole ring carbon atoms in the <sup>13</sup>C NMR spectra served for the assignment of two main forms (**A** and **B**) of the triazole ring (three potential tautomers). The assignments were confirmed with the literature data,<sup>6-8,10</sup> and the equilibrium constant ( $K_T$ ) and corresponding Gibbs free energy ( $\Delta G_{300}$ ) values were estimated (Table 3).

N-NH



Scheme 3. Theoretically possible tautomers of polymethylene-bis(1H-1,2,4-triazol-5(3)-amines).

4

et fuutomerie com	position porjine		, $1 \text{ tild 201 } \mathcal{O}(\mathcal{O})$ to		
			$H_2N \xrightarrow{N-N}_2 (CH_2)$	$ \begin{array}{c} N - NH \\ H_2 N \\ N \\ B \\ 2 \end{array} $	
Compounds	n	Composition (%)			A Gase (k Impl <sup>-1</sup> )
		Α	В	κ <sub>[</sub>	
12	2	72	28	2.57	2.35
13	3	68	32	2.13	1.89
14	4	67	33	2.03	1.77
15	5	66	34	1.94	1.65
16	6	66	34	1.94	1.65
17	7	65	35	1.86	1.55
6	8	65	35	1.86	1.55
11	10	65	35	1.86	1.55

Table 3. Tautomeric composition polymethylene-bis(1,2,4-triazol-5(3)-amines) in DMSO-d6.ª

<sup>a</sup> Assigned by NMR spectroscopy performed at 27 °C.

In DMSO- $d_6$  solution, the tautomeric equilibria was shifted mainly towards the 5-amino-1*H*-form (**A**) of the triazole ring with the 3-amino-1*H*-form (**B**) being less prominent and the 4*H*form (**C**) undetectable. The expansion of the polymethylene link between the triazole rings led to an increase in the 3-amino-1*H*form (**B**) in the equilibrium. This trend can be explained by the diminishing distance electron-withdrawing effect of one triazole ring on another. However, for compounds possessing seven or more methylenic groups in the linking chain, the equilibrium constant remained the same. It should be noted that the tautomer signals were indistinguishable when the NMR spectra of the compounds were recorded in CDCl<sub>3</sub>.

In crystals, the tautomerizable 5(3)-amino-1,2,4-triazoles usually appear in the 5-amino-1*H*-tautomer form.<sup>11</sup> An example of 5-amino-1*H*- and 3-amino-1*H*-tautomers crystallising together has also been reported.<sup>12</sup> With the objective to investigate the tautomeric preference in crystals of the prepared compounds, we performed single crystal X-ray crystallography for representative compound **14**.<sup>13</sup> The molecular structure of **14**, shown in Fig. 2(a), revealed that this compound crystallised with the 3-amino-1*H*-1,2,4-triazole rings on both sides of the molecule. This tautomeric form is not typical for crystals of tautomerizable 5(3)-amino-1,2,4-triazoles and represents a rather unique case of finding 3-amino-1*H*-1,2,4-triazoles in crystals.



**Figure 2.** Molecular structure of **14** showing atom labelling and 70% probability displacement ellipsoids; unlabelled atoms are related by the symmetry operation -x, 2-y, -z.

In conclusion, we have developed an efficient and practical one-pot, microwave-promoted synthesis of polymethylenebis(1H-1,2,4-triazol-5(3)-amines) from readily available dicarboxylic acids. The method was found to be scalable and transferable to another microwave reactor. The tautomeric preferences in the products were determined in solution (by NMR spectroscopy) and solid state (X-ray crystallography). The tautomeric equilibrium in the solution depended on the length of the chain connecting the triazole rings. A unique crystallisation of the 5-substituted 3-amino-1H-1,2,4-triazole tautomeric form was also observed.

#### Acknowledgments

The Research Centre for Crystalline Materials (Sunway University) is thanked for the X-ray intensity data. This work is supported by the Ministry of Higher Education, Malaysia under Fundamental Research Grant Scheme, grant number FRGS/1/2015/SG01/MUSM/03/1.

#### **References and notes**

- (a) Venkatraj, M.; Salado, I. G.; Heeres, J.; Joossens, J.; Lewi, P. J.; Caljon, G.; Maes, L.; Van der Veken, P.; Augustyns, K. Eur. J. Med. Chem. 2018, 143, 306-319; (b) Schmidt, I.; Pradel, G.; Sologub, L.; Golzmann, A.; Ngwa, C. J.; Kucharski, A.; Schirmeister, T.; Holzgrabe, U. Bioorg. Med. Chem. 2016, 24, 3636-3642; (c) Fielding, A. J.; Lukinović, V.; Evans, P. G.; Alizadeh-Shekalgourabi, S.; Bisby, R. H.; Drew, M. G. B.; Male, V.; Del Casino, A.; Dunn, J. F.; Randle, L. E.; Dempster, N. M.; Nahar, L.; Sarker, S. D.; Cantú Reinhard, F. G.; de Visser, S. P.; Dascombe, M. J.; Ismail, F. M. D. Chem. Eur. J. 2017, 23, 6811-6828; (d) Panozzo-Zénere, E. A.; Porta, E. O. J.; Arrizabalaga, G.; Fargnoli, L.; Khan, S. I.; Tekwani, B. L.; Labadie, G. R. Eur. J. Med. Chem. 2018, 143, 866-880.
- (a) Suzuki, K.; Nomura, I.; Ninomiya, M.; Tanaka, K.; Koketsu, M. Bioorg. Med. Chem. Lett. 2018, in press, doi: 10.1016/j.bmcl.2018.06.050; (b) Hua, X.; Zhang, G.; Zhang, D.; Wu, Y. J. Heterocycl. Chem. 2018, 55, 1504-1508; (c) Trang, T. T. T; Dieltjens, L.; Hooyberghs, G.; Waldrant, K.; Ermolat'ev, D. S.; Van der Eycken, E. V.; Steenackers, H. P. L. Bioorg. Med. Chem. 2018, 26, 1470-1480.

#### Tetrahedron Letters

, North

- (a) Kumar, A.; Banerjee, S.; Roy, P.; Sondhi, S. M.; Sharma, A. Bioorg. Med. Chem. Lett. 2017, 27, 501-504; (b) Kopsida, M.; Barron, G. A.; Bermano, G.; Kong Thoo Lin, P.; Goua, M. Org. Biomol. Chem. 2016, 14, 9780-9789; (c) Sosic, A.; Zuravka, I.; Schnitt, N.-K.; Miola, A.; Goettlich, R.; Fabris, D.; Gatto, B. ChemMedChem 2017, 12, 1471-1479; (d) Santiago-Vazquez, Y.; Das, S.; Das, U.; Robles-Escajeda, E.; Ortega, N. M.; Lema, C.; Varela-Ramirez, A.; Aguilera, R. J.; Balzarini, J.; De Clercq, E.; Dimmock, S. G.; Gorecki, D. K. J.; Dimmock, J. R. Eur. J. Med. Chem. 2014, 77, 315-322; (e) Islam, M. K.; Jackson, P. J. M.; Thurston, D. E.; Rahman, K. M. Org. Biomol. Chem. 2018, 16, 4424-4428; (f) I El Azab, I. H.; Elkanzi, N. A. A. Curr. Org. Synth. 2017, 14, 309-320.
- Burmistrov, V.; Morisseau, C.; Pitushkin, D.; Karlov, D.; Fayzullin, R. R.; Butov, G. M.; Hammock, B. D. Bioorg. Med. Chem. Lett. 2018, 28, 2302-2313.
- Fujisaki, F.; Usami, H.; Nakashima, S.; Nishida, S.; Fujioka, T.; Kashige, N.; Miake, F.; Sumoto, K. *Heterocycles* 2013, 87, 665-675
- Dolzhenko, A. V.; Pastorin, G.; Dolzhenko, A. V.; Chui, W. K. Tetrahedron Lett. 2009, 50, 2124-2128.
- Lim, F. P. L.; Tan, L. Y.; Tiekink, E. R. T.; Dolzhenko, A. V. RSC Adv. 2018, 8, 22351-22360.
- Chernyshev, V. M.; Chernysheva, A. V.; Starikova, Z. A. Heterocycles 2010, 81, 2291-2311.
- 9 General method for the microwave-assisted synthesis of bis(5(3)amino-1,2,4-triazol-3(5)-yl)polymethylenes (6, 11-17). A mixture of the corresponding dicarboxylic acids (3.4 mmol) and aminoguanidine hydrochloride (1.88 g, 17 mmol) in water (2 mL) was irradiated in a 10 mL seamless pressure vial using the microwave system operating at maximum microwave power (up to 300 W) at 220 °C for 15 min. After cooling to ambient temperature, of 17 M aq. NaOH (1 mL) was added to the vial and the reaction mixture was irradiated again at 200  $^{\circ}\!\mathrm{C}$  for 10 min. After cooling to ambient temperature, the precipitated solid was filtered, washed with cold water and recrystalised from water to give desired product. The reaction was also replicated on an increased scale of the dicarboxylic acids (17 mmol) and aminoguanidine hydrochloride (9.40 g, 85 mmol) in water (10 mL).
- Buzykin, B. I.; Mironova, E. V.; Nabiullin, V. N.; Gubaidullin, A. T.; Litvinov, I. A. *Russ. J. Gen. Chem.* **2006**, *76*, 1471-1486.
- (a) Garcia, E.; Lee, K. Y. Acta Crystallogr., Sect. C: Cryst. Struct. Commun. 1992, C48, 1682-1683; (b) Borbulevych, O. Y.; Shishkin, O. V.; Desenko, S. M.; Chernenko, V. N.; Orlov, V. D. Acta Crystallogr., Sect. C: Cryst. Struct. Commun. 1998, C54, 442-444; (c) Dolzhenko, A. V.; Tan, G. K.; Koh, L. L.; Dolzhenko, A. V.; Chui, W. K. Acta Crystallogr., Sect. E: Struct. Rep. Online 2009, 65, o125; (d) Zhang, S.; Liu, P.-J.; Ma, D.-S.; Hou, G.-F. Acta Crystallogr., Sect. E: Struct. Rep. Online 2012, 68, o83; (e) Liu, B.; Tome, J. P. C.; Cunha-Silva, L.; Almeida Paz, F. A. Acta Crystallogr., Sect. E: Struct. Rep. Online 2012, 68, o2700-o2701; (f) Hernandez-Gil, J.; Ferrer, S.; Ballesteros, R.; Castineiras, A. Acta Crystallogr., Sect. E: Struct. Rep. Online 2013, 69, o227-o228.
- Dolzhenko, A. V.; Tan, G. K.; Koh, L. L.; Dolzhenko, A. V.; Chui, W. K. Acta Crystallogr., Sect. E: Struct. Rep. Online 2009, 65, o126.
- 13. Crystal data for  $C_8$ H<sub>14</sub>N<sub>8</sub> (14): M = 222.27, monoclinic,  $P_{21}/n$ , a = 5.00860(10), b = 5.1190(2), c = 19.5646(6) Å,  $\beta = 93.306(3)^\circ$ , V = 500.78(3) Å<sup>3</sup>, Z = 2,  $D_x = 1.474$  g cm<sup>3</sup>, F(000) = 236,  $\mu = 0.844$  mm<sup>-1</sup>. CCDC 1851291.

#### **Supplementary Material**

Supplementary data associated with this article can be found, in the online version, at http://dx.doi.org/10.1016/j.tetlet.2018.XX.XXX. These data include experimental details, copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra of the prepared compounds, and X-ray crystallography data for compound **14**.

6

- Acctinition

7