

Note

Nickel/N-Heterocyclic Carbene-Catalyzed Suzuki-Miyaura Type Cross-Coupling of Aryl Carbamates

Akimichi Ohtsuki, Kousuke Yanagisawa, Takayuki Furukawa, Mamoru Tobisu, and Naoto Chatani

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.6b01627 • Publication Date (Web): 07 Sep 2016

Downloaded from <http://pubs.acs.org> on September 7, 2016

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3 **Nickel/N-Heterocyclic Carbene-Catalyzed Suzuki-Miyaura Type Cross-Coupling of Aryl**
4
5 **Carbamates**

6
7
8 Akimichi Ohtsuki, Kousuke Yanagisawa, Takayuki Furukawa, Mamoru Tobisu,^{*,†} and Naoto

9
10
11 Chatani*

12
13 *Department of Applied Chemistry, Faculty of Engineering, Osaka University, Suita, Osaka 565-*
14
15 *0871, Japan*

16
17
18 [†]*Center for Atomic and Molecular Technologies, Graduate School of Engineering, Osaka*
19
20 *University, Suita, Osaka 565-0871, Japan*

21
22 *tobisu@chem.eng.osaka-u.ac.jp; chatani@chem.eng.osaka-u.ac.jp*
23
24
25
26
27
28

29
30 **Abstract:** The utility of N-heterocyclic carbene ligands in nickel-catalyzed Suzuki-Miyaura cross-
31
32 coupling of aryl esters and carbamates is investigated. Imidazol-2-ylidene bearing 2-adamantyl
33
34 groups at its nitrogen atoms generates the most active nickel species among the ligands examined,
35
36 allowing cross-coupling of a range of aryl carbamates and pivalates. Unlike the previously reported
37
38 system using tricyclohexylphosphine, this protocol is suitable for the cross-coupling using
39
40 arylboronic esters in addition to arylboronic acids.
41
42
43
44
45
46
47
48

49 The Suzuki-Miyaura reaction has been recognized as a powerful carbon-carbon bond formation
50
51 method because of the favorable features of organoboron reagents, such as high functional group
52
53 compatibility and ease of handling.¹ The Suzuki-Miyaura reaction is particularly useful for the
54
55 synthesis of (hetero)biaryls, in which various aryl halides and their equivalents are coupled with
56
57 arylboronic acid derivatives. Recent efforts have been directed toward the use of inert phenol
58
59 derivatives,² such as aryl ethers, esters and carbamates, as aryl halide surrogates for Suzuki-
60

Miyaura reactions. These efforts will not only provide cost-effective, halogen-free processes but also lead to diverse synthetic strategies such as orthogonal cross-coupling and late-stage functionalization. In this context, we have reported nickel-catalyzed Suzuki-Miyaura coupling of methoxyarenes via the cleavage of C(aryl)-O bonds.³ Although we initially used tricyclohexylphosphine (PCy₃) as a suitable ligand for this cross-coupling,^{3a} a subsequent study led us to identify 1,3-dicyclohexylimidazol-2-ylidene (ICy) as a superior ligand with broadened scope of methoxyarenes.^{3c} The outstanding activity of the Ni/ICy system⁴ for C(aryl)-O bond activation of methoxyarenes naturally encouraged us to consider whether this system is able to promote the C-O bond activation of other inert phenol derivatives. Aryl esters and carbamates have attracted substantial attention as a class of inert phenol derivatives for use in nickel-catalyzed cross-coupling reactions.² Although several groups have already revealed that the Ni/PCy₃ system is able to catalyze Suzuki-Miyaura type reactions of aryl esters⁵ and carbamates,⁶ the utility of an NHC ligand in these processes has not been verified.^{7,8} Herein, we report Ni/NHC-catalyzed cross-coupling of aryl esters and carbamates with arylboron reagents.


Table 1. Reported Conditions for Nickel-Catalyzed Suzuki-Miyaura Cross-Coupling of Aryl Esters and Carbamates

	catalyst	boron sources	reference
Garg	NiCl ₂ (PCy ₃) ₂	ArB(OH) ₂	5a, 6a
Shi	NiCl ₂ (PCy ₃) ₂	(ArBO) ₃ + 0.88 H ₂ O	5b, 6c
Snieckus	NiCl ₂ (PCy ₃) ₂	ArB(OH) ₂ + 0.1 (ArBO) ₃	6b


Reliable catalytic conditions used for the cross-coupling of aryl esters and carbamates with arylboron reagents are summarized in Table 1. Close inspection of the reported boron sources employed indicates that these reactions apparently need precise control of the amount of water

1
2
3 present in the reaction mixture. For example, Garg used arylboronic acid ArB(OH)_2 ,^{5a,6a} which
4
5 generates water by the reversible formation of arylboroxine $(\text{ArBO})_3$ at elevated temperature. Shi
6
7 used a mixture of $(\text{ArBO})_3$ and 0.88 equiv of H_2O ,^{5b,6c} while Snieckus employed a 10:1 mixture of
8
9 ArB(OH)_2 and $(\text{ArBO})_3$.^{6b} Although an excess amount of water is thought to be detrimental to
10
11 Ni(0) catalysis,^{6c} a suitable amount of water is essential, presumably to generate active Ni(0)
12
13 species by the reaction of $\text{NiCl}_2(\text{PCy}_3)_2$ with ArB(OR)_2 . To avoid the potential complexity
14
15 associated with the effect of water during ligand screening experiments, we decided to use
16
17 anhydrous conditions with Ni(cod)_2 as a catalyst and an arylboronic ester as a nucleophile. Thus, 2-
18
19 naphthyl carbamate (**1a**) was reacted with phenylboronic ester **2a** in the presence of Ni(cod)_2 and a
20
21 ligand using K_3PO_4 as a base at 120 °C (Table 2). When PCy_3 was used as the ligand under these
22
23 conditions, cross-coupling product **3aa** was formed in only 9% yield (entry 1).⁹ A series of NHC
24
25 ligands was then examined. *N*-Aryl substituted carbenes such as **L1** (entry 2) and **L2** (entry 3) were
26
27 found to be ineffective for this coupling. In contrast to our expectation, the use of **L3**, which
28
29 represents the most effective ligand for nickel-catalyzed Suzuki-Miyaura reaction of
30
31 methoxyarenes, afforded **3aa** in only 11% yield (entry 4). NHC ligands bearing adamantyl groups
32
33 led to better results. Whereas 1-adamantyl derivative **L4** provided a slight increase in the yield of
34
35 **3aa** to 27% (entry 5), isomeric 2-adamantyl ligand **L5**^{4c,10} delivered **3aa** in 88% yield (entry 6).
36
37 Having optimized ligand **L5** in hand, we next examined a less reactive substrate **1b**¹¹ using these
38
39 conditions, which resulted in a lower yield of the coupling product (entry 7). Further optimization
40
41 revealed that conducting the reaction at 150 °C with a lower catalyst loading (5 mol%) increased
42
43 the yield of **3ba** to 71% (entry 8). These conditions were used to explore the scope of this reaction.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Nickel/NHC-Catalyzed Cross-Coupling of Carbamate **1** with Phenylboronic Ester **2a**^a



entry	substrate	ligand	GC yield of 3aa or 3ba (%)
1 ^b	1a	PCy ₃	9
2	1a	L1	0
3	1a	L2	3
4	1a	L3	11
5	1a	L4	27
6	1a	L5	88
7	1b	L5	40
8 ^c	1b	L5	71




^aReaction conditions: **1** (0.30 mmol), **2a** (0.45 mmol), Ni(cod)₂ (0.030 mmol), NHC·HCl (0.12 mmol), NaO'Bu (0.12 mmol) and K₃PO₄ (1.4 mmol) in toluene (1.0 mL) at 120 °C for 24 h.

^bConducted in the absence of NaO'Bu. ^cPerformed using Ni(cod)₂ (0.015 mmol), **L5** (0.045 mmol) and NaO'Bu (0.045 mmol) at 150 °C for 20 h.

The scope of the reaction with respect to the boronic ester component was examined using **1b** as the substrate (Scheme 1). Both electron-rich and -deficient aryl groups can be introduced with comparable efficiency. Similar to the standard Suzuki-Miyaura reaction, functional groups such as amines (**3be**), esters (**3bf**), fluorides (**3bg**) and acetals (**3bi**) were well tolerated. In addition, heteroarylboronic esters including **2j** and **2k** successfully underwent the cross-coupling with **1b**.

Scheme 1. Scope of Arylboron Reagents^a




23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

^aReaction conditions: **1b** (0.30 mmol), arylboron reagent (0.45 mmol), Ni(cod)₂ (0.015 mmol), **L5** (0.045 mmol) NaO^tBu (0.045 mmol) and K₃PO₄ (1.4 mmol) in toluene (1.0 mL) at 150 °C for 20 h. ^b**1a** was used as the substrate. ^cConducted using Ni(cod)₂ (0.030 mmol), **L5** (0.090 mmol) and NaO^tBu (0.090 mmol) at 160 °C.

The scope of aryl carbamates was next investigated using **2b** as a coupling partner (Scheme 2). Although the Ni/**L5** system can catalyze the reactions of an electronically diverse set of aryl carbamates, electron-rich substrates such as **1d** required higher catalyst loading to obtain satisfactory yields of the products. When ortho substituted derivative **1g** was used as the substrate, the yield was decreased markedly because of the formation of an undesired reduced product (i.e., biphenyl with ca. 75% yield). The yield of **3gb** was improved by switching the ligand to **L3** and increasing the amount of boronic ester **2b**. Carbamate is known as a versatile directing group for ortho C-H functionalization reactions.¹² Therefore, the compatibility of the ortho substituent allows sequential carbamate-directed ortho functionalization/arylation of the carbamate directing group.¹³ Heteroaryl carbamates (i.e., **1i**) and aryl pivalates (i.e., **1j**) also underwent cross-coupling with **2b** under these conditions. Phenyl *p*-toluenesulfonate and *p*-tert-butylanisole were found to be much

less reactive under these conditions (34% and 0% yield, respectively, for the reaction with **2b**). Moreover, arylation at the benzylic position is also possible, as demonstrated by the reaction of **1k**.

Scheme 2. Scope of Aryl Carbamates^a



^aReaction conditions: substrate (0.30 mmol), **2b** (0.45 mmol), $\text{Ni}(\text{cod})_2$ (0.015 mmol), **L5** (0.045 mmol), NaO^tBu (0.045 mmol) and K_3PO_4 (1.4 mmol) in toluene (1.0 mL) at 150 °C for 20 h.

^bConducted using $\text{Ni}(\text{cod})_2$ (0.030 mmol), **L5** (0.090 mmol) and NaO^tBu (0.090 mmol) at 160 °C.

^cPerformed using **2b** (1.8 mmol), $\text{Ni}(\text{cod})_2$ (0.030 mmol), **L3** (0.090 mmol) and NaO^tBu (0.090 mmol) at 160 °C. ^dPerformed using **2e** (1.8 mmol), $\text{Ni}(\text{cod})_2$ (0.030 mmol), **L3** (0.090 mmol) and NaO^tBu (0.090 mmol) at 160 °C.

One advantage of using an NHC ligand over PCy_3 in the nickel-catalyzed Suzuki-Miyaura reaction is its tolerance of a protecting group of the boronic acid component (Table 3). When the Ni/**L5** system was used, neopentylglycolate **2b**, pinacolate **2b'** and free boronic acid **2b''** were all successfully coupled with aryl carbamate **1b** (entries 1,3 and 5, respectively). In stark contrast, boronic esters **2b'** and **2b''** did not form an appreciable amount of **3bb** under the conditions using PCy_3 (entries 2 and 4, respectively). The use of boronic esters is preferred over the parent boronic

acids when boronic acids are unstable¹⁴ or the boron compounds are prepared by C-H borylation.¹⁵ In such a situation, our catalytic system should offer a useful protocol for the arylation of aryl esters and carbamates.¹⁶ ArBF₃K was unreactive under these nickel/**L5**-catalyzed conditions.

Table 3. Effect of Ligands on the Reactivity of Boron Reagents^a

entry	boron reagent	ligand	yield of 2b (%)
1	2b	L5	70
2	2b	PCy ₃	trace
3	2b'	L5	39 (61) ^b
4	2b'	PCy ₃	trace
5	2b''	L5	81
6	2b''	PCy ₃	75

^aReaction conditions: **1b** (0.30 mmol), organoboron reagent (0.45 mmol), Ni(cod)₂ (0.015 mmol), **L5** (0.045 mmol), NaO^tBu (0.045 mmol) and K₃PO₄ (1.4 mmol) in toluene (1.0 mL) at 150 °C for 20 h. ^bConducted using Ni(cod)₂ (0.030 mmol), **L5** (0.090 mmol) and NaO^tBu (0.045 mmol) at 160 °C.

In conclusion, we revealed that 2-adamantyl-substituted NHC ligand **L5** serves as an active ligand for nickel-catalyzed Suzuki-Miyaura cross-coupling of aryl esters and carbamates. Use of **L5** as the ligand allows boronic esters to participate in the cross-coupling reaction, which is in sharp contrast to the previously reported catalytic system using PCy₃, in which boronic acids are essential. Although the Ni/PCy₃ system dominates in catalytic transformations involving the activation of

1
2
3 inert phenol derivatives,² we anticipate that this work will stimulate the application of NHC ligands
4
5
6 to such reactions.
7
8
9

10 11 12 13 **Experimental**

14
15 **General Information.** ¹H NMR and ¹³C NMR spectra were recorded at 400 and 100 MHz,
16
17 respectively, in CDCl₃ with tetramethylsilane as an internal reference standard. High resolution
18
19 mass spectra (HRMS) were recorded by EI using a double-focusing mass spectrometer. Column
20
21 chromatography was performed with SiO₂ (230-400 mesh). Unless otherwise noted, all of the
22
23 reagents used in this study were obtained from commercial suppliers and used as received without
24
25 further purification. Aryl carbamates used in this study are all known compounds and were
26
27 prepared by the reaction of the corresponding phenols with ClCONEt₂. Arylboronic esters used in
28
29 this study are also known compounds and were prepared by the reaction of the corresponding
30
31 arylboronic acids with 2,2-dimethylpropane-1,3-diol. Ni(cod)₂, NaO^tBu, K₃PO₄, **L1** and **L2** were
32
33 purchased from the commercial suppliers and used as received. **L3**,¹⁷ **L4**¹⁸ and **L5**^{4c} were prepared
34
35 according to the literature procedure.
36
37
38
39

40
41 *4-(Dimethylamino)phenyl diethylcarbamate (1d).* In a dried round-bottom flask, NaH (60%
42
43 dispersion in mineral oil, 230 mg, 5.8 mmol) was dissolved in DME (20 mL) and then the mixture
44
45 was cooled to 0 °C. 4-(Dimethylamino)phenol (660 mg, 4.8 mmol) was added portionwise. After
46
47 the mixture was stirred at 0 °C for 10 min, diethylcarbonyl chloride (781 mg, 5.8 mmol) was
48
49 added dropwise. The mixture was warmed to rt and then stirred for another 1.5 h. The resulting
50
51 mixture was quenched with H₂O and an aqueous solution of NaOH (1.0 M) was added. The
52
53 resulting mixture was extracted with Et₂O. The organic phase was washed with brine, dried over
54
55 MgSO₄ and concentrated in vacuo. The crude mixture was purified by flash column
56
57 chromatography over silica gel (eluted with hexane/AcOEt = 3/1) to give the target product as a
58
59
60

white solid (916 mg, 81%); Rf 0.20 (hexane/ AcOEt = 5/1); Mp: 37.5-38.0 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.15-1.25 (m, 6H), 2.91 (s, 6H), 3.45-3.55 (m, 4H), 6.72 (dt, *J* = 8.8, 2.2 Hz, 2H), 6.98 (dt, *J* = 8.8, 2.2 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 13.6, 14.4, 41.4, 41.9, 42.3, 113.6, 122.3, 142.8, 148.5, 155.1; IR(ATR): 3431 m, 2968 m, 2940 w, 2889 w, 2254 m, 1650 m, 1460 m, 1416 s, 1303 m, 1224 m, 1088 s, 1048 s, 907 s, 857 m, 805 w. HRMS (EI+) *m/z* [M]⁺ calcd for C₁₃H₂₀N₂O₂⁺ 236.1525, Found 236.1527.

Ethyl 3-((diethylcarbamoyl)oxy)benzoate (1f). In a dried round-bottom flask, NaH (60% dispersion in mineral oil, 480 mg, 12 mmol) was dissolved in DME (25 mL) and then the mixture was cooled to 0 °C. Ethyl 3-hydroxybenzoate (1.66 g, 10 mmol) was added portionwise. After the mixture was stirred at 0 °C for 10 min, diethylcarbamoyl chloride (1.63 g, 12 mmol) was added dropwise. The mixture was warmed to rt and then stirred for another 1.5 h. The resulting mixture was quenched with H₂O and an aqueous solution of NaOH (1.0 M) was added. The organic phase was washed with brine, dried over MgSO₄ and concentrated in vacuo. The crude mixture was purified by flash column chromatography over silica gel (eluted with hexane/AcOEt = 3/1) to give the target product as a clear oil (2.61 g, 98%); Rf 0.54 (hexane/ AcOEt = 1/1). ¹H NMR (400 MHz, CDCl₃) δ 1.18-1.28 (m, 6H), 1.39 (t, *J* = 7.2 Hz, 3H), 3.38-3.47 (m, 4H), 4.37 (q, *J* = 7.2 Hz, 2 H), 7.33-7.35 (m, 1H), 7.42 (t, *J* = 7.8 Hz, 1H), 7.73 (t, *J* = 1.8 Hz, 1H), 7.88 (dt, *J* = 8.0 Hz, 1.4 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 13.5, 14.38, 14.44, 42.1, 42.5, 61.3, 123.0, 126.4, 126.6, 129.2, 131.9, 151.6, 154.0, 166.0; IR(ATR): 2978w, 2936 w, 2904 w, 2874 w, 1713 s, 1589 w, 1473 w, 1455 w, 1445 w, 1415 m 1381 w, 1367 w, 1283 m, 1254 s, 1223 m, 1120 s, 1150 s, 1097 s, 1075 m, 1042 m, 1021 m, 965 m, 912 w, 866 w, 826 w, 784 w, 750 s, 684 m. HRMS (EI+) *m/z* [M]⁺ calcd for C₁₄H₁₉NO₄ 265.1314, Found 265.1312.

Typical Procedure. Ni(cod)₂ (4.2 mg, 0.015 mmol, 5 mol%), **L5** (17 mg, 0.045 mmol), NaO^tBu (4.3 mg, 0.045 mmol), and toluene (0.4 mL) were added to a 10 mL-sample vial with a Teflon-

1
2
3 sealed screwcap in a glove box with N₂-atmosphere. The resulting mixture was stirred at room
4
5 temperature for 5 min. K₃PO₄ (287 mg, 1.35 mmol), **1b** (58 mg, 0.30 mmol), **2b** (92 mg, 0.45
6
7 mmol), and toluene (0.6 mL) were added to the vial and the cap was closed. The contents of the
8
9 vial were stirred at 150 °C for 20 h. The reaction mixture was cooled to room temperature, and then
10
11 the crude mixture was filtered through a pad of silica gel, and analyzed by GC. The filtrate was
12
13 concentrated *in vacuo* to give a residue, which was purified by column chromatography over silica
14
15 gel (eluent: hexane/EtOAc =9/1) to give **3bb** as a white solid (35 mg, 70% yield).
16
17
18
19
20
21

22 *2-Phenylnaphthalene (3aa)*.¹⁹ The general procedure was used, except that the reaction was
23
24 conducted at 120 °C for 24 h. The yield was only determined by GC since this is a model
25
26 compound used for optimization studies (88% GC yield); ¹H NMR (400 MHz, CDCl₃) δ 7.39-7.41
27
28 (m, 1H), 7.47-7.52 (m, 4H), 7.72-7.77 (m, 3H), 7.86-7.94 (m, 3H), 8.05 (s, 1H); ¹³C NMR (100
29
30 MHz, CDCl₃) δ 125.7, 125.9, 126.1, 126.4, 127.5, 127.6, 127.8, 128.3, 128.6, 129.0, 132.8, 133.8,
31
32 138.7, 141.3. HRMS (EI+) m/z [M]⁺ calcd for C₁₆H₁₂ 204.0939, Found 204.0937.
33
34
35
36

37 *Biphenyl (3ba)*.²⁰ The general procedure was used, except that the reaction was conducted at
38
39 120 °C for 24 h. The yield was only determined by GC since this is a model compound used for
40
41 optimization studies (71% GC yield); ¹H NMR (400 MHz, CDCl₃) δ 7.35-7.39 (m, 2H), 7.44-7.48
42
43 (m, 4H), 7.61-7.63 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 127.3, 127.4, 128.9, 141.4. HRMS
44
45 (EI+) m/z [M]⁺ calcd for C₁₂H₁₀ 154.0783, Found 154.0780.
46
47
48

49 *4-Methyl-1,1'-Biphenyl (3bb)*.²¹ The general procedure was used; white solid (35 mg, 70% yield);
50
51 ¹H NMR (400 MHz, CDCl₃) δ 2.40 (s, 3H), 7.24-7.28 (m, 2H), 7.33 (t, *J* = 7.2 Hz, 1H), 7.43 (t, *J* =
52
53 7.8 Hz, 2H), 7.49 (dd, *J* = 6.4, 2.0 Hz, 2H), 7.58 (dd, *J* = 8.4, 1.2 Hz, 2H); ¹³C NMR (100 MHz,
54
55 CDCl₃) δ 21.2, 127.1, 128.9, 129.6, 137.2, 138.5, 141.3. HRMS (EI+) m/z [M]⁺ calcd for C₁₃H₁₂
56
57 168.0939, Found 168.0935.
58
59
60

4-(1,1-Dimethylethyl)-1,1'-Biphenyl (3bc).²² The general procedure was used; white solid (43 mg,

68% yield); ^1H NMR (400 MHz, CDCl_3) δ 1.37 (s, 9H), 7.33 (t, $J = 7.4$ Hz, 1H), 7.41-7.49 (m, 4H), 7.54 (d, $J = 8.8$ Hz, 2H), 7.59 (d, $J = 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 31.5, 34.7, 125.9, 126.9, 127.1, 127.2, 128.8, 138.5, 141.2, 150.4. HRMS (EI+) m/z $[\text{M}]^+$ calcd for $\text{C}_{16}\text{H}_{18}$ 210.1409, Found 210.1405.

1,1':4',1''-Terphenyl (3bd).²³ The general procedure was used; white solid (53 mg, 76% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.37 (dt, $J = 7.6, 1.2$ Hz, 2H), 7.49 (t, $J = 7.8$ Hz, 4H), 7.64-7.67 (m, 4H), 7.69 (s, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ 127.2, 127.5, 127.6, 129.0, 140.3, 140.9. HRMS (EI+) m/z $[\text{M}]^+$ calcd for $\text{C}_{18}\text{H}_{14}$ 230.1096, Found 230.1095.

N,N-Dimethyl-[1,1'-biphenyl]-4-amine (3be).²⁴ The general procedure was used; white solid (46 mg, 77% yield); ^1H NMR (400 MHz, CDCl_3) δ 3.02 (s, 6H), 6.84 (d, $J = 8.8$ Hz, 2H), 7.28 (t, $J = 7.6$ Hz, 1H), 7.43 (t, $J = 7.8$ Hz, 2H), 7.54 (dd, $J = 6.8, 2.4$ Hz, 2H), 7.59 (dd, $J = 7.2, 1.6$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 40.7, 112.9, 126.1, 126.4, 127.8, 128.8, 129.4, 141.4, 150.1. HRMS (EI+) m/z $[\text{M}]^+$ calcd for $\text{C}_{14}\text{H}_{15}\text{N}$ 197.1204, Found 197.1200.

Methyl [1,1'-biphenyl]-4-carboxylate (3bf).²⁵ The general procedure was used; white solid (43 mg, 67% yield); ^1H NMR (400 MHz, CDCl_3) δ 3.95 (s, 3H), 7.40 (tt, $J = 7.4, 1.6$ Hz, 1H), 7.48 (tt, $J = 7.4, 1.6$ Hz, 2H), 7.63 (dt, $J = 8.4, 1.2$ Hz, 2H), 7.67 (dt, $J = 8.2, 1.8$ Hz, 2H), 8.11 (dt, $J = 8.2, 1.8$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 52.3, 127.2, 127.4, 128.3, 129.1, 130.2, 140.2, 145.8, 167.2. HRMS (EI+) m/z $[\text{M}]^+$ calcd for $\text{C}_{14}\text{H}_{12}\text{O}_2$ 212.0837, Found 212.0839.

2-(4-Fluorophenyl)-naphthalene (3ag).²⁶ The general procedure was used except that **1a** was used as the substrate; white solid (36 mg, 53% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.16-7.21 (m, 2H), 7.51-7.56 (m, 2H), 7.67-7.72 (m, 3H), 7.89-7.93 (m, 3H), 8.00-8.02 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 115.8 (d, $J = 22.0$ Hz), 125.5, 125.8, 126.1, 126.5, 127.8, 128.3, 128.7, 129.1 (d, $J = 7.67$ Hz), 132.7, 133.8, 137.4 (d, $J = 2.88$ Hz), 137.7, 162.7 (d, $J = 246.3$ Hz). HRMS (EI+) m/z $[\text{M}]^+$ calcd for $\text{C}_{16}\text{H}_{11}\text{F}$ 222.0845, Found 222.0844.

1
2
3 *1-Phenylnaphthalene (3bh)*.²⁷ The general procedure was followed used; white solid (41 mg, 67%
4 yield); ¹H NMR (400 MHz, CDCl₃) δ 7.56-7.44 (m, 9H) ,7.86-7.93 (m, 3H); ¹³C NMR (100 MHz,
5 CDCl₃) δ 125.5, 125.9, 126.2, 127.1, 127.4, 127.8, 128.4, 130.2, 131.7, 133.9, 140.4, 140.9. HRMS
6
7
8
9
10
11 (EI+) m/z [M]⁺ calcd for C₁₆H₁₂ 204.0939, Found 204.0938.

12
13 *5-Phenylbenzo[d][1,3]dioxole (3bi)*.²⁸ The general procedure was used; white solid (45 mg, 75%
14 yield); ¹H NMR (400 MHz, CDCl₃) δ 6.00 (s, 2H), 6.89 (dd, *J* = 8.0, 0.8 Hz, 1H), 7.08-7.05 (m,
15 2H), 7.32 (tt, *J* = 7.2, 1.4 Hz, 1H), 7.42 (t, *J* = 7.6 Hz, 2H), 7.52 (dd, *J* = 7.6, 1.0 Hz, 2H); ¹³C NMR
16
17
18
19
20
21
22 (100 MHz, CDCl₃) δ 101.3, 107.8, 108.7, 120.8, 127.0, 127.1, 128.9, 135.8, 141.1, 147.2, 148.3.
23
24 HRMS (EI+) m/z [M]⁺ calcd for C₁₃H₁₀O₂ 198.0681, Found 198.0682.

25
26 *1-Methyl-5-phenyl-1H-indole (3bj)*.²⁹ The general procedure was used; pale yellow solid (47mg,
27 76% yield); ¹H NMR (400 MHz, CDCl₃) δ 3.83 (s, 3H), 6.55 (d, *J* = 2.4 Hz, 1H), 7.09 (d, *J* = 3.2
28 Hz, 1H), 7.31 (t, *J* = 7.2 Hz, 1H), 7.36-7.53 (m, 4H), 7.66 (dd, *J* = 8.0, 0.8 Hz, 2H), 7.85 (d, *J* = 1.2
29 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 33.1, 101.5, 109.5, 119.6, 121.5, 126.4, 127.5, 128.8,
30
31
32
33
34
35
36
37
38
39 129.1, 129.6, 133.0, 136.4, 142.8. HRMS (EI+) m/z [M]⁺ calcd for C₁₅H₁₃N 207.1048, Found
207.1048.

40
41 *3-Phenylpyridine (3bk)*.³⁰ The general procedure was used; white solid (32 mg, 68% yeild); ¹H
42 NMR (400 MHz, CDCl₃) δ 7.20 (dd, *J* = 7.6, 4.8 Hz, 1H), 7.29-7.31 (m, 1H), 7.33-7.38 (m, 2H),
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60 7.44-7.47 (m, 2H), 7.71(dt, *J* = 8.0, 1.6 Hz, 1 H), 8.50 (dd, *J* = 4.8, 1.6 Hz, 1H), 8.78 (d, *J* = 2.4 Hz,
1H); ¹³C NMR (100 MHz, CDCl₃) δ 123.2, 126.8, 127.8, 128.8, 133.9, 136.2, 137.4, 148.0, 148.2.
HRMS (EI+) m/z [M+H]⁺ calcd for C₁₁H₁₀N⁺ 156.0808, Found 156.0814.

Ethyl 4'-methyl-[1,1'-biphenyl]-4-carboxylate (3cb).³¹ The general procedure was used; white solid
(43 mg, 60% yield); ¹H NMR (400 MHz, CDCl₃) δ 1.42 (t, *J* = 7.4 Hz, 3H), 2.42 (s, 3H), 4.41 (q, *J*
= 7.0 Hz, 2H), 7.28 (d, *J* = 8.4 Hz, 2H), 7.53 (d, *J* = 7.6 Hz, 2H), 7.65 (d, *J* = 8.4 Hz, 2H), 8.11 (d, *J*
= 8.2 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.5, 21.3, 61.1, 126.9, 127.2, 129.1, 129.8, 130.2,

1
2
3 137.3, 138.2, 145.6, 166.7. HRMS (EI+) m/z [M]⁺ calcd for C₁₆H₁₆O₂ 240.1150, Found 240.1148.

4
5 *N,N,4'-Trimethyl-[1,1'-biphenyl]-4-amine (3db)*.³² The general procedure was used, except that the
6
7
8 reaction was conducted at 160 °C using 10 mol% of Ni(cod)₂, 30 mol% of **L5** and NaO^tBu; white
9
10 solid (38 mg, 60% yield); ¹H NMR (400 MHz, CDCl₃) δ 2.38 (s, 3H), 3.00 (s, 6H), 6.81 (d, *J* = 8.0
11
12 Hz, 2H), 7.22 (d, *J* = 8.4 Hz, 2H), 7.47 (d, *J* = 7.6 Hz, 2H), 7.50 (d, *J* = 8.0 Hz, 2H); ¹³C NMR (100
13
14 MHz, CDCl₃) δ 21.2, 40.8, 113.0, 126.3, 127.0, 127.7, 129.5, 135.8, 138.5, 149.9. HRMS (EI+) m/z
15
16 [M]⁺ calcd for C₁₅H₁₇N 211.1361, Found 211.1360.

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
4'-Methyl-3-(trifluoromethyl)-1,1'-biphenyl (3eb).³³ The general procedure was used; white solid
(46 mg, 65% yeild); ¹H NMR (400 MHz, CDCl₃) δ 2.42 (s, 3H), 7.29 (d, *J* = 8.4 Hz, 2H), 7.49-7.59
(m, 4H), 7.76 (d, *J* = 7.2 Hz, 1H), 7.83 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 21.3, 123.8 (q, *J* =
3.9 Hz), 124.4 (q, *J* = 271 Hz), 127.2, 129.3, 129.9, 130.3, 131.2 (q, *J* = 31.87 Hz), 137.0, 138.1,
142.1. HRMS (EI+) m/z [M]⁺ calcd for C₁₄H₁₁F₃ 236.0813, Found 236.0809.

Ethyl 4'-methyl-[1,1'-biphenyl]-3-carboxylate (3fb).³⁴ The general procedure was used; white solid
(51 mg, 71%); ¹H NMR (400 MHz, CDCl₃) δ 1.34 (t, *J* = 7.4 Hz, 3H), 2.34 (s, 3H), 4.34 (q, *J* = 7.2
Hz, 2H), 7.20 (d, *J* = 8.4 Hz, 2H), 7.42 (t, *J* = 7.8 Hz, 1H), 7.46 (d, *J* = 8.0 Hz, 2H), 7.68 (dt, *J* =
7.8, 1.6 Hz, 1H), 7.92 (dt, *J* = 7.8, 1.4 Hz, 1H), 8.19 (t, *J* = 1.8 Hz, 1H); ¹³C NMR (100 MHz,
CDCl₃) δ 14.5, 21.3, 61.2, 127.2, 128.2, 128.9, 129.7, 131.1, 131.4, 137.5, 137.7, 141.5, 166.8.
HRMS (EI+) m/z [M]⁺ calcd for C₁₆H₁₆O₂ 240.1150, Found 240.1153.

4-Methyl-1,1':2',1''-terphenyl (3gb).³⁵ The general procedure was used, except that the reaction was
conducted at 160 °C using Ni(cod)₂ (10 mol%), **L3** (30 mol%), NaO^tBu (30 mol%) and **2b** (6.0
equiv); colorless oil (43 mg, 59%). ¹H NMR (400 MHz, CDCl₃) δ 2.32 (s, 3H), 6.94-7.05 (m, 4H),
7.18-7.15 (m, 2H), 7.26-7.22 (m, 3H), 7.41-7.43 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 21.2,
126.5, 127.4, 127.6, 128.0, 128.7, 129.9, 130.0, 130.8, 136.2, 138.7, 140.7, 141.8. HRMS (EI+) m/z
[M]⁺ calcd for C₁₉H₁₆ 244.1252, Found 244.1251.

N,N,2'-Trimethyl[1,1'-biphenyl]-4-amine (3he).³⁶ The general procedure was used, except that the

1
2
3 reaction was conducted at 160 °C using Ni(cod)₂ (10 mol%), **L3** (30 mol%), NaO^tBu (30 mol%)
4 and **2e** (6.0 equiv); colorless oil (47 mg, 75%). ¹H NMR (400 MHz, CDCl₃) δ 2.33 (s, 3H), 3.00 (s,
5 6H), 6.79-6.81 (m, 2H), 7.21-7.26 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 20.8, 40.8, 112.3, 125.9,
6 7 8 9 10 11 12 13 14
126.7, 130.06, 130.08, 130.4, 135.6, 142.1, 149.4. HRMS (EI+) m/z [M]⁺ calcd for C₁₅H₁₇N⁺
211.1361, Found 211.1357.

15 2-(*p*-Tolyl)pyridine (**3ib**).³⁷ The general procedure was used; white solid (36 mg, 70%); ¹H NMR
16 17 18 (400 MHz, CDCl₃) δ 2.41 (s, 3H), 7.20 (ddd, *J* = 6.4, 4.8, 1.8 Hz, 1H), 7.29 (d, *J* = 8.2 Hz, 2H),
19 20 21 22 23 24 25 26
7.68-7.78 (m, 2H), 7.90 (d, *J* = 8.4 Hz, 2H), 8.68 (d, *J* = 5.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃)
δ 21.4, 120.4, 121.9, 126.9, 129.6, 136.8, 139.1, 149.7, 157.6. HRMS (EI+) m/z [M]⁺ calcd for
C₁₂H₁₁N 169.0891, Found 169.0880.

27 2-(*p*-Tolyl)naphthalene (**3jb**).³⁸ The general procedure was used; white solid (40 mg, 61%); ¹H
28 29 30 31 32 33 34 35 36 37 38 39
NMR (400 MHz, CDCl₃) δ 2.42 (s, 3H), 7.30 (d, *J* = 7.8 Hz, 2H), 7.56-7.42 (m, 2H), 7.63 (d, *J* =
8.2 Hz, 2H), 7.74 (dd, *J* = 8.7, 1.8 Hz, 1H), 7.79-7.97 (m, 3H), 8.02 (d, *J* = 1.4 Hz, 1H); ¹³C NMR
(100 MHz, CDCl₃) δ 21.2, 125.5, 125.6, 125.8, 126.3, 127.3, 127.7, 128.2, 128.4, 129.7, 132.6,
133.8, 137.2, 138.3, 138.6. HRMS (EI+) m/z [M]⁺ calcd for C₁₇H₁₄ 218.1096, Found 218.1097.

40 2-(4-Methylbenzyl)naphthalene (**3kb**).³⁹ The general procedure was used; pale-yellow oil (58 mg,
41 42 43 44 45 46 47 48 49 50 51 52 53 54 55
83%); ¹H NMR (400 MHz, CDCl₃) δ 2.36 (s, 3H), 4.14 (s, 2H), 7.07-7.25 (m, 4H), 7.35 (dd, *J* =
8.5, 1.6 Hz, 1H), 7.40-7.58 (m, 2H), 7.67 (s, 1H), 7.73-7.94 (m, 3H); ¹³C NMR (100 MHz, CDCl₃)
δ 21.2, 41.8, 125.4, 126.1, 127.1, 127.68, 127.75, 127.77, 128.2, 129.0, 129.3, 132.2, 133.8, 135.8,
138.1, 139.0. HRMS (EI+) m/z [M]⁺ calcd for C₁₈H₁₆ 232.1252, Found 232.1250.

56 **Acknowledgements.** This work was supported by ACT-C (J1210B2576) from JST, Japan and
57 Scientific Research on Innovative Area "Precisely Designed Catalysts with Customized
58 Scaffolding" (16H01022) from MEXT, Japan. We also thank the Instrumental Analysis Center,
59 60

1
2
3 Faculty of Engineering, Osaka University, for their assistance with HRMS.
4
5
6
7

8 **Supporting Information Available:** ^1H and ^{13}C NMR spectra for **1d**, **1f** and all the cross-coupling
9 products is available free of charge via the Internet at <http://pubs.acs.org>.
10
11
12
13
14
15
16
17

18 **References and Notes**

19
20 (1) (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457. (b) Suzuki, A. *Angew. Chem., Int. Ed.*
21
22 **2011**, *50*, 6722.
23

24
25 (2) (a) Rosen, B. M.; Quasdorf, K. W.; Wilson, D. A.; Zhang, N.; Resmerita, A.-M.; Garg, N. K.;
26 Percec, V. *Chem. Rev.* **2011**, *111*, 1346-1416. (b) Li, B.-J.; Yu, D.-G.; Sun, C.-L. ; Shi, Z.-J. *Chem.–*
27
28 *Eur. J.* **2011**, *17*, 1728-1759. (c) Yamaguchi, J.; Muto, K.; Itami, K. *Eur. J. Org. Chem.* **2013**, *2013*,
29
30 19-30. (d) Cornella, J.; Zarate, C.; Martin, R. *Chem. Soc. Rev.* **2014**, *43*, 8081-8097. (e) Tobisu, M.;
31
32 Chatani, N. *Acc. Chem. Res.* **2015**, *48*, 1717-1726.
33
34

35
36 (3) (a) Tobisu, M.; Shimasaki, T.; Chatani, N. *Angew. Chem., Int. Ed.* **2008**, *47*, 4866. (b)
37
38 Shimasaki, T.; Konno, Y.; Tobisu, M.; Chatani, N. *Org. Lett.* **2009**, *11*, 4890. (c) Tobisu, M.;
39
40 Yasutome, A.; Kinuta, H.; Nakamura, K.; Chatani, N. *Org. Lett.* **2014**, *16*, 5572.
41
42

43
44 (4) Other catalytic transformations enabled by the Ni/ICy catalytic system: (a) Tobisu, M.;
45
46 Takahira, T.; Ohtsuki, A.; Chatani, N. *Org. Lett.* **2015**, *17*, 680. (b) Tobisu, M.; Takahira, T.;
47
48 Chatani, N. *Org. Lett.* **2015**, *17*, 4352. (c) Tobisu, M.; Morioka, T.; Ohtsuki, A.; Chatani, N. *Chem.*
49
50 *Sci.* **2015**, *6*, 3410. (d) Furukawa, T.; Tobisu, M.; Chatani, N. *Chem. Commun.* **2015**, *51*, 6508. (e)
51
52 Morioka, T.; Nishizawa, A.; Nakamura, K.; Tobisu, M.; Chatani, N. *Chem. Lett.* **2015**, *44*, 1729. (f)
53
54 Nakamura, K.; Tobisu, M.; Chatani, N. *Org. Lett.* **2015**, *17*, 6142. (g) Hu, J.; Sun, H.; Cai, W.; Pu,
55
56 X.; Zhang, Y.; Shi, Z. *J. Org. Chem.* **2016**, *81*, 14. (h) Basch, C. H.; Cobb, K. M.; Watson, M. P.
57
58
59 *Org. Lett.* **2016**, *18*, 136.
60

- 1
2
3 (5) (a) Quasdorf, K. W.; Tian, X.; Garg, N. K. *J. Am. Chem. Soc.* **2008**, *130*, 14422. (b) Guan, B.-
4 T.; Wang, Y.; Li, B.-J.; Yu, D.-G.; Shi, Z.-J. *J. Am. Chem. Soc.* **2008**, *130*, 14468. (c) Molander, G.
5 T.; Wang, Y.; Li, B.-J.; Yu, D.-G.; Shi, Z.-J. *J. Am. Chem. Soc.* **2008**, *130*, 14468. (c) Molander, G.
6 T.; Wang, Y.; Li, B.-J.; Yu, D.-G.; Shi, Z.-J. *J. Am. Chem. Soc.* **2008**, *130*, 14468. (c) Molander, G.
7 T.; Wang, Y.; Li, B.-J.; Yu, D.-G.; Shi, Z.-J. *J. Am. Chem. Soc.* **2008**, *130*, 14468. (c) Molander, G.
8 A.; Beaumard, F. *Org. Lett.* **2010**, *12*, 4022. (d) Xu, M.; Li, X.; Sun, Z.; Tu, T. *Chem. Commun.*
9 **2013**, *49*, 11539.
- 10
11
12 (6) (a) Quasdorf, K. W.; Riener, M.; Petrova, K. V.; Garg, N. K. *J. Am. Chem. Soc.* **2009**, *131*,
13 17748. (b) Antoft-Finch, A.; Blackburn, T.; Snieckus, V. *J. Am. Chem. Soc.* **2009**, *131*, 17750. (c)
14 Xu, L.; Li, B.-J.; Wu, Z.-H.; Lu, X.-Y.; Guan, B.-T.; Wang, B.-Q.; Zhao, K.-Q.; Shi, Z.-J. *Org.*
15 *Lett.* **2010**, *12*, 884. (d) Baghbanzadeh, M.; Pilger, C.; Kappe, C. O. *J. Org. Chem.* **2011**, *76*, 1507.
16
17 (e) Quasdorf, K. W.; Antoft-Finch, A.; Liu, P.; Silberstein, A. L.; Komaromi, A.; Blackburn, T.;
18 Ramgren, S. D.; Houk, K. N.; Snieckus, V.; Garg, N. K. *J. Am. Chem. Soc.* **2011**, *133*, 6352.
- 19
20 (7) The use of NHC ligands in other nickel-catalyzed transformations of aryl esters and carbamates:
21
22 (a) Shimasaki, T.; Tobisu, M.; Chatani, N. *Angew. Chem., Int. Ed.* **2010**, *49*, 2929. (b) Mesganaw,
23 T.; Silberstein, A. L.; Ramgren, S. D.; Nathel, N. F. F.; Hong, X.; Liu, P.; Garg, N. K. *Chem. Sci.*
24 **2011**, *2*, 1766. (c) Hie, L.; Ramgren, S. D.; Mesganaw, T.; Garg, N. K. *Org. Lett.* **2012**, *14*, 4182.
- 25
26 (8) During the preparation of this manuscript, nickel/NHC-catalyzed cross-coupling of aryl
27 pivalates with alkyl(9-BBN) was reported: Guo, L.; Hsiao, C.-C.; Yue, H.; Liu, X.; Rueping, M.
28 *ACS Catal.* **2016**, *6*, 4438.
- 29
30 (9) Lower reactivity of arylboronic esters than the parent arylboronic acids in nickel-catalyzed
31 Suzuki-Miyaura cross-coupling of phenol derivatives has been reported: Leowanawat, P.; Zhang,
32 N.; Percec, V. *J. Org. Chem.* **2012**, *77*, 1018.
- 33
34 (10) Nakamura, K.; Yasui, K.; Tobisu, M.; Chatani, N. *Tetrahedron* **2015**, *71*, 4484.
- 35
36 (11) Phenol derivatives are less reactive than those of π -extended phenols, such as naphthol, in
37 many nickel-catalyzed cross-coupling reactions. See ref 2e.
- 38
39 (12) Snieckus, V. *Chem. Rev.* **1990**, *90*, 879.
- 40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (13) A review on convertible directing groups: Zhang, F.; Spring, D. R. *Chem. Soc. Rev.* **2014**, *43*,
4
5 6906.
6
7
8 (14) (a) Yamamoto, Y.; Takizawa, M.; Yu, X.-Q.; Miyaura, N. *Angew. Chem., Int. Ed.* **2008**, *47*,
9
10 928. (b) Billingsley, K. L.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 4695. (c) Knapp, D.
11
12 M.; Gillis, E. P.; Burke, M. D. *J. Am. Chem. Soc.* **2009**, *131*, 6961.
13
14
15 (15) Mkhaliid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. *Chem. Rev.*
16
17 **2010**, *110*, 890.
18
19
20 (16) It was mentioned that Ni/PCy₃ catalyst system can promote the cross-coupling of *naphthyl*
21
22 carbamates with arylboronic esters. (a) Ramgren, S. D.; Hie, L.; Ye, Y.; Garg, N. K. **2013**, *1*, 4974.
23
24 (b) Malineni, J.; Jezorek, R. L.; Zhang, N.; Percec, V. *Synthesis* **2016**, *48*, *in press*; DOI:
25
26 10.1055/s-0035-1562343. See also ref 9.
27
28
29 (17) Mistryukov, E.A. *Mendeleev Commun.* **2006**, *16*, 258.
30
31
32 (18) Medina, F., Michon, C., Agbosson-Niederccen, A. *Eur. J. Org. Chem.* **2012**, *31*, 6218.
33
34
35 (19) Li, B.-J.; Li, Y.-Z.; Lu, X.-Y.; Liu, J.; Guan, B.-T.; Shi, Z.-J. *Angew. Chem. Int. Ed.* **2008**, *47*,
36
37 10124.
38
39
40 (20) Cepanec, I.; Litvić, M.; Udiković, J.; Pogorelić, I.; Lovrić, M. *Tetrahedron* **2007**, *63*, 5614.
41
42
43 (21) Old, D. W.; Wolfe, J. P.; Buchwald, S. L. *J. Am. Chem. Soc.* **1998**, *120*, 9722.
44
45
46 (22) Susanto, W.; Chu, C.-Y.; Ang, W. J.; Chou, T.-C.; Lo, L.-C.; Lam, Y. *J. Org. Chem.* **2012**, *77*,
47
48 2729.
49
50
51 (23) Bai, L.; Wang, J. X. *Adv. Synth. Catal.* **2008**, *350*, 315.
52
53
54 (24) Chen, X.; Ke, H.; Chen, Y.; Guan, C.; Zou, G. *J. Org. Chem.* **2012**, *77*, 7572.
55
56
57 (25) Martinez-Solorio, D.; Melillo, B.; Sanchez, L.; Liang, Y.; Lam, E.; Houk, K. N.; Smith III, A.
58
59 B. *J. Am. Chem. Soc.* **2016**, *138*, 1836.
60
61
62 (26) Chen, H.; Huang, Z.; Hu, X.; Tang, G.; Xu, P.; Zhao, Y.; Cheng, C.-H. *J. Org. Chem.* **2011**,
63
64 76, 2338.

- 1
2
3 (27) Zhou, C.; Liu, Q.; Li, Y.; Zhang, R.; Fu, X.; Duan, C. *J. Org. Chem.* **2012**, *77*, 10468.
4
5
6 (28) Peng, Z.; Hu, G.; Qiao, H.; Xu, P.; Gao, Y.; Zhao, Y. *J. Org. Chem.* **2014**, *79*, 2733.
7
8 (29) Quasdorf, K. W.; Riener, M.; Petrova, K. V.; Garg, N. K. *J. Am. Chem. Soc.* **2009**, *131*, 17748.
9
10 (30) Zeng, J.; Liu, K. M.; Duan, X. F., *Org. Lett.* **2013**, *15*, 5342
11
12 (31) Zhu F.; Wang, Z.-X. *J. Org. Chem.* **2014**, *79*, 4285.
13
14 (32) Guan, B.-T.; Lu, X.-Y.; Zheng, Y.; Yu, D.-G.; Wu, T.; Li, K.-L.; Li, B.-J.; Shi, Z.-J. *Org. Lett.*
15
16 **2010**, *12*, 396.
17
18 (33) Ackermann, L.; Althammer, A. *Org. Lett.* **2006**, *8* 3457.
19
20 (34) Badone, D.; Baroni, M.; Cardamone, R.; Ielmini, A.; Guzzi, U. *J. Org. Chem.* **1997**, *62*, 7170.
21
22 (35) Kuriyama, M.; Shimazawa, R.; Shirai, R. *Tetrahedron* **2007**, *63*, 9393.
23
24 (36) Bolliger, J. L.; Frech, C. M. *Adv. Synth. Catal.* **2010**, *352*, 1075.
25
26 (37) Luzung, M. R.; Patel, J. S.; Yin, J. *J. Org. Chem.* **2010**, *75*, 8330.
27
28 (38) Ackermann, L.; Kapdi, A. R.; Fenner, S.; Kornhaab, C.; Schulzke, C. *Chem.-Eur. J.* **2011**, *17*,
29
30 2965.
31
32 (39) Cao, Z.-C.; Yu, D.-G.; Zhu, R.-Y.; Weia, J.-B.; Shi, Z.-J. *Chem. Commun.* **2015**, *51*, 2683.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(Graphic for Table of Contents)

