# **Chemical Science**

## **Accepted Manuscript**

## **Chemical Science**



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This Accepted Manuscript will be replaced by the edited and formatted Advance Article as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard **Terms & Conditions** and the **ethical guidelines** that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

## RSCPublishing

www.rsc.org/chemicalscience

Registered Charity Number 20789

View Online

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/chemicalscience

## EDGE ARTICLE

### Organocatalytic Conversion of Cellulose into a Platform Chemical<sup>†</sup>

Benjamin R. Caes,<sup>a</sup> Michael J. Palte,<sup>b</sup> and Ronald T. Raines<sup>a,c,\*</sup>

Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX DOI: 10.1039/b000000x

The search for a source of fuels and chemicals that is both abundant and renewable has become of paramount importance. The polysaccharide cellulose meets both criteria, and methods have been developed for its transformation into the platform chemical 5-(hydroxymethyl)furfural (HMF). These methods employ harsh reaction conditions or toxic heavy metal catalysts, deterring large-scale implementation. Here, we describe a low-temperature, one-pot route that uses *ortho*-carboxyl-substituted phenylboronic acids as organocatalysts in conjunction with hydrated magnesium chloride and mineral acids to convert cellulose and cellulose-rich municipal waste to HMF in yields comparable to processes that use toxic heavy metal catalysts. Isotopic labeling studies indicate that the key aldose-to-ketose transformation occurs via an enediol intermediate. The route, which also allows for facile catalyst recovery and recycling, provides a green prototype for cellulose conversion.

#### Introduction

Published on 03 October 2012 on http://pubs.rsc.org | doi:10.1039/C2SC21403B

10

Downloaded by University of Massachusetts - Amherst on 03 October 2012

The dependence on fossil fuels for energy and chemicals has <sup>15</sup> become unsustainable.<sup>1</sup> Abundant, renewable biomass resources could meet the fuel and chemical demands of the future,<sup>2</sup> but must recapitulate the wide array of products now derived from fossil fuels. The six-carbon furanic 5-(hydroxymethyl)furfural (HMF) has the potential to meet this challenge.<sup>3</sup> HMF can be transformed into a variety of useful products, such as common polyester building blocks<sup>4</sup> and the promising liquid fuel 2,5dimethylfuran.<sup>5</sup> The carbon skeleton of HMF is identical to those of the hexose sugars that are the primary components of cellulose

and hemicelluloses in biomass. Still, accessing this resource requires a process that efficiently transforms these carbohydrates into HMF.

The conversion of cellulose to HMF can proceed via three steps: hydrolysis of cellulose to glucose, isomerization of glucose to fructose, and dehydration of fructose to HMF (Fig. 1).

- <sup>30</sup> Although several methods exist for transforming glucose and fructose into HMF,<sup>6</sup> few are capable of producing HMF in high yields directly from cellulose.<sup>3</sup> Solid acids<sup>7</sup> and heavy metals<sup>6b,8</sup> are the best extant catalysts for this conversion. The inefficiency of solid acid catalysts and the toxicity of heavy metals could
- <sup>35</sup> diminish their impact.<sup>9</sup> Considering the ever-growing need for green chemistry, we sought a conversion process that uses benign and recyclable reagents, catalysts, and solvents, as well as mild reaction conditions. Here, we report on our discovery of such a process.

#### 40 Results and discussion

Our initial experiments focused on discovering an alternative to heavy metals for accessing HMF from glucose. As a solvent, we choose dimethylacetamide (DMA), a polar aprotic solvent that has served as the medium for other biomass conversions.<sup>8b,10</sup> We

<sup>45</sup> screened a variety of metal chlorides (Table S1) and found that <sup>50</sup> magnesium, nickel, zinc, cerium, or lanthanum, like chromium,



**Fig. 1.** Putative route for the conversion of cellulose to 5-(hydroxymethyl)furfural, mediated by the depicted boronic acids. Labeled hydrogens in glucose (green) and water (blue) have the indicated fates.

35





				111111	40
		Т	time	yield	40
metal chloride, mol%	boronic acid, wt%	(°C)	(h)	(%)	
_	—	100	6	0	-
CrCl <sub>2</sub> , 25	—	100	6	58	
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	—	120	6	29	
NiCl <sub>2</sub> ·6H <sub>2</sub> O, 300	—	120	3	32	
ZnCl <sub>2</sub> , 300	—	120	6	21	45
CeCl <sub>3</sub> ·7H <sub>2</sub> O, 300	—	120	3	22	
LaCl <sub>3</sub> ·7H <sub>2</sub> O, 300	—	120	6	22	
—	2-carboxyphenyl, 100	120	6	2	
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	2-carboxyphenyl, 100	120	4	54	
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	2-carboxyphenyl, 50	120	6	48	
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	2-carboxyphenyl, 25	120	6	46	50
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	2-carboxyphenyl, 10	120	6	31	50
NiCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-carboxyphenyl, 100	105	4	24	
$ZnCl_2 + 6H_2O, 300$	2-carboxyphenyl, 100	105	4	18	
CeCl <sub>3</sub> ·7H <sub>2</sub> O, 300	2-carboxyphenyl, 100	120	3	51	
LaCl <sub>3</sub> ·7H <sub>2</sub> O, 300	2-carboxyphenyl, 100	120	3	49	
—	2-methoxycarbonylphenyl, 100	100	6	22	
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	2-methoxycarbonylphenyl, 100	120	4	57	55
—	2-ethoxycarbonylphenyl, 100	100	6	15	
MgCl <sub>2</sub> ·6H <sub>2</sub> O, 200	2-ethoxycarbonylphenyl, 100	120	4	52	

<sup>*a*</sup> Glucose was at 10 wt% relative to DMA. Mol% and HMF yield (HPLC) are relative to glucose.

provides appreciable yields of HMF (Table 1). We chose to employ magnesium due to its being one of the most abundant elements in Earth's crust<sup>11</sup> and the human body.<sup>12</sup>

Next, we attempted to increase the efficiency of the conversion while maintaining environmental benignancy. To do so, we turned to organocatalysis,<sup>13</sup> which had not been employed previously in a biomass conversion process.<sup>2,3</sup> We were aware that boronic acids readily form boronate esters with 1,2- and 1,3diols, and bind to furanose sugars with greater affinity than to pyranose sugars.<sup>14</sup> We reasoned that these organoboranes would alter the equilibrium between glucose and fructose (which exist primarily in the pyranose and furanose forms, respectively), and thereby enhance HMF production. The precedents were discouraging, however, as phenylboronic acid had been reported to inhibit the hydrolysis of cellulose and the dehydration of sugar

monomers.<sup>15,16</sup> Phenylboronic acids have an affinity for carbohydrates that is

tunable based on substituents.<sup>14</sup> We assessed the ability of twenty derivatives of phenylboronic acid to promote the conversion of glucose to HMF (Tables S2–S4 in the Supporting Information). Our data corroborated the earlier work with phenylboronic acid itself,<sup>15</sup> which is not a catalyst in our hands, but revealed that adding an *ortho* carboxyl group engenders catalysis, especially in

- the presence of MgCl<sub>2</sub>·6H<sub>2</sub>O (Table 1). Further optimization revealed prospects for catalytic turnover. For example, adding 25 mol% of 2-carboxyphenylboronic acid to the magnesium increases the HMF yield from 29% to 46%.
- Water can counter the dehydration of fructose (Fig. 1). Accordingly, we attempted to increase our HMF yields by using anhydrous MgCl<sub>2</sub>. We found, however, that anhydrous MgCl<sub>2</sub> <sub>90</sub> provides lower HMF yields than does the hydrated salt (Table S4). To determine how much water is beneficial for HMF

Page 2 of 4

production, we added water to reaction mixtures containing the anhydrous salt (Table S5 in the Supporting Information) and found that HMF yields increase steadily up to an H<sub>2</sub>O:Mg(II) ratio of ~6 (Fig. S1). Other anhydrous metal chlorides that work synergistically with 2-carboxyphenylboronic also have a similar reliance on water (Table S5).

Encouraged by the successful conversion of glucose to HMF, we next attempted to convert cellulose to HMF using boronic acids in 1-ethyl-3-methylimidazolium chloride ([EMIM]Cl), an ionic liquid that dissolves cellulose<sup>17</sup> and enables a mineral acid to catalyze its hydrolysis.<sup>18</sup> We observed that phenylboronic acids with an ortho carboxylic acid or ester promote the conversion of cellulose to HMF (Tables S6 and S7 in the Supporting Information), and that MgCl<sub>2</sub>·6H<sub>2</sub>O is necessary to maximize HMF production (Table 2). The observed yields of HMF (e.g., 41% in 1 h at 105 °C with 2-methoxycarbonylphenyl boronic acid) are comparable to those attainable with a heavy metal.8 Boronic acid/MgCl<sub>2</sub>·6H<sub>2</sub>O also promote the conversion of cotton, paper towels, and newspaper (Table 2), which are cellulose-rich components of municipal waste. The HMF yields correlate with the purity of the cellulose in these materials. Thus, HMF can be produced efficiently from cellulosic material in a single, one-pot process that is devoid of heavy metals.

The high boronate loadings necessary to turnover cellulose rapidly ( $\leq 2$  h) led us to search for a recovery method that enables boronate recovery. Our initial separation strategy focused on isolating 2-carboxyphenylboronic acid from a reaction mixture using an anion-exchange resin. The reaction mixture was diluted with water, filtered to remove any humins (which are insoluble byproducts from aldol addition and condensation<sup>19</sup>), extracted with ethyl acetate to remove HMF, and passed through a column of resin. The anionic boronate was retained on the resin and eluted with aqueous NH<sub>4</sub>HCO<sub>3</sub> (1 M) to yield purified boronate, which retained catalytic activity (Fig. S2). This strategy was, however, inapplicable to the *o*-esterphenylboronic acids, which have low solubility in water and high solubility in ethyl acetate.

Boronic acid moieties become anionic at high pH and partition completely into an aqueous phase. Hence, we added basic water to reaction mixtures containing 2-ethoxycarbonylphenylboronic acid, filtered to remove any humins, and extracted with ethyl acetate (Fig. S3). Removing solvent from the organic phase provided HMF. Evaporating water from the aqueous phase recovered the boronic acid and MgCl<sub>2</sub>. The recovered catalysts provided HMF in comparable yield through four reaction cycles.

Additional data provided insight on the chemical mechanism of our conversion (Fig. 1). Chromium is known to mediate the isomerization of aldose to ketose sugars by a 1,2-hydride shift.10,20 To probe the mechanism of our conversion, we performed two deuterium-labeling experiments. In the first, glucose-2-d converted HMF was into hν 2-ethoxycarbonylphenylboronic acid and MgCl<sub>2</sub>·6H<sub>2</sub>O. <sup>1</sup>H NMR spectroscopy revealed that virtually no deuterium was retained in the HMF product. In the second, unlabeled glucose was converted in the presence of D<sub>2</sub>O, and a substantial quantity of deuterium was found at C-1 of HMF (Fig. S4). These results are compatible with a mechanism that avails an enediol intermediate (Fig. 1). A similar mechanism is used by the enzyme

15

 Table 2
 Conversion of cellulosic substrates into HMF in ionic liquids<sup>a</sup>

ОН

		HOOOF	HO OH cataly HO OH	HO			
		(	cellulose	HMF			
					Т	time	HMF yield
substrate	ionic liquid	acid, wt%	metal chloride, mol%	boronic acid, mol%	(°C)	(h)	(%)
cellulose	[EMIM]Cl	HCl, 0.61	—	—	105	2	10
cellulose	[EMIM]Cl	HCl, 0.61	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	—	105	3	15
cellulose	[EMIM]Cl	HCl, 0.61	—	2-methoxycarbonylphenyl, 120	105	2	12
cellulose	[EMIM]Cl	HCl, 0.61	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-methoxycarbonylphenyl, 120	105	2	39
cellulose	[EMIM]Cl	H <sub>2</sub> SO <sub>4</sub> , 0.88	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-methoxycarbonylphenyl, 120	105	1	41
cellulose	[EMIM]Cl	HCl, 0.61	—	2-ethoxycarbonylphenyl, 160	105	1	21
cellulose	[EMIM]Cl	HCl, 0.61	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-ethoxycarbonylphenyl, 160	105	2	38
cellulose	[EMIM]Cl	H <sub>2</sub> SO <sub>4</sub> , 0.88	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-ethoxycarbonylphenyl, 160	105	1	36
cotton	[BMIM]Cl	_	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-ethoxycarbonylphenyl, 160	105	2	40
paper towel	[EMIM]Cl	HCl, 0.61	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-ethoxycarbonylphenyl, 160	105	2	31
newspaper	[BMIM]Cl	—	MgCl <sub>2</sub> ·6H <sub>2</sub> O, 300	2-ethoxycarbonylphenyl, 160	105	2	18

<sup>*a*</sup> Substrates were at 5 wt% relative to the ionic liquid. Wt% is relative to the ionic liquid. Mol% and HMF yield (HPLC) are relative to glucose monomers within the substrate, which was assumed to be pure cellulose.

50

phosphoglucose isomerase, which catalyzes the interconversion of glucose 6-phosphate and fructose 6-phosphate.<sup>21</sup>

Boronate ester-formation is known to be more favorable with fructose than glucose.<sup>14</sup> We found that boronic acids also serve <sup>40</sup> by catalyzing fructose dehydration (Table S8). We suspect that the organocatalyst relies on an *ortho* carboxyl group because its oxygen can donate electron density into the empty *p*-orbital of boron, thereby decreasing the strength of the fructose–boronate complex in the nearly water-free medium.<sup>14b</sup> Finally, we propose <sup>45</sup> that water attenuates the reactivity of Mg(II), allowing its participation in catalysis, but deterring reaction pathways that can lead to humins.<sup>22</sup>

#### **Concluding Remarks**

- <sup>15</sup> In summary, we have discovered novel reactivity mediated by a simple boronic acid composed of hydrogen, boron, carbon, and oxygen—four of the first eight elements in the periodic table. <sup>55</sup> This organocatalyst mediates the transformation of fructose, glucose, and cellulose into the platform chemical HMF. Although
- <sup>20</sup> boronate loadings for rapid ( $\leq$ 2-h) conversions are high, this liability is overcome by the facility of its recovery (or, <sup>60</sup> potentially, by its immobilization). This discovery, which is the first to link two highly active areas of research: organocatalysis and biomass conversion, could facilitate the transition from
- <sup>25</sup> fossil-based fuels and chemicals to a more sustainable biomassbased society.

#### Acknowledgements

We are grateful to Prof. S. S. Stahl and Dr. N. A. McGrath for contributive discussions. This work was supported by the DOE <sup>70</sup>
 <sup>30</sup> Great Lakes Bioenergy Research Center (DOE BER Office of Science DE-FC02-07ER64494), a DOE Bioenergy Research Center. M.J.P. was supported by pre-doctoral fellowship 09PRE2260125 (American Heart Association). This work made use of the National Magnetic Resonance Facility at Madison, <sup>72</sup>

<sup>35</sup> which is supported by NIH grants P41RR02301 (BRTP/NCRR) and P41GM66326 (NIGMS). Additional equipment was purchased with funds from the University of Wisconsin, the NIH (RR02781, RR08438), the NSF (DMB-8415048, OIA-9977486, BIR-9214394), and the USDA.

#### Notes and references

<sup>a</sup>Department of Chemistry, University of Wisconsin-Madison, 1101 University Avenue, Madison, WI 53706, USA <sup>b</sup>Medical Scientist Training Program and Molecular & Cellular Pharmacology Graduate Training Program, University of Wisconsin-Madison, 1300 University Avenue, Madison, WI 53706, USA <sup>c</sup>Department of Biochemistry, University of Wisconsin-Madison, 433 Babcock Drive, Madison, WI 53706, USA. E-mail: rtraines@wisc.edu † Electronic Supplementary Information (ESI) available: Experimental procedures, Tables S1–S8, and Figs. S1–S4. See DOI: 10.1039/b000000x/

- 1 US National Petroleum Council, *Facing the Hard Truths about Energy*, Washington, DC, 2007.
- (a) D. Tilman, R. Socolow, J. A. Foley, J. Hill, E. Larson, L. Lynd, S. Pacala, J. Reilly, T. Searchinger, C. Somerville and R. Williams, *Science*, 2009, 325, 270; (b) C. O. Tuck, E. Pérez, I. T. Horváth, R. A. Sheldon and M. Poliakoff, *Science*, 2012, 337, 695-699; (c) R. A. Sheldon, *Chem. Soc. Rev.*, 2012, 41, 1437-1451.
- 3 (a) J. N. Chheda, G. W. Huber and J. A. Dumesic, *Angew. Chem. Int. Ed.*, 2007, 46, 7164-7183; (b) M. E. Zakrewska, E. Bogel-Łukasik and R. Bogel-Łukasik, *Chem. Rev.*, 2011, 111, 397-417.
- 4 J. Lewkowski, ARKIVOC, 2001, 17-54.
- 5 (a) S. Zhong, R. Daniel, H. Xu, J. Zhang, D. Turner, M. L. Wyszynski and P. Richards, *Energy Fuels*, 2010, 24, 2891-2899; (b) G. Tian, R. Daniel, H. Li, H. Xu, S. Shuai and P. Richards, *Energy Fuels*, 2010, 24, 3898-3905; (c) R. Daniel, G. Tian, H. Xu, M. L. Wyszynski, X. Wu and Z. Huang, *Fuel*, 2011, 90, 449-458; (d) D. A. Rothamer and J. H. Jennings, *Fuel*, 2012, 98, 203-212.
- 6 (a) Y. Román-Leshkov, J. N. Chheda and J. A. Dumesic, *Science*, 2006, **312**, 1933-1937; (b) H. Zhao, J. E. Holladay, H. Brown and Z. C. Zhang, *Science*, 2007, **316**, 1597-1600.
- 7 (a) C. V. McNeff, D. T. Nowlan, L. C. McNeff, B. Yan and R. L. Fedie, *Appl. Catal. A-Gen.*, 2010, **384**, 65-69; (b) S. Zhao, M. Cheng, J. Li, J. Tian and X. Wang, *Chem. Commun.*, 2011, **47**, 2176-2178.
- 8 (a) Y. Su, H. M. Brown, X. Huang, X. Zhou, J. E. Amonette and C. Z. Zhang, *Appl. Catal. A-Gen.*, 2009, **361**, 117-122;

(b) J. B. Binder and R. T. Raines, J. Am. Chem. Soc., 2009, 131, 1979-1985; (c) B. Kim, J. Jeong, D. Lee, S. Kim, H.-J. Yoon, Y.-S. Lee and J. K. Cho, Green Chem., 2011, 13, 1503-1506; (d) P. Wang, H. Yu, S. Zhan and S. Wang, Bioresource Technol., 2011, 102, 4179-4183.

- 9 (a) S. Langrrd, Am. J. Ind. Med., 1990, 17, 189-214; (b) A. Levina and P. A. Lay, Chem. Res. Toxicol., 2008, 21, 563-571; (c) R. Marin, Y. Ahuja and R. N. Bose, J. Am. Chem. Soc., 2010, 132, 10617-10619.
- (a) J. B. Binder, A. V. Cefali, J. J. Blank and R. T. Raines, 10 Energy Environ. Sci., 2010, 3, 765-771; (b) J. B. Binder, A. V. Cefali, J. J. Blank and R. T. Raines, ChemSusChem, 2010, 3, 1268-1272.
- G. B. Haxel, J. B. Hedrick and G. J. Orris, U.S. Geological 11 Survey, 2002, Fact Sheet 087-002.

15

20

25

35

45

50

Published on 03 October 2012 on http://pubs.rsc.org | doi:10.1039/C2SC21403B

Downloaded by University of Massachusetts - Amherst on 03 October 2012

- M. J. Laires, C. P. Monteiro and M. Bicho, Front. Biosci., 12 2004, 9, 262-276.
- (a) S. C. Pan and B. List, Ernst Schering Found. Symp. 13 Proc., 2007, 1-43; (b) D. W. C. MacMillan, Nature, 2008, 455, 304-308; (c) E. N. Jacobsen and D. W. C. MacMillan, Proc. Natl. Acad. Sci. USA, 2010, 107, 20618-20619.
- 14 For examples, see: (a) G. Springsteen and B. H. Wang, Tetrahedron, 2002, 58, 5291-5300; (b) M. Berube, M. Dowlut and D. G. Hall, J. Org. Chem., 2008, 73, 6471-6479; (c) M. G. Chudzinski, Y. Chi and M. S. Taylor, Aust. J. Chem., 2011, 64, 1466-1469; (d) G. A. Ellis, M. J. Palte and R. T. Raines, J. Am. Chem. Soc., 2012, 134, 3631-3634.
- H. Kawamoto, S. Saito and S. Saka, J. Anal. Appl. Pyrol., 15 2008, 82, 78-82.
- 16 Boric acid is known to assist an enzyme-catalyzed transformation of glucose to fructose (Huang, R.; Qi, W.; Su, R.; He, Z. Chem. Commun. 2010, 1115-1117) but has varied effects on the other steps in Figure 1 (ref. 15; Kawamoto, H.; Saito, S.; Shiro, S. Carbohydr. Res. 2008, 343, 249-255; Ståhlberg, T.; Rodriguez-Rodriguez, S.; Fristrup, P.; Riisager, A. Chem. Eur. J. 2011, 17, 1456-1464). Moreover, the reactivity of boric acid cannot be tuned by chemical modification, nor can boric acid be immobilized readily.
- 17 R. P. Swatloski, S. K. Spear, J. D. Holbrey and R. D. 40 Rogers, J. Am. Chem. Soc., 2002, 124, 4974-4975.
  - 18 J. B. Binder and R. T. Raines, Proc. Natl. Acad. Sci. U.S.A., 2010, 107, 4516-4521.
  - 19 S. K. R. Patil and C. R. F. Lund, Energy Fuels, 2011, 25, 4745-4755.
  - 20 Y. Román-Leshkov, M. Moliner, J. A. Labinger and M. E. Davis, Angew. Chem. Int. Ed., 2010, 49, 8954-8957.
  - 21 (a) Y. J. Topper, J. Biol. Chem., 1957, 225, 419-426; (b) I. A. Rose and E. L. O'Connell, J. Biol. Chem., 1961, 236, 3086-3092.
- 22 C. Sievers, I. Musin, T. Marzialetti, M. B. V. Olarte, P. K. Agrawal and C. W. Jones, ChemSusChem, 2009, 2, 665-671.

Page 4 of 4