This article was downloaded by: [University of Sydney] On: 04 May 2015, At: 20:50 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Synthesis of 1,3-Bis[(3aryl)-S-triazolo[3,4-b]-[1,3,4]thiadiazole-6yl]benzenes

Dejiang Li^a, Deqing Long^a & Heqing Fu^b ^a Department of Chemistry, Yunyang Teachers College, Danjiangkou, Hubei, P. R. China

^b Research Institute of Chemical Engineering, South China University of Technology, Guangzhou, P. R. China

Published online: 01 Feb 2007.

To cite this article: Dejiang Li , Deqing Long & Heqing Fu (2006) Synthesis of 1,3-Bis[(3-aryl)-S-triazolo[3,4-b]-[1,3,4]thiadiazole-6-yl]benzenes, Phosphorus, Sulfur, and Silicon and the Related Elements, 181:3, 519-526, DOI: <u>10.1080/10426500500265008</u>

To link to this article: http://dx.doi.org/10.1080/10426500500265008

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Synthesis of 1,3-Bis[(3-aryl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzenes

Dejiang Li

Deqing Long

Department of Chemistry, Yunyang Teachers College, Danjiangkou, Hubei, P. R. China

Heqing Fu

Research Institute of Chemical Engineering, South China University of Technology, Guangzhou, P. R. China

1,3-bis[(3-aryl)-s-triazolo[3,4-b]-[1,3,4]thiadiazole-6-yl]benzenes ${\bf 2}$ were synthesized in high yields by the reaction of 3-aryl 4-amino-5-mercapto-1,2,4-triazole ${\bf 1}$ with m-phthalic acid.

 $\label{eq:keywords} \begin{array}{ll} 1,3-Bis[(3-aryl)-s-triazolo-[3,4-b]-[1,3,4] thiadiazole-6-yl] benzenes; \\ 2,4-triazole; 3-aryl-4-amino-5-mercapto-1; synthesis \end{array}$

S-triazolo[3,4-*b*]-[1,3,4]thiadiazole derivatives were found to possess significant biological activities such as antiflammatory, antiviral, antifungal, antineoplastic, and antidepressant effects¹⁻⁴ and electrochemical properties.⁵ They are highly important heterocycles and have been used in the research and development of agrochemicals and in the pharmaceutical chemistry. Most derivatives that have been reported only contain one s-triazolo[3,4-*b*]-[1,3,4]thiadiazole in one moleclule.⁶⁻¹¹ As part of our current studies on the synthesis of the biologically active s-triazolo[3,4-*b*]-[1,3,4]thiadiazole derivatives, we now report an efficient synthesis of some novel fused heterocyclic compounds combining two s-triazolo[3,4-*b*]-[1,3,4]thiadiazole rings in one framework from 3-aryl 4-amino-5-mercapto-1,2,4-triazoles with the hope to obtain compounds of better biological activities.

Received March 28, 2005; accepted May 1, 2005.

We gratefully acknowledge financial support of this work by the Science Foundation of Hubei Province Education Committee, China (Project No. 2004D001).

Address correspondence to Dejiang Li, Department of Chemistry, Yunyang Teachers College Danjiangkou, Hubei 442700, P. R. China. E-mail: lidejiang999@yahoo.com.cn

RESULTS AND DISCUSSION

The aroylhydrazides were prepared by the esterification and hydrazinolysis of corresponding aromatic carboxylic acids (Scheme 1). The reaction of aroylhydrazides with CS_2/KOH in absolute ethanol gave potassium aroyldithiocarbazates, and then the hydrazinolysis of potassium aroyldithiocarbazates with hydrazine hydrate afforded 3-aryl-4-amino-5-mercapto-1,2,4-triazoles **1** (Scheme 2).¹²

Ar-COOH
$$\xrightarrow{\text{H}_2\text{SO}_4}$$
 Ar-COOCH₂CH₃ $\xrightarrow{\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}}$ Ar-CONHNH₂

SCHEME 1



SCHEME 2

The synthesis of **2** was accomplished in one step with good yields by condensing 3-aryl-4-amino-5-mercapto-1,2,4-triazoles **1** with *m*phthalic acid in the presence of POCl₃ and the phase transfer catalyst tetrabutylammonium iodide (Scheme 3, Table I).



SCHEME 3

The structures of all of compounds **2** was established on the basis of elemental analysis and spectral data. The IR spectral data of compound **2** showed bands at 1600–1630 cm⁻¹, 1235–1260 cm⁻¹, and 700 cm⁻¹, due to -C=N, N-N=C, and -C-S-C, respectively. The ¹H NMR spectra of **2** exhibited multiple signals in the δ 7.00–8.70 range accounting for hydrogens of the aryl group. With compound **2g**, as an example, multiple signals appeared in the range of δ 8.59–8.44, δ 8.30–8.24, δ 8.07–8.05, and δ 7.59–7.56, which accounted for the 12 hydrogens of the aryl group. A single signal at δ 2.57 accounted for 6 hydrogens of 2 methyl group. The EI-MS for compounds **2**, except for **2b**, **2c**, and **2d**, exhibited strong

Entry	Ar	Condition	Yield (%) ^a	m.p. (°)
2a	Ph	115–120°C/11 h	76	>300
2b	2-Cl-Ph	115–120°C/12 h	62	>300
2c	3-Cl-Ph	115–120°C/13 h	60	> 300
2d	4-Cl-Ph	115–120°C/11 h	65	> 300
2e	2-CH ₃ -Ph	115–120°C/14 h	56	>300
2f	3-CH ₃ -Ph	115–120°C/13 h	61	> 300
2g	4-CH ₃ -Ph	115–120°C/14 h	70	> 300
2 h	3-Br-Ph	115–120°C/12 h	63	> 300
2i	4-Br-Ph	115–120°C/11 h	68	>300
2j	2-I-Ph	115–120°C/11 h	60	> 300
2k	3-I-Ph	115–120°C/11 h	67	>300
21	4-I-Ph	115–120°C/10 h	70	>300
2m	4-OCH ₃ -Ph	115–120°C/15 h	66	>300
2n	4-Pyridyl	115–120°C/12 h	53	>300
2o	3-Pyridyl	115–120°C/13 h	51	>300
2p	2-Furyl	115–120°C/10 h	45	>300

TABLE I Preparation of 1,3-Bis[(3-aryl)-s-triazolo[3,4-b]-[1,3,4]thiadiazole-6-yl]benzenes 2 from 3-aryl-4-amino-5-mercapto-1,2,4-triazoles 1

^aPurified yields of **2a-2p** based on *m*-ophthalmic acid.

molecular ion peaks, including some base peaks, for example, **2g** and **2o**. Compounds **2b**, **2c**, and **2d** showed molecular ion peaks corresponding to $[M^++1]$ at m/z 548 and $[M^+-1]$ m/z 546.

EXPERIMENTAL

Melting points were determined on an X_4 melting point apparatus and were uncorrected. The IR spectra were recorded on a Nicolet Nexus 470 FT-IR spectrophotometer using KBr discs in the range 4000–400 cm⁻¹. ¹H NMR spectra were recorded on a Varian Mercury-Plus 400 NMR spectrometer in CF₃COOD or pyridine- d_5 solution using TMS as an internal reference. MS spectra were recorded on a Finnigan Trace GC-MS spectrometer. Elemental analyses were taken on a Perkin-Elemer-2400-CHN elemental analysis instrument.

General Procedure for the Preparation of 3-Aryl-4-amino-5-mercapto-1,2,4-triazole 1¹²

A mixture of potassium aroyldithiocarbazate (30 mmol) and hydrazine hydrate (80%, 36 mL) was refluxed with stirring for 4–5 h at 120°C. The color of the reaction mixture changed to green, and a homogeneous

solution resulted. The acidification with HCl (3 mol/L) resulted in the precipitation of a white solid. The product was filtered, washed with cold water, and recrystallized (ethanol) to give the pure products **1**.

3-p-Methylphenyl-4-amino-5-mercapto-1,2,4-triazole (1 g)

Needle crystals, yield 55%, m.p. $217-218^{\circ}$. ¹H NMR (DMSO- d_6 , 400 MHz) δ 12.7 (s, 1H, S-H), 7.51–8.24 (m, 4H, Ar-H), 5.63 (s, 2H, NH₂), 2.54 (s, 3H, CH₃); IR (cm⁻¹): 3270, 1621, 1600, 1548, 1251. Elemental anal. calcd. for C₉H₁₀N₄S: C, 52.41; H, 4.89; N, 27.16. Found: C, 52.30; H, 4.97; N, 27.03.

General Procedure for the Preparation of 1,3-bis[(3-aryl)-S-triazolo[3,4-b]-[1,3,4]thiadiazole-6-yl]benzenes 2

A mixture of compound 3-aryl-4-amino-5-mercapto-1,2,4-triazole (2.2 mmol), *m*-phthalic acid (1.0 mmol), the phase transfer catalyst tetrabutylammonium iodide (0.5 mmol), and POCl₃ (7 mL) was stirred for 6 h at 55–60°C, and then refluxed for 11–15 h at 115–120°C. Excess POCl₃ was removed under reduced pressure. The concentrated mass was cooled and poured into crushed ice and neutralized with potassium carbonate. The separated solid was filtered, washed with water and ethanol, and then dried. The crude material was recrystallized (ethanol-pyridine), giving the pure products **2**.

1,3-Bis[(3-phenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2a)

Yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.52–8.42 (m, 5H, Ar-H), 8.19–8.11 (m, 4H, Ar-H), 7.83–7.58 (m, 5H, Ar-H); IR (KBr, cm⁻¹): 1614, 1253, 702; MS (*m*/*z*): 478 (M⁺, 2%), 321 (2%), 303 (3%), 146 (86%), 128 (100%). elemental anal. calcd. for C₂₄H₁₄N₈S₂: C, 60.24; H, 2.95; N, 23.42. Found: C, 60.15; H, 2.88; N, 23.61.

1,3-Bis[(3-*o*-chlorophenyl)-S-triazolo[3, 4-*b*]-[1,3,4] thiadiazole-6-yl]benzene (2b)

Yellow powder, ¹H NMR (CF₃COOD, 400 MHz) 8.53–8.47 (m, 4H, Ar-H), 8.31–8.26 (m, 3H, Ar-H), 8.20–8.14 (m, 2H, Ar-H), 7.83–7.51 (m, 3H, Ar-H); IR (KBr, cm⁻¹): 1623, 1251, 701; MS (*m*/*z*): 548 (M⁺+1, 54%), 546 (M⁺-1, 100%), 356 (1%), 357 (22%), 146 (53%). Elemental anal. calcd. for $C_{24}H_{12}N_8S_2Cl_2$: C, 52.66; H, 2.21; N, 20.47. Found: C, 52.81; H, 2.13; N, 20.56.

1,3-Bis[(3-*m*-chlorophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene(2c)

Pale yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.71–8.46 (m, 3H, Ar-H), 8.38–8.24 (m, 2H, Ar-H), 8.17–8.10 (m, 4H, Ar-H), 7.81–7.52 (m, 3H, Ar-H); IR (KBr, cm⁻¹): 1609, 1243, 700; MS (*m/z*): 548 (M⁺+1, 65%), 546 (M⁺-1, 78%), 356 (4%), 357 (12%), 146 (100%). Elemental anal. calcd. for C₂₄H₁₂N₈S₂Cl₂: C, 52.66; H, 2.21; N, 20.47. Found: C, 52.51; H, 2.30; N, 20.67.

1,3-Bis[(3-*p*-chlorophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl] benzene (2d)

Yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.64–8.34 (m, 7H, Ar-H), 8.05–7.71 (m, 5H, Ar-H); IR (KBr, cm⁻¹): 1627, 1254, 701; MS (m/z): 548 (M⁺+1, 71%), 546 (M⁺-1, 100%), 356(3%), 357 (16%), 146 (45%). Elemental anal. calcd. for C₂₄H₁₂N₈S₂Cl₂: C, 52.66; H, 2.21; N, 20.47. Found: C, 52.78; H, 2.30; N, 20.41.

1, 3-Bis[(3-*o*-methylphenyl)-S-triazolo[3, 4-*b*]-[1,3,4]thiadiazole-6-yl]benzene(2e)

Pale yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.46–8.41 (m, 3H, Ar-H), 8.35–8.29 (m, 2H, Ar-H), 8.12–8.05 (m, 4H, Ar-H), 7.57–7.49 (m, 3H, Ar-H), 2.56 (s, 6H, 2CH₃); IR (KBr, cm⁻¹): 1627, 1261, 695; MS (*m/z*): 506 (M⁺, 94%), 335 (9%), 317 (32%), 146 (21%), 117 (100%). Elemental anal. calcd. for C₂₆H₁₈N₈S₂: C, 61.64; H, 3.58; N, 22.12. Found: C, 61.76; H, 3.60; N, 22.25.

1,3-Bis[(3-*m*-methylphenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2f)

Pale yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.41–8.35 (m, 5H, Ar-H), 8.29–8.01 (m, 4H, Ar-H), 7.51–7.42 (m, 3H, Ar-H), 2.55 (s, 6H, 2CH₃); IR (KBr, cm⁻¹): 1621, 1252, 705; MS (*m*/*z*): 506 (M⁺, 100%), 335 (12%), 317 (52%), 146 (13%), 145 (67%), 117 (41%). Elemental anal. calcd. for C₂₆H₁₈N₈S₂: C, 61.64; H, 3.58; N, 22.12. Found: C, 61.52; H, 3.49; N, 22.27.

1,3-Bis[(3-*p*-methylphenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2g)

Yellow powder, 1H NMR (CF_3COOD, 400 MHz) δ 8.59–8.44 (m, 3H, Ar-H), 8.30–8.24 (m, 4H, Ar-H), 8.07–8.05 (m, 1H, Ar-H), 7.59–7.56 (m, 4H,

Ar-H), 2.57 (s, 6H, 2CH₃); IR (KBr, cm⁻¹): 1615, 1252, 701; MS (*m/z*): 506 (M⁺, 100%), 335 (12%), 317 (28%), 146 (32%), 145 (51%), 117 (32%), 116 (91%). Elemental anal. calcd. for $C_{26}H_{18}N_8S_2$: C, 61.64; H, 3.58; N, 22.12. Found: C, 61.52; H, 3.69; N, 22.07.

1,3-Bis[(3-*m*-bromophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2h)

Yellow powder, ¹H NMR (Pyridine- d_5 , 400 MHz) δ 8.52–8.46 (m, 3H, Ar-H), 8.37–8.30 (m, 3H, Ar-H), 8.12–8.07 (m, 2H, Ar-H), 7.57–7.52 (m, 4H, Ar-H); IR (KBr, cm⁻¹): 1621, 1263, 704; MS (*m*/*z*): 636 (M⁺, 46%), 555 (8%), 399 (26), 381 (23%), 146 (100%), 101 (43%). Elemental anal. calcd. for C₂₄H₁₂N₈S₂Br₂: C, 45.31; H, 1.90; N, 17.61. Found: C, 45.42; H, 2.11; N, 17.49.

1,3-Bis[(3-*p*-bromophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2i)

Yellow powder, ¹H NMR (Pyridine- d_5 , 400 MHz) δ 8.48–8.41 (m, 4H, Ar-H), 8.31–8.25 (m, 3H, Ar-H), 8.09–8.04 (m, 2H, Ar-H), 7.54–7.50 (m, 3H, Ar-H); IR (KBr, cm⁻¹): 1614, 1254, 700; MS (*m*/*z*): 636 (M⁺, 55%), 555 (7%), 399 (15), 381 (37%), 146 (63%), 145 (100%), 101 (61%). Elemental anal. calcd. for C₂₄H₁₂N₈S₂Br₂: C, 45.31; H, 1.90; N, 17.61. Found: C, 45.42; H, 2.11; N, 17.49.

1,3-Bis[(3-*o*-lodophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2j)

Yellow powder, ¹H NMR (CF₃COOD, 400MHz) δ 8.71–8.68 (m, 4H, Ar-H), 8.42–8.34 (m, 2H, Ar-H), 8.21–8.09 (m, 3H, Ar-H), 7.50–7.42 (m, 3H, Ar-H); IR (KBr, cm⁻¹): 1615, 1258, 703; MS (*m*/*z*): 730 (M⁺, 52%), 604 (2%), 447 (13), 429 (51%), 276 (100%), 229 (26%), 146 (38%), 145 (63%), 101 (71%). Elemental anal. calcd. for C₂₄H₁₂N₈S₂I₂: C, 39.47; H, 1.66; N, 15.34. Found: C, 39.59; H, 1.58; N, 15.42.

1,3-Bis[(3-*m*-lodophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2k)

Pale yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.76–8.72 (m, 3H, Ar-H), 8.35–8.45 (m, 4H, Ar-H), 8.15–8.07 (m, 3H, Ar-H), 7.49–7.45 (m, 2H, Ar-H); IR (KBr, cm⁻¹): 1610, 1239, 710; MS (*m*/*z*): 730 (M⁺, 44%), 604 (4%), 447 (10), 429 (55%), 276 (100%), 229 (41%), 146 (50%), 145

 $(56\%),\,101\,(84\%).$ Elemental anal. calcd. for $C_{24}H_{12}N_8S_2I_2;\,C,\,39.47;\,H,\,1.66;\,N,\,15.34.$ Found: C, 39.59; H, 1.58; N, 15.42.

1,3-Bis[(3-*p*-lodophenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6yl]benzene (2l)

Pale yellow powder, ¹H NMR (CF₃COOD, 400 MHz) δ 8.45–8.42 (m, 2H, Ar-H), 8.13–7.81 (m, 10H, Ar-H); IR (KBr, cm⁻¹): 1620, 1247, 700; MS (*m*/*z*): 730 (M⁺, 88%), 604 (4%), 447 (19), 429 (85%), 302 (20%), 229 (39%), 146 (41%), 145 (100%), 101 (61%). Elemental anal. calcd. for C₂₄H₁₂N₈S₂I₂: C, 39.47; H, 1.66; N, 15.34. Found: C, 39.38; H, 1.52; N, 15.27.

1,4-Bis[(3-*p*-methoxyphenyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2m)

Pale yellow powder, ¹H NMR (CF₃COOD, 400 MHz) 8.56–8.42 (m, 4H, Ar-H), 8.31–8.25 (m, 3H, Ar-H), 8.11–8.04 (m, 3H, Ar-H), 7.42–7.40 (m, 2H, Ar-H), 3.89 (s, 6H, 2OCH₃); IR (KBr, cm⁻¹): 1614, 1254, 700; MS (*m*/*z*): 538 (M⁺, 99%), 351 (3%), 333 (32), 146 (30%), 145 (32%), 132 (100%), 101 (22%). Elemental anal. calcd. for $C_{26}H_{18}N_8O_2S_2$: C, 57.98; H, 3.37; N, 20.80. Found: C, 57.83; H, 3.42; N, 20.73.

1,3-Bis[(3-4'-pyridyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (2n)

Pale yellow powder, ¹H NMR (Pyridine- d_5 , 400 MHz) 8.51–8.38 (m, 3H, Ar-H), 8.33–8.24 (m, 4H, Ar-H), 8.17–8.00 (m, 2H, Ar-H), 7.38-7.26 (m, 3H, Ar-H); IR (KBr, cm⁻¹): 1626, 1242, 707; MS (*m*/*z*): 480 (M⁺, 100%), 322 (26), 304 (10), 146 (23%), 145 (51%), 104 (11%). Elemental anal. calcd. for C₂₂H₁₂N₁₀S₂: C, 54.99; H, 2.52; N, 29.15. Found: C, 54.81; H, 2.42; N, 29.26.

1,3-Bis[(3-3'-pyridyl)-S-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]benzene (20)

Yellow powder, ¹H NMR (Pyridine- d_5 , 400 MHz) 8.57-8.43 (m, 4H, Ar-H), 8.35–8.22 (m, 3H, Ar-H), 8.14–8.01 (m, 3H, Ar-H), 7.31–7.25 (m, 2H, Ar-H); IR (KBr, cm⁻¹): 1614, 1254, 700; MS (*m*/*z*): 480 (M⁺, 100%), 324 (7%), 322 (42), 304 (12), 146 (27%), 145 (40%), 104 (18%). Elemental anal. calcd. for C₂₂H₁₂N₁₀S₂: C, 54.99; H, 2.52; N, 29.15. Found: C, 54.83; H, 2.69; N, 29.06.

1,3-Bis[(3-2'-furyl)-S-triazolo[3, 4-b]-[1,3,4]thiadiazole-6-yl]benzene (2p)

Pale yellow powder, ¹H NMR (Pyridine- d_5 , 400 MHz) 8.45–8.34 (m, 2H, Ar-H), 8.27–8.16 (m, 2H, Ar-H), 7.12–6.37 (m, 6H, furyl-H); IR (KBr, cm⁻¹): 1614, 1254, 700; MS (*m*/*z*): 458 (M⁺, 48%), 311 (9%), 293 (64), 146 (32%), 145 (100%), 93 (27%). Elemental anal. calcd. for C₂₀H₁₀N₈O₂S₂: C, 52.39; H, 2.20; N, 24.40. Found: C, 52.48; H, 2.11; N, 24.31.

REFERENCES

- [1] N. F. Ewiss and A. A. Bahajaj, J. Heterocylic Chem., 24, 1173 (1987).
- [2] M. Jan and G. S. R. Anjaneyulu, Indian J. Chem., 27B, 128(1987).
- [3] X. Xiong, L. X. Zhang, A. J. Zhang, and X. Sh. Lu, Synth. Commun., 32, 3455 (2002).
- [4] X. C. Wang, L. Zheng, Z. J. Quang, and D. J. Xu, Synth.Commun., 33, 2891 (2003).
- [5] M. Chen and T. Xiao, Chem. Res. Chinese U., 62, 20 (2004)
- [6] S. M. El-Khawass and N. S. Habib, J. Heterocylic Chem., 26, 177 (1989).
- [7] Q. M. Bano and S. G. Tiwari, *Indian J. Chem.*, **31B**, 714 (1992).
- [8] F. M. Liu, J. X. Yu, and W. J. Lu, Chin. J. Chem., 17, 62 (1999).
- [9] L. X. Zhang, A. J. Zhang, and G. Q. Zhou, Chinese J. Org. Chem., 22, 663 (2002).
- [10] M. Li, L. R. Wen, and W. J. Fu, Chinese J. Org. Chem., 23, 678 (2003).
- [11] G. Q. Hu, Z. Q. Zhang, and W. L. Huang, Chinese Acta Chim. Sinica., 62, 204 (2004).
- [12] R. Reid and N. D. Heindel, J. Heterocylic Chem., 13, 925 (1976).