

pubs.acs.org/JACS

Copper-Mediated Radical—Polar Crossover Enables Photocatalytic Oxidative Functionalization of Sterically Bulky Alkenes

Nicholas L. Reed, Grace A. Lutovsky, and Tehshik P. Yoon*

Cite This: J. Ar	n. Chem. Soc. 2021, 143, 6065–6070	Read Online	
ACCESS	III Metrics & More	E Article Recommendations	Supporting Information

ABSTRACT: Oxidative heterofunctionalization reactions are among the most attractive methods for the conversion of alkenes and heteroatomic nucleophiles into complex saturated heterocycles. However, the state-of-the-art transition-metal-catalyzed methods to effect oxidative heterofunctionalizations are typically limited to unhindered olefins, and different nucleophilic partners generally require quite different reaction conditions. Herein, we show that Cu(II)-mediated radical—polar crossover allows for highly efficient and exceptionally mild photocatalytic oxidative heterofunctionalization reactions between bulky tri- and tetrasubstituted alkenes and a wide variety of nucleophilic partners. Moreover, we demonstrate that the broad scope of this transformation arises from photocatalytic alkene activation and thus complements existing transition-metal-catalyzed methods for oxidative heterofunctionalizations and that the combination of photocatalytic substrate activation and Cu(II)-mediated radical oxidation can address long-standing challenges in catalytic oxidation chemistry.

xidative alkene functionalizations¹ are powerful methods for the rapid synthesis of saturated heterocycles commonly found in many important natural products, pharmaceuticals, and agrochemicals.² These reactions are synthetically attractive for a number of reasons. First, they enable the direct oxidative coupling of alkenes with heteronucleophiles without the need to prepare preoxidized and often unstable heteroatom donors as internal oxidants.³⁻⁶ Moreover, and in contrast to redox-neutral alkene hydro-functionalization reactions,^{7–9} oxidative alkene heterofunctionalizations conserve an olefin functionality that can be valuable for further synthetic elaborations. The state-of-the-art methods are Pd(II)-catalyzed processes that can proceed via either of two mechanistically distinct pathways (Scheme 1):^{10,11} synnucleopalladation involves migratory insertion of an alkene across a Pd-heteroatom bond,¹² while *anti*-nucleopalladation involves attack of a heteroatomic nucleophile on a Pd(II)coordinated alkene. The details of these mechanisms, however, engender several notable limitations. First, both mechanistic manifolds rely upon the coordination of the alkene to the Pd(II) center. Highly substituted alkenes that coordinate weakly, therefore, are poor reaction partners in all but a handful of highly specialized cases.^{13,14} Second, relatively subtle changes to the substrate structures and reaction conditions can switch the operative mechanistic pathway,¹⁵ requiring reoptimization of reaction conditions for heteroatom nucleophiles with different metal-binding propensities. Thus, the identification of conditions for the oxidative functionalization of sterically hindered alkenes with diverse heteronucleophiles remains an unsolved challenge that Pd(II) catalysis seems poorly positioned to address.

We imagined that photoredox catalysis might offer a complementary strategy toward oxidative alkene heterofunctionalization that would be free of these long-standing

Scheme 1. Oxidative Heterofunctionalization of Alkenes



Received: March 12, 2021 **Published:** April 15, 2021



Communication

Journal of the American Chemical Society

limitations. Electron-rich tri- and tetrasubstituted alkenes that are generally poor substrates for Pd(II)-catalyzed oxidative heterofunctionalizations are easily activated by photoredox catalysts. Moreover, Nicewicz and others have shown that the resulting alkene radical cations are susceptible to highly regioselective nucleophilic attack, and because this step does not involve a discrete nucleophile-catalyst interaction, they are relatively agnostic toward the identity of the nucleophilic reaction partner. $^{16-22}$ Thus, if an appropriate terminal oxidant could be identified to intercept the resulting organoradical intermediate and promote oxidative elimination, this process could constitute a new, alternative platform for oxidative alkene heterofunctionalization. Lei and co-workers recently demonstrated the feasibility of a similar approach in the oxidative coupling of olefins with alcohols or azoles using a dual photoredox/hydrogen evolution system.²³ However, these reactions were limited to activated styrenic olefins and structurally simple nucleophiles, and thus the construction of complex saturated heterocyclic motifs has yet to be realized.

Inexpensive Cu(II) salts are ideal terminal oxidants for photoredox applications.²⁴ Many electron-rich radical species are efficiently oxidized by Cu(II), which offers a means to divert the prototypical organoradical chemistry enabled by photoredox catalysis toward formally cationic reactivity via radical-polar crossover.²⁵⁻²⁷ We have leveraged this combination to design a photocatalytic protocol for the alkoxylation of benzylic C–H bonds²⁸ and the photocatalytic oxyamination of electron-rich olefins.²⁹ In both instances, oxidation of a photogenerated organoradical was followed by substitution with a heteroatomic nucleophile. To achieve oxidative heterofunctionalization, however, the radical mechanism would have to be terminated by an alternative oxidative elimination process. Seminal investigations by Kochi showed that thermally generated organoradicals could be diverted either toward oxidative substitution or oxidative elimination by tuning of reaction conditions,³⁰⁻³³ and more recent studies by Glorius and Tunge have demonstrated that the Cu(II)mediated oxidative elimination pathway is operative for radicals produced by photochemical decarboxylation.^{34,35}

We therefore propose a new mechanism for copper-enabled photoredox oxidative heterofunctionalization (Scheme 2). Photoexcitation of MesAcrPh⁺ affords a highly oxidizing excited state MesAcrPh⁺* (${}^{*}E_{1/2}^{\text{red}} = +2.12$ V vs SCE).²¹ Rapid reductive quenching by alkene 1 would furnish radical cation 2 and MesAcrPh[•]. Radical cation 2 is a potent electrophile and should be readily trapped by a pendant heteroatomic nucleophile to generate carbon-centered radical 3. Radical oxidative addition of 3 by an appropriate Cu(II) oxidant would afford an organocopper(III) intermediate. Key to the successful realization of this strategy would be the optimization of the elimination step to favor contrathermodynamic elimination to 4. The alternate regiochemistry of oxidative elimination would afford an electron-rich enamine that would not survive the highly oxidizing conditions of this transformation. Finally, oxidation of MesAcrPh[•] by Cu(I) generated over the course of the reaction would close the photoredox catalytic cycle and regenerate ground-state MesAcrPh⁺.

We began our investigations by examining the oxidative cyclization of trisubstituted alkene 1 upon irradiation in the presence of 2.5 mol % *N*-phenylacridinium tetrafluoroborate (MesAcrPh⁺), 1 equiv of trifluoroacetic acid (TFA), and a variety of Cu(II) terminal oxidants (Table 1, entries 1–6). The

Scheme 2. Proposed Reaction Design for Photocatalytic Oxidative Heterofunctionalization

pubs.acs.org/JACS



Table 1. Optimization of Oxidative Amination



^{*a*}Reactions conducted using 1 (0.1 mmol), oxidant (2 equiv), TFA (1 equiv), and photocatalyst (2.5 mol %) in degassed 1,2-DCE and irradiated with a 15 W blue LED flood lamp for 16 h. ^{*b*}Yields were determined by ¹H NMR analysis of the unpurified reaction mixtures using phenanthrene as an internal standard. ^{*c*}Reaction conducted in absence of 1 equiv TFA. ^{*d*}Reaction vessel was covered in aluminum foil.

formation of cyclized product 4 is observed in most cases, but as expected, the identity of the Cu(II) oxidant proved to be crucial. We hypothesize that sterically bulky carboxylate ligands improve the regioselectivity of alkene formation during Cu(II)-mediated oxidative elimination.^{36–38} Other highly oxidizing



Table 2. Scope of Photocatalytic Oxidative Heterofunctionalization^a



photocatalysts afford **4** in moderate yields (entries 7 and 8), but MesAcrPh⁺ proved optimal for this transformation. Empirically, we also observed that the addition of 1 equiv TFA improved the rate and reproducibility of oxidative cyclization (entry 9). This is most readily rationalized by the increased solubility of Cu(II) salts in organic media upon the addition of trifluoroacetic acid.^{39–41} Thus, optimized conditions were found to be 2.5 mol % MesAcrPh⁺, 2 equiv of copper(II) 2-ethylhexanoate (Cu(EH)₂), 1 equiv of **1**, and 1 equiv of TFA in 1,2-DCE with irradiation by two 15 W blue LED flood lamps (entry 6). Control reactions verified the photocatalytic nature of this process: omitting photocatalyst,

Cu(II) oxidant, or light resulted in no formation of pyrrolidine 4 (entries 10-12).

The degree of generality exhibited by this transformation is notable (Table 2). A wide variety of arylsulfonamides undergo efficient oxidative cyclization. Electron-rich and electrondeficient aryl sulfonamides react smoothly (4-8), including sterically hindered *ortho*-substituted sulfonamides (8). Unfortunately, 2-nosylsulfonamide was not tolerated (9), presumably due to the redox activity of the nitroarene. Alkyl sulfonamides are excellent reaction partners (10 and 11); these include a sterically hindered *tert*-butylsulfonamide, which gives a 67% yield of 12. Sulfamoyl ureas (13), alcohols (14), and carboxylic acids (15) give high yields of cyclized product, offering efficient access to a diverse array of heterocyclic scaffolds. Importantly, no alteration of the catalyst system or reaction conditions is necessary in these cases despite the disparate nucleophiles examined.

Modifications to the length and identity of the tethering group are readily accommodated, and the functional group tolerance of this method is excellent. Common organic functional groups including esters (16), alcohols (17), silvl ethers (18), and acetals (19) are tolerated. These reactions exhibit excellent chemoselectivity as only cyclization of the sulfonamide is observed in the presence of nucleophilic free alcohols (17). While cyclizations affording five-membered rings are most efficient (20), six-membered rings are also readily accessible (21). Given the oxidizing nature of these conditions, we were pleased to find that tertiary amines (22) and anilines (23) are well tolerated. Lewis basic heterocycles that might coordinate to Cu(II) also do not interfere with the desired transformation (24); however, substrates containing Lewis basic functionalities react most efficiently with 2 equiv of p-TsOH acid in place of TFA, presumably because the basic nitrogen is protonated under these conditions. These functionalities are also particularly noteworthy because they are common poisons for transition metal catalysts, further demonstrating the unique complementarity of this photocatalytic system to its Pd(II)-catalyzed counterparts. A sterically hindered tertiary nucleophile undergoes cyclization, delivering a densely functionalized spirocyclic tetrahydrofuran (25) in 57% yield. Azetidine 26 was prepared in 54% yield, showcasing the potential utility of this method in the synthesis of medicinally desirable structures.⁴² Bicyclic heterocycles could also be synthesized (27), demonstrating the application of this method in the construction of higher-order molecular architectures.

Trisubstituted alkenes are excellent substrates in this reaction, and in all cases, we observe exclusive anti-Markovnikov selectivity in the initial bond-forming event.⁴³⁻⁴⁵ Thus, cyclization can proceed in endo fashion, consistent with generation of a more stable radical intermediate upon nucleophilic trapping (28). In this experiment, the oxidative elimination gives a moderate preference for the exocylic terminal olefin (4:1), in line with the regioselectivites observed by Glorius for decarboxylative olefination.³⁴ Additionally, oxidative elimination affording endocyclic olefins can be conducted without issue (29). Styrenic olefins afford 1,1-disubstituted styrenes (30) as the sole products. Importantly, and in contrast to many Pd(II)catalyzed oxidative amination methods, no alkene isomerization is observed.^{46–49} Short reaction times were crucial to obtain good yields of cyclized product, however, as 30 undergoes slow oxidative decomposition under the reaction conditions. To our delight, tetrasubstituted alkenes, typically among the most challenging substrates for Pd(II)-catalyzed heterofunctionalization methods, react smoothly, and consistent with the absence of a discrete catalyst-nucleophile interaction, the identity of the heteroatomic nucleophile has little effect on the efficiency of cyclization. Sulfonamides (31), carboxylic acids (32), and alcohols (33) all afford good to excellent yields of the desired heterocycles.

Importantly, the scope of this new oxidative heterofunctionalization reaction provides a synthetic capacity that state-ofthe-art Pd(II)-catalyzed methods do not. To demonstrate this complementarity, we subjected trisubstituted alkene 1 and tetrasubstituted alkene 34 to several known, highly active Pd(II)-based catalyst systems for oxidative amination (Table 3a and 3b). $^{50-53}$ These systems uniformly afforded unsat-

Table 3. Survey of Oxidative Amination Methods

A Oxidative Amination of Trisubstituted Alkene 1

Me Me	Me Me	•	4 Me
entry	catalyst system	% yield	% RSM
1 ^a	MesAcrPh ⁺ /h _V , Cu(EH) ₂	87	0
2 ^b	Pd(OAc) ₂ /DMSO, O ₂	trace	70
3c	Pd(OAc) ₂ /pyridine, O ₂	13	59
4 ^d	Pd(TFA) ₂ /(–)-sparteine, O ₂	trace	69

B Oxidative Amination of Tetrasubstituted Alkene 34



^{*a*}Reaction conducted using MesAcrPh⁺ (2.5 mol %), Cu(EH)₂ (2 equiv), TFA (1 equiv), and 1,2-DCE. Irradiated for 16–18 h. ^{*b*}Reaction conducted using Pd(OAc)₂ (5 mol %), NaOAc (2 equiv), O₂, DMSO, rt, 72 h. ^{*c*}Reaction conducted using Pd(OAc)₂ (5 mol %), pyridine (10 mol %), O₂, toluene (0.1 M), 80 °C, 24 h. ^{*d*}Reaction conducted using Pd(TFA)₂ (10 mol %), (–)-sparteine (40 mol %), DIPEA (2 equiv), MS 3 Å, O₂, toluene (0.1 M), 80 °C, 26 h.

isfactory yields of oxidative cyclization with significant decomposition of the starting alkenes. The highest yields were obtained with the catalyst system reported by Stahl $(Pd(OAc)_2/pyridine)^{52}$ but only after extended reaction times at elevated temperatures (13% and 22% yield of 4 and 31, respectively). In contrast, our photocatalytic protocol effects rapid and efficient oxidative cyclization of both 1 and 34 to their corresponding heterocycles.

The origin of this complementary reactivity can readily be rationalized upon examination of each reaction component by cyclic voltammetry (Figure 1). Trisubstituted alkenes are oxidized at significantly less positive potentials than sulfonamides, alcohols, or carboxylic acids in 1,2-DCE. Moreover, the oxidation of functionalized alkene 1 occurs at significantly less positive potentials than either alkenes or sulfonamides alone. As detailed by Moeller in studies of the reactions of heteronucleophiles with electrochemically generated alkene radical cations, ^{54,55} this observation would be consistent with a relatively slow alkene oxidation followed by a rapid cyclization event. Thus, the activation of the polysubsituted alkene moiety by photochemical oxidation rather than by transition metal



Figure 1. Cyclic voltammetry experiments.

coordination is responsible for the complementary reactivity profile we observe. 56

In summary, we have shown that a photocatalytic system consisting of a highly oxidizing photocatalyst and a Cu(II) terminal oxidant enables the oxidative heterofunctionalization of highly substituted alkenes with a diverse range of nucleophilic partners. This reaction addresses several longstanding synthetic limitations of state-of-the-art oxidative alkene heterofunctionalizations. The mechanism of photoactivation involving the generation of an alkene radical cation intermediate obviates the need for discrete metal-alkene or metal-nucleophile interactions, and the scope of this photocatalytic method thus complements those of existing Pd(II)catalyzed reactions. More generally, this reaction adds to the growing evidence that Cu(II) salts are ideal terminal oxidants for photoredox oxidation reactions. While we have previously shown that a similar system enables oxidative substitution reactions of photochemically generated organoradical intermediates, here we show that their reactivity can be diverted toward oxidative elimination reactions. Together, these results suggest a powerful and potentially general strategy for the development of a wide range of novel oxidative functionalization reactions that exploit the versatility of photoredox activation.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/jacs.1c02747.

Experimental procedures; characterization data; spectra for all new compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

Tehshik P. Yoon – Department of Chemistry, University of Wisconsin—Madison, Madison, Wisconsin 53706, United States; o orcid.org/0000-0002-3934-4973; Email: tyoon@chem.wisc.edu

Authors

Nicholas L. Reed – Department of Chemistry, University of Wisconsin—Madison, Madison, Wisconsin 53706, United States; Occid.org/0000-0003-4234-5464 pubs.acs.org/JACS

Communication

Grace A. Lutovsky – Department of Chemistry, University of Wisconsin—Madison, Madison, Wisconsin 53706, United States; orcid.org/0000-0002-6040-546X

Complete contact information is available at: https://pubs.acs.org/10.1021/jacs.1c02747

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Funding for this project was provided by the NIH (GM095666). G.L. is grateful to 3M for a 3M Science & Technology Fellowship. NMR and MS facilities at UW–Madison are funded by the NIH (1S10 OD020022-1) and a generous gift from the Paul J. and Margaret M. Bender Fund.

REFERENCES

(1) For a review, see: Nakamura, I.; Yamamoto, Y. Transition-Metal-Catalyzed Reactions in Heterocyclic Synthesis. *Chem. Rev.* **2004**, *104*, 2127–2198.

(2) Heterocycles in Natural Product Synthesis; Majumdar, K. C., Chattopadhyay, S. K., Eds.; Wiley-VCH Verlag & Co. KGaA: Weinheim, 2011.

(3) Stahl, S. S. Palladium Oxidase Catalysis: Selective Oxidation of Organic Chemicals via Direct Dioxygen-Coupled Catalytic Turnover. *Angew. Chem., Int. Ed.* **2004**, *43*, 3400–3420.

(4) Stahl, S. S. Palladium-Catalyzed Oxidation of Organic Chemicals with O₂. *Science* **2005**, *309*, 1824–1826.

(5) Kitamura, M.; Narasaka, K. Synthesis of Aza-Heterocycles from Oximes by Amino-Heck Reaction. *Chem. Rec.* **2002**, *2*, 268–277.

(6) Race, N. J.; Hazelden, I. R.; Faulkner, A.; Bower, J. F. Recent developments in the use of aza-Heck cyclizations for the synthesis of chiral N-heterocycles. *Chem. Sci.* **201**7, *8*, 5248–5260.

(7) Muller, T. E.; Hultzsch, K. C.; Yus, M.; Foubelo, F.; Tada, M. Hydroamination: Direct Addition of Amines to Alkenes and Alkynes. *Chem. Rev.* **2008**, *108*, 3795–3892.

(8) Crossley, S. W. M.; Obradors, C.; Martinez, R. M.; Shenvi, R. A. Mn-, Fe-, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev.* 2016, 116, 8912–9000.

(9) Margrey, K. A.; Nicewicz, D. A. A General Approach to Catalytic Alkene Anti-Markovnikov Hydrofunctionalization Reactions via Acridinium Photoredox Catalysis. *Acc. Chem. Res.* **2016**, *49*, 1997–2006.

(10) McDonald, R. I.; Liu, G.; Stahl, S. S. Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications. *Chem. Rev.* **2011**, *111*, 2981–3019.

(11) Kočovský, P.; Bäckvall, J.-E. The syn/anti-Dichotomy in the Palladium-Catalyzed Addition of Nucleophiles to Alkenes. *Chem. - Eur. J.* **2015**, *21*, 36–56.

(12) Hanley, P. S.; Hartwig, J. F. Migratory Insertion of Alkenes into Metal–Oxygen and Metal–Nitrogen Bonds. *Angew. Chem., Int. Ed.* **2013**, *52*, 8510–8525.

(13) Yang, G.; Shen, C.; Zhang, W. An Asymmetric Aerobic Aza-Wacker-Type Cyclization: Synthesis of Isoindolinones Bearing Tetrasubstituted Carbon Stereocenters. *Angew. Chem., Int. Ed.* **2012**, *51*, 9141–9145.

(14) Kou, X.; Shao, Q.; Ye, C.; Yang, G.; Zhang, W. Asymmetric Aza-Wacker-Type Cyclization of N-Ts Hydrazine-Tethered Tetrasubstituted Olefins: Synthesis of Pyrazolines Bearing One Quaternary or Two Vicinal Stereocenters. J. Am. Chem. Soc. 2018, 140, 7587–7597. (15) Trend, R. M.; Ramtohul, Y. K.; Stoltz, B. M. Oxidative Cyclizations in a Nonpolar Solvent Using Molecular Oxygen and Studies on the Stereochemistry of Oxypalladation. J. Am. Chem. Soc. 2005, 127, 17778–17788.

Journal of the American Chemical Society

pubs.acs.org/JACS

(16) Gassman, P. G.; Bottorff, K. J. Anti-Markovnikov Addition of Nucleophiles to a Non-conjugated Olefin *via* Single Electron Transfer Photochemistry. *Tetrahedron Lett.* **1987**, *28*, 5449–5452.

(17) Mizuno, K.; Nakanishi, I.; Ichinose, N.; Otsuji, Y. Structural Dependence on Photoaddition of Methanol to Arylalkenes. Solvent and Additive Effects on Photoinduced Electron Transfer Reaction. *Chem. Lett.* **1989**, *18*, 1095–1098.

(18) Hamilton, D. S.; Nicewicz, D. A. Direct Catalytic Anti-Markovnikov Hydroetherification of Alkenols. J. Am. Chem. Soc. 2012, 134, 18577–18580.

(19) Nguyen, T. M.; Nicewicz, D. A. Anti-Markovnikov Hydroamination of Alkenes Catalyzed by an Organic Photoredox System. 2013, *135*, 9588–9591.

(20) Perkowski, A. J.; Nicewicz, D. A. Direct Catalytic Anti-Markovnikov Addition of Carboxylic Acids to Alkenes. J. Am. Chem. Soc. 2013, 135, 10334–10337.

(21) Wilger, D. J.; Grandjean, J. M.; Lammert, T.; Nicewicz, D. A. The Direct Anti-Markovnikov Addition of Mineral Acids to Styrenes. *Nat. Chem.* **2014**, *6*, 720–72.

(22) Wang, H.; Man, Y.; Xiang, Y.; Wang, K.; Li, N.; Tang, B. Regioselective Intramolecular Markovnikov and anti-Markovnikov hydrofunctionalization of alkenes *via* photoredox catalysis. *Chem. Commun.* **2019**, *55*, 11426–11429.

(23) Yi, H.; Niu, L.; Song, C.; Li, Y.; Dou, B.; Singh, A. K.; Lei, A. Photocatalytic Dehydrogenative Cross-Coupling of Alkenes with Alcohols or Azoles without External Oxidant. *Angew. Chem., Int. Ed.* **2017**, *56*, 1120–1124.

(24) Reed, N. L.; Yoon, T. P. Oxidase reactions in photoredox catalysis. *Chem. Soc. Rev.* 2021, *50*, 2954–2967.

(25) Lampard, C.; Murphy, J. A.; Lewis, N. Tetrathiafulvalene as a catalyst for radical-polar crossover reactions. J. Chem. Soc., Chem. Commun. 1993, 295–297.

(26) Murphy, J. A.; Rasheed, F.; Roome, S. J.; Lewis, N. Termination of radical-polar crossover reactions by intramolecular nucleophiles. *Chem. Commun.* **1996**, 737–738.

(27) Murphy, J. A.; Rasheed, F.; Gastaldi, S.; Ravishanker, T.; Lewis, N. Synthesis of functionalized indolines by radical-polar crossover reactions. *J. Chem. Soc., Perkin Trans.* 1 **1997**, 1549–1558.

(28) Lee, B. J.; DeGlopper, K. S.; Yoon, T. P. Site-Selective Alkoxylation of Benzylic C–H Bonds by Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2020**, *59*, 197–202.

(29) Reed, N. L.; Herman, M. I.; Miltchev, V. P.; Yoon, T. P. Photocatalytic Oxyamination of Alkenes: Copper(II) Salts as Terminal Oxidants in Photoredox Catalysis. *Org. Lett.* **2018**, *20*, 7345–7350.

(30) Kochi, J. K. Oxidation of Allylic Radicals by Electron Transfer: Effect of Complex Copper Salts. J. Am. Chem. Soc. **1962**, 84, 3271–3277.

(31) Kochi, J. K. The Decomposition of Peroxides Catalyzed by Copper Compounds and the Oxidation of Alkyl Radicals by Cupric Salts. J. Am. Chem. Soc. **1963**, 85, 1958–1968.

(32) Kochi, J. K.; Bemis, A. Carbonium Ions from Alkyl Radicals by Electron Transfer. J. Am. Chem. Soc. **1968**, 90, 4038–4051.

(33) Kochi, J. K.; Bemis, A.; Jenkins, C. L. Mechanism of Electron Transfer Oxidation of Alkyl Radicals by Copper(II) Complexes. *J. Am. Chem. Soc.* **1968**, *90*, 4616–4625.

(34) Tlahuext-Aca, A.; Candish, L.; Garza-Sanchez, R. A.; Glorius, F. Decarboxylative Olefination of Activated Aliphatic Acids Enabled by Dual Organophotoredox/Copper Catalysis. *ACS Catal.* **2018**, *8*, 1715–1719.

(35) Cartwright, K. C.; Lang, S. B.; Tunge, J. A. Photoinduced Kochi Decarboxylative Elimination for the Synthesis of Enamides and Enecarbamates from N-Acyl Amino Acids. *J. Org. Chem.* **2019**, *84*, 2933–2940.

(36) Faulkner, A.; Race, N. J.; Scott, J. S.; Bower, J. F. Copper catalyzed Heck-like cyclizations of oxime esters. *Chem. Sci.* 2014, *5*, 2416–2421.

(37) Wu, X.; Riedel, J.; Dong, V. M. Transforming Olefins into γ,δ -Unsaturated Nitriles through Copper Catalysis. *Angew. Chem., Int. Ed.* **2017**, *56*, 11589–11593.

(38) Wang, C.; Liu, R.-H.; Tian, M.-Q.; Hu, X.-H.; Loh, T.-P. Regioselective Copper-Catalyzed Oxidative Coupling of α -Alkylated Styrenes with Tertiary Alkyl Radicals. *Org. Lett.* **2018**, *20*, 4032–4035.

(39) Antilla, J. C.; Buchwald, S. L. Copper-Catalyzed Coupling of Arylboronic Acids and Amines. *Org. Lett.* 2001, *3*, 2077–2079.

(40) Yoo, W.-J.; Tsukamoto, T.; Kobayashi, S. Visible-Light-Mediated Chan–Lam Coupling Reactions of Aryl Boronic Acids and Aniline Derivatives. *Angew. Chem., Int. Ed.* **2015**, *54*, 6587–6590. (41) Brewer, A. C.; Hoffman, P. C.; Martinelli, J. R.; Kobierski, M.

E.; Mullane, N.; Robbins, D. Development and Scale-Up of a Continuous Aerobic Oxidative Chan–Lam Coupling. *Org. Process Res. Dev.* **2019**, 23, 1484–1498.

(42) Marson, C. M. New and unusual scaffolds in medicinal chemistry. *Chem. Soc. Rev.* 2011, 40, 5514–5533.

(43) Neunteufel, R. A.; Arnold, D. R. Radical Ions in Photochemistry. I. 1,1-Diphenylethylene Cation Radical. J. Am. Chem. Soc. 1973, 95, 4080–4081.

(44) Arnold, D. R.; Du, X.; Chen, J. The Effect of Meta- or Para-Cyano Substitution on the Reactivity of the Radical Cations of Arylalkenes and Alkanes. Radical Ions in Photochemistry, Part 34. *Can. J. Chem.* **1995**, 73, 307–318.

(45) Arnold, D. R.; Chan, M. S.; McManus, K. A. Photochemical Nucleophile-Olefin Combination, Aromatic Substitution (photo-NOCAS) Reaction, Part 12. Factors Controlling the Regiochemistry of the Reaction with Alcohol as the Nucleophile. *Can. J. Chem.* **1996**, *74*, 2143–2166.

(46) Kotov, V.; Scarborough, C. C.; Stahl, S. S. Palladium-Catalyzed Aerobic Oxidative Amination of Alkenes: Development of Intra- and Intermolecular Aza-Wacker Reactions. *Inorg. Chem.* **2007**, *46*, 1910–1923.

(47) Kohler, D. G.; Gockel, S. N.; Kennemur, J. L.; Waller, P. J.; Hull, K. L. Palladium-catalyzed *anti*-Markovnikov selective oxidative amination. *Nat. Chem.* **2018**, *10*, 333–340.

(48) Allen, J. R.; Bahamonde, A.; Furukawa, Y.; Sigman, M. S. Enantioselective *N*-Alkylation of Indoles via an Intermolecular Aza-Wacker-Type Reaction. *J. Am. Chem. Soc.* **2019**, *141*, 8670–8674.

(49) Bahamonde, A.; Al Rifaie, B.; Martin-Heras, V.; Allen, J. R.; Sigman, M. S. Enantioselective Markovnikov Addition of Carbamates to Allylic Alcohols for the Construction of α -Secondary and α -Tertiary Amines. J. Am. Chem. Soc. **2019**, 141, 8708–8711.

(50) Liu, G.; Stahl, S. S. Two-Faced Reactivity of Alkenes: cis- versus trans-Aminopalladation in Aerobic Pd-Catalyzed Intramolecular Aza-Wacker Reactions. J. Am. Chem. Soc. 2007, 129, 6328–6335.

(51) Larock, R. C.; Hightower, T. R.; Hasvold, L. A.; Peterson, K. P. Palladium(II)-Catalyzed Cyclization of Olefinic Tosylamides. *J. Org. Chem.* **1996**, *61*, 3584–3585.

(52) Fix, S. R.; Brice, J. L.; Stahl, S. S. Efficient Intramolecular Oxidative Amination of Olefins through Direct Dioxygen-Coupled Palladium Catalysis. *Angew. Chem., Int. Ed.* **2002**, *41*, 164–166.

(53) Yip, K.-T.; Yang, M.; Law, K.-L.; Zhu, N.-Y.; Yang, D. Pd(II)-Catalyzed Enantioselective Oxidative Tandem Cyclization Reactions. Synthesis of Indolines through C–N and C–C Bond Formation. J. Am. Chem. Soc. 2006, 128, 3130–3131.

(54) Moeller, K.; Tinao, L. V. Intramolecular Anodic Olefin Coupling Reactions: The Use of Bis Enol Ether Substrates. J. Am. Chem. Soc. **1992**, 114, 1033–1041.

(55) Xu, H.-C.; Moeller, K. D. Intramolecular Anodic Olefin Coupling Reactions: Use of the Reaction Rate to Control Substrate/ Product Selectivity. *Angew. Chem., Int. Ed.* **2010**, *49*, 8004–8007.

(56) When a 1,2-disubstituted alkene was subjected to our photocatalytic conditions only unreacted starting material was observed (see Supporting Information). This result is inconsistent with an alternate pathway involving initial oxidation of the heteroatomic nucleophile by the excited state photocatalyst.