

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.138$
 $S = 0.884$
 10 384 reflections
 584 parameters
 H-atom parameters
 constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0513P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$
 Extinction correction:
 SHELXTL (Sheldrick,
 1997)
 Extinction coefficient:
 0.0063 (10)
 Scattering factors from
 International Tables for
 Crystallography (Vol. C)

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$)

O1A—C9A	1.377 (2)	O1B—C9B	1.373 (2)
O1A—C14	1.450 (2)	O1B—C11B	1.453 (2)
O2A—C12A	1.442 (2)	O2B—C11B	1.440 (2)
O2A—C11A	1.443 (2)	O2B—C12B	1.446 (2)
O3A—C20A	1.368 (2)	O3B—C20B	1.366 (2)
O3A—C21A	1.386 (2)	O3B—C21B	1.408 (2)
O4A—C27A	1.376 (2)	O4B—C27B	1.375 (2)
O4A—C28A	1.397 (3)	O4B—C28B	1.393 (2)
C2A—C3A	1.325 (2)	C2B—C3B	1.325 (2)
C7A—C10A	1.508 (2)	C7B—C10B	1.510 (2)
C8A—C13A	1.501 (2)	C8B—C13B	1.501 (2)
C10A—C11A	1.486 (3)	C10B—C11B	1.490 (3)
C11A—C12A	1.456 (3)	C11B—C12B	1.445 (3)
C12A—C13A	1.483 (3)	C12B—C13B	1.491 (2)
C12A—O2A—C11A	60.6 (1)	C11B—O2B—C12B	60.1 (1)
C21A—O3A—C20A—C19A	−14.4 (3)		
C28A—O4A—C27A—C26A	−30.6 (3)		
C21B—O3B—C20B—C19B	6.3 (3)		
C28B—O4B—C27B—C26B	−1.4 (3)		

Table 2. Hydrogen-bonding geometry ($\text{\AA}$, $^\circ$)

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C12A—H12A $\cdots$ O2B	0.98	2.52	3.261 (2)	132

The data collection covered over a hemisphere of reciprocal space by a combination of three sets of exposures; each set had a different φ angle (0, 88 and 180°) for the crystal and each exposure of 30 s covered 0.3° in ω . The crystal-to-detector distance was 3 cm and the detector swing angle was -20° . Crystal decay was monitored by repeating 30 initial frames at the end of data collection and analysing the intensity of duplicate reflections, and was found to be negligible. The structure was solved by direct methods and refined by full-matrix least-squares techniques. After checking their presence in the difference map, all the H atoms were placed at geometrically calculated positions and a riding model was used for their refinement; rotating group refinement was used for the methyl groups.

Data collection: *SMART* (Siemens, 1996). Cell refinement: *SAINT* (Siemens, 1996). Data reduction: *SAINT*. Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1997). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL* and *PARST* (Nardelli, 1995).

The authors would like to thank the Malaysian Government and Universiti Sains Malaysia for research grant R&D No. 190-9609-2801. KC thanks the Universiti Sains Malaysia for a Visiting Post-Doctoral Fellowship.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: NA1398). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Boyd, D. R. & Sharma, N. D. (1986). *Chem. Soc. Rev.* **25**, 289–296.
 Nardelli, M. (1983). *Acta Cryst.* **C39**, 1141–1142.
 Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
 Sheldrick, G. M. (1997). *SHELXTL. Structure Determination Software Programs*. Version 5.10. Bruker Analytical X-ray Systems Inc., Madison, Wisconsin, USA.
 Siemens (1996). *SMART and SAINT. Area Detector Control and Integration Software*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Smith, J. G. (1984). *Synthesis*, pp. 629–656.

Acta Cryst. (1999). **C55**, 770–772

4-(*p*-Methoxyphenyl)-3,5-bis(2-pyridyl)-4*H*-1,2,4-triazole

HOONG-KUN FUN,^a KANDASAMY CHINNAKALI,^{a†} SICHANG SHAO,^{b‡} DUNRU ZHU^b AND XIAO ZENG YOU^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and
^bCoordination Chemistry Institute, State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing 210093, People's Republic of China. E-mail: hkfun@usm.my

(Received 4 January 1999; accepted 28 January 1999)

Abstract

In the title molecule, C₁₉H₁₅N₅O, the dihedral angle between the phenyl and the triazole rings is $59.30(7)^\circ$. The triazole ring forms dihedral angles of $47.90(8)$ and $26.91(8)^\circ$ with the two pyridyl rings. In the crystal, the molecules exist as centrosymmetrically C—H $\cdots$ N intermolecular hydrogen-bonded dimers. The glide-related molecules translated a unit along the *b* axis are linked by weak C—H $\cdots$ N hydrogen bonds to form an infinite chain along the *c* axis which is further stabilized by C—H $\cdots$ π interactions.


Comment

Extensive studies have been carried out on the substituted 1,2,4-triazole ligands (Cornelissen *et al.*, 1992; Gupta & Bhargava, 1978; Kunkeler *et al.*, 1996). It

† On leave from: Department of Physics, Anna University, Chennai 600 025, India.

‡ Visiting Scholar from: Fuyang Normal College, 236032 Anhui, People's Republic of China.

is of interest that some iron(II) complexes containing substituted 1,2,4-triazole ligands are the spin-crossover materials which could be used as molecular-based memory devices, displays and optical switches (Garcia *et al.*, 1997; Kahn & Martinez, 1998). We have recently synthesized the title molecule, (I), which can act as a potentially dinucleating ligand. The X-ray structure determination was carried out in order to elucidate the molecular conformation.



Bond lengths and angles in the structure are comparable with those reported for related structures (Wang *et al.*, 1998; Chen *et al.*, 1998). The substituted pyridyl groups and the phenyl ring lie in a propeller arrangement around the central 1,2,4-triazole ring. The dihedral angle between the phenyl and triazole rings is $59.30(7)^\circ$. The two pyridyl rings form dihedral angles of $47.90(8)$ and $26.91(8)^\circ$ with the triazole ring. In the crystal, the molecules exist as centrosymmetrically $C7-H7 \cdots N2(-x, 1-y, 1-z)$ hydrogen-bonded dimers (Table 2). The triazole and one of the pyridyl rings (N5) of the glide-related molecules translated a unit along the *b* axis are stacked along the *c* direction, with a perpendicular distance of $3.301(2)$ Å. The possibility of π - π stacking interactions involving these rings is reduced as these planes are inclined at an angle of 27° . However, these molecules are linked by weak $C12-H12 \cdots N4(x, \frac{3}{2}-y, z-\frac{1}{2})$ hydrogen bonds to form an infinite chain along the *c* axis. This arrangement is further stabilized by $C-H \cdots \pi$ interactions involving (i) H14 and the phenyl ring at $(x, \frac{3}{2}-y, z+\frac{1}{2})$, and (ii) H18 and a pyridyl ring (N5) at $(x, \frac{3}{2}-y, z-\frac{1}{2})$. The geometry of these interactions are listed in Table 2.




Fig. 1. The structure of (I) showing 50% probability displacement ellipsoids and the atom-numbering scheme.

$z - \frac{1}{2}$) hydrogen bonds to form an infinite chain along the *c* axis. This arrangement is further stabilized by $C-H \cdots \pi$ interactions involving (i) H14 and the phenyl ring at $(x, \frac{3}{2}-y, z+\frac{1}{2})$, and (ii) H18 and a pyridyl ring (N5) at $(x, \frac{3}{2}-y, z-\frac{1}{2})$. The geometry of these interactions are listed in Table 2.

Experimental

The title compound was synthesized by the reaction of equivalent amounts of 4,4'-methoxyphenylphosphazone and *N,N'*-dipyridylhydrazine in *N,N*-dimethylaniline for 3 h at 463–473 K (Grimmel *et al.*, 1946; Klingsberg, 1958). Single crystals suitable for X-ray diffraction analysis were obtained from acetone.

Crystal data

$C_{19}H_{15}N_5O$
 $M_r = 329.36$
 Monoclinic
 $P2_1/c$
 $a = 11.5436(4)$ Å
 $b = 18.2820(6)$ Å
 $c = 8.2935(3)$ Å
 $\beta = 107.141(1)^\circ$
 $V = 1672.5(1)$ Å³
 $Z = 4$
 $D_x = 1.308$ Mg m⁻³
 D_m not measured

Mo $K\alpha$ radiation
 $\lambda = 0.71073$ Å
 Cell parameters from 6971 reflections
 $\theta = 2.16$ – 38.60°
 $\mu = 0.086$ mm⁻¹
 $T = 293(2)$ K
 Plate
 $0.50 \times 0.32 \times 0.14$ mm
 Colourless

Data collection

Siemens SMART CCD area-detector diffractometer
 ω scans
 Absorption correction: none
 13 978 measured reflections
 4831 independent reflections
 3310 reflections with $I > 2\sigma(I)$

$R_{int} = 0.032$
 $\theta_{max} = 30^\circ$
 $h = -15 \rightarrow 15$
 $k = 0 \rightarrow 25$
 $l = 0 \rightarrow 11$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.120$
 $S = 1.029$
 4831 reflections
 287 parameters
 All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0457P)^2 + 0.2459P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} < 0.001$
 $\Delta\rho_{max} = 0.19$ e Å⁻³
 $\Delta\rho_{min} = -0.19$ e Å⁻³
 Extinction correction: *SHELXTL* (Sheldrick, 1997)
 Extinction coefficient: $0.017(2)$
 Scattering factors from *International Tables for Crystallography* (Vol. C)

Table 1. Selected geometric parameters (Å, °)

O1—C16	1.369 (2)	N2—N3	1.384 (2)
O1—C19	1.406 (2)	N3—C2	1.316 (2)
N1—C1	1.369 (2)	N4—C4	1.336 (2)
N1—C2	1.373 (2)	N4—C3	1.341 (2)
N1—C13	1.442 (1)	N5—C12	1.335 (2)
N2—C1	1.311 (2)	N5—C8	1.339 (2)
N2—C1—C3—C7	48.1 (2)	C1—N1—C13—C18	57.1 (2)
N3—C2—C8—C9	26.8 (2)		

Table 2. Hydrogen-bonding geometry (Å, °)

D—H...A	D—H	H...A	D...A	D—H...A
C7—H7...N2 ⁱ	0.94 (2)	2.56 (2)	3.318 (2)	137 (1)
C12—H12...N4 ⁱⁱ	1.01 (2)	2.60 (2)	3.382 (2)	134 (2)
C14—H14... π (phenyl ⁱⁱⁱ)	0.98 (2)	3.06 (2)	3.876 (2)	141 (2)
C18—H18... π (pyridyl ⁱⁱ)	0.99 (2)	2.87 (2)	3.682 (2)	140 (2)

Symmetry codes: (i) $-x, 1-y, 1-z$; (ii) $x, \frac{1}{2}-y, z-\frac{1}{2}$; (iii) $x, \frac{1}{2}-y, \frac{1}{2}+z$.

The data collection covered a hemisphere of reciprocal space by a combination of three sets of exposures; each set had a different φ angle (0, 88 and 180°) for the crystal and each exposure of 30 s covered 0.3° in ω . The crystal-to-detector distance was 3 cm and the detector swing angle was -40° . Crystal decay was monitored by repeating 30 initial frames at the end of data collection and analysing the intensity of duplicate reflections, and was found to be negligible.

The structure was solved by direct methods and refined by full-matrix least-squares techniques. All H atoms were located from a difference map and refined isotropically.

Data collection: *SMART* (Siemens, 1996). Cell refinement: *SAINT* (Siemens, 1996). Data reduction: *SAINT*. Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1997). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 1990).

This work was financially supported by the State Science and Technology Commission and the National Nature Science Foundation of China. The authors would like to thank the Malaysian Government and Universiti Sains Malaysia for research grant R & D No. 190-9609-2801. KC thanks the Universiti Sains Malaysia for a Visiting Postdoctoral Fellowship.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK1269). Services for accessing these data are described at the back of the journal.

References

- Chen, W., Wang, Z. X., Jian, F. F., Bai, Z. P. & You, X. Z. (1998). *Acta Cryst. C* **54**, 851–852.
- Cornelissen, J. P., Diemen, J. H., Groeneveld, L. R., Haasnoot, J. G., Spek, A. L. & Reedijk, J. (1992). *Inorg. Chem.* **31**, 198–202.
- Garcia, Y., Koningsbruggen, P. J., Codjovi, E., Lapouyade, R., Kahn, O. & Rabardel, L. (1997). *J. Mater. Chem.* **7**, 857–858.
- Grimmel, H. W., Guenther, A. & Morgan, J. F. (1946). *J. Am. Chem. Soc.* **68**, 539–542.
- Gupta, A. K. & Bhargava, K. P. (1978). *Pharmazie*, **23**, 430–431.
- Kahn, O. & Martinez, C. J. (1998). *Science*, **279**, 44–48.
- Klingsberg, E. (1958). *J. Org. Chem.* **23**, 1086–1087.
- Kunkeler, P. J., Koningsbruggen, P. J., Cornelissen, J. P., Horst, A. N., Kraan, A. M., Spek, A. L., Haasnoot, J. G. & Reedijk, J. (1996). *J. Am. Chem. Soc.* **118**, 2190–2197.
- Sheldrick, G. M. (1997). *SHELXTL. Structure Determination Software Programs*. Version 5.10. Bruker Analytical X-ray Systems Inc., Madison, Wisconsin, USA.
- Siemens (1996). *SMART and SAINT. Area Detector Control and Integration Software*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Spek, A. L. (1990). *Acta Cryst. A* **46**, C-34.
- Wang, Z. X., Bai, Z. P., Yang, J. X., Okamoto, K. I. & You, X. Z. (1998). *Acta Cryst. C* **54**, 438–439.

Acta Cryst. (1999). **C55**, 772–774

2-Ethyl-2-(3-oxobutyl)cycloheptane-1,3-dione

KANDASAMY CHINNAKALI,^{a†} HOONG-KUN FUN,^b DESIKAN RAJAGOPAL^c AND SAMBASIVAM SWAMINATHAN^c

^aDepartment of Physics, Anna University, Chennai 600 025, India, ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^cDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India. E-mail: hkfun@usm.my


(Received 26 November 1998; accepted 22 December 1998)

Abstract

In the title compound, C₁₃H₂₀O₃, the cycloheptane ring adopts a twist-chair conformation. The molecules are packed as layers parallel to the (202) plane. The planar oxobutyl group forms a dihedral angle of 48.3 (1)° with the best plane through the cycloheptane ring.

Comment

The title compound, (I), is the first solid prochiral triketone system in carbocyclic chemistry. It serves as an important precursor for the synthesis of homo-Wieland Miescher ketone which is a potential building block for the synthesis of terpenes (Ireland & Aristoff, 1979) and other natural-product intermediates. It can also be used for the study of oxy-cope rearrangement (Selvarajan, 1966) and synthesis of optically active carbocyclic intermediates (Rajagopal & Swaminathan, 1998). The X-ray structure determination of the compound was carried out to study the molecular conformation.



Some of the bond lengths and intra-annular valence angles in the cycloheptane ring that deviate significantly from the reported mean values (Allen *et al.*, 1993) may be due to the presence of the extended oxobutyl substituent at C1, and Csp² atoms C2 and C7. The cycloheptane ring adopts a twist-chair conformation with a twofold axis running through C1 and the mid-point of the C4—C5 bond. The Cremer & Pople puckering parameters (Cremer & Pople, 1975) are $q_2 = 0.540$ (2) Å,

† Visiting Post-Doctoral Fellow, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia.