

Asymmetric Catalysis Hot Paper

How to cite:

International Edition: doi.org/10.1002/anie.202010484 German Edition: doi.org/10.1002/ange.202010484

## **Dynamic Kinetic Resolution of Alcohols by Enantioselective Silylation Enabled by Two Orthogonal Transition-Metal Catalysts**

Jan Seliger and Martin Oestreich\*

Abstract: A nonenzymatic dynamic kinetic resolution of acyclic and cyclic benzylic alcohols is reported. The approach merges rapid transition-metal-catalyzed alcohol racemization and enantioselective Cu-H-catalyzed dehydrogenative Si-O coupling of alcohols and hydrosilanes. The catalytic processes are orthogonal, and the racemization catalyst does not promote any background reactions such as the racemization of the silyl ether and its unselective formation. Often-used ruthenium halfsandwich complexes are not suitable but a bifunctional ruthenium pincer complex perfectly fulfills this purpose. By this, enantioselective silylation of racemic alcohol mixtures is achieved in high yields and with good levels of enantioselection.

**D**ynamic kinetic resolution (DKR) is a powerful tool for the preparation of enantiomerically enriched chiral alcohols.<sup>[1]</sup> As an advantage over conventional kinetic resolution (KR) processes, the theoretical limit of 50% yield for each enantiomer is overcome by rapid in situ racemization of the starting material. Tracing back to an early example by Williams and co-workers,<sup>[2]</sup> this is typically achieved by catalytic transfer hydrogenation with only a few exceptions.<sup>[1a,3,4]</sup> A particularly versatile racemization catalyst is ruthenium complex 1 developed by Bäckvall and co-workers.<sup>[3a]</sup> In combination with Candida antarctica lipase B and isopropenyl acetate as the acyl source, efficient chemoenzymatic DKR of a number of structurally unbiased secondary alcohols was achieved (Scheme 1, top). Later, Fu and coworkers succeeded in developing a DKR of secondary alcohols using Bäckvall's complex 1 and a ferrocene-based, planar chiral DMAP derivative as a nucleophilic activator towards an acyl source (Scheme 1, middle).<sup>[5]</sup> This is the only nonenzymatic approach to date.

The acyl group introduced by these DKRs is usually cleaved to liberate the free alcohol or occasionally used further as a protecting group. Silyl ethers excel at the protection of hydroxy groups owing to the ease of their installation and removal as well as their tuneable stability and orthogonality.<sup>[6]</sup> Unlike the numerous (D)KR processes by enzymatic and nonenzymatic enantioselective acylation,<sup>[1,5]</sup> corresponding silylation-based methods had been unknown

[*]	J. Seliger, Prof. Dr. M. Oestreich
	Institut für Chemie, Technische Universität Berlin
	Strasse des 17. Juni 115, 10623 Berlin (Germany)
	E-mail: martin.oestreich@tu-berlin.de
	Homepage: http://www.organometallics.tu-berlin.de
	Supporting information and the ORCID identification number(s) for
Ā	the author(s) of this article can be found under:

https://doi.org/10.1002/anie.202010484.

Angew. Chem. Int. Ed. 2020, 59, 1-6

until approximately twenty years ago.<sup>[7]</sup> Starting from an initial report by Ishikawa,<sup>[8]</sup> asymmetric alcohol silylation has undergone tremendous advances.<sup>[7]</sup> Today, both organocatalytic<sup>[9]</sup> and transition-metal-catalyzed<sup>[10]</sup> approaches for the KR of alcohols are available. Conversely, suitable conditions for a related DKR have remained elusive (Scheme 1, bottom). For this, the resolving system and the racemization catalyst must be compatible, and racemization must occur significantly faster than the conversion of the slow-reacting enantiomer to the product silyl ether. Additionally, the racemization catalyst must not promote any racemic background reactions, that is the racemization of the silyl ether and the coupling of the alcohol and the silylation reagent.

Our study commenced with an investigation of several ruthenium and osmium complexes as potential racemization catalysts with (S)-1-phenylethanol [(S)-2a] as the model substrate and benzene- $d_6$  as the solvent (Figure 1 and Table 1). As expected, fast racemization of (S)-2a was seen using ruthenium complex 1 as the precatalyst and NaOtBu as the base (entry 1). The situation changed dramatically under our established KR conditions. The racemization was substantially slowed down in the presence of 6.0 mol% of (R,R)-Ph-BPE and stopped almost entirely when 1.3 equiv of

Chemoenzymatic DKR by enantioselective acylation (Bäckvall 2005, Ref. [3a])



Nonenzymatic DKR by enantioselective acylation (Fu 2012, Ref. [5])



Nonenzymatic DKR by enantioselective silylation (this work)



**Scheme 1.** Representative DKRs of alcohols by acylation (top) and planned silylation approach (bottom). DMAP=4-dimethylaminopyridine, (R,R)-Ph-BPE=1,2-bis[(2R,5R)-2,5-diphenylphospholan-1-yl]ethane.

© 2020 Wiley-VCH GmbH

### Wiley Online Library These are not the final page numbers!

Communications



Figure 1. Candidates for catalytic alcohol racemization.

Table 1: Identification of a suitable racemization catalyst.<sup>[a]</sup>

∧ ↓	<i>w/o coj</i> catalys NaO <i>t</i> B ( <i>R</i> , <i>R</i> )-Ph-I <i>n</i> Bu <sub>3</sub> Si	oper catalyst it (5.0 mol%) u (5.0 mol%) BPE (6.0 mol%) H (1.3 equiv)	он		o A
	/le bei RT t	nzene-d <sub>6</sub> for 30 min	Me	+	Me
(S)- <b>2a</b>			<i>rac-<b>2a</b> &gt;95</i>	:	<5 <sup>[b]</sup>
Entry	Catalyst	Variation			ee [%] <sup>[c</sup>
1	1	w/o (R,R)-	Ph-BPE, <i>n</i> Bu₃S	iН	0
2	1	w/o nBu₃S	SiH		32
3	1	w/o ( <i>R</i> , <i>R</i> )-	Ph-BPE		93
4	1	none			95
5	3 a	none			93
6	3 b	none			94
7	3 c	none			90
8	4a	none			0
9	4 b	none			0
10	5	none			0

[a] All reactions were performed on a 0.1 mmol scale. [b] Estimated by in situ <sup>1</sup>H NMR analysis; ratio based on the integration of baseline-separated resonance signals. [c] Determined by HPLC analysis on a chiral stationary phase.

*n*Bu<sub>3</sub>SiH were added (entries 2–4).<sup>[12]</sup> A similar result was obtained with the structurally related, more robust ruthenium complex 3a where one carbonyl ligand is replaced by triphenylphosphine (entry 5).<sup>[3b]</sup> The electronic modification of this complex using electron-poorer and -richer phosphines as in **3b** and **3c** had no effect (entries 6 and 7). We then turned our attention towards bifunctional catalysts bearing a metalbound amino group that have been employed in transfer hydrogenation with great success.<sup>[13]</sup> In contrast to the halfsandwich complexes 1 and 3a-c operating through innersphere mechanisms,<sup>[14]</sup> these catalysts likely operate through outer-sphere mechanisms involving hydrogen-bonding networks between the substrate and the NH<sub>2</sub> functionality as a crucial feature.<sup>[15]</sup> In recent years, transition-metal pincer complexes have emerged as highly active and robust catalysts, with [M(CNN)(dppb)Cl] 4a (M = Ru) and 4b (M = Os) as well as [Ru(PNP)(CO)HCl] 5 achieving particularly high turnover frequencies.<sup>[16]</sup> Also, ruthenium complex 5 was reported to facilitate the catalytic hydrogenolysis of chlorosilanes to produce hydrosilanes,<sup>[17]</sup> thereby making catalyst poisoning under our setup unlikely. Indeed, fast racemization of (S)-1-phenylethanol [(S)-2a] in the presence of both (R,R)- Ph-BPE and  $nBu_3SiH$  was accomplished with all of the bifunctional catalysts **4a**, **4b**, and **5** (entries 8–10) and merely trace amounts of acetophenone were observed.

With three promising candidates as racemization catalysts in hand, we next investigated their activity in the aforementioned racemic background reactions (Scheme 2). Importantly, none of the pincer complexes catalyzed the racemization of the silyl ether (S)-6a in basic medium (Scheme 2, top). Additionally, both the Ru- and the Os-CNN complexes 4a and 4b did not promote the dehydrogenative coupling of 1phenylethanol (2a) and  $nBu_3SiH$  (Scheme 2, bottom). Conversely, when Ru-PNP complex 5 was employed as the precatalyst, substantial amounts of silyl ether 6a were formed within 14 h, therefore thwarting its suitability for our endeavor. Without any obvious differences in catalytic activity between 4a and 4b, we continued with Ru-CNN complex 4a as the designated racemization precatalyst.

We then tried the DKR of racemic 1-phenylethanol (2a) in the presence of a copper catalyst, employing ruthenium complex 4a and our established resolving system consisting of CuCl, NaOtBu, (R,R)-Ph-BPE, and  $nBu_3SiH$  in toluene. With 1.1 equiv of the hydrosilane, quantitative conversion of the alcohol was achieved after stirring for 14 h at room temperature, and the corresponding silyl ether (S)-6a was isolated with 86% *ee.* It is worth noting that the reaction also proceeded smoothly in polar protic solvents such as *tert*-amyl alcohol with unchanged reactivity and enantioselectivity (see the Supporting Information for details). However, reducing the reaction temperature to 0°C did not improve the enantiomeric excess although good reactivity was maintained.

With suitable conditions established, we explored the scope of this DKR (Scheme 3). Various sterically and electronically modified benzylic alcohols  $2\mathbf{a}-\mathbf{z}$  were tested, and the corresponding silyl ethers (*S*)- $6\mathbf{a}-\mathbf{y}$  were generally obtained in almost quantitative yields with good to high levels of enantioselection. Both electron-donating (Me and OMe) and electron-withdrawing substituents (aryl, heteroaryl, halogens, CF<sub>3</sub>, and CO<sub>2</sub>/Bu) were well tolerated but a trend regarding the reactivity or selectivity was not apparent. A detailed survey of the aryl group's steric effects revealed that monosubstitution of the parent 1-phenylethanol ( $2\mathbf{a}$ ) with

#### Potential racemization of the silyl ether



rac-6a

**Scheme 2.** Interrogation of potential background reactions. All reactions were performed on a 0.2 mmol scale. Enantiomeric excesses were determined by HPLC analysis on a chiral stationary phase after cleavage of the silyl ether. Conversions were determined by GLC analysis using tetracosane as an internal standard.

#### www.angewandte.org

© 2020 Wiley-VCH GmbH

rac-2a

**K** These are not the final page numbers!



**Scheme 3.** DKR of acyclic benzylic alcohols. All reactions were performed on a 0.2 mmol scale. Yields are isolated yields after flash chromatography on silica gel. Enantiomeric excesses were determined by HPLC analysis on chiral stationary phases after cleavage of the silyl ether. [a] The reaction time was 48 h. [b] Estimated enantiomeric excess since no baseline separation was achieved. [c] The reaction time was 36 h. [d] The reaction time was 84 h. [e] The reaction time was 60 h. Boc = tert-butoxycarbonyl.

methyl groups in the *ortho-, meta-* and *para-*positions as in **2b**, **2d**, and **2f** was compatible with only little influence on the enantiomeric excess. However, when a bulkier bromine atom was installed in either of these positions a more pronounced effect was observed. Here, *ortho-*substituted (*S*)-**6c** was obtained with considerably lower *ee*, pointing towards a possible limitation of the racemization catalyst **4a**. Conversely, the enantioselectivities observed for *meta-* and *para*substituted derivatives (*S*)-**6e** and (*S*)-**6j** were in the expected range, yet noticeably higher in the former case. Accordingly, disubstitution in the *meta-*positions led to the silyl ethers (*S*)-**6n-p** with improved enantioselectivities while *ortho,ortho*dimethylated (*S*)-**6q** was isolated with drastically reduced enantiomeric excess and after prolonged reaction time. The latter result stands in stark contrast to the outstanding selectivity factor of 170 observed in the KR of 2q with the same copper catalyst.<sup>[10e]</sup> Consequently, steric bulk in the ortho-position exerts a strong effect on the racemization ability of catalyst **4a**. In line with these findings, both  $\alpha$ - and  $\beta$ naphthyl-substituted derivatives 2r and 2s as well as alcohols bearing heterocyclic furan-2-yl and thien-2-yl units as in 2t and 2u exhibited high reactivity and good enantioselectivity. Although 1-(pyridin-4-yl)ethanol (2v) proved substantially less reactive, the enantiodifferentiation stayed in the same range. As expected from our previous work,<sup>[10e]</sup> both the enantiomeric excess and the reactivity dwindled with increasing size of the alkyl group Me < Et < Bn  $\ll i$ Pr [2a and 2w- $\mathbf{y} \rightarrow (S)$ -6a and (S)-6w–y], and synthetically useful 2-chloro-1-phenylethanol (2z) was entirely unreactive. This trend was further corroborated in an application to 7, a precursor of duloxetine (gray box, left). The thien-2-yl unit had been shown to be compatible before [cf.  $2\mathbf{u} \rightarrow (S)$ - $6\mathbf{u}$ ] but the ethylene-tethered, Boc-protected amino group was detrimental, affording (S)-9 with moderate 62% ee; the coordination ability of the Boc group could however contribute to this outcome. In another application, carbamate-containing 8 participated in the DKR with good efficiency to yield the rivastigmine precursor (S)-10 in 93% ee (gray box, right).

Selected aliphatic alcohols were also included into this study (Figure 2), especially to compare 1-phenylethanol (2a) with fully saturated 1-cyclohexylethanol. The corresponding silyl ether (S)-11a (79% *ee*) did form with slightly diminished *ee* compared to (S)-6a (86% *ee*), and the reaction had a significantly smaller rate (days versus hours). Both substitution of the cyclohexyl moiety with a cyclopentyl group (as in (S)-11b) and slight chain elongation (i.e. Me  $\rightarrow$  Et as in (S)-11c) paired low reactivity with further deteriorated enantioselectivity, and full conversion was not even reached in the latter case.

Finally, we investigated cyclic benzylic alcohols 12a-h with different ring sizes as substrates for our reaction (Scheme 4). Again, yields were nearly quantitative and enantioselectivities were usually slightly better than for their acyclic counterparts. Hence, five- to seven-membered carbocyclic alcohols 12a-c smoothly underwent the reaction, although the benzosuberol-derived silyl ether (S)-13c was obtained with notably lower *ee.* Substitution with a heteroa-



**Figure 2.** DKR of aliphatic alcohols. All reactions were performed on a 0.2 mmol scale. Yields are isolated yields after flash chromatography on silica gel. Enantiomeric excesses were determined by HPLC analysis on a chiral stationary phase after cleavage of the silyl ether and derivatization as the *p*-nitrobenzoate. [a] The reaction time was 84 h. [b] The reaction time was 120 h after which 65% conversion of 1cyclohexylpropanol was detected by GLC analysis using tetracosane as an internal standard. [c] Estimated enantiomeric excess since no baseline separation was achieved. Communications



**Scheme 4.** DKR of cyclic benzylic alcohols. All reactions were performed on a 0.2 mmol scale. Yields are isolated yields after flash chromatography on silica gel. Enantiomeric excesses were determined by HPLC analysis on chiral stationary phases after cleavage of the silyl ether.

tom in the saturated ring was also tolerated, and the highly functionalized (thio)chromanol-derived silyl ethers (S)-13d-f and (S)-13g were amenable with high *ee*. Furthermore, the position of the heteroatom had no significant effect on the selectivity of our reaction as evident from the tetrahydro-(iso)quinolinol-derived silyl ethers (S)-13h and (R)-14 (gray box).

DKR of secondary alcohols is typically achieved by enantioselective acylation techniques for which several chemoenzymatic protocols are available.<sup>[1a]</sup> Fu's nonenzymatic process had remained an isolated example<sup>[5]</sup> and is also a beautiful example of orthogonal tandem catalysis.<sup>[18]</sup> A cognate tool relying on the stereoselective silylation of alcohols has been unprecedented so far. We reported here a DKR of secondary alcohols by means of a Cu-H-catalyzed dehydrogenative coupling with the simple hydrosilane  $nBu_3SiH$ . The choice of the racemization catalyst is crucial as commonly used ruthenium half-sandwich complexes fail under our previously established, commercially available resolving system.<sup>[10e]</sup> In turn, bifunctional ruthenium and osmium pincer complexes were identified as suitable alternatives and were successfully combined with our KR procedure.<sup>[10e]</sup> Our method is applicable to a broad range of acyclic and cyclic benzylic alcohols and exhibits good functionalgroup tolerance. An extension of this approach to allylic<sup>[10e, 19]</sup> and propargylic<sup>[10f]</sup> alcohols will require further catalyst design and is currently under investigation in our laboratory.

#### Acknowledgements

This research was supported by the Deutsche Forschungsgemeinschaft (Oe 249/14-1) and the Fonds der Chemischen Industrie (predoctoral fellowship to J.S., 2018–2020). M.O. is indebted to the Einstein Foundation Berlin for an endowed professorship. We thank Takuya Kinoshita of Osaka University for his experimental contributions. Open access funding enabled and organized by Projekt DEAL.

#### Conflict of interest

J.S. and M.O. filed a patent application of the reported method.

**Keywords:** asymmetric catalysis · copper · dehydrogenative coupling · dynamic kinetic resolution · silicon

- For reviews of the (D)KR of alcohols by enzymatic and nonenzymatic acylation, see: a) O. Verho, J.-E. Bäckvall, J. Am. Chem. Soc. 2015, 137, 3996-4009; b) E. Vedejs, M. Jure, Angew. Chem. Int. Ed. 2005, 44, 3974-4001; Angew. Chem. 2005, 117, 4040-4069; c) J. M. Keith, J. F. Larrow, E. N. Jacobsen, Adv. Synth. Catal. 2001, 343, 5-26.
- [2] P. M. Dinh, J. A. Howarth, A. R. Hudnott, J. M. J. Williams, W. Harris, *Tetrahedron Lett.* **1996**, *37*, 7623–7626.
- Selected examples of alcohol racemization catalysts operating by transfer hydrogenation mechanisms. Ruthenium(II)-based: a) B. Martín-Matute, M. Edin, K. Bogár, F. B. Kaynak, J.-E. Bäckvall, J. Am. Chem. Soc. 2005, 127, 8817-8825; b) S.-B. Ko, B. Baburaj, M.-J. Kim, J. Park, J. Org. Chem. 2007, 72, 6860-6864; c) G. Bossi, E. Putignano, P. Rigo, W. Baratta, Dalton Trans. 2011, 40, 8986-8995; d) J. A. Fernández-Salas, S. Manzini, S. P. Nolan, Chem. Eur. J. 2014, 20, 13132-13135; aluminum(III)-based: e) A. Berkessel, M. L. Sebastian-Ibarz, T. N. Müller, Angew. Chem. Int. Ed. 2006, 45, 6567-6570; Angew. Chem. 2006, 118, 6717-6720; iridium(III)-based: f) R. M. Haak, F. Berthiol, T. Jerphagnon, A. J. A. Gayet, C. Tarabiono, C. P. Postema, V. Ritleng, M. Pfeffer, D. B. Janssen, A. J. Minnaard, B. L. Feringa, J. G. de Vries, J. Am. Chem. Soc. 2008, 130, 13508-13509; iron(II)-based: g) O. El-Sepelgy, N. Alandini, M. Rueping, Angew. Chem. Int. Ed. 2016, 55, 13602-13605; Angew. Chem. 2016, 128, 13800-13803.
- [4] Examples of alcohol racemization through dehydration-hydration mechanisms. Under zeolite catalysis: a) S. Wuyts, K. De Temmerman, D. De Vos, P. Jacobs, *Chem. Commun.* 2003, 1928–1929; b) Y. Zhu, K.-L. Fow, G.-K. Chuah, S. Jaenicke, *Chem. Eur. J.* 2007, *13*, 541–547; under vanadium(V) catalysis: c) M. Egi, K. Sugiyama, M. Saneto, R. Hanada, K. Kato, S. Akai, *Angew. Chem. Int. Ed.* 2013, *52*, 3654–3658; *Angew. Chem.* 2013, *125*, 3742–3746.
- [5] S. Y. Lee, J. M. Murphy, A. Ukai, G. C. Fu, J. Am. Chem. Soc. 2012, 134, 15149–15153.
- [6] a) P. G. M. Wuts, Greene's Protective Groups in Organic Synthesis, 5th ed., Wiley, Hoboken, 2014, pp. 201–270; b) P. J. Kocieńsky, Protecting Groups, 3rd ed., Thieme, Stuttgart, 2005, pp. 188–230.
- [7] a) A. H. Hoveyda, M. L. Snapper in Organosilicon Chemistry– Novel Approaches and Reactions (Eds.: T. Hiyama, M. Oestreich), Wiley-VCH, Weinheim, 2019, pp. 459–493; b) J. Seliger, M. Oestreich, Chem. Eur. J. 2019, 25, 9358–9365; c) L.-W. Xu, Y. Chen, Y. Lu, Angew. Chem. Int. Ed. 2015, 54, 9456–9466; Angew. Chem. 2015, 127, 9590–9601.

www.angewandte.org

© 2020 Wiley-VCH GmbH

These are not the final page numbers!

- [8] T. Isobe, K. Fukuda, Y. Araki, T. Ishikawa, *Chem. Commun.* 2001, 243–244.
- [9] Selected examples of KRs of alcohols through organocatalytic silylation: a) Y. Zhao, A. W. Mitra, A. H. Hoveyda, M. L. Snapper, Angew. Chem. Int. Ed. 2007, 46, 8471-8474; Angew. Chem. 2007, 119, 8623-8626; b) C. I. Sheppard, J. L. Taylor, S. L. Wiskur, Org. Lett. 2011, 13, 3794-3797; c) A. D. Worthy, X. Sun, K. L. Tan, J. Am. Chem. Soc. 2012, 134, 7321-7324; d) R. W. Clark, T. M. Deaton, Y. Zhang, M. I. Moore, S. L. Wiskur, Org. Lett. 2013, 15, 6132-6135; e) L. Wang, R. K. Akhani, S. L. Wiskur, Org. Lett. 2015, 17, 2408-2411; f) S. Y. Park, J.-W. Lee, C. E. Song, Nat. Commun. 2015, 6, 7512.
- [10] Selected examples of KRs of alcohols through transition-metalcatalyzed silylation: a) S. Rendler, G. Auer, M. Oestreich, *Angew. Chem. Int. Ed.* 2005, 44, 7620-7624; *Angew. Chem.* 2005, 117, 7793-7797; b) H. F. T. Klare, M. Oestreich, *Angew. Chem. Int. Ed.* 2007, 46, 9335-9338; *Angew. Chem.* 2007, 119, 9496-9499; c) B. Karatas, S. Rendler, R. Fröhlich, M. Oestreich, *Org. Biomol. Chem.* 2008, 6, 1435-1440; d) A. Weickgenannt, M. Mewald, T. W. T. Muesmann, M. Oestreich, *Angew. Chem. Int. Ed.* 2010, 49, 2223-2226; *Angew. Chem.* 2010, 122, 2269-2272; e) X. Dong, A. Weickgenannt, M. Oestreich, *Nat. Commun.* 2017, 8, 15547; f) J. Seliger, X. Dong, M. Oestreich, *Angew. Chem. Int. Ed.* 2019, 58, 1970-1974; *Angew. Chem.* 2019, 131, 1991-1996.
- [11] X. Dong, PhD thesis, Technische Universität Berlin, Berlin, 2018.
- [12] In a separate experiment, we observed the formation of a Ru-H species by <sup>1</sup>H NMR spectroscopy that failed to deprotonate (*S*)-1-phenylethanol [(*S*)-**2a**]. This observation is in good agreement with studies conducted by Bäckvall and co-workers regarding

the racemization mechanism of **1**. These authors found that  $C_5Ph_5(CO)_2RuH$  was completely inactive in the racemization of (*S*)-**2a** and therefore excluded this species as a relevant intermediate. See Ref. [3a] for details.

- [13] D. Wang, D. Astruc, Chem. Rev. 2015, 115, 6621-6686.
- [14] M. C. Warner, J.-E. Bäckvall, Acc. Chem. Res. 2013, 46, 2545– 2555.
- [15] P. A. Dub, J. C. Gordon, Nat. Rev. Chem. 2018, 2, 396-408.
- [16] a) W. Baratta, G. Chelucci, S. Gladiali, K. Siega, M. Toniutti, M. Zanette, E. Zangrando, P. Rigo, *Angew. Chem. Int. Ed.* 2005, *44*, 6214–6219; *Angew. Chem.* 2005, *117*, 6370–6375; b) W. Baratta, M. Ballico, G. Chelucci, K. Siega, P. Rigo, *Angew. Chem. Int. Ed.* 2008, *47*, 4362–4365; *Angew. Chem.* 2008, *120*, 4434–4437; c) M. Bertoli, A. Choualeb, A. J. Lough, B. Moore, D. Spasyuk, D. G. Gusev, *Organometallics* 2011, *30*, 3479–3482.
- [17] A. Glüer, J. I. Schweizer, U. S. Karaca, C. Würtele, M. Diefenbach, M. C. Holthausen, S. Schneider, *Inorg. Chem.* 2018, 57, 13822–13828.
- [18] For reviews on tandem catalysis, see: a) T. L. Lohr, T. J. Marks, *Nat. Chem.* 2015, *7*, 477–482; b) J.-C. Wasilke, S. J. Obrey, R. T. Baker, G. C. Bazan, *Chem. Rev.* 2005, *105*, 1001–1020.
- [19] A complex mixture was obtained when (E)-4-phenylbut-3-en-2ol was subjected to the reaction conditions. Another experiment revealed that the allylic alcohol was isomerized to the corresponding ketone, 4-phenylbutan-2-one, within seconds in the presence of ruthenium complex **4a**.

Manuscript received: July 31, 2020

Revised manuscript received: September 28, 2020

Version of record online:



## Communications

# Asymmetric Catalysis

Dynamic Kinetic Resolution of Alcohols by Enantioselective Silylation Enabled by Two Orthogonal Transition-Metal Catalysts



The combination of a chiral copper catalyst and a bifunctional ruthenium pincer complex enables the nonenzymatic dynamic kinetic resolution of acyclic and cyclic benzylic alcohols. The enantioselective Cu-H-catalyzed dehydrogenative Si-O coupling and the rapid transitionmetal-catalyzed alcohol racemization are perfectly orthogonal. By this, high yields and good levels of enantioselection are achieved.

6 www.angewandte.org
These are not the final page numbers!

© 2020 Wiley-VCH GmbH