

A Journal of the Gesellschaft Deutscher Chemiker

# Angewandte Chemie

GDCh

International Edition

www.angewandte.org

## Accepted Article

**Title:** Ni-Catalyzed Regioselective Alkenylarylation of  $\gamma,\delta$ -Alkenyl Ketones via Carbonyl Coordination

**Authors:** Ramesh Giri, Roshan K Dhungana, Vivek Aryal, Doleshwar Niroula, Rishi R Sapkota, and Margaret G Lakomy

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Angew. Chem. Int. Ed.* 10.1002/anie.202104871

**Link to VoR:** <https://doi.org/10.1002/anie.202104871>


## COMMUNICATION

Ni-Catalyzed Regioselective Alkenylarylation of  $\gamma,\delta$ -Alkenyl Ketones via Carbonyl Coordination

Roshan K. Dhungana, Vivek Aryal, Doleshwar Niroula, Rishi R. Sapkota, Margaret G. Lakomy, and Ramesh Giri\*

**Abstract:** We disclose a nickel-catalyzed reaction, which enabled us to difunctionalize unactivated  $\gamma,\delta$ -alkenes in ketones with alkenyl triflates and arylboronic esters. The reaction was made feasible by the use of 5-chloro-8-hydroxyquinoline as a ligand along with  $\text{NiBr}_2\cdot\text{DME}$  as a catalyst and  $\text{LiOtBu}$  as base. The reaction proceeded with a wide range of cyclic, acyclic, endocyclic and exocyclic alkenyl ketones, and electron-rich and electron-deficient arylboronate esters. The reaction also worked with both cyclic and acyclic alkenyl triflates. Control experiments indicate that carbonyl coordination is required for the reaction to proceed.

Metal-catalyzed alkene dicarbofunctionalization<sup>[1]</sup> is an emerging synthetic method for complex molecule synthesis<sup>[2]</sup> with a promise to reduce a multistep process to a few-step process.<sup>[3]</sup> This reaction integrates two carbon sources into alkenes with high regiofidelity, and provides branched molecules in one step.<sup>[4]</sup> However, its development has been met with formidable challenges largely due to the formation of Heck products by  $\beta$ -H elimination (Scheme 1).<sup>[5]</sup> The reaction selectivity toward alkene difunctionalization and the Heck product depend on the difference in kinetic barriers between a single-step  $\beta$ -H elimination and a two-step transmetalation/reductive elimination process. In the majority of catalytic processes,  $\beta$ -H elimination has been shown to occur with a much lower kinetic barrier than both the transmetalation and reductive elimination steps.<sup>[6]</sup>



**Scheme 1.** Alkene 1,2-dicarbofunctionalization and complications by  $\beta$ -H elimination

[\*] Prof. R. Giri, Mr. R. K. Dhungana, Mr. Vivek Aryal, Dr. Doleshwar Niroula, Dr. Rishi R. Sapkota, Ms. Margaret G. Lakomy  
Department of Chemistry  
Pennsylvania State University,  
University Park, Pennsylvania 16802, USA  
E-mail: rkg5374@psu.edu


Supporting information for this article and crystallographic data (CCDC number: 2059802) for compound **33** are given via a link at the end of the document.

Early work focused on difunctionalizing 1,1-disubstituted terminal<sup>[7]</sup> and bicyclic<sup>[8]</sup> alkenes to avoid  $\beta$ -H elimination. Recently, a broad class of substrates were functionalized using a coordination approach<sup>[1a-f]</sup> in which alkylmetal intermediates were stabilized as transient metallacycles.<sup>[9]</sup> However, this strategy requires installation and removal of strong coordinating groups like imines,<sup>[10]</sup> pyridines,<sup>[11]</sup> 2-aminopyrimidines,<sup>[12]</sup> and 8-aminoquinolines<sup>[13]</sup> for stabilizing metallacycles.<sup>[14]</sup> Additionally, this strategy generally works with substrates that generate rigid and planar five-membered metallacycles. Limited examples of reactions involving six-membered metallacycles are known but they either require strongly coordinating bidentate groups<sup>[15]</sup> or are derived from vinylarenes containing activated alkenes<sup>[10]</sup> to planarize and rigidify metallacycles. Alkene difunctionalization reactions that involve nonplanar and fluxional six-membered metallacycles form rearranged products through contraction of metallacycles by  $\beta$ -H elimination.<sup>[5a]</sup>

We recently introduced a cationic catalysis<sup>[16]</sup> to suppress the contraction of metallacycles. While the reaction was successful, the cationization of organometallic intermediates is also known to promote  $\beta$ -H elimination.<sup>[6a, 6b, 17]</sup> As such, when weakly coordinating groups are used, the cationization of metallacycles promotes  $\beta$ -H elimination faster than the transmetalation/reductive elimination steps.<sup>[5b]</sup> For that reason, regioselective alkene dicarbofunctionalization in weakly coordinating ketone substrates are not known.<sup>[18]</sup> Herein, we disclose a new Ni-catalyzed reaction in the presence of 5-chloro-8-hydroxyquinoline that addresses the issue of  $\beta$ -H elimination, and materializes alkenylarylation of unactivated  $\gamma,\delta$ -alkenes in simple ketones. To the best of our knowledge, the current reaction represents the first example of keto carbonyl-assisted dicarbofunctionalization of unactivated alkenes.<sup>[19]</sup>

In our initial studies, we examined various ancillary ligands (**L1-L7**) for alkenylarylation of  $\gamma,\delta$ -alkene in 2-allylcyclohexanone (Scheme 2). We were pleased to discover that a combination of 20 mol % 8-hydroxyquinoline (**L7**) and 10 mol %  $\text{NiBr}_2\cdot\text{DME}$ , along with  $\text{LiOtBu}$  as a base, generated the alkenylarylation product **3** with cyclohexenyl triflate and phenylboronic ester **B3** in 48% yield. Control experiments conducted in the absence of **L7** and in the presence of **L8**, a neutral variant of **L7**, produced no product, confirming that the anionic ancillary ligand played a critical role in the reaction. Upon further evaluation (**L9-L12**), we found that 5-chloro-8-hydroxyquinoline (**L9**) performed optimally and furnished the product **3** in 75% yield. Examination of different arylboron reagents showed that phenylboronic ester **B3** afforded the product **3** in best yield. Replacing 2-allylcyclohexanone with 2-vinyl- and 2-butenylketones also did not furnish expected products (Scheme 2).

## COMMUNICATION



<sup>a</sup>Reactions run at 0.10 mmol scale in 0.5 mL THF. <sup>1</sup>H NMR yields with pyrene as a standard. <sup>b</sup>The yield of isolated product from a 0.50 mmol scale reaction in 2.5 mL THF in parenthesis. dr is 1.1:1.

**Scheme 2.** Examination of reaction parameters<sup>a</sup>

**Table 1.** Further reaction optimization<sup>a</sup>

entry	deviation in reaction condition	yields of <b>3</b> (%) <sup>b</sup>
1	none	75 (72) <sup>c</sup>
2	Lithium-1-adamantanolate (LiOAd) instead of LiOtBu	75
3	LiOMe instead of LiOtBu	23
4	LiOPr instead of LiOtBu	15
5	1 equiv LiOtBu instead of 1.5 equiv	42
6	2 equiv LiOtBu instead of 1.5 equiv	65
7	NiI <sub>2</sub> or NiCl <sub>2</sub> ·DME instead of NiBr <sub>2</sub> ·DME	70-72
8	Ni(TMHD) <sub>2</sub> instead of NiBr <sub>2</sub> ·DME	53
9	Ni(PPh <sub>3</sub> ) <sub>4</sub> instead of NiBr <sub>2</sub> ·DME	12
10	Ni(cod) <sub>2</sub> instead of NiBr <sub>2</sub> ·DME	52
11	other metals instead of NiBr <sub>2</sub> ·DME <sup>d</sup>	0
12	room temperature instead of 80 °C	10
13	60 °C instead of 80 °C	62
14	2-Me-THF instead of THF	70
15	1,4-dioxane instead of THF	60
16	MeCN or toluene instead of THF	35-38
17	NMP, DMF, DMSO or dioxane instead of THF	0-25

<sup>a</sup>Reactions run at 0.10 mmol scale in 0.5 mL solvent. <sup>b1</sup>H NMR yields with pyrene as a standard. <sup>c</sup>The yield of isolated product from a 0.50 mmol scale reaction in 2.5 mL THF in parenthesis. dr is 1.1:1. <sup>d</sup>Pd(OAc)<sub>2</sub>, CuI, Co(OAc)<sub>2</sub>, FeCl<sub>2</sub>.

Examination of different alkoxide bases indicated that the reaction was best performed with LiOtBu or LiOAd (Table 1, entries 1-6). NiBr<sub>2</sub>·DME could be replaced with NiCl<sub>2</sub>·DME or NiI<sub>2</sub> (entry 7). Other Ni-catalysts furnished the product in lower yields (entries 8-10) and other metal-catalysts such as FeCl<sub>2</sub>, Co(OAc)<sub>2</sub>, CuI and Pd(OAc)<sub>2</sub> generated no product at all (entry 11). The reaction was optimally performed at 80 °C in THF since other solvents and lower temperatures either decreased the yield or formed no product (entries 12-17).

After optimization of the reaction conditions, we explored the scope of the alkenylarylation reaction. Examination of different arylboronic esters indicated that the reaction tolerated both electron-rich and electron-deficient arenes, and the products were formed in good

yields with moderate diastereoselectivity (Table 2, A). The reaction was compatible with various substituents like Me, dioxanyl, OMe, OCF<sub>3</sub> and F, and transition metal-sensitive functional groups such as Br, nitrile and ester (**4-11**). This method can be applied to couple arylboronic esters having polyaromatic hydrocarbons including naphthyl, methoxynaphthyl and binaphthyl (**11-13**). The reaction also tolerates heterocyclic arylboronic esters, such as thiazolyl and carbazolylboronic esters (**14-15**), and alkenylboronic esters (**16**).

Moreover, we evaluated reaction scope with regard to  $\gamma,\delta$ -alkenyl ketones and alkenyl triflates (Table 2, B). Various  $\gamma,\delta$ -alkenyl endocyclic ketones containing both monocyclic and bicyclic scaffolds, such as cyclohexenonyl, tetralonyl and acenaphthylenonyl, could be implemented as substrates along with different alkenyl triflates and arylboronic esters (**17-22**). The reaction could also be performed with  $\gamma,\delta$ -alkenyl exocyclic ketones bearing both cyclic and acyclic aliphatic backbones such as cyclohexyl and isopropyl (**23-25**). Surprisingly,  $\gamma,\delta$ -alkenyl arylketones furnished products in low yields when the reactions were conducted in THF (20-30%). Unlike  $\gamma,\delta$ -alkenyl alkyketones for which THF was the solvent of choice at 80 °C,  $\gamma,\delta$ -alkenyl arylketones furnished products in best yields in MeCN at 60 °C.  $\gamma,\delta$ -Alkenyl arylketones with *p*-CF<sub>3</sub>, *o*-Me and *p*-Cl substituents and a naphthyl group afforded dicarbofunctionalized products in good yields with different alkenyl triflates and arylboronic esters (**26-33**). Reactions of these diverse  $\gamma,\delta$ -alkenyl ketones could be conducted with carbocyclic alkenyl triflates containing cyclopentenyl, cyclohexenyl, cyclooctenyl and 4-methylcyclohexenyl rings, and heterocyclic alkenyl triflates, such as 3,6-dihydro-2H-thiopyran-4-yl (**18**) and 3,6-dihydro-2H-pyran-4-yl triflates (**22** and **27**). The reaction condition was also amenable for difunctionalizing acyclic alkenyl triflates (**34-35**). These types of product derivatives form the cores of pharmaceutical molecules and natural products.<sup>[20]</sup>



We have further demonstrated synthetic applications by difunctionalizing unactivated alkenes in natural products, pharmaceuticals and complex spirocyclic molecules, and by using alkenyl triflates and arylboronic esters derived from complex molecules (Table 2, C). For example, an alkenyl triflate derived from a steroid, cholestenone, was incorporated into  $\gamma,\delta$ -alkene in cyclohexanone (**37**). In addition,  $\gamma,\delta$ -alkenes contained in complex molecules such as cholestenone, probenecid (a uricosuric drug) and naproxen (a NSAID drug) could also be difunctionalized with cyclohexenyl triflate and various arylboronic esters (**38-40**). Arylboronic ester derived from 9,9'-spirobifluorene also afforded products (**41-42**) with alkenylcyclohexanone and alkenylarylketone. Furthermore, we examined an internal  $\gamma,\delta$ -alkene contained in a spirobicyclic ketone, which reacted to furnish a *cis*-difunctionalized product (**43**).

Control experiments with an alkene lacking carbonyl group support carbonyl coordination during reaction (Scheme 3). Moreover, alkenyl ketones bearing less sterically bulky substituents on the carbonyl group predominantly produced alkenylarylated products **24** and **29** along with trace amounts of Heck products **44** and **45** (Scheme 4). In contrast, a sterically demanding *tert*-butyl group on the carbonyl group generated the Heck product **47** in significant amounts (22%) in addition to the alkenylarylation product **46** in 35% yield. These experiments indicate that carbonyl coordination to nickel is required to generate alkenylarylated products and that the Heck products are generated due to equilibrium between the cyclic and acyclic alkylnickel species **48** and **49** (Scheme 4).<sup>[21]</sup> Intimate analyses of regioselectivity in products indicated that the reaction should be catalyzed by a Ni(I) catalyst and transmetalation should precede the reaction of alkenyl triflates (see SI for details).<sup>[22]</sup>


## COMMUNICATION

Table 2. Reaction scope<sup>a</sup>

## A. scope with organoboron reagents

B. Scope with  $\gamma,\delta$ -alkenyl ketones and alkenyl triflates


## C. Scope with complex substrates derived from natural products, pharmaceuticals and materials






## COMMUNICATION

<sup>a</sup>Reactions run in 0.50 mmol scale in 2.5 mL solvent. Reactions for alkylketones (products **4-22**, **34-36**) run in THF at 80 °C while those for arylketones (products **26-33**, **39** and **42**) run in MeCN 60 °C for 6 h. <sup>b</sup>Reaction run under MeCN conditions. <sup>c</sup>Reactions run under THF conditions at 60 °C for 6 h. <sup>d</sup>20 mol % NiBr<sub>2</sub>·DME. Alkenyl triflate E/Z: 2:1. <sup>e</sup>A mixture of 4 diastereomers based on <sup>13</sup>C NMR (see SI for details).



**Scheme 3.** Carbonyl coordination effect



**Scheme 4.** Carbonyl steric effect

In summary, we report a nickel-catalyzed reaction, which addresses the issue of  $\beta$ -H elimination in alkene difunctionalization in carbonyl-assisted alkene dicarbofunctionalization reaction. The success of the reaction relied upon the use of a combination of 5-chloro-8-hydroxyquinoline and Ni(cod)<sub>2</sub> in the presence of LiOtBu. This catalysis enabled us to difunctionalize unactivated  $\gamma,\delta$ -alkenes in ketones with alkenyl triflates and arylboronic esters. The reaction proceeded with a wide range of cyclic, acyclic, endocyclic and exocyclic alkenyl ketones along with electron-rich and electron-deficient arylboronate esters, and cyclic and acyclic alkenyl triflates. Control experiments with a substrate lacking a carbonyl group and with a sterically hindered carbonyl group indicated that the carbonyl coordination was required for the reaction to proceed.

## Acknowledgements

We gratefully acknowledge the NIH NIGMS (R35GM133438) and The Pennsylvania State University for support of this work, and the PSU NMR facility for NMR support. We thank Dr. Christy George for help with stereochemistry determination of compound **43** by NMR.

**Keywords:** Alkenyl ketone • alkenylarylation • dicarbofunctionalization • nickel-catalyzed • 8-hydroxyquinoline

- [1] For reviews, see: a) R. K. Dhungana, S. KC, P. Basnet, R. Giri, *Chem. Rec.* **2018**, *18*, 1314-1340; b) R. Giri, S. KC, *J. Org. Chem.* **2018**, *83*, 3013-3022; c) J. Derosa, O. Apolinar, T. Kang, V. T. Tran, K. M. Engle, *Chem. Sci.* **2020**, *11*, 4287-4296; d) S. O. Badir, G. A. Molander, *Chem* **2020**, *6*, 1327-1339; e) H.-Y. Tu, S. Zhu, F.-L. Qing, L. Chu, *Synthesis* **2020**, *52*, 1346-1356; f) X. Qi, T. Diao, *ACS Catal.* **2020**, *10*, 8542-8556; g) A. Herath, W. Li, J. Montgomery, *J. Am. Chem. Soc.* **2008**, *130*, 469-471; h) J. Montgomery, *Acc. Chem. Res.* **2000**, *33*, 467-473.
- [2] K. C. Nicolaou, S. Rigol, *Nat. Prod. Rep.* **2020**.
- [3] a) S. R. Chemler, S. D. Karyakarte, Z. M. Khoder, *J. Org. Chem.* **2017**, *82*, 11311-11325; b) J. P. Wolfe, *Synlett* **2008**, *2008*, 2913-2937; c) W. Ahmed, S. Zhang, X. Yu, X. Feng, Y. Yamamoto, M. Bao, *Angew. Chem. Int. Ed.* **2019**, *58*, 2495-2499; d) J.-B. Peng, F.-P. Wu, D. Li, H.-Q. Geng, X. Qi, J. Ying, X.-F. Wu, *ACS Catal.* **2019**, *9*, 2977-2983.
- [4] a) T. Qin, J. Cornella, C. Li, L. R. Malins, J. T. Edwards, S. Kawamura, B. D. Maxwell, M. D. Eastgate, P. S. Baran, *Science* **2016**, *352*, 801-805; b) L. Zhang, G. J. Lovering, E. K. Edelstein, A. A. Szymaniak, M. P. Chierchia, J. P. Morken, *Science* **2016**, *351*, 70-74; c) M. Nakamura, A. Hirai, E. Nakamura, *J. Am. Chem. Soc.* **2000**, *122*, 978-979; d) S. R. Neufeldt, M. S. Sanford, *Org. Lett.* **2013**, *15*, 46-49.
- [5] For selected examples on three-component alkene dicarbofunctionalization, see: a) P. Basnet, R. K. Dhungana, S. Thapa, B. Shrestha, S. KC, J. M. Sears, R. Giri, *J. Am. Chem. Soc.* **2018**, *140*, 7782-7786; b) R. K. Dhungana, S. KC, P. Basnet, V. Aryal, L. J. Chesley, R. Giri, *ACS Catal.* **2019**, *9*, 10887-10893; c) J. Jeon, H. Ryu, C. Lee, D. Cho, M.-H. Baik, S. Hong, *J. Am. Chem. Soc.* **2019**, *141*, 10048-10059; d) L. Liao, R. Jana, K. B. Urkalan, M. S. Sigman, *J. Am. Chem. Soc.* **2011**, *133*, 5784-5787; e) V. Saini, M. S. Sigman, *J. Am. Chem. Soc.* **2012**, *134*, 11372-11375; f) M. Orlandi, M. J. Hilton, E. Yamamoto, F. D. Toste, M. S. Sigman, *J. Am. Chem. Soc.* **2017**, *139*, 12688-12695; g) H.-M. Huang, P. Bellotti, P. M. Pflüger, J. L. Schwarz, B. Heidrich, F. Glorius, *J. Am. Chem. Soc.* **2020**, *142*, 10173-10183; h) Y. Li, H. Wei, D. Wu, Z. Li, W. Wang, G. Yin, *ACS Catal.* **2020**, *10*, 4888-4894.
- [6] a) V. H. Menezes da Silva, A. A. C. Braga, T. R. Cundari, *Organometallics* **2016**, *35*, 3170-3181; b) A. N. Campbell, M. R. Gagné, *Organometallics* **2007**, *26*, 2788-2790; c) P. Veerakumar, P. Thanasekaran, K.-L. Lu, K.-C. Lin, S. Rajagopal, *ACS Sustain. Chem. Eng.* **2017**, *5*, 8475-8490.
- [7] For alkene dicarbofunctionalization by cyclization/coupling, see: a) B. Burns, R. Grigg, P. Ratananukul, V. Sridharan, P. Stevenson, S. Sukirthalingam, T. Worakun, *Tetrahedron Lett.* **1988**, *29*, 5565-5568; b) P. Fretwell, R. Grigg, J. M. Sansano, V. Sridharan, S. Sukirthalingam, D. Wilson, J. Redpath, *Tetrahedron* **2000**, *56*, 7525-7539; c) R. Grigg, J. Sansano, V. Santhakumar, V. Sridharan, R. Thangavelanthum, M. Thornton-Pett, D. Wilson, *Tetrahedron* **1997**, *53*, 11803-11826; d) R. Grigg, E. Mariani, V. Sridharan, *Tetrahedron Lett.* **2001**, *42*, 8677-8680; e) J. E. Wilson, *Tetrahedron Lett.* **2012**, *53*, 2308-2311; f) P. Fan, Y. Lan, C. Zhang, C. Wang, *J. Am. Chem. Soc.* **2020**, *142*, 2180-2186; g) Y. Jin, C. Wang, *Angew. Chem. Int. Ed.* **2019**, *58*, 6722-6726; h) K. Wang, Z. Ding, Z. Zhou, W. Kong, *J. Am. Chem. Soc.* **2018**, *140*, 12364-12368; i) L. Zhou, S. Li, B. Xu, D. Ji, L. Wu, Y. Liu, Z.-M. Zhang, J. Zhang, *Angew. Chem. Int. Ed.* **2020**, *59*, 2769-2775; j) A. D. Marchese, E. M. Larin, B. Mirabi, M. Lautens, *Acc. Chem. Res.* **2020**, *53*, 1605-1619.
- [8] a) M. Kosugi, H. Tamura, H. Sano, T. Migita, *Tetrahedron* **1989**, *45*, 961-967; b) K. Masanori, K. Tomoyuki, O. Hiroshi, M. Toshihiko, *Bull. Chem. Soc. Jpn.* **1993**, *66*, 3522-3524; c) M. Kosugi, H. Tamura, H. Sano, T. Migita, *Chem. Lett.* **1987**, *16*, 193-194; d) M. Catellani, G. P. Chiusoli, S. Concar, *Tetrahedron* **1989**, *45*, 5263-5268; e) S.-K. Kang, J.-S. Kim, S.-C. Choi, K.-H. Lim, *Synthesis* **1998**, *1998*, 1249-1251; f) K. M. Shaulis, B. L. Hoskin, J. R. Townsend, F. E. Goodson, C. D. Incavito, A. L. Rheingold, *J. Org. Chem.* **2002**, *67*, 5860-5863; g) A. A. Kadam, T. L. Metz, Y. Qian, L. M. Stanley, *ACS Catal.* **2019**, *9*, 5651-5656.
- [9] For examples of alkene dicarbofunctionalization without a coordinating group, see: a) A. García-Domínguez, Z. Li, C. Nevado, *J. Am. Chem. Soc.* **2017**, *139*, 6835-6838; b) P. Gao, L.-A. Chen, M. K. Brown, *J. Am. Chem. Soc.* **2018**, *140*, 10653-10657; c) S. KC, R. K. Dhungana, B. Shrestha, S. Thapa, N. Khanal, P. Basnet, R. W. Lebrun, R. Giri, *J. Am. Chem. Soc.* **2018**, *140*, 9801-9805; d) L. Liu, W. Lee, C. R. Youshaw, M. Yuan, M. B. Geherty, P. Y. Zavali, O. Gutierrez, *Chem. Sci.* **2020**, *11*, 8301-8305; e) D. Anthony, Q. Lin, J. Baudet, T. Diao, *Angew. Chem. Int. Ed.* **2019**, *58*, 3198-3202; f) M. W. Campbell, J. S. Compton, C. B. Kelly, G. A. Molander, *J. Am. Chem. Soc.* **2019**, *141*, 20069-20078; g) W. Zhang, S. Lin, *J. Am. Chem. Soc.* **2020**, *142*, 20661-20670; h) J. Jeon, Y.-T. He, S. Shin, S. Hong, *Angew. Chem. Int. Ed.* **2020**, *59*, 281-285; i) K. Lee, S. Lee, N. Kim, S. Kim, S. Hong, *Angew. Chem. Int. Ed.* **2020**, *59*, 13379-13384; j) S. Sakurai, A. Matsumoto, T. Kano, K. Maruoka, *J.*


## COMMUNICATION

- Am. Chem. Soc.* **2020**, *142*, 19017-19022; k) D. Huang, D. Olivieri, Y. Sun, P. Zhang, T. R. Newhouse, *J. Am. Chem. Soc.* **2019**, *141*, 16249-16254; l) X. Yong, Y.-F. Han, Y. Li, R.-J. Song, J.-H. Li, *Chem. Commun.* **2018**, *54*, 12816-12819; m) X.-H. Ouyang, M. Hu, R.-J. Song, J.-H. Li, *Chem. Commun.* **2018**, *54*, 12345-12348; n) M. Chierchia, P. Xu, G. J. Lovinger, J. P. Morken, *Angew. Chem. Int. Ed.* **2019**, *58*, 14245-14249; o) J. A. Myhill, C. A. Wilhelmsen, L. Zhang, J. P. Morken, *J. Am. Chem. Soc.* **2018**, *140*, 15181-15185; p) T. Ishii, K. Ota, K. Nagao, H. Ohmiya, *J. Am. Chem. Soc.* **2019**, *141*, 14073-14077; q) X.-G. Wang, Y. Li, H.-C. Liu, B.-S. Zhang, X.-Y. Gou, Q. Wang, J.-W. Ma, Y.-M. Liang, *J. Am. Chem. Soc.* **2019**, *141*, 13914-13922.
- [10] B. Shrestha, P. Basnet, R. K. Dhungana, S. Kc, S. Thapa, J. M. Sears, R. Giri, *J. Am. Chem. Soc.* **2017**, *139*, 10653-10656.
- [11] S. Thapa, R. K. Dhungana, R. T. Magar, B. Shrestha, S. Kc, R. Giri, *Chem. Sci.* **2018**, *9*, 904-909.
- [12] W. Li, J. K. Boon, Y. Zhao, *Chem. Sci.* **2018**, *9*, 600-607.
- [13] a) J. Derosa, V. T. Tran, M. N. Boulous, J. S. Chen, K. M. Engle, *J. Am. Chem. Soc.* **2017**, *139*, 10657-10660; b) T. Yang, Y. Jiang, Y. Luo, J. J. H. Lim, Y. Lan, M. J. Koh, *J. Am. Chem. Soc.* **2020**, *142*, 21410-21419; c) Z. Bai, Z. Bai, F. Song, H. Wang, G. Chen, G. He, *ACS Catal.* **2020**, *10*, 933-940.
- [14] For radical-related alkene dicarbofunctionalization and related reactions, see: a) C. Xu, R. Cheng, Y.-C. Luo, M.-K. Wang, X. Zhang, *Angew. Chem. Int. Ed.* **2020**, *59*, 18741-18747; b) J.-W. Gu, Q.-Q. Min, L.-C. Yu, X. Zhang, *Angew. Chem. Int. Ed.* **2016**, *55*, 12270-12274; c) L. Wang, C. Wang, *Org. Lett.* **2020**, *22*, 8829-8835; d) O. Apolinar, V. T. Tran, N. Kim, M. A. Schmidt, J. Derosa, K. M. Engle, *ACS Catal.* **2020**, *10*, 14234-14239; e) L. Guo, M. Yuan, Y. Zhang, F. Wang, S. Zhu, O. Gutierrez, L. Chu, *J. Am. Chem. Soc.* **2020**, *142*, 20390-20399; f) K. Semba, N. Ohta, Y. Nakao, *Org. Lett.* **2019**, *21*, 4407-4410; g) S. Zheng, Z. Chen, Y. Hu, X. Xi, Z. Liao, W. Li, W. Yuan, *Angew. Chem. Int. Ed.* **2020**, *59*, 17910-17916; h) M. Koy, P. Bellotti, F. Katzenburg, C. G. Daniliuc, F. Glorius, *Angew. Chem. Int. Ed.* **2020**, *59*, 2375-2379; i) N. A. Weires, Y. Slutskyy, L. E. Overman, *Angew. Chem. Int. Ed.* **2019**, *58*, 8561-8565; j) J. Liu, S. Wu, J. Yu, C. Lu, Z. Wu, X. Wu, X.-S. Xue, C. Zhu, *Angew. Chem. Int. Ed.* **2020**, *59*, 8195-8202; k) D. R. White, E. M. Hinds, E. C. Bornowski, J. P. Wolfe, *Org. Lett.* **2019**, *21*, 3813-3816; l) E. C. Bornowski, E. M. Hinds, D. R. White, Y. Nakamura, J. P. Wolfe, *Org. Process Res. Dev.* **2019**, *23*, 1610-1630.
- [15] a) T. Yang, X. Chen, W. Rao, M. J. Koh, *Chem* **2020**, *6*, 738-751; b) J. Derosa, V. A. van der Puyl, V. T. Tran, M. Liu, Keary M. Engle, *Chem. Sci.* **2018**, *9*, 5278-5283.
- [16] P. Basnet, S. Kc, R. K. Dhungana, B. Shrestha, T. J. Boyle, R. Giri, *J. Am. Chem. Soc.* **2018**, *140*, 15586-15590.
- [17] L. Chen, D. R. Sanchez, B. Zhang, B. P. Carrow, *J. Am. Chem. Soc.* **2017**, *139*, 12418-12421.
- [18] For dicarbofunctionalization of carboxylates and carboxamides, see: a) X. Zhao, H.-Y. Tu, L. Guo, S. Zhu, F.-L. Qing, L. Chu, *Nat. Commun.* **2018**, *9*, 3488; b) H.-Y. Tu, F. Wang, L. Huo, Y. Li, S. Zhu, X. Zhao, H. Li, F.-L. Qing, L. Chu, *J. Am. Chem. Soc.* **2020**, *142*, 9604-9611; c) J. Derosa, T. Kang, V. T. Tran, S. R. Wisniewski, M. K. Karunananda, T. C. Jenkins, K. L. Xu, K. M. Engle, *Angew. Chem. Int. Ed.* **2020**, *59*, 1201-1205; d) J. Derosa, R. Kleinmans, V. T. Tran, M. K. Karunananda, S. R. Wisniewski, M. D. Eastgate, K. M. Engle, *J. Am. Chem. Soc.* **2018**, *140*, 17878-17883.
- [19] While our manuscript was under review, one example of ketone-directed alkene diarylation reaction appeared in a preprint. See: DOI: 10.26434/chemrxiv.14150174.
- [20] a) E. Cini, L. Banfi, G. Barreca, L. Carcone, L. Malpezzi, F. Manetti, G. Marras, M. Rasparini, R. Riva, S. Roseblade, A. Russo, M. Taddei, R. Vitale, A. Zanotti-Gerosa, *Org. Process Res. Dev.* **2016**, *20*, 270-283; b) Y.-P. Li, Y.-C. Yang, Y.-K. Li, Z.-Y. Jiang, X.-Z. Huang, W.-G. Wang, X.-M. Gao, Q.-F. Hu, *Fitoterapia* **2014**, *95*, 214-219.
- [21] Quaternarization of  $\alpha$ -C proximal to alkenes suppressed the reaction regardless of presence or absence of  $\alpha$ -H in alkenylketones. Therefore, the role of  $\alpha$ -H in the reaction could not be established unambiguously.
- [22] For mechanisms of Ni-catalyzed alkene difunctionalization reactions, see: a) Q. Lin, T. Diao, *J. Am. Chem. Soc.* **2019**, *141*, 17937-17948; b) J. Diccianni, Q. Lin, T. Diao, *Acc. Chem. Res.* **2020**, *53*, 906-919. For general mechanisms of Ni-catalyzed C-C bond-forming reactions, see: c) M. Yuan, Z. Song, S. O. Badir, G. A. Molander, O. Gutierrez, *J. Am. Chem. Soc.* **2020**, *142*, 7225-7234; d) G. D. Jones, J. L. Martin, C. McFarland, O. R. Allen, R. E. Hall, A. D. Haley, R. J. Brandon, T. Konovalova, P. J. Desrochers, P. Pulay, D. A. Vivic, *J. Am. Chem. Soc.* **2006**, *128*, 13175-13183; e) D. J. Weix, *Acc. Chem. Res.* **2015**, *48*, 1767-1775; f) Y. Li, Y. Luo, L. Peng, Y. Li, B. Zhao, W. Wang, H. Pang, Y. Deng, R. Bai, Y. Lan, G. Yin, *Nat. Commun.* **2020**, *11*, 417.

## COMMUNICATION

TOC

## COMMUNICATION



A nickel-catalyzed alkenylarylation reaction of  $\gamma,\delta$ -alkenylketones with alkenyl triflates and arylboronic esters is described. The reaction was made feasible by the use of 5-chloro-8-hydroxyquinoline as a ligand along with  $\text{NiBr}_2 \cdot \text{DME}$  as a catalyst and  $\text{LiOtBu}$  as base, and works with cyclic, acyclic, endocyclic and exocyclic alkenyl ketones. Control experiments indicate that carbonyl coordination is required for the reaction to proceed.

Roshan K. Dhungana, Vivek Aryal,  
Doleshwar Niroula, Rishi R. Sapkota,  
Margaret G. Lakomy, and Ramesh Giri\*

Page No. – Page No.

**Ni-Catalyzed Regioselective  
Alkenylarylation of  $\gamma,\delta$ -Alkenyl  
Ketones via Carbonyl Coordination**