

Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Nucleosides 5^{15} : Synthesis of Novel 1,2,4-Triazolo[3,4-c]-1,2,4-Triazole Nucleosides

M. A. N. Mosselhi ^a & R. Neidlein ^b

^a Chemistry Department, Faculty of Science, Cairo University, Giza, Egypt

^b Pharmazeutisch-Chemisches Institut, Heidelberg, Germany

Version of record first published: 24 Nov 2009.

To cite this article: M. A. N. Mosselhi & R. Neidlein (2009): Nucleosides 5^{15} : Synthesis of Novel 1,2,4-Triazolo[3,4-c]-1,2,4-Triazole Nucleosides, Nucleosides, Nucleotides and Nucleic Acids, 28:11-12, 1095-1103

To link to this article: <http://dx.doi.org/10.1080/15257770903368377>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.


The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

NUCLEOSIDES 5¹⁵: SYNTHESIS OF NOVEL 1,2,4-TRIAZOLO[3,4-c]-1,2,4-TRIAZOLE NUCLEOSIDES

M. A. N. Mosselhi¹ and R. Neidlein²

¹Chemistry Department, Faculty of Science, Cairo University, Giza, Egypt

²Pharmazeutisch-Chemisches Institut, Heidelberg, Germany



Ribosylation of 3-methylthio-5-phenyl-1,2,4-triazole (1) with ribose derivative 2 gave the protected 1,2,4-triazole-nucleoside 3, which reacted with hydrazine hydrate to afford the 3-hydrazino-1,2,4-triazole derivative 5. Reaction of 5 with aromatic aldehydes yielded the corresponding hydrazones 6, which cyclized under bromination in acetic acid to give 8. Debenzoylation of 8 afforded novel 1,2,4-triazolo[3,4-c]-1,2,4-triazole nucleosides 9.

Keywords Ribosylation; 3-methylthio-5-phenyl-1,2,4-triazole; 1,2,4-triazole-nucleoside; 1-O-acetyl-2,3,5-tri-O-benzoyl-β-D-ribofuranose; 1,2,4-triazolo[3,4-c]-1,2,4-triazole nucleosides

Received 12 May 2009; accepted 6 October 2009.

Present address for M.A.N.M.: Taif University, Faculty of Science, Department of Chemistry Taif, Saudi Arabia.

The authors thank den Deutschen Forschungsgemeinschaft (DFG) for the financial support. The authors also thank Prof. Dr. Helmut Duddeck, Institut fuer Organische Chemie, Hannover, Germany for measuring the NOE-¹H NMR spectrum of compound 3.

Address correspondence to M.A.N. Mosselhi, Chemistry Department, Faculty of Science, Cairo University, Giza 12613, Egypt. E-mail: mosselhi2008@hotmail.com

INTRODUCTION


1,2,4-Triazoles represent a class of heterocyclic compounds of significant importance in agriculture and medicine.^[1,2] Compounds with 1,2,4-triazole moiety also have received considerable attention among medicinal chemists because molecules with these structural features have been found to display a wide range of potent biological activities, such as antihypertensive,^[2] antifungal,^[3] and antibacterial^[4] activities. Appropriately functionalized 1,2,4-triazoles, such as 4-[3,5-bis(2-hydroxyphenyl)-1,2,4-triazole-1-yl]benzoic acid, have been shown to selectively form a complex with iron (III), which is useful in iron overload therapy.^[5]

Considering the important biological properties of 1, 2, 4-triazole compounds, several efficient triazole syntheses have been reported.^[6,7,8] According to these interesting biological activities, and as a part of our research program on the nucleoside chemistry, we report here a synthesis of two novel fused 1,2,4-triazole nucleosides. So far in the literature the few examples of 1,2,4-triazolo-[1,2,4]triazole, include 1,2,4-triazolo[1,2-a],^[9] 1,2,4-triazolo[3,4-c],^[10-12] and 1,2,4-triazolo[3,4-b] [1,2,4]triazoles.^[13]

RESULTS AND DISCUSSION


The starting material, 3-methylthio-5-phenyl-1(2H) [1,2,4]triazole (**1**), was prepared according to the literature reported method.^[14] The ribosylation of **1** with 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (**2**) was carried out in two different methods to give the benzoylated nucleoside; 3-methylthio-5-phenyl-2-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)-(2*H*)-1,2,4-triazole **3** or 3-methylthio-5-phenyl-4-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)-(4*H*)-1,2,4-triazole **4** (see Scheme 1). The first method was silylation of **1** by reflux in hexamethyl disilazane (HMDS) with ammonium sulfate as a catalyst and then stirring of the silylated product with ribose derivative **2** in dry acetonitrile and TMS-triflate (CF₃SO₂OSiMe₃) at room temperature for 48 hours. This method afforded a product in 60% yield.

The second method was stirring of the lithium salt of compound **1** with **2** in tetrahydrofuran and TMS-triflate at -78°C for 5 hours and followed by stirring at room temperature for 24 hours. This method yielded a product in 90%. The structure of the latter product was established and confirmed on the bases of its elemental analyses and spectral data. Thus, the analytical data for it revealed a molecular formula C₃₅H₂₉N₃O₇S (m/z 635). Its ¹H NMR spectrum showed a doublet at δ 6.1 assigned to the anomeric proton of the ribose moiety with a spin-spin coupling constant equal to 4.1 Hz, which corresponds to a diaxial orientation for the 1'- and 2'- H protons; that is, the β -configuration.^[15-19] The ultraviolet (UV) spectra of the above product revealed λ_{max} (ethanol) at 249 nm, which is similar to that of 5-methyl-4-phenyl-3-(tri-O-acetyl- β -D-xylopyranosylmercapto)-1,2,4-

SCHEME 1 X = a, H, b, CH_3 .

triazole [λ_{max} (ethanol) 245 nm].^[19] Furthermore, the irradiation of the anomeric proton of **3** gave a clear nuclear Overhauser effect (NOE) response from the *ortho* protons of the 5-phenyl group due to spatial neighborhood. These findings confirmed the structure of **3** and the structure **4** was discarded.

The reaction of compound **3** with hydrazine hydrate under reflux in ethanol afforded 3-hydrazino-2-(2',3',5'-tri-O-benzoyl)- β -D-ribo-furanozyl-5-phenyl-(2*H*)-1,2,4-triazole (**5**). Furthermore compound **5** reacted with aromatic aldehydes in presence of acetic acid to give the corresponding hydrazones (**6**) in good yields. Next, we examined the bromination of **6** in acetic acid and anhydrous sodium acetate at room temperature. Thus, it afforded directly the triazolotriazole nucleosides (**8**) via in situ formed hydrazonoyl halide derivatives (**7**) that were not isolated (see Scheme 2). However, the debenzoylation of compounds **8** in methanol containing sodium methoxide was successful and gave the free 1,2,4-triazolo[3,4-*c*]-1,2,4-triazole nucleosides (**9**) (Scheme 2). Structural assignments of the



SCHEME 2

products **8** and **9** isolated were made on the basis of their mass spectrometry (MS), ^1H NMR, ^{13}C NMR, infrared (IR) spectra, and elemental analyses (see Experimental section). For example, the absence of the methine and nitrogen protons of hydrazones **6** in the ^1H NMR spectra of **8** or **9** confirmed the structure.

EXPERIMENTAL

All chemicals were supplied by Sigma-Aldrich and Merck (Germany).

IR spectra were recorded for KBr discs on Testscan Shimadzu FTIR 8000 Series and Bruker IFS 113V spectrophotometers (Germany), ^1H -NMR and ^{13}C -spectra were recorded on a Bruker Ac 250 MHz and on a Varian Gemini 200 MHz NMR spectrometer (Germany) using trimethylsilane (TMS) as an internal standard; NOE-diffraction was recorded on a Bruker AM 400 MHz spectrometer (Germany); UV spectra were recorded on a Perkin Elmer spectrophotometer Lambda 5 (USA); MS spectra were recorded on Varian Mat 711 spectrometer (electron ion [EI] and fast atom bombardment [FAB] in nitrobenzyl alcohol [NBA]). Thin layer chromatography (TLC) was performed on silica gel sheets F 1550 LS 254 of Schleicher & Schull and column chromatography on Merck silica gel 60 (particle size 0.063–0.20 mm). Melting points were measured on Gallenkamp melting point apparatus (UK) and are uncorrected. Elemental analyses were carried out at Pharmazeutisch-Chemisches Institut (Heidelberg, Germany) and at Microanalytical Centre (Cairo University, Egypt). 3-Methylthio-5-phenyl-1,2,4-triazole^[10] was prepared as previously described.

3-Methylthio-5-phenyl-2-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)-(2H)-1,2,4-triazole **3**

Method A. A mixture of **1** (20 mmol) and dry hexamethyldisilazane (100 ml) was heated under reflux for 10 hours with a catalytic amount of ammonium sulfate (100 mg). After the clear solution was cooled, it was evaporated to dryness under anhydrous condition to give the silylated derivative, which directly was dissolved in 50 ml of dry 1,2-dichloroethane. To this a solution of 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (**2**) (9.6 g, 19.6 mmol) in dry 1,2-dichloroethane (50 ml) was added. The mixture was cooled in ice bath and a solution of trimethylsilyl trifluoromethanesulfonate (4 ml, 20 mmol) in dry 1,2-dichloroethane (20 ml), which was added dropwise. It was stirred at room temperature for 24 hours, diluted with chloroform (500 ml), washed with a saturated solution of aqueous sodium bicarbonate (200 ml) and water (3 \times 150 ml), and dried over anhydrous sodium sulfate. The solvent was removed in vacuo and the residue was chromatographed on silica gel with chloroform as eluent to afford white

solid, which was crystallized from ethanol to yield colorless crystals of the corresponding nucleoside derivative **3**.

Method B. A mixture of **1** (20 mmol) and sodium hydride (20 mmol) was dissolved in dry tetrahydrofuran (100 ml) at -70°C and stirred for 1 hour. To this mixture a solution of 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (**2**) (9.6 g, 19.6 mmol) in dry tetrahydrofuran (40 ml) was then added and the mixture was stirred at -70°C for 5 hours. The excess solvent was evaporated and the residue was extracted from chloroform and purified by chromatography as in the above method to give **3**, which is identical in all physical constants with that obtained from the above method.

Compound **3**: Yield (60%, method A and 90% method B); m.p. $158\text{--}159^{\circ}\text{C}$; ν (cm^{-1}) (KBr) 1750 (CO); λ_{max} (EtOH) (249 nm); δ_{H} (CDCl_3): 2.61 (s, 3H, SCH₃), 4.6–4.9 (m, 3H, 2 H-5', H-4'), 6.1 (d, 1H, H-1', $J_{1',2'} = 4.1$ Hz), 6.25–6.35 (pt, 1H, H-3'), 6.4–6.5 (pt, 1H, H-2'), 7.3–8.1 (m, 20H, Ar-H); δ_{C} (CDCl_3): 18, 60, 71, 78, 84, 93, 121, 125–140, 151, 159, 168 (CO); m/z (M^+): 635, 445, 323, 201, 105, 77; Anal. Found: C, 65.70; H, 4.60; N, 6.72; S, 4.88% $\text{C}_{34}\text{H}_{26}\text{N}_2\text{O}_8\text{S}$ requires; C, 66.14; H, 4.57; N, 6.61; S, 5.04%.

3-Hydrazino-5-phenyl-2-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)-(2H)-1,2,4-triazole **5**

A mixture of the protected nucleoside **3** (15 mmol), absolute ethanol (50 ml), and hydrazine hydrate (20 ml) was heated under reflux for 2 hours. Evaporation of the solvent under vacuum gave a colorless solid, which was crystallized from ethanol/dioxane to give **5** as colorless crystals.

Compound **5**: Yield (75%); m.p. $222\text{--}224^{\circ}\text{C}$; ν (cm^{-1}) (KBr) 3310, 3190 (NH_2 , NH), 1750 (CO); δ_{H} (DMSO-d_6): 4.65–4.90 (m, 3H, 2 H-5', H-4'), 5.9 (s, 2H, NH_2), 6.2 (d, 1H, H-1', $J_{1',2'} = 4.1$ Hz), 6.30–6.35 (pt, 1H, H-3'), 6.45–6.50 (pt, 1H, H-2'), 7.25–8.3 (m, 20H, Ar-H), 9.2 (s, 1H, NH); m/z (M^+): 603, Anal. Found: C, 69.5; H, 4.8; N, 6.6%. $\text{C}_{35}\text{H}_{29}\text{N}_3\text{O}_7$ requires; C, 69.64; H, 4.84; N, 6.96%.

Benzaldehyde and p-Methylbenzaldehyde-[(5-phenyl-2-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)-(2H)-1,2,4-triazol-3-yl] Hydrazones **6**

General method. A mixture of **5** (0.01 mol) and benzaldehyde or p-methylbenzaldehyde (5 ml each) in ethanol (50 ml) and acetic acid (0.5 ml) was heated under reflux for 1 hour. The mixture was evaporated to half and then cooled. The solid formed was filtered and crystallized from ethanol to give **6** as colorless crystals.

Compound **6a**: Yield (48%); m.p. $192\text{--}193^{\circ}\text{C}$, ν (cm^{-1}) (KBr) 3311 (NH), 1755 (CO); δ_{H} (DMSO-d_6): 4.6–4.7 (m, 1H, 4'-H), 4.85–4.95 (m, 2H, 5', 5''-H; 6.0 (d, 1H, $J_{1',2'} = 4.0$ Hz, 1'-H), 6.2–6.25 (m, 1H, 3'-H); 6.35–6.45

(m, 1H, 2'-H); 7.2–8.5 (m, 26H, Ar-H & = CH), 9.3 (s, 1H, NH); m/z (M⁺): 707; Anal. Found: C, 69.2; H, 4.7; N, 10.1%. C₄₁H₃₃N₅O₇ requires; C, 69.58; H, 4.70; N, 9.90%.

Compound **6b**: Yield (50%); m.p. 180–182°C, ν (cm⁻¹) (KBr) 3315 (NH), 1755 (CO); δ_{H} (DMSO-d₆): 2.2 (s, 3H, CH₃), 4.5–4.6 (m, 1H, 4'-H); 4.75–4.8 (m, 2H, 5',5''-H); 5.95 (d, 1H, J_{1',2'} = 4.0 Hz, 1'-H), 6.1–6.2 (m, 1H, 3'-H); 6.3–4.4 (m, 1H, 2'-H); 7.3–8.2 (m, 25H, Ar-H & = CH), 9.1 (s, 1H, NH); m/z (M⁺): 721; Anal. Found: C, 69.5; H, 4.5; N, 10.0%. C₄₂H₃₅N₅O₇ requires; C, 69.89; H, 4.89; N, 9.70%.

1-(2',3',5'-Tri-O-benzoyl- β -D-ribofuranosyl)-5-aryl-3-phenyl-1,2,4-triazolo[3,4-c]-1,2,4-triazole Nucleosides 8

General method. A suspension of **6** (5 mmol) in acetic acid (10 ml) was treated with bromine (5 ml) in acetic acid (2 ml) dropwise and the reaction mixture was stirred for 24 hours. It was then diluted with water and the product was filtered and crystallized from ethanol/dioxane to give **8**.

Compound **8a**: Yield (65%); m.p. 220–222°C, ν (cm⁻¹) (KBr) 1765 (CO); λ_{max} (EtOH) (253 nm); δ_{H} (DMSO-d₆) 4.6–4.7 (m, 1H, 4'-H); 4.8–4.85 (m, 2H, 5',5''-H); 6.0 (d, 1H, J_{1',2'} = 2.0 Hz, 1'-H), 6.1–6.2 (m, 1H, 3'-H); 6.3–4.4 (m, 1H, 2'-H); 7.2–8.3 (m, 25H, Ar-H); δ_{C} (DMSO-d₆): 65, 73, 75, 82, 90, 123, 128–135, 162, 170 (CO); m/z (M⁺): 705; Anal. Found: C, 68.0; H, 4.5; N, 9.8%. C₄₁H₃₁N₅O₇ requires; C, 69.78; H, 4.43; N, 9.92%.

Compound **8b**: Yield (70%); m.p. 212–213°C, ν (cm⁻¹) (KBr) 1760 (CO); λ_{max} (EtOH) (250 nm); δ_{H} (DMSO-d₆) 2.1 (s, 3H, CH₃), 4.5–4.6 (m, 1H, 4'-H); 4.75–4.8 (m, 2H, 5',5''-H); 6.0 (d, 1H, J_{1',2'} = 2.0 Hz, 1'-H), 6.15–6.25 (m, 1H, 3'-H); 6.3–4.4 (m, 1H, 2'-H); 7.2–8.3 (m, 24H, Ar-H); δ_{C} (DMSO-d₆) 24, 63, 71, 78, 84, 95, 121, 126–136, 160, 169 (CO); m/z (M⁺): 719; Anal. Found: C, 69.6; H, 4.4; N, 9.5%. C₄₂H₃₃N₅O₇ requires; C, 70.09; H, 4.62; N, 9.73%.

1-(β -D-ribofuranosyl)-5-aryl-3-phenyl-1,2,4-triazolo[3,4-c]-1,2,4-triazole Nucleosides 9

General method. A mixture of the protected nucleoside **8** (1 mmol), absolute methanol (20 ml), and sodium methoxide (60 mg, 1.1 mol) was stirred at room temperature for 24 hours. Evaporation of the solvent under vacuum gave a colorless solid, which was dissolved in hot water and neutralized with acetic acid. The precipitate was filtered off and afforded upon crystallization from water, the nucleosides **9** as colorless crystals.

Compound **9a**: Yield (40%); m.p. 245–246°C, ν (cm⁻¹) (KBr) 3356 (OH); λ_{max} (EtOH) (265 nm); δ_{H} (DMSO-d₆) 3.5–3.6 (m, 2H, 5',5''-H); 3.9–4.0 (m, 1H, 4'-H); 4.15–4.2 (m, 1H, 3'-H); 4.35–4.45 (m, 1H, 2'-H); 4.7–4.8 (t, 1H, 5'-OH); 5.1–5.2 (d, 1H, 3'-OH); 5.4–5.5 (d, 1H, 2'-OH);

6.1–6.2 (d, 1H, $J_{1',2'} = 2.0$ Hz, 1'-H); 7.2–8.0 (m, 10H, Ar-H); m/z (M^+): 393; Anal. Found: C, 60.0; H, 4.5; N, 17.4%. $C_{20}H_{19}N_5O_4$ requires; C, 61.06; H, 4.87; N, 17.8%.

Compound **9b**: Yield (45%); m.p. 233–234°C, ν (cm^{-1}) (KBr) 3360 (OH); λ_{max} (EtOH) (260nm); δ_H (DMSO- d_6) 2.0 (s, 3H, CH_3), 3.45–3.55 (m, 2H, 5',5''-H); 3.7–3.9 (m, 1H, 4'-H); 4.15–4.25 (m, 1H, 3'-H); 4.3–4.4 (m, 1H, 2'-H); 4.65–4.75 (t, 1H, 5'-OH); 5.1–5.2 (d, 1H, 3'-OH); 5.4–5.5 (d, 1H, 2'-OH); 6.1–6.2 (d, 1H, $J_{1',2'} = 2.0$ Hz, 1'-H); 7.2–7.8 (m, 9 H, Ar-H); m/z (M^+): 407; Anal. Found: C, 62.0; H, 5.4; N, 16.8%. $C_{21}H_{21}N_5O_4$ requires; C, 61.91; H, 5.2; N, 17.19%.

REFERENCES

1. Mavrova, A.Ts.; Wesselinova, D.; Tsenov, Y.A.; Denkova, P. Synthesis, cytotoxicity and effects of some 1,2,4-triazole and 1,3,4-thiadiazole derivatives on immunocompetent cells. *Eur. J. Med. Chem.* **2009**, 44, 63.
2. Al-Masoudi, I.A.; Al-Soud, Y.A.; Al-Salihi, N.J.; Al-Masoudi, N.A. 1,2,4-Traizoles: Synthetic approaches and pharmacological importance. *Chem. Heterocycl. Compd.* **2006**, 42, 1377.
3. Kaku, Y.; Tsuruoka, A.; Kakinuma, H.; Tsukada, I.; Yanagisawa, M.; Natio, T. A novel route for chiral synthesis of the triazole antifungal ER-30346. *Chem. Pharm. Bull.* **1998**, 46, 1125.
4. Sui, Z.; Guan, J.; Hlasta, D.J.; Macielag, M.J.; Foleno, B.D.; Goldschmidt, R.M.; Loeloff, M.J.; Webb, G.C.; Barrett, J.F. Sar studies of diaryl triazoles against bacterial two-component regulatory systems and their antibacterial activities. *Bioorg. Med. Chem. Lett.* **1998**, 8, 1929.
5. Heiz, U.; Hegetschweiber, K.; Acklin, P.; Faller, B.; Lattmann, R.; Schnebli, H.P. 4-[3,5-Bis(2-hydroxyphenyl)-1,2,4-triazol-1-yl]-benzoic acid: A novel efficient and selective iron(III) complexing agent. *Angew. Chem. Int. Ed. Engl.* **1990**, 38, 2568.
6. Neidlein, R.; Sui, Z. The 1,3-Dipolar Cycloadditions of Nitrile Oxides and Nitrile Imines to Alkyl Dicyanoacetates. *Helv. Chim. Acta* **1991**, 74, 501.
7. Mosselhi, M.A.N.; Abdallah, M.A.; Riyadh, S.M.; Harhash, A.E.; Shawali, A.S. Tautomerism and reactions of 1H-1,2,4-triazole-5-thiones with hydrazonoyl halides. *J. Prakt. Chem.* **1998**, 30(2), 160.
8. Kumarapperuma, S.C.; Sun, Y.; Jeselnik, M.; Chung, K.; Parker, W.B.; Jonsson, C.B.; Arterburna, J.B. Synthesis effects on the phosphorylation of 3-substituted 1- β -D-ribofuranosyl-1,2,4-triazoles by human adenosine kinase. *Bioorg. Med. Chem. Lett.* **2007**, 17, 3203.
9. Korotkikh, N.I.; Pekhtereva, T.M.; Suikov, S. Yu; Smirnov, Yu.I. Synthesis and structure of the glycidyl derivatives of two-nuclear and annealed triazolones. *Russ. J. Org. Chem.* **1999**, 35(4), 591.
10. a) Padhy, A.K.; Nug, V.L.; Panda, C.S. Studies on the synthesis and bio-activity of some thiadiazole derivatives. *Indian J. Chem.* **1999**, 38B(8), 998; b) Wolkoff, P. 1-Substituted 3,5-diaryl-s-triazolo[3,4-c]-s-triazoles. *Acta Chem. Scan.* **1976**, B30, 463.
11. Hegarty, A.F.; Quain, P.; O'Mahony, T.A.F.; Scott, F.L. Mechanism of cyclization of N-(1,2,4-triazol-3-yl)hydrazonoyl bromides to mixtures of isomeric triazolotriazoles. *J. Chem. Soc., Perkin Trans.* **1972**, 2, 997.
12. Shawali, A.S.; Abdallah, M.A.; Abbas, I.M.; Eid, G.M. New regioselective synthesis and biological activity of substituted 1H-[1,2,4]triazolo[3,4-c][1,2,4]triazoles. *J. Chin. Chem. Soc.* **2004**, 51(2), 351.
13. Jois, Y.H.R.; Kwong, C.D.; Riordan, J.M.; Montgomery, J.A.; Secrist III, J.A. Synthesis and antiviral evaluation of some novel [1,2,4]triazolo[4,3-b][1,2,4]-triazole nucleoside analogs. *J. Heterocycl. Chem.* **1993**, 30, 1289.
14. Hoggarth, E. Compounds related to thiosemicarbazide. Part II. 1-Benzoylthiosemicarbazides. *J. Chem. Soc.* **1949**, 1163.
15. Break, L.M.; Shmiss, N.A.M.M.; Mosselhi, A.N.M. Nucleosides 4: Synthesis of some new S-Nucleoside derivatives of 2-thioxo and 2,4-dithioxo-5,6,7,8-tetrahydrobenzo-thieno[2,3-d]pyrimidin-4(3H) ones. *Phosphorus, Sulfur, Silicon & the Related Elements* (in press, 2010).

16. Mosselhi, M.A.N. Nucleoside 1: Ribosylation of 8-substituted theophylline derivatives. *Nucleosides Nucleotides Nucleic Acids* **1993**, 12(5), 431.
17. Mosselhi, M.A.N. Nucleosides 2: Synthesis and properties of 3,4-diaryl-4,5-Dihydro-1-(β -D-ribofuranosyl)-1,2,4-triazole-5-thiones. *Nucleosides Nucleotides Nucleic Acids* **1999**, 18(9), 2043.
18. Mosselhi, M.A.N. Nucleosides 3: Synthesis of Novel 1,3,4-Oxadiazole Nucleosides, Seliger H; *J. Chem. Res. (S)* **2001**, 359.
19. Wagner, G., Valz, G., Dietzsch, B., Fischer, G. Glucosides and xylosides of 1-phenyl-5-mercaptotetrazole and 3-methyl-4-phenyl-5-mercapto-1,2,4-triazole. 49. Glycosides of heterocyclic compounds. *Pharmazie* **1974**, 29(2), 90.