This article was downloaded by: [University of Guelph] On: 04 January 2015, At: 12:41 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

Novel Regioselective Hydroxyl-Alkylation of 4,5-Diphenylimidazole-2-Thione and A Competitive Intramolecular Ring Closure of the S-Hydroxyalkyl-Imidazoles to Imidazo[2,1-b]Thiazines and Thiazoles. Role of Catalyst, Microwave Irradiation, and Solid Support

E. S. H. El Ashry $^{\rm a}$, N. Rashed $^{\rm a}$, L. F. Awad $^{\rm a}$, E. Ramadan $^{\rm a}$, S. M. Abdel-Maggeed $^{\rm a}$ & N. Rezki $^{\rm a}$

^a Chemistry Department, Faculty of Science, Alexandria University, Alexandria, Egypt Published online: 19 Jun 2007.

To cite this article: E. S. H. El Ashry, N. Rashed, L. F. Awad, E. Ramadan, S. M. Abdel-Maggeed & N. Rezki (2007) Novel Regioselective Hydroxyl-Alkylation of 4,5-Diphenylimidazole-2-Thione and A Competitive Intramolecular Ring Closure of the S-Hydroxyalkyl-Imidazoles to Imidazo[2,1-b]Thiazines and Thiazoles. Role of Catalyst, Microwave Irradiation, and Solid Support, Nucleosides, Nucleotides and Nucleic Acids, 26:5, 423-435, DOI: <u>10.1080/15257770701426179</u>

To link to this article: http://dx.doi.org/10.1080/15257770701426179

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

Nucleosides, Nucleotides, and Nucleic Acids, 26:423–435, 2007 Copyright © Taylor & Francis Group, LLC ISSN: 1525-7770 print / 1532-2335 online DOI: 10.1080/15257770701426179



NOVEL REGIOSELECTIVE HYDROXYL-ALKYLATION OF 4,5-DIPHENYLIMIDAZOLE-2-THIONE AND A COMPETITIVE INTRAMOLECULAR RING CLOSURE OF THE S-HYDROXYALKYL-IMIDAZOLES TO IMIDAZO[2,1-*b*]THIAZINES AND THIAZOLES. ROLE OF CATALYST, MICROWAVE IRRADIATION, AND SOLID SUPPORT

E. S. H. El Ashry, N. Rashed, L. F. Awad, E. Ramadan, S. M. Abdel-Maggeed, and N. Rezki
Chemistry Department, Faculty of Science, Alexandria University, Alexandria, Egypt

□ Under both conventional method (CM) and microwave (MW) irradiation (MWI) conditions, alkylation of 4,5-diphenylimidazole-2-thione (1) with halogeno-alkanols 2 or 5, chloroglycerol 11 and 2,3-O-isopropylidene-1-O-(p-tolylsulfonyl)-glycerol (8) in presence of sodium ethoxide or sodium acetate in alcohol afforded regioselectively the corresponding S-alkylated analogues 3, 6, 9, and 12; they also were obtained using MW in absence and presence of bentonite as solid support with no change in regioselectivity. In the presence of potassium carbonate in DMF, the bisalkylated analogues 4, 7, 10, and 13 were obtained except in case of compound 13 where it was accompanied with the imidazothiazine 14. A convenient approach for imidazo-[2,1-b]thiazines and thiazoles 14–16 could be achieved by intramolecular dehydrative ring closure of the S-hydroxyalkylated imidazoles 3, 6, and 12 using potassium carbonate in DMF under both conventional and microwave methods. Isopropylidenation of 12 and 13 and deprotection of 9 and 10 also were investigated.

Keywords Imidazole; imidazo[2,1-*b*][1,3]thiazine; Imidazo[2,1-*b*][1,3]thiazole; acyclonucleoside; alkylation; microwave irradiation; solid support

INTRODUCTION

Many acyclonucleoside analogues^[1] such as acyclovir^[2] (ACV), ganciclovir^[3] (GCV) and penciclovir^[4] (PCV), and (*S*)-9-(2,3-dihydroxy-propyl)adenine^[5] posses potent antiviral activity. Extensive efforts have

Received 25 July 2006; accepted 5 March 2007.

E. S. H. El-Ashry is on leave to International Center for Chemical and Biological Sciences, HEJ Research Institute, University of Karachi, Karachi, Pakistan.

N. Rezki is on leave from the Ministry of High Education and Scientific Research, Algeria.

Address correspondence to E. S. H. El Ashry, Chemistry Department, Faculty of Science, Alexandria University, Alexandria, Egypt. E-mail: eelashry60@hotmail.com

been directed to synthesize acyclonucleoside analogues of the natural ones by modification of the heterocyclic base and/or the acyclic side chain.^[1,6–11]

Valuable medicinal applications such as antiinflammatory,^[12] antiasthmatic,^[13] antiulcerative^[14] and antithrombotic^[15] activities have been associated with 2-thiosubstituted imidazoles and diaryl imidazoles.

Microwave technology has opened up new horizons for chemists and increasingly used to accelerate organic reactions.^[16–20] The solvent free conditions also have been adopted to organic synthesis as a tool for green chemistry^[21–24] particularly when coupled to microwave irradiation.

On the basis of these findings, it was of interest to report on the regioselectivity encountered and the intramolecular anellation to form imidazothiazine and imidazothiazole during the synthesis of acyclonucle-oside analogues from the 4,5-diphenylimidazole-2-thione^[25] under both conventional method (CM) and microwave (MW) as a continuation of a program directed to use microwave technology in our laboratory.^[16,17,26–36] The competitive intramolecular cyclization with the N-hydroxylalkylation can be explained by the semiemperical AM1 theoritical calculations.

RESULTS AND DISCUSSION

This work describes the regioselective alkylation of 4.5diphenylimidazole-2-thione (1) by reaction with one and two equivalents of different hydroxyl-alkylating agents including halogeno-ethanol 2, propanol 5, 2,3-dihydroxychloropropanol 11 and 2,3-O-isopropylidene-1-O-(*p*-tolylsulfonyl)-glycerol (8) under both conventional and microwave methods. Reaction of 1 with 1.1 equivalents of 2, 3, 8, and 11 in a boiling solution of sodium ethoxide in ethanol for 2-6 hours or in boiling methanol in the presence of NaOAc for 4-14 hours afforded only the S-alkyl derivatives 3, 6, 9, and 12, respectively in 62–74%. Deprotection of 9 with 70% acetic acid at reflux temperature gave the corresponding 2-(2,3-dihydroxyprop-1-ylthio)-4,5-diphenylimidazole (12) whose reaction with acetone in sulfuric acid gave back compound 9 (Schemes 1 and 2). When the above reactions were carried out under MW, 1–5 minutes were required to give the same products in higher yields (Table 1), and no change in regioselectivity was noticed. A higher yield (79-88%) was achieved under MW irradiation for 1–3 minutes of a mixture of 1 with the alkylating agent adsorbed on the surface of the bentonite in a closed Teflon vessel.

The structures of the S-alkylated compounds have been established on the basis of their elemental analyses and spectral data. The IR spectra of **3**, **6**, and **12** showed characteristic absorption band at 3314—3413 cm⁻¹ due to OH of the acyclic side chains. Their ¹H NMR spectra indicated the



a) NaOEt, EtOH, reflux, CM; b) NaOEt, EtOH, MW;

- c) NaOAc, MeOH, reflux, CM; d) NaOAc, MeOH, MW;
- e) Bentonite, MW;
- f) K₂CO₃, DMF, reflux, CM; g) K₂CO₃, DMF, MW

SCHEME 1 Reaction of imidazole thione with halogeno-ethanol and propanol.

	Conditions					Conventional method (CM)		Microwave method (MW)	
Compd.	Alkylating				Solid	Time	Yield	Time	Yield
No.	agent	Entry*	Base	Solvent	support	(hr)	(%)	(min)	(%)
3	2	a, b	NaOEt	EtOH	_	2.0	74	1.0	89
3	2	c, d	NaOAc	MeOH	—	4.0	70	2.0	84
3	2	e	—	—	Bentonite	—	_	1.0	88
4	2	f, g	K_2CO_3	DMF	—	8.0	66	4.0	81
6	3	a, b	NaOEt	EtOH	—	3.0	72	2.0	88
6	3	c, d	NaOAc	MeOH	—	6.0	69	2.5	82
6	3	e	_	_	Bentonite	_	_	1.5	86
7	3	f, g	K_2CO_3	DMF	—	10.0	65	4.5	76
9	5	a, b	NaOEt	EtOH	_	06.0	67	3.0	82
9	5	c, d	NaOAc	MeOH	—	14.0	62	5.0	76
9	5	e	_	_	Bentonite	—	_	3.0	79
10	5	f, g	K_2CO_3	DMF	—	20.0	62	5.0	75
12	4	a, b	NaOEt	EtOH	_	05.0	69	3.0	81
12	4	c, d	NaOAc	MeOH	—	10.0	65	4.0	76
12	4	e	_	_	Bentonite	—	_	3.0	79
13, 14	4	f, g	K ₂ CO ₃	DMF	_	14.0	68	6.0	79
14	—	f, g	K_2CO_3	DMF	—	48.0	60	4.0	70
15	—	f, g	K_2CO_3	DMF	—	96.0	65	6.0	78
16	—	f, g	K_2CO_3	DMF	—	72.0	55	4.0	65

TABLE 1 Comparative results from conventional method (CM) and microwave (MW) method

*See Scheme.



a) NaOEt, EtOH, boil, CM; b) NaOEt, EtOH, MW;

c) NaOAc, MeOH, boil, CM;d) NaOAc, MeOH, MW;

e) Bentonite, MW; f) K₂CO₃, DMF, boil, CM;

g) K₂CO₃, DMF, MW; h) AcOH, H₂O; i) Acetone, H₂SO₄

SCHEME 2 Reaction of imidazolethione with glycerol derivatives.

formation of the S-alkylated products by the assignment of one exchangeable proton at $\delta_{\rm H}$ 12.54–12.61 due to a one NH proton of the imidazole ring. Moreover, the OH protons of **3** and **6** were assigned to the singlets at $\delta_{\rm H}$ 5.13 and 4.72, respectively. The spectrum of **12** showed two doublet of doublets of the SCH₂ at $\delta_{\rm H}$ 3.14 and 3.31 ppm, which were correlated with the multiplet at $\delta_{\rm H}$ 3.7–3.7 ppm of CHO whereas their respective carbons were assigned at $\delta_{\rm c}$ 37.2 and 71.8, respectively. The multiplet at $\delta_{\rm H}$ 3.35–3.43 ppm for CH₂O was correlated with its corresponding carbon at $\delta_{\rm c}$ 64.5. The spectrum of **12** also showed a triplet at $\delta_{\rm H}$ 4.98 and a singlet at $\delta_{\rm H}$ 5.45, which were assigned to the OH groups. The ¹H NMR spectrum of compound **9** has two doublet of doublets at $\delta_{\rm H}$ 3.20 and 3.27 ppm corresponding to the SCH₂ protons, whereas the two doublet of doublets at $\delta_{\rm H}$ 3.80 and 4.16 were assigned to the CH₂O protons. Its ¹³C NMR spectrum showed signals at $\delta_{\rm c}$ 36.8, 68.4, 77.1, and 110.1 corresponding to SCH₂, CHO, -CH₂O, and Me₂C, respectively.

Attempted preparation of the acyclonucleosides 4, 7, 10, and 13 from 1 by using 2.2 equivalents of the alkylating agents 2, 5, and 8 in presence of NaOEt or NaOAc in alcoholic solvent was unsuccessful. On the other hand, in the presence of K_2CO_3 in boiling DMF as base for 8–20 hours the corresponding acyclonucleosides analogues 4, 7, and 12 were successfully obtained in 62–66% yield. The use of 1.1 equivalent of the alkylating agent in presence of K_2CO_3 in boiling DMF gave a mixture of the Salkylated and S,N-bis(alkylated) derivatives as well as some starting material. Isopropylidenation of 13 gave 10 whose deprotection gave 13.

Unexpectedly, reaction of 1 with 11 under the same condition did not only afford the S,N-bis(dihydroxypropyl) 13, but an additional product was obtained and identified as imidazothiazine 14 (Scheme 2). The identity of such cyclized product indicated that its precursor would be the respective monoalkylated derivative 12. The formation of the two products has reflected that a competitive reaction, following the preliminary common S-alkylation, between the intramolecular dehydrative cyclization reaction and the further alkylation on the nitrogen-imidazole atom has taken place. Such tendency of intramolecular ring closure has prompted us to apply these conditions, K_2CO_3 in DMF, to the S-alkylated analogue 12 whereby the cyclized product was formed as a sole product. Moreover, the cyclization of 3 or 6 gave 15 and 16, respectively, under similar conditions, indicating the generality of these conditions to achieve such intramolecular cyclization and it is favorable that the S-alkylated derivatives were the intermediates. Although, the cyclizations required heating for long periods of times, much shorter times were only needed under the MW to give even better yields (Table 1). Mitsunobu reaction condition^[37,38] was used to achieve the cyclization of 6-amino-9-(3-hydroxypropyl)-7H-purine-8(9H)-thione to the anhydro analogue.

The structures of the fused heterocyles were deduced from the analysis of their spectra. The IR and the ¹H NMR spectra showed the disappearance of NH proton of the imidazole ring. The ¹H NMR spectrum of **14** indicated that the terminal OH of its precursor **12** disappeared upon the cyclization and the multiplet of the terminal methylene protons of **12** was shifted to the downfield region as two doublets of doublets at $\delta_{\rm H}$ 3.55 and 3.76 ppm.

In order to explain the observed difference in the mode of reaction of **1** with the different alkylating agents and the competitive intramolecular cyclization with the formation of acyclonucleoside analogues, the semiemperical AM1 theoretical methods, carried out with MOPAC7 program pack-



EQUATION 1 Synthesis of imadazo-thiazones and thaizoles.

age has been used. Thus, the heat of formation for the mono S-alkylated derivatives **3**, **4**, and **12** and the S,N-bis(alkylated) products **4**, **7**, and **13** as well as the respective fused rings **14–16** have been calculated. The higher stability of the S-alkylated product **12** ($\Delta H = -2.95 \text{ Kcal} \cdot \text{mol}^{-1}$) compared to other analogues **3** and **6** ($\Delta H = 48.06$ and 41.38 Kcal \cdot mol^{-1}, respectively) allowed a more chance of intramolecular dehydrative cyclization to take place to give **14** in addition to the alkylation to give **13**. Moreover, the lower heat of formation of imidazo-thiazine **14** ($\Delta H = 59.69 \text{ Kcal} \cdot \text{mol}^{-1}$) than that of the other fused imidazo derivative **15** and **16** ($\Delta H = 119.15$ and 106.26 Kcal \cdot mol⁻¹, respectively) facilitated its formation (Table 2). The imidazo-thiazines **14** and **16** were found to be more stable than the imidazo-thiazole which is reflected in the higher yields of **14** and **16** compared to **15** which was accompanied with a recovered starting material.

The energy difference ($\Delta E_1 - 71.09$ and $\Delta E_3 - 64.88$ Kcal·mol⁻¹) between the mono-S-alkylated products **3** and **6** and their corresponding fused rings were higher than that between the mono alkylated derivatives and the bis-alkylated analogues ($\Delta E_2 = 46.67$ and $\Delta E_4 = 51.54$ Kcal·mol⁻¹,

Compd. No.	Heat of formation ΔH (Kcal. mol ⁻¹)	Energy difference ΔE (Kcal. mol ⁻¹)
3	48.06	
4	1.39	$\Delta E_1 = \Delta H(3) - \Delta H(4) = 46.67$
15	119.15	$\Delta E_2 = \Delta H(3) - \Delta H(15) = -71.09$
6	41.38	
7	-10.16	$\Delta E_3 = \Delta H(6) - \Delta H(7) = 51.54$
16	106.26	$\Delta E_4 = \Delta H(6) - \Delta H(16) = -64.88$
12	-2.95	
13	-98.02	$\Delta \mathbf{E}_5 = \Delta \mathbf{H}(12) - \Delta \mathbf{H}(13) = 95.07$
14	59.69	$\Delta \mathbf{E}_6 = \Delta \mathbf{H}(12) - \Delta \mathbf{H}(14) = -62.64$

TABLE 2 Heat of formation of products

respectively). These results explain the formation of the bis-alkylated **4** and **7** as sole products from the reaction of **1** or **3** and **6** with bromoethanol (**2**) or chloropropanol (**5**) in presence of potassium carbonate in boiling DMF.

In contrast, the energy difference between the mono-alkylated **12** and its corresponding imidazo-thiazine ($\Delta E_6 = -62.64 \text{ Kcal.mol}^{-1}$) is lower than that between **12** and the bis-alkylated **13** ($\Delta E_5 = 95.07 \text{ Kcal.mol}^{-1}$). On the basis of these results, one could predict that the imidazo-thiazine **14** was favored to be formed rather than the bis-alkylated product but the higher stability of **12** has altered the energetic behavior of this reaction where both of the bis-alkylated analogue **13** and the imidazo-thiazine **14** were formed. These conclusions were in agreement with the experimental results.

CONCLUSIONS

A regioselective alkylation of 4,5-diphenylimidazole-2-thiol (1) with hydroxyl-alkylating agents in the presence of NaOEt or NaOAc has been achieved successfully to give only the respective S-alkylated derivatives, via the formation of the sodium salt of the thiolate anion generated by proton abstraction from the thiol group. Further alkylation of the S-alkylated analogues in boiling DMF and in the presence of potassium carbonate afforded the acyclonucleoside analogues. This could presumably be due to the ability of K_2CO_3 to abstract the proton from NH group which is not the case with NaOEt or NaOAc. These conditions were used to develop a general method for the synthesis of the fused imidazo-thiazines or thiazolines via the cyclization of **4**, **6**, and **12** with K_2CO_3 /DMF.

A comparison of the results from using MWI with that under conventional method revealed that higher yields were obtained in shorter times. Better results were obtained when bentonite was used as solid support. Thus, economical and environmental impacts can be achieved when the MW irradiation was combined with the solid support. The results were supported by the semiemoerical AM1 theoritical calculations.

EXPERIMENTAL

General Procedures

Melting points were determined with a Melt-Temp apparatus and are uncorrected. TLC was performed on Baker-Flex silica gel (Merck) using ethyl acetate-hexane as developing solvents, and the spot were detected by UV light absorption. Irradiation was done in a domestic microwave oven EM-230M (1200 watt output power). IR spectra were recorded with Perkin-Elmer 1430 spectrometer. The NMR spectra were recorded on Jeol spectrometer (500 MHz). Chemical shifts δ are given in ppm relative to the signal for TMS as internal standard. The double quatum filteredcorrelation spectroscopy DQF-COSY and heteronuclear multiple quantum correlation HMQC experiments were used to confirm the spectral analysis. The elemental analyses were performed by the microanalysis unit at the Faculty of Science, Cairo University.

General Procedure for the Alkylation 4,5-Diphenylimidazole-2-thione (1)

Conventional Method (CM). To a solution of compound 1, 3, 6, 9, or 12 (1 mmol) in a solvent (15 mL) and base (1.1 or 2.2 mmol), the appropriate hologeno-ethanols 2 or 3, chloroglycerol 11, and 2,3-O-isopropylidene-1-O-(p-tolylsulfonyl)-glycerol (8) (1.1 mmol or 2.2 mmol) were added with stirring. The conditions of the reaction are shown in Table 1. The reaction mixture was poured onto crushed ice. The products were washed with water, dried, and recrystallized from ethanol.

Microwave Method (MW). A mixture of compound 1, 3, 6, 9, or 12 (0.36 mmol), solvent (3 mL), base (0.36 or 0.73 mmol), and the appropriate alkylating agents (0.33 mmol or 0.73 mmol) were placed in a closed Teflon vessel and then irradiated by MW. The reaction mixture was processed as described above (Table 1).

Microwave and Solid Support Method. A mixture of 1, 3, 6, 9, or 12 (0.36 mmol) and an appropriate alkylating agent (0.36 mmol or 0.73 mmol) and bentonite (0.3 g) were mixed uniformly. The mixture was placed in a closed Teflon vessel and then irradiated by MW for 1–3 minutes. After cooling the products were extracted by boiled ethanol and were recrystallized from ethanol.

2-[(2-Hydroxyeth-1-yl)thio]-4,5-diphenyl-1*H***-imidazole (3).** This compound was obtained as colorless crystals, m.p: 168–170°C, IR (KBr): 3087 (CH-Ar), 3413 (OH). ¹H NMR (DMSO-d₆) δ : 3.17 (dd, 2H, *J* = 6.1 Hz, *J* = 6.9 Hz, SCH₂), 3.67 (ddd, 2H, *J* = 6.1 Hz, *J* = 6.9 Hz, *J* = 11.9 Hz, CH₂O), 5.13 (br, 1H, D₂O exchangeable, OH), 7.15–7.43 (m, 10H, Ar-H), 12.54 (s, 1H, D₂O exchangeable, NH). *Anal. Calcd.* for C₁₇H₁₆N₂OS (296.10): C, 68.89; H, 5.44; N, 9.45. Found: C, 68.79; H, 5.65; N, 9.25.

1-(2-Hydroxyeth-1-yl)-2-[(2-hydroxyeth-1-yl)thio]-4,5-diphenylimidazole (4). This compound was obtained as colorless crystals; m.p: 119–120°C, IR (KBr): 3410 (OH). ¹H NMR (DMSO-d₆) δ : 3.12 (br, 1H, D₂O exchangeable, OH), 3.24 (t, 2H, J = 4.6 Hz, SCH₂), 3.65 (dd, 2H, J = 5.3 Hz, J = 6.1 Hz, NCH₂), 3.97 (dd, 2H, J = 5.3 Hz, J = 6.1 Hz, CH₂O), 4.05 (t, 2H, J = 4.6 Hz, CH₂O), 4.42 (br, 1H, D₂O exchangeable, OH), 7.15–7.43 (m, 10H, Ar-H). *Anal. Calcd.* for C₁₉H₂₀N₂O₂S (340.12): C, 67.03; H, 5.92; N, 8.23. Found: C, 66.97; H, 5.61; N, 8.03.

2-[(3-Hydroxyprop-1-yl)thio]-4,5-diphenyl-1*H***-imidazole (6).** This compound was obtained as colorless crystals; m.p: 158–160°C, IR (KBr): 1585

(C = C), 3059 (CH-Ar), 3390 cm⁻¹ (OH). ¹H NMR (DMSO-d₆) δ : 1.74–1.79 (m, 2H, CH₂CH₂CH₂), 3.12 (t, 2H, J = 6.9 Hz, SCH₂), 3.49 (t, 2H, J = 6.1 Hz, CH₂O), 4.72 (br, 1H, D₂O exchangeable, OH), 7.20–7.32 (m, 10H, Ar-H), 12.55 (s, 1H, D₂O exchangeable, NH). *Anal. Calcd.* for C₁₈H₁₈N₂OS (310.11): C, 69.65; H, 5.84; N, 9.02. Found: C, 69.76; H, 6.03; N, 8.67.

1-(3-Hydroxyprop-1-yl)-2-[(3-hydroxyprop-1-yl)thio]-4,5-diphenylimidazole (7). This compound was obtained as yellow crystals, m.p: 145°C; IR (KBr): 3438 (OH). ¹H NMR (DMSO-d₆) δ: 1.54–1.59 (m, 2H, SCH₂CH₂CH₂) 1.82–1.85 (m, 2H, NCH₂CH₂CH₂), 3.19–3.22 (m, 4H, SCH₂, NCH₂), 3.50–3.52 (m, 2H, CH₂O), 3.76–3.79 (m, 2H, CH₂O), 4.46–4.49 (br, 1H, D₂O exchangeable, OH), 4.68–4.71 (br, 1H, D₂O exchangeable, OH), 7.05–7.47 (m, 10H, Ar-H). *Anal. Calcd.* for C₂₁H₂₄N₂O₂S (368.16): C, 68.45; H, 6.56; N, 7.60. Found: C, 68.32; H, 6.43; N, 7.89.

2-[(**2**,**3**-*O*-**Isopropylidene-2**,**3**-dihydroxyprop-1-yl)thio]-4,5-diphenyl-1*H*imidazole (9). This compound was obtained as colorless crystals; m.p: 159–160°C, IR (KBr): 3022 cm⁻¹ (C-H Ar). ¹H NMR (CDCl₃) δ : 1.40, 1.44 (2 s, 6H, 2 × CH₃), 3.20 (dd, 1H, J = 6.9 Hz, J = 14.5 Hz, SCH₂), 3.27 (dd, 1H, J = 4.6 Hz, J = 14.5 Hz, SCH₂), 3.80 (dd, 1H, J = 7.7 Hz, J = 8.4 Hz, CH₂O), 4.16 (dd, 1H, J = 6.1 Hz, J = 8.4 Hz, CH₂O), 4.48–4.50 (m, 1H, CHO), 7.26–7.48 (m, 10H, Ar-H) 12.60 (s, 1H, D₂O exchangeable, NH). ¹³C NMR (CDCl₃) δ : 25.7 (CH₃), 26.8 (CH₃), 36.8 (SCH₂), 68.4 (*CH*₂O), 77.1 (*CHO*), 110.1 (*C*(CH₃)₂), 127.6, 128.7, 140.14 (Ar-C). *Anal. Calcd.* for C₂₁H₂₂N₂O₂S (366.14): C, 68.82; H, 6.05; N, 7.64. Found: C, 68.50; H, 6.22; N, 7.98.

1-(2,3-*O*-Isopropylidene-2,3-dihydroxyprop-1-yl)-2-[(2,3-*O*-isopropylidene-2,3-dihydroxyprop-1-yl)thio]-4,5-diphenylimidazole (10). This compound was obtained as colorless crystals, m.p: 180–182°C; ¹H NMR (DMSO-d₆) δ : 1.19, 1.20, 1.21, 1.27 (4 s, 12H, 4 × CH₃), 3.39 (dd, 2H, J = 3.1 Hz, J = 9.2Hz, SCH₂), 3.73 (dd, 1H, J = 2.3 Hz, J = 5.3 Hz, J = 9.2, CH₂O(a)), 3.90 (dd, 1H, J = 5.3Hz, J = 9.2, CH₂O(b)), 3.99–4.09 (m, 2H, 2 × CH₂O(a), CH₂O(b)), 4.21 (dd, 1H, J = 5.3 Hz, J = 14.5 Hz, NCH₂), 4.28–4.32 (m, 1H, NCH₂), 4.33–4.46 (m, 2H, 2 × CHO), 7.18–7.41 (m, 10H, Ar-H). *Anal. Calcd.* for C₂₇H₃₂N₂O₄S (480.21): C, 67.47; H, 6.71; N, 5.83. Found: C, 67.68; H, 6.91; N, 5.90.

2-[(2,3-Dihydroxyprop-1-yl)thio]-4,5-diphenyl-1*H***-imidazole** (12). This compound was obtained as colorless crystals; m.p: 148°C, IR (KBr): 3096 (C-H Ar), 3314 (br OH). ¹H NMR (DMSO-d₆) δ : 3.14 (dd, 1H, J = 6.1 Hz, J = 13.7 Hz, SCH₂), 3.31 (dd, 1H, J = 6.1 Hz, J = 13.7 Hz, SCH₂), 3.35–3.43 (m, 2H, CH₂O), 3.72–3.75 (m, 1H, CHO), 4.98 (t, 1H, J = 5.3 Hz, D₂O exchangeable, OH), 5.45 (br, 1H, D₂O exchangeable, OH), 7.15–7.43 (m, 10H, Ar-H), 12.61 (s, 1H, D₂O exchangeable, NH). ¹³C NMR (DMSO-d₆) δ : 37.2 (SCH₂), 64.5 (CH₂O), 71.8 (CHO), 127.2, 127.4, 128.2, 128.8, 129.2,

131.1, 135.1, 137.2 (Ar-C). *Anal. Calcd.* for C₁₈H₁₈N₂O₂S (326.11): C, 66.23; H, 5.56; N, 8.58. Found: C, 65.98; H, 5.86; N, 8.41.

1-(2,3-Dihydroxyprop-1-yl)-2-[(2,3-dihydroxyprop-1-yl)thio-4,5-diphenylimidazole (13). This compound was obtained as colorless needless after purification by column chromatography using EtOAc-Hexane (40–60) as eluant; m.p: 125–126°C; IR (KBr): 1565 (C = C), 1610 (C = N), 3463 (OH). ¹H NMR (DMSO-d₆) δ : 3.20 (ddd, 1H, J = 5.4 Hz, J = 7.7 Hz, J = 10.7Hz, SCH₂), 3.33–3.43 (m, 5H, SCH₂, 2 × CH₂O), 3.68–3.75 (m, 1H, CHO), 3.84–3.90 (m, 1H, CHO), 4.05 (dd, 1H, J = 8.4 Hz, J = 14.5 Hz, NCH₂), 4.24 (dd, 1H, J = 3.8 Hz, J = 14.5 Hz, NCH₂), 4.80 (t, 1H, J = 6.1 Hz, D₂O exchangeable, OH), 4.90 (t, 1H, J = 6.2 Hz, D₂O exchangeable, OH), 5.17 (d, 1H, J = 4.5 Hz, D₂O exchangeable, OH), 5.21 (dd, 1H, J = 3.1 Hz, J =5.3 Hz, D₂O exchangeable, OH), 7.16–7.45 (m, 10H, Ar-H). Anal. Calcd. for C₂₁H₂₄N₂O₄S (400.15): C, 62.98; H, 6.04; N, 6.99. Found: C, 63.19; H, 5.95; N, 6.68.

Deisopropylidenation of 9 and 10. General procedure. The isopropylidenes **9** and **10** (5 mmol) were dissolved in 70% AcOH (5 mL). The mixture was heated under reflux for 2 hours. The solvent was evaporated under reduced pressure and the resulting products were collected and crystallized from ethanol to give **12** and **13**, respectively. The products were identical with those prepared by the above method.

Isopropylidenation of 12 and 13. General procedure. Compound 12 and 13 (0.25 mmol) were stirred vigorously with dry acetone (10 mL) and 96% H₂SO₄ (3 drops) for 2 hours, and then kept for overnight at room temperature. The resulting mixture was neutralized by Na₂CO₃, filtered, and the inorganic salts were well washed with dry acetone. The filtrate was evaporated under reduced pressure and the resulting products 9 and 10 were crystallized from ethanol or purified by column chromatography to give identical products with those obtained by the above alkylation method.

General Procedure for the Synthesis of Imidazothiazines and Thiazole

Conventional method (CM). A mixture of **3**, **7**, or **12** (1 mmol) and potassium carbonate (1.5 mmol) in DMF (15 mL) was heated under reflux. The reaction mixture was then poured onto crushed ice (10 mL). The product was filtered out and recrystallized from ethanol.

Microwave Method (MW). A mixture of **6**, **7**, or **12** (0.33 mmol) and potassium carbonate (0.5 mmol) in DMF (3 mL) in a closed Teflon vessel was irradiated by MWI. The obtained reaction mixture was treated as described above (Table 1).

6,7-Diphenyl-2*H*,3*H*,4*H*-3-hydroxy-tetrahydroimidazo[2,1-*b*][1,3]thiazine (14). This compound was obtained as colorless crystals, m.p: 210–212°C; IR (KBr): 3269 cm⁻¹ (OH). ¹H NMR (DMSO-d₆) δ : 3.11 (dd, 1H, *J* = 2.3 Hz, J = 12.6 Hz, SCH₂), 3.26 (dd, 1H, J = 6.3 Hz, J = 13.0 Hz, SCH₂), 3.55 (dd, 1H, J = 6.5 Hz, J = 12.6 Hz, NCH₂), 3.76 (dd, 1H, J = 3.4 Hz, J = 12.6 Hz, NCH₂), 4.26–4.30 (m, 1H, CHO), 5.61 (br, 1H, D₂O exchangeable OH), 7.18–7.41 (m, 10H, Ar-H). *Anal. Calcd.* for C₁₈H₁₆N₂S (308.40): C, 70.10; H, 5.23; N, 9.08. Found: C, 69.92; H, 5.39; N, 9.36.

5,6-Diphenyl-2,3-dihydroimidazo[2,1-*b***][1,3]thiazole (15).** This compound was obtained as colorless plates, m.p: 232–234°C; ¹H NMR (DMSO-d₆) δ : 3.82 (t, 2H, J = 8.4, SCH₂), 4.10 (t, 2H, J = 8.4, NCH₂), 7.15–7.23 (m, 2H, Ar-H), 7.34–7.40 (m, 5H, Ar-H), 7.49–7.54 (m, 3H, Ar-H). *Anal. Calcd.* for C₁₇H₁₄N₂S (278.37): C, 73.35; H, 5.07; N, 10.06. Found: C, 73.21; H, 4.93; N, 10.19.

6,7-Diphenyl-2*H*,3*H*,4*H*-tetrahydroimidazo[2,1-*b*][1,3]thiazine (16). This compound was obtained as colorless plates, m.p. 272°C; ¹H NMR (DMSO-d₆) δ : 1.92–1.95 (m, 2H, CH₂CH₂CH₂), 3.46 (t, 2H, *J* = 6.1, SCH₂), 3.83 (t, 2H, *J* = 5.3, NCH₂), 7.15–7.23 (m, 2H, Ar-H), 7.34–7.40 (m, 5H, Ar-H), 7.49–7.54 (m, 3H, Ar-H). *Anal. Calcd.* for C₁₈H₁₆N₂S (292.40): C, 73.94; H, 5.52; N, 9.58. Found: C, 74.07; H, 5.83; N, 9.39.

REFERENCES

- (a) El Ashry, E.S.H.; El Kilany, Y. Acyclonucleosides: Part 1. Seco-Nucleosides. Adv. Heterocyc. Chem. 1996, 67, 391–438; (b) El Ashry, E.S.H.; El Kilany, Y. Acyclonucleosides: Part 2. diseco-Nucleosides. Adv. Heterocyc. Chem. 1997, 68, 1–88; (c) El Ashry, E.S.H.; El Kilany, Y. Acyclonucleosides: Part 3. tri-,tetra-, and pentaseco-Nucleosides. Adv. Heterocyc. Chem. 1998, 9, 129–215.
- Shaeffer, H.J.; Beauchamp, L.; de Miranda, P.; Elion, G.B.; Bauer, D.J.; Collins, P. 9-(2-Hydroxyethoxymethyl) guanine activity against viruses of the herpes group. *Nature* (London) 1978, 272, 583–585.
- Martin, J.C.; Dvorak, C.A.; Smee, D.F.; Matthews, T.R.; Julien, P.H.; Verheyden, J.P. 9-[(1,3-Dihydroxy-2-propoxy)methyl]guanine: A new potent and selective antiherpes agent. *J. Med. Chem.* 1983, 26, 759–761.
- Hardern, M.R.; Jarvest, R.L.; Bacon, T.H.; Boyd, M.R. Synthesis and antiviral activity of 9-[4-hydroxy-3-(hydroxymethyl)but-1-yl]purines. J. Med. Chem. 1987, 39, 1636–1642.
- De Clercq, E.; De Somer, P.; Holý, A.; Descampus, J. (S)-9-(2,3-Dihydroxypropyl) adenine: an aliphatic nucleoside analog with broad-spectrum antiviral activity. *Science* 1978, 200, 563–565.
- Chu, C.K.; Cutler, S.J. Chemistry and antiviral activities of acyclonucleosides. J. Heterocyc. Chem. 1986, 23, 289–319.
- El Ashry, E.S.H.; Rashed, N.; Awad, L.F.; Abdel-Rahman, A.A.-H.; Rasheed, H.A. Synthesis of AZT analogues: 7-(3-azido-2-hydroxpropyl)-, 7-(3-Amino-2-hydroxypropyl)-, 7-(3-triazolyl-2-hydroxypropyl) theophyllines. *Nucleosides Nucleoides Nucleic Acids* 2006, 25, 299–305.
- (a) Lin, T.-S.; Liu, M.-C. Synthesis of 9-(2,3-dihydroxy-1-propoxymethyl)guanine-a new potential antiviral agent. *Tetrahedron Lett.* 1984, 25, 905–906; (b) Ashton, W.T.; Canning, L.F.; Reynolds, G.F.; Tolman, R.L.; Karkas, J.D.; Liou, R.; Davies, M.E.M.; Dewitt, C.M.; Perry, H.C.; Field, A.K. Synthesis and antiherpetic activity of (S)-, (R)-, and (±)-9-[(2,3-dihydroxy-1-propoxy)methyl]guanine, linear Isomers of 2'-N or-2'-deoxyguanosine. *J. Med. Chem.* 1985, 28, 926–933.
- El Ashry, E.S.H.; Rashed, N.; Awad, L.F.; Abdel-Rahman, A.A.-H.; Rasheed, H.A. Synthesis of new 7-alkylated theophyllines by chemical modification of dyphylline. *Journal of Chemical Research (S)* 2001, 2, 129–130. *J. Chem. Res. (M)* 2001, 440–450.
- El Ashry, E.S.H.; Abdel Rahman, A.A.-H.; Rashed, N.; Rasheed, H.A. Homoacylovir analogues of unnatural bases and their activity against hepatitis B virus. *Pharmazie* 1999, 54, 893–897.
- 11. El Ashry, E.S.H.; Abdel-Rahman, A.A.H.; Rashed, N.; Rasheed, H.A. Synthesis and anti-hepatitis B

virus activity of some 2,3-dihydroxyprop-1-yl unnatural hetaryls. Arch. Pharm. Med. Chem. 1999, 332, 327-330.

- Fenske, D.C.; Kuo, E.A.; Tully, R.W. Preparation and formulation of imidazoles, imidazopyridines, and imidazopyrimidines as inflammation inhibitors. U.K. Patent 1989, 2,193,962; *Chem. Abstr.* 1988, 109, 170425d.
- Renth, E.O.; Schromm, K.; Anderskewitz, R.; Birke, F.; Fuegner, A.; Heuer, H. Heterocyclecontaining amidine derivatives, their preparation and use as LTB₄ antagonist. German Offen Patent 1994, 4,309,285; *Chem. Abstr.* 1995, 122, 81416c.
- Chiesi, P.; Servadio, V.; Razzetti, R.. Thiomethyl derivatives having gastric acid antisecretory activity, processes for their preparation, and pharmaceutical compositions containing them. European Patent 1989, 301,422; *Chem. Abstr.* 1989, 111, 225320q.
- Nishi, T.; Uno, T.; Koga, Y.; Chu, G.N. Preparation of heterocyclic carbostyril derivatives as Inhibitors of thrombocyte adhesion. European Patent 1987, 240,015; *Chem. Abstr.* 1988, 108, 167318v.
- El Ashry, E.S.H.; Ramadan, E.; Kassem, A.A.; Haggar, M. Microwave irradiation for accelerating organic reactions. Part 1: Three, four and five membered heterocycles. *Adv. Heterocyc. Chem.* 2005, 88, 1–110.
- (a) El Ashry, E.S.H.; Kassem, A.A.; Ramadan, E. Microwave irradiation for accelerating organic reactions. Part 2: six, seven fused and spiro heterocyclic ring Systems. *Adv. Heterocyc. Chem.* 2006, 90, 1–127; (b) El Ashry, E.S.H.; Kassem, A.A. Account of microwave irradiation for accelerating organic reactions. *Arkivoc* 2006, ix, 1–15.
- (a) Kappe, C.O. High-speed combinatorial synthesis utilizing microwave irradiation. *Curr. Opin. Chem. Biol.* 2002, 6, 314–320; (b) Stadler, A.; Kappe, C.O. High-speed couplings and cleavages in microwave-heated, solid-phase reactions at high temperatures. *Eur. J. Org. Chem.* 2001, 919–925.
- Hamelin, J.; Bazureau, J.P.; Texier-Boullet, F. In *Microwave in Organic Synthesis*. ed., A. Loupy, Wiley-VCH, Weinheim, 2002, 253–293.
- Perreux, L.; Loupy, A.A. Tentative rationalisation of microwave effects in organic synthesis according to the reaction medium, and mechanistic considerations. *Tetrahedron* 2001, 57, 9199–9223.
- Lidström, P.; Tierney, J.; Wathey, B.; Westman, J. Microwave assisted organic synthesis. *Tetrahedron* 2001, 57, 9225–9283.
- Perreux, L.; Cotterill, Ic.; Usyatinsky, I.Y.; Arnold, J.M.; Clark, D.S.; Dordick, J.S.; Michels, P.C.; Khmelnitsky, Y.L. Microwave assisted combinatorial chemistry. Synthesis of substituted pyridines. *Tetrahedron Lett.* **1998**, 39, 1117–1120.
- Larhed, M.; Hallberg, A. Microwave-assisted high-speed chemistry a new techniques in drug discovery. *Drug Disc. Today* 2001, 6, 406–416.
- Bougrin, K.; Loupy, A.; Soufiaoui, M. Microwave-assisted solvent-free heterocyclic synthesis. J. Photochem. Photobiol. C 2005, 6, 139–167.
- Abdel-Rahman, A.-H.; El Ashry, E.S.H. Efficient synthesis of 5-hydroxymethyl pyrimidines and their nucleosides using microwave irradiation. *Synlett* 2002, 2043–2044.
- Abdel Hamid, H.M.; Ramadan, E.; Hagar, M.; El Ashry, E.S.H. Synthesis of aryloxyacetic acids, esters and hydrazides assisted by microwave irradiation. *Synth. Comm.* 2004, 34, 377–382.
- El Ashry, E.S.H.; Ramadan, E.; Abdel Hamid, H.M.; Hagar, M. Synthesis of azalactones, phenylpyruvic acid and 1,2,4-triazine derivatives under microwave irradiation. *Lett. Org. Chem.* 2004, 2, 415–418.
- El Ashry, E.S.H.; Ramadan, E.; Abdel Hamid, H.; Haggar, M. Microwave irradiation for enhancing the synthesis of quinoline derivatives from isatin. *Synth. Comm.* 2005, 35, 2243–2250.
- El Ashry, E.S.H.; Ramadan, E.; Abdel-Hamid, H.; Hagar, M. Microwave irradiation for enhancing the regioselective synthesis of 6*H*-indolo[2,3-*b*]quinoxaline. *J. Chem. Re.* 2005, 229–232.
- El Ashry, E.S.H.; Ramadan, E.; Abdel-Hamid, H.; Hagar, M. Microwave irradiation for accelerating the synthesis of 1,2,4-triazino[5,6-b]indole-2-thiols and their derivatives from isatin and 5chloroisatin. Synlett 2004,, 723–725.
- El Ashry, E.S.H.; Awad, L.F.; Abdel-Hamid, H.; Atta, A.I. Synthesis of interglycosidically S-linked 1-thio-oligosaccharides under microwave irradiation. J. Carb. Chem. 2005, 24, 745–753.
- El Ashry, E.S.H.; Awad, L.F.; Abdel-Hamid, H.; Atta, A.I. Microwave-assisted organic synthesis of 3-(D-gluco-pentitol-1-yl)-1*H*-1,2,4-triazole. *Nucleosides Nucleotides Nucleic Acids* 2005, 25, 325–335.
- El Ashry, E.S.H.; Awad, L.F.; Abdel-Hamid, H.; Atta, A.I. Int. Soc. For Nucleosides Nucleotides Nucleotides Nucleotides XVIInt. Round Table, Minneaoplis, Minnesota, USA, (2004) PS. Microwave Irradiation for accelerating the synthesis of acyclonucleosides of 1,2,4-triazole. *Nucleosides Nucleotides Nucleic Acids* 2005, 24, 427–429.

- 34. El Ashry, E.S.H.; Rashed, N.; Awad, L.F.; Ramadan, E., Abdel-Mageed, S.M.; Rezki, N. MAOS Vversus conventional synthesis of 4,5-di- and 3,4,5-triphenylimidazole-2-thione and their derivatives. *ARKIVOC* 2007, 7, 30–40.
- 35. El Ashry, E.S.H.; Kassem, A.A.; Abdel-Hamid, H.; Louis, F.F.; Khattab, Sh.A.N. Aouad, M. R. Synthesis and alkylation of 5-(3-chloro benzo[b]thien-2-yl)-4H-1,2,4-triazole-3-thiol under classical and microwave conditions. AM1 Semiemperical calculations for investigating the regioselectivity of alkylation. J. Heterocyc. Chem. 2006, 43, 1421–1429.
- 36. (a) Gürsoy, A.; Demirayak, Ş.; Cesur, Z.; Reisch, J.; Ötük, G. Synthesis of some new hydrazinehydrazone, thiosemicarbazide, thiadiazoles, triazoles and their derivatives as possible antimicrobials. Pharmazie **1990**, *45*, 246–250; (b) Biltz, H.; Krebs, P. 4,4',5.5'-Tetrasubstituierte di-2-imidazolylderivative. *Liebegs Annalen Chemie* **1912**, 391, 191–194.
- 37. Mitsunobu, O. The use of diethyl azodicaboxylate and triphenylphosphine in synthesis and transformation of natural products. *Synthesis* **1981**, 1–28.
- Janeba, Z.; Holy, A. Masojidkova, M. Synthesis of acyclic nucleoside and nucleotide analogs derived from 6-amino-7*H*-purine-8(9*H*)-thione. *Coll. Czech Chem. Comm.* 2000, 65, 1698–1712.