

## *Exo,Exo*-2,3-Diaminoborneol-Derived Imidazolidinone as Chiral Auxiliary for Asymmetric Alkylations

Claudio Palomo<sup>a\*</sup>, Mikel Oiarbide<sup>a</sup>, Alberto González<sup>b</sup>, Jesús M. García<sup>b</sup> and Fabienne Berrée<sup>a</sup>

<sup>a</sup>Dpto. de Química Orgánica. Facultad de Química. Universidad del País Vasco. Apartado: 1072. 20080-San Sebastián. Spain.

<sup>b</sup>Departamento de Química. Universidad Pública de Navarra. Campus de Arrosadía. 31006-Pamplona. Spain


**Abstract:** A new camphor-derived imidazolidinone type auxiliary is described. Its acylated derivatives under Na enolate formation and treatment with primary halides give alkylated products in highly diastereoselective manner. Copyright © 1996 Elsevier Science Ltd

The use of the Evans  $\alpha$ -amino acid-derived *N*-acyloxazolidinones for asymmetric C-C bond construction is one of the most straightforward and efficient strategies in modern organic synthesis.<sup>1</sup> Despite exhibiting high levels of diastereoselectivity in numerous transformations, the absolute stereochemical control in reactions of particularly poor diastereoface selection continues to be an important challenge. A recent report from this laboratory has addressed this issue and high levels of asymmetric induction were observed in reactions involving *N*-acylimides derived from *exo,exo*-2-amino-3-borneol **1**.<sup>2</sup> The design of **1** was inspired by the results achieved with the *tert*-leucine derived oxazolidinone **2**, and by the advantageous topological bias of camphor skeleton to form conformationally rigid derivatives.<sup>3</sup> Herein we wish to report on the preparation and synthetic utility of the imidazolidinone **5**, Scheme I, in asymmetric enolate alkylations.<sup>4</sup> The reagent **5**, which is closely related to **1**, has been delineated with the hope that the phenyl chromophore present in the imidazolidinone ring could facilitate reaction monitoring and/or product isolation.




The imidazolidinone **5** was easily prepared in simple steps from the monoimine **3** derived from (1*R*)-(-)-camphorquinone.<sup>5</sup> When **3** was treated with benzylamine and TiCl<sub>4</sub> followed by reduction of the

resulting diimine with  $\text{NaBH}_4$  the diamine **4** was obtained in 70% overall yield.<sup>6</sup> Subsequent *N*-debenzylation and further treatment with triphosgene produced the desired *exo,exo*-imidazolidinone **5** in 80% yield.<sup>7</sup>



Scheme I

To test the chemical and stereochemical behaviour of imidazolidinone **5** for asymmetric synthesis, we evaluated the alkylation of *N*-acylimides **6-8** with representative alkyl halides. The enolates **9-11** were generated by addition of  $\text{NaHMDS}$  to a solution of the corresponding *N*-acylimide under usual conditions<sup>8</sup> and after complete enolization, 1 h at  $-60^\circ\text{C}$ , a threefold excess of the alkylating agent was added at the same temperature. After the disappearance of the starting material which could be checked by TLC (1-3 h), the mixture was quenched with saturated  $\text{NH}_4\text{Cl}$  and product isolated.



Scheme II

As the results in Table 1 reveal methyl iodide, benzyl bromide and allyl bromide reacted with these enolates to give the respective alkylated products with excellent yields. In every case the analysis of the crude products by  $^{13}\text{C}$  NMR showed a unique set of signals. This observation was confirmed by HPLC analysis, where absorption bands appeared well resolved and typically showed a retention time gap larger than 20 seconds. On the other hand, although a somewhat lower yield was produced in the alkylation of **9** (entry a) with ethyl iodide to give **12**, the diastereoselection level was found to be virtually complete. In an effort to enhance the reactivity of imide enolate **9** the use of HMPA as cosolvent (entry b) was found to be

effective albeit in lower diastereoselectivity.<sup>9</sup> Nevertheless, the most striking feature of the chiral auxiliary **5** is that the sterically undemanding methylations and ethylations, which often are difficult to control, proceeded with remarkable diastereoselectivity. From these results it is clear that auxiliary **5** gives rise to the same excellent level of diastereoselection than does **1** in alkylation reactions, including some monitoring advantages as mentioned before.

Table 1. Diastereoselective Alkylations of Chiral Enolates **9-11**<sup>a</sup>

Entry	Enolate	R	R <sup>1</sup> X	Product	d.r. <sup>b</sup>	Yield, % <sup>c</sup>	mp °C	[ $\alpha$ ] <sub>D</sub> <sup>25</sup> <sup>d</sup>
a	<b>9</b>	CH <sub>3</sub>	EtI	<b>12</b>	>99:1	41	120-122	+20.4
b	<b>9</b>		EtI(HMPA)	<b>12</b>	92:8	73		
c	<b>9</b>		PhCH <sub>2</sub> Br	<b>13</b>	>99:1	88	oil	+11.8
d	<b>10</b>	CH <sub>2</sub> CH <sub>3</sub>	MeI	<b>14</b>	>99:1	90	73-75	+74.5
e	<b>10</b>		CH <sub>2</sub> =CHCH <sub>2</sub> Br	<b>15</b>	>99:1	79	98-100	+50.4
f	<b>11</b>	CH <sub>2</sub> Ph	MeI	<b>16</b>	>99:1	78	oil	+32.1

<sup>a</sup> Reactions conducted on a 1 mmol scale. Enolate formation at -60°C in THF as solvent during 1 h and alkylations performed at -60°C using 3 equiv. of alkyl halide, except for MeI (5 equiv.). <sup>b</sup> Diastereomeric ratio determined by HPLC using a Shimadzu LC-8A instrument (column Hibar Lichrosorb Si 60 7 $\mu$ m, Supelco). <sup>c</sup> Yields of isolated pure products, after flash-column chromatography (Silica gel, Hex:EtOAc 4:1). <sup>d</sup> Measured in CHCl<sub>3</sub> at c 1.0.

The sense of asymmetric induction can be interpreted by assuming alkylations occur predominantly at the relatively unhindered  $\alpha$ -face of the corresponding chelated enolates. The stereochemical course of these reactions was unambiguously confirmed by lithium aluminium hydride reduction of both **13** and **16** to afford carbinols **17** {[ $\alpha$ ]<sub>D</sub><sup>25</sup> +11.1° (c=1.15, C<sub>6</sub>H<sub>6</sub>); lit.<sup>10</sup> [ $\alpha$ ]<sub>D</sub><sup>25</sup> +11.0° (c=1.15, C<sub>6</sub>H<sub>6</sub>)} and **18** {[ $\alpha$ ]<sub>D</sub><sup>25</sup> -10.6° (c=1.15, C<sub>6</sub>H<sub>6</sub>)} in 87% and 85% yield respectively, along with the imidazolidinone **5** which could be separated by column chromatography and recycled.

#### ACKNOWLEDGEMENTS

This work was supported by Gobierno Vasco (Project PI9386) and in part by Diputación Foral de Gipuzkoa. A grant from MEC to F.Berrée is gratefully acknowledged.

#### REFERENCES AND NOTES

- For some comprehensive reviews, see: (a) Evans, D.A.; Nelson, J.V.; Taber, T.R. *Top. Stereochem.* **1982**, *31*, 1. (b) Evans, D.A. *Aldrichimica Acta* **1982**, *15*, 23. (c) Evans, D.A. in *Asymmetric Synthesis*; Morrison J.D. Ed.; Academic: New York, vol. 3, **1984**; p. 1; (d) Masamune, S.; Choy, W.; Petersen, J.S.; Sita, L.R. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 1. (e) Seebach, D. *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 1320. (f) Yamamoto, Y.; Sasaki, N. in *Stereochemical Control, Bonding and*

- Steric Rearrangements*; Bernal I., Ed.; Elsevier: Amsterdam, **1990**; p. 1. (g) Franklin, A.S.; Paterson, I. *Contemp.Org. Synth.* **1994**, *1*, 317.
- Palomo, C.; Berré, F.; Linden, A.; Villalgorido, J.M. *J. Chem. Soc.; Chem. Commun.* **1994**, 1861.
  - For a review on *tert*-leucine, see: Bonmarius, A.S.; Schwarm, M.S.; Stingl, K.; Kottenhanhn, M.; Huthmacher, K.; Drauz, K. *Tetrahedron Asymmetry* **1995**, *6*, 2851. For a review on camphor-derived chiral auxiliaries, see: Oppolzer, W. *Tetrahedron* **1987**, *43*, 1969.
  - For asymmetric enolate alkylations on N-acyl imidazolidin-2-ones, see, for example: (a) Cardillo, G.; Orena, M.; Romero, M.; Sandri, S. *Tetrahedron* **1989**, *45*, 1501. (b) Cardillo, G.; D'Amico, A.; Orena, M.; Sandri, S. *J. Org. Chem.* **1988**, *53*, 2354. (c) Orena, M.; Porzi, G.; Sandri, S. *Tetrahedron Lett.* **1992**, *33*, 3797. (d) Lotz, G.A.; Palacios, S.M.; Rossi, R.A. *Tetrahedron Lett.* **1994**, *35*, 7711. (e) Davies, S.G.; Evans, G.B.; Mortlock, A.A. *Tetrahedron Asymmetry* **1994**, *5*, 585. For an extensive list of references concerning other chiral auxiliaries, see: (f) Suder, A.; Hintermann, T.; Seebach, D. *Helv. Chim. Acta* **1995**, *78*, 1185.
  - (a) Saigo, K.; Yamamoto, M.; Morimura, K.; Nohira, H. *Chem. Lett.* **1979**, 545. (b) Denmark, S.E.; Rivera, I. *J. Org. Chem.* **1994**, *59*, 6887.
  - Spectroscopic data for compound **4** (CDCl<sub>3</sub>, δ ppm): <sup>1</sup>H NMR 7.41-6.50 (m, 10H, arom), 5.48 (s<sub>b</sub>, 1H, NH), 3.73 (s, 2H, CH<sub>2</sub>Ph), 3.29 (dd, 1H, J=8.4Hz, J'=3.8Hz, CHN), 2.88 (d, 1H, J=8.4Hz, CHN), 1.90 (d, 1H, J=4.5Hz, CH bridgehead), 1.80-1.70 (m, 1H, CH<sub>a</sub>H<sub>b</sub>), 1.79-1.55 (m, 1H, CH<sub>a</sub>H<sub>b</sub>), 1.42 (s<sub>b</sub>, 1H, NH), 1.40-1.15 (m, 2H, CH<sub>a</sub>H<sub>b</sub>), 1.09, 0.96 and 0.79 (s, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR 148.2, 140.4, 129.2, 128.5, 128.1, 127.0, 115.7, 112.4, 69.4, 60.2, 55.8, 48.9, 48.8, 47.0, 36.7, 25.6, 21.8, 21.0, 12.1
  - Preparation of **5**: H<sub>2</sub> was supplied by balloon for 12 h to a solution of **4** (18.3 g, 54 mmol) in MeOH (150 mL) in the presence of 10% Pd/C (1.83 g, 10% w/w). After filtration of the catalyst and evaporation of the solvent, to the oily residue dissolved in dry THF (350 mL) and cooled at 0°C were added successively under a N<sub>2</sub> atmosphere Et<sub>3</sub>N (37.8 mL, 270 mmol) and dropwise during 1 h a solution of (Cl<sub>3</sub>CO)<sub>2</sub>CO (8.02 g, 27 mmol) in THF (50 mL). The resulting solution was stirred at r.t. for 1h, solvent evaporated, the residue dissolved in CH<sub>2</sub>Cl<sub>2</sub> (300 mL), washed with H<sub>2</sub>O (2 x 100 mL) and brine (100 mL), dried over MgSO<sub>4</sub>, and solvent removed. The pale yellow solid thus obtained was purified by flash column chromatography (EtOAc/Hex 1:3) to afford pure compound **5** (8.37 g, 80%) which recrystallized from Et<sub>2</sub>O/Hex: mp 184-185°C; [α]<sub>D</sub><sup>25</sup> -9.9° (CH<sub>2</sub>Cl<sub>2</sub>, c 1.0); <sup>1</sup>H NMR (CDCl<sub>3</sub>, δ ppm) 7.51-7.00 (m, 5H, arom), 5.73 (s, 1H, NH), 4.31 (d, 1H, J=9.3Hz, CH), 3.59 (dd, 1H, J=9.3Hz, J'=2.2Hz, CH), 2.10 (d, 1H, J=4.9Hz, CH bridgehead), 1.84-1.71 (m, 1H, CH), 1.61-1.50 (m, 1H, CH), 1.17-1.03 (m, 2H, CH<sub>a</sub>H<sub>b</sub>), 1.07, 0.96 and 0.81 (s, 3H, CH<sub>3</sub>); <sup>13</sup>C NMR 159.3, 139.1, 128.8, 122.6, 119.0, 65.0, 62.3, 48.2, 46.7, 45.9, 33.4, 25.0, 23.3, 18.8, 10.9. Anal. Calcd. for C<sub>17</sub>H<sub>22</sub>N<sub>2</sub>O: C, 75.52%; H, 8.20%; N, 10.36%. Found: C, 75.47%; H, 8.67%; N, 10.20%.
  - Evans, D.A.; Ennis, M.D.; Mathre, D.J. *J. Am. Chem. Soc.* **1982**, *104*, 1737.
  - Less reactive alkyl bromides such as n-propyl bromide and isobutyl iodide afforded very low yields of the expected products. For a solution to this problem which is general for imide enolates, see: Evans, D.A.; Dow, R.L.; Shih, L.; Takacs, J.M.; Zahler, R. *J. Am. Chem. Soc.* **1990**, *112*, 5290.
  - Ahn, K.H.; Lim, A.; Lee, S. *Tetrahedron Asymmetry* **1993**, *4*, 2435.

(Received in UK 25 March 1996; revised 30 April 1996; accepted 3 May 1996)