## Synthesis and reactivity of triethylborane adduct of N-heterocyclic carbene: versatile synthons for synthesis of N-heterocyclic carbene complexes<sup>†</sup>

Yoshitaka Yamaguchi,\*<sup>a</sup> Taigo Kashiwabara,<sup>a</sup> Kenichi Ogata,<sup>a</sup> Yumiko Miura,<sup>a</sup> Yoshiyuki Nakamura,<sup>b</sup> Kimiko Kobayashi<sup>c</sup> and Takashi Ito\*<sup>a</sup>

- <sup>a</sup> Department of Materials Chemistry, Graduate School of Engineering, Yokohama National University, Tokiwadai, Hodogaya-ku, Yokohama 240-8501, Japan
- <sup>b</sup> Chemical Resources Laboratory, Tokyo Institute of Technology, Nagatsuda, Midori-ku, Yokohama 226-8503, Japan
- <sup>c</sup> The Institute of Physical and Chemical Research (RIKEN), Hirosawa, Wako, Saitama 351-0198, Japan

Received (in Cambridge, UK) 14th April 2004, Accepted 5th July 2004 First published as an Advance Article on the web 20th August 2004

The reaction of an imidazolium salt with  $LiBEt_3H$  afforded triethylborane adduct of imidazol-2-ylidene, which can act as a carbene precursor for the synthesis of a transition metal complex as well as a main group element complex.

N-Heterocyclic carbenes (NHCs) are an increasingly useful class of ligands for not only transition metals but also main group elements.<sup>1</sup> One of the most widely used methods is the thermal cleavage of the enetetramines, so-called electron-rich olefins, in the presence of metal species.<sup>2</sup> However the metal precursor and/or product of the NHC are occasionally decomposed due to the high temperature requirement for enetetramine cleavage. The successful isolation of the free NHCs by Arduengo and co-workers in 1991<sup>3</sup> afforded the most common method of direct complexation of the free NHC, either isolated or generated in situ, produced by deprotonation of the corresponding salt. However manipulation of the free carbenes was often difficult due to their highly reactive nature. Recently the employment of NHC adducts as "protected" forms of the free carbene attracted much attention for synthesis of the metal complexes bearing these ligands. Grubbs and co-workers reported the utilization of NHC-alcohol or -chloroform adducts for saturated NHCs (imidazolin-2-ylidenes),<sup>4</sup> and Lin *et al.* reported the carbene transmetallation of silver complexes for unsaturated NHCs (imidazol-2-ylidenes).<sup>5</sup> Recent studies have revealed that the latter method is reliable for the preparation of the unsaturated NHC complexes.

During the course of our research on NHC complexes, a BEt<sub>3</sub> adduct of NHC was obtained unexpectedly by the reaction of an imidazolium salt with  $LiBEt_3H$ ; the latter acts not only as a base for the generation of a carbene but also the protecting group for it. We herein report the preparation, characterization, and utilization of NHC·BEt<sub>3</sub> compounds.

Treatment of 1,3-diisopropylimidazolium chloride (I<sup>i</sup>Pr·HCl, 1a) with one equivalent of LiBEt<sub>3</sub>H in THF at -78 °C yielded a homogeneous reaction mixture from which pale yellow crystals of 2a were isolated in 80% yield. In the case of 1b (IMes·HCl), the 2,4,6-trimethylphenyl (mesityl) analogue of 1a, a similar reaction took place to give a pale yellow solid 2b (46%). Elemental analyses, and <sup>1</sup>H, <sup>13</sup>C, and <sup>11</sup>B NMR spectra established the formation of BEt<sub>3</sub>-adduct NHC compounds (2a and 2b) (Scheme 1). In the <sup>1</sup>H NMR spectra, the BEt<sub>3</sub> coordinated carbene carbon in both compounds displayed a triplet and a quartet for the methyl and methylene groups, respectively, in a region of high magnetic field (*ca.* -0.1 to 0.6 ppm). The <sup>13</sup>C NMR spectra showed the broad signals assignable to the carbene carbons at 174.0 ppm for 2a and at 180.4 ppm for 2b, respectively. The broadening may be caused by the influence of the binding boron. A singlet signal assignable to

† Electronic supplementary information (ESI) available: full experimental data. See http://www.rsc.org/suppdata/cc/b4/b405459h/

BEt<sub>3</sub> was observed at -12.6 ppm for **2a** and at -13.3 ppm for **2b**, respectively, in the <sup>11</sup>B NMR spectra. These chemical shifts are consistent with that of the reported carbene–BEt<sub>3</sub> compound.<sup>7</sup>

The X-ray analyses of 2a and 2b were undertaken.<sup>‡</sup> The structures of 2a and 2b are shown in Fig. 1 and are consistent with their spectroscopic data. The B–C(carbene) bond distances are 1.683(5) Å for 2a and 1.678(6) Å for 2b. The B–C bond distances in 2 are remarkably longer than those in the BH<sub>3</sub>-adduct of imidazol-2-ylidene compounds (1.596(4) Å for IMes·BH<sub>3</sub> and 1.603(3) Å for 1,3-diethyl-4,5-dimethylimidazol-2-ylidene·BH<sub>3</sub>),<sup>8</sup> and are slightly longer than those for the BF<sub>3</sub>-adduct of imidazol-2-ylidene (1.635(5) Å for IMes·BF<sub>3</sub> and 1.669(6) Å for 1,3-dimesityl-4,5-dichloroimidazol-2-ylidene·BF<sub>3</sub>).<sup>9</sup>

Recent study on the carbene adducts with Group 13 elements such as B, Al, Ga, and In revealed that these adducts have the remarkable thermal stability because of the formation of the stable acid–base adduct.<sup>10</sup> However, we focused attention on the elongated B–C(carbene) bond distance in the compounds 2, because, if the bond cleavage would occur under mild conditions, compounds 2 could be utilized as the precursor for the HNCs. So, we investigated the reaction of compounds 2 with some metal compounds (Scheme 2).

On treating 2a with BH3 THF complex, a clean substitution



Fig. 1 Molecular structures of 2a (left) and 2b (right). ORTEP diagrams drawn at 30% probability (upper) and space filling diagrams (bottom).

10.1039/b405459h

ä



reaction took place to give the BH<sub>3</sub>-adduct of I<sup>1</sup>Pr (3a). However, **3a** was so highly soluble toward hydrocarbons that a low isolated yield (31%) resulted. The similar reaction employing **2b** in the place of 2a also afforded the BH<sub>3</sub>-adduct 3b as a white powder in a good yield (93%).11 11B NMR spectra of 3a and 3b displayed quartet signals at -36.4 ppm ( ${}^{1}J_{BH} = 88.0$  Hz) and at -36.8 ppm ( ${}^{1}J_{BH} =$ 88.0 Hz), respectively, whose data are consistent with the reported data for the related compound.<sup>8</sup> Other spectroscopic data supported the formation of BH<sub>3</sub>-adducts 3. Secondly the reaction of BF3 ·OEt2 with 2b was examined, and thus the BF3 adduct of IMes (4b) was formed as a colorless solid. The characterization of 4b was done by <sup>1</sup>H and <sup>11</sup>B NMR measurements and these spectral data are consistent with the reported data by Arduengo.9 Treatment of 2a with Mo(CO)<sub>6</sub> in toluene under reflux conditions afforded a carbene complex of molybdenum 5a as a pale yellow solid in moderate yield (67%). In the case of 2b, Mo(CO)<sub>5</sub>(IMes) (5b) was also obtained as a pale yellow solid in 67% yield. These complexes were characterized by elemental analyses, IR, and <sup>1</sup>H and <sup>13</sup>C NMR spectra.<sup>12</sup>

It is of interest to compare the behavior of BH<sub>3</sub>- and/or BF<sub>3</sub>adducts toward Mo(CO)<sub>6</sub> to that of the BEt<sub>3</sub>-adduct with a view to find possible reagents capable of carbene transfer to transition metals. Treatment of BH<sub>3</sub> adduct **3b** with Mo(CO)<sub>6</sub> under the same conditions, *i.e.*, refluxing in toluene, afforded the carbene complex **5b**, accompanied by the formation of Mo( $\eta$ -toluene)(CO)<sub>3</sub> (6) (the ratio **5b/6** being *ca.* 1/1). On the other hand, from the reaction of Mo(CO)<sub>6</sub> with BF<sub>3</sub> adduct **4b**, the carbene complex **5b** was not formed and 82% of **4b** was recovered unreacted. These results clearly show that the BEt<sub>3</sub> adduct of NHC (**2**) effectively acts as a carbene precursor for a metal complex, which might be due to the weak Lewis acidity<sup>13</sup> and the steric bulk of BEt<sub>3</sub> compared with BF<sub>3</sub> and BH<sub>3</sub>.

In order to investigate whether this method is applicable to the bidentate system or not, we examined the reaction of xylyl bridged bis(imidazolium) salts (7a: *ortho*-form, 7b: *meta*-form) with two equivalents of LiBEt<sub>3</sub>H. In both reactions, BEt<sub>3</sub>-adducts of bis(imidazol-2-ylidene) (8a and 8b) were obtained as white powders in good yield (Scheme 3).

In summary, we showed the one-pot synthesis of BEt<sub>3</sub> adduct of NHC (2) by the reaction of imidazolium salt (1) with LiBEt<sub>3</sub>H. The manipulation of BEt<sub>3</sub> adducts of NHC (2) is very easy due



Scheme 3

to their stability toward air and moisture. We also demonstrated the utilization of these adducts for the preparation of metal complexes.

The authors are grateful to Professor Kohtaro Osakada and Dr Yasushi Nishihara (Tokyo Institute of Technology), and Dr Noriyuki Suzuki (RIKEN) for the kind help in the elemental analyses.

## Notes and references

‡ Crystal data for **2a**: C<sub>15</sub>H<sub>31</sub>BN<sub>2</sub>, M = 250.23, orthorhombic, a = 16.033(8), b = 9.904(8), c = 10.58(1) Å, U = 1680(2) Å<sup>3</sup>, T = 223 K, space group *Pnma* (no. 62), Z = 4,  $\mu$ (Mo-K $\alpha$ ) = 0.56 cm <sup>-1</sup>, 2788 measured reflections, 2575 unique reflections. *R*1 = 0.082, *wR*2 = 0.212, for 1754 reflections. For **2b**: C<sub>27</sub>H<sub>39</sub>BN<sub>2</sub>, M = 402.43, orthorhombic, a = 16.929(10), b = 17.29(1), c = 8.511(8) Å, U = 2490(3) Å<sup>3</sup>, T = 173 K, space group *P*<sub>2121</sub>(no. 19), Z = 4,  $\mu$ (Mo-K $\alpha$ ) = 0.61 cm<sup>-1</sup>, 4085 measured reflections, 4058 unique reflections. *R*1 = 0.062, *wR*2 = 0.173, for 3219 reflections. CCDC numbers: **2a**: 236673; **2b**: 236674. See http:// www.rsc.org/suppdata/cc/b4/b405459h/ for crystallographic data in .cif format.

- For a review of *N*-heterocyclic carbene complexes, see: W. A. Herrmann and C. Köcher, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2162–2187; D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39–91; T. Weskamp, V. P. W. Böhm and W. A. Herrmann, *J. Organomet. Chem.*, 2000, **600**, 12–22; W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1290–1309; A. H. Cowley, *J. Organomet. Chem.*, 2001, **617–618**, 105–109.
- M. F. Lappert, J. Organomet. Chem., 1988, 358, 185–214; M. F. Lappert,
  P. L. Pye and G. M. McLaughlin, J. Chem. Soc., Dalton Trans., 1977, 1272–1282; M. F. Lappert and P. L. Pye, J. Chem. Soc., Dalton Trans., 1977, 1283–1291; M. F. Lappert, P. L. Pye, A. J. Rogers and
  G. M. McLaughlin, J. Chem. Soc., Dalton Trans., 1981, 701–704;
  M. F. Lappert, J. Organomet. Chem., 1975, 100, 139–159.
- 3 A. J. Arduengo, III, R. L. Harlow and M. Kline, J. Am. Chem. Soc., 1991, **113**, 361–363; A. J. Arduengo, III, Acc. Chem. Res., 1999, **32**, 913–921.
- 4 M. Scholl, S. Ding, C. W. Lee and R. H. Grubbs, *Org. Lett.*, 1999, 1, 953–956; T. M. Trnka, J. P. Morgan, M. S. Sanford, T. E. Wilhelm, M. Scholl, T.-L. Choi, S. Ding, M. W. Day and R. H. Grubbs, *J. Am. Chem. Soc.*, 2003, 125, 2546–2558.
- 5 H. M. J. Wang and I. J. B. Lin, Organometallics, 1998, 17, 972-975.
- 6 A. R. Chianese, X. Li, M. C. Janzen, J. W. Faller and R. H. Crabtree, *Organometallics*, 2003, 22, 1663–1667; M. Frøseth, A. Dhindsa, H. Røise and M. Tilset, *Dalton Trans.*, 2003, 4516–4524; A. M. Magill, D. S. McGuinness, K. J. Cavell, G. J. P. Britovsek, V. C. Gibson, A. J. P. White, D. J. Williams, A. H. White and B. W. Skelton, *J. Organomet. Chem.*, 2001, 617–618, 546–560.
- 7 A. Wacker, H. Pritzkow and W. Siebert, *Eur. J. Inorg. Chem.*, 1998, 843–849.
- 8 T. Ramnial, H. Jong, I. D. McKenzie, M. Jennings and J. A. C. Clyburne, *Chem. Commun.*, 2003, 1722–1723; N. Kuhn, G. Henkel, T. Kratz, J. Kreutzberg, R. Boese and A. H. Maulitz, *Chem. Ber.*, 1993, **126**, 2041–2045.
- 9 A. J. Arduhengo, III, F. Davidson, R. Krafczyk, W. J. Marshall and R. Schmutzler, *Monatsh. Chem.*, 2000, **131**, 251–265.
- 10 R. J. Baker, M. L. Cole, C. Jones and M. F. Mahon, J. Chem. Soc., Dalton Trans., 2002, 1992; R. J. Baker, A. J. Davies, C. Jones and M. Kloth, J. Organmet. Chem., 2002, 656, 203; M. D. Francis, D. E. Hibbs, M. B. Hursthoues, C. Jones and N. A. Smithies, J. Chem. Soc., Dalton Trans., 1998, 3249.
- 11 Jennings and Clyburne also reported that the direct reaction of **1b** with NaBH<sub>4</sub> resulted in the formation of **3b**. See reference 8.
- 12 A related compound was reported, see: K. Öfele, W. A. Herrmann, D. Mihalios, M. Elison, E. Herdtweck, W. Scherer and J. Mink, *J. Organomet. Chem.*, 1993, **459**, 177–184; H. G. Raubenheimer, Y. Stander, E. K. Marais, C. Thompson, G. J. Kruger, S. Cronje and M. Deetlefs, *J. Organomet. Chem.*, 1999, **590**, 158–168.
- 13 N. Frafán and R. Contreras, J. Chem. Soc., Perkin Trans. 2, 1987, 771–773.