

Bassam A. Abu Thaher [a], Jalal A. Zahra [b] and Mustafa M. El-Abadelah* [b]

[a] Chemistry Department, Faculty of Science, Islamic University of Gaza, Gaza Strip.

[b] Chemistry Department, Faculty of Science, University of Jordan, Amman, Jordan.

Received November 19, 2001

2-Aminobenzoic acid reacts readily, in the presence of triethylamine, with hydrazonoyl chlorides (**5a-c**) (precursors of the reactive nitrile imine 1,3-dipolar species) to afford high yields of the corresponding acyclic amidrazone adducts (**6a-c**). The latter adducts undergo, in THF in presence of 1,1'-carbonyldiimidazole, smooth intramolecular cyclization involving the activated carboxyl and the NH- termini to deliver unequivocally the respective dihydro-1,3,4-benzotriazepin-5-ones (**7a-c**).

J. Heterocyclic Chem., **39**, 901(2002).

Introduction.

1,4-benzodiazepines, exemplified by chlorbenzodiazepoxide (Librium, **1**) and diazepam (Valium, **2**) (Chart 1), constitute a unique class of CNS drugs that are widely used as tranquilizers in several anxiety states [1]. Extensive research in the field of benzodiazepines has stimulated interest in the synthesis and pharmacologic action of their aza-homologues, the benzotriazepines. Amongst these, the 1,3,4-benzotriazepin-5-ones (*e.g.* **3**, Chart 1) [2-7] have been attended to, but research efforts in this area were comparatively modest. Their preparation was reported at large *via* the reaction of 2-aminobenzoylhydrazides with orthoesters (**3a-d**) [2], or *via* the reaction of 3,1-benzoxazin-4-ones with hydrazine (**3e,f**) [3], and once from the reaction of isatoic anhydride with 2-pyridylamidrazone and cyclizing the product with acid (**3g**) [4]. Some 1,3,4-benzotriazepin-6-ones (**4**) (Chart 1) were reported as useful antihypertensives, cardiotonics and fungicides [5].

It is worth mentioning that considerable confusion has surrounded these literature methods, which could have easily generated varying amounts of the isomeric 3-aminoquinazolin-4-ones and/or 1,3,4-oxadiazoles. In fact, several of the earlier reported syntheses of presumed 1,3,4-benzotriazepines [6] were later proven erroneous, and the alternate isomeric structures established [2a,4,7]. The formation of those intervening isomeric five- and six-membered heterocycles was occasionally suppressed by the proper choice, adjustment of, and monitoring of the reaction conditions [2c], and was excluded by the use of 1-(2-aminobenzoyl)-1-methylhydrazides [8,9].

In light of these uncertainties, we thought it was worthwhile to explore alternate preparative avenues starting from easily accessible compounds. Herein we wish to report on a convenient and versatile synthetic route toward new 1,3,4-benzotriazepin-5-ones (**7**), utilizing hydrazonoyl chlorides (**5**) (Scheme 1).


Results and Discussion.

Synthesis.

2-Aminobenzoic acid, acting as nitrogen nucleophile in basic medium, adds readily onto nitrile imines (the reactive 1,3-dipolar intermediates, generated *in situ* from their hydrazonoyl chloride precursors **5a-c** in presence of triethylamine) to furnish the corresponding amidrazone adducts **6** (Scheme 1). This mode of nucleophilic addition reaction of various nucleophiles onto 1,3-dipoles is well-documented, and several amidrazone adducts, related to **6**, were obtained from the reaction of primary and secondary amines with hydrazonoyl chlorides (such as **5**) [10,11]. The required hydrazonoyl chlorides **5a** [12-14], **5b** [12,13] and **5c** [12-14] were previously characterized, and are made accessible *via* the Japp-Klingemann reaction [12,15,16] involving coupling of the appropriate arenediazonium chloride with 3-chloro-2,4-pentanedione in aqueous pyridine.

In a subsequent step, these acyclic adducts **6** in tetrahydrofuran in presence of the coupling reagent, 1,1'-carbonyldiimidazole, underwent smooth cyclocondensation involving the activated carboxyl group and the hydrazone-NH terminus to deliver the requisite benzotriazepinones **7** (Scheme 1).

Chart 1



a: R = Ph (R' = H)

b: R = Me (R' = H)

c: R = R' = H

d: R = H; Me; Et; Bu
(R' = H; Me; Cl; Br; NO₂)e: R' = H; R = C₆H₄-X(p)
(X = Me; OMe; Cl; NO₂)


f: R' = H; R = COX (X = allylamino)

g: R = 2-Pyridyl (R' = H)

R = NH₂; alkylthio

R' = Furoyl

R'' and R''' = H; alkoxy



Spectral Data.

The ir, ms, and nmr spectral data and microanalyses of the new compounds **6a-c** and **7a-c** conform to the suggested structures, and are given in the experimental section. Thus, their ms spectra display the correct molecular ions for which the measured high-resolution data are in good agreement with the calculated values. ^1H -signal assignments are straightforward, and ^{13}C -assignments follow from DEPT and 2D (COSY, HMQC and HMBC) experiments. Thus, long range correlation is observed between H-6 and each of C-9a and C-5 in HMBC experiments for compounds **7a-c**. Likewise, H-8 is correlated with C-5a, while H-3'/H-5' show correlation with C-1'.

In conclusion, a new synthetic route toward 1,3,4-benzotriazepin-5-ones, described in the present work, utilizes readily available and inexpensive reactants (anthranilic acid and hydrazoneyl chlorides). This versatile and efficient route involves two-step reactions that are conveniently conducted in a short time, at or below room temperature, to give unequivocally the desired benzotriazepinones in good overall yield and high purity. In essence, this viable method competes favorably with ambiguous reported methods which also suffered from inherent drawbacks (*vide supra*).

EXPERIMENTAL

Melting points (uncorrected) were determined on an electrothermal Mel-Temp. Apparatus. ^1H - and ^{13}C nmr spectra were measured on a Bruker DPX-300 instrument with TMS as internal reference. Electron-impact mass spectra were obtained using a Finnigan MAT TSQ-70 spectrometer at 70 eV; ion source temperature = 200 °C. Ir spectra were recorded as KBr discs on a Nicolet Impact-400 FT-IR spectrophotometer. Microanalyses

were performed at the Microanalytical Laboratory-Inorganic Chemistry Department, Tübingen University, Germany.

1-Arylhyaazono-1-chloropropanones (**5a-c**).

These hydrazoneyl chlorides were prepared *via* direct coupling of the appropriate arenediazonium chloride with 3-chloro-2,4-pentanedione in aqueous pyridine, following standard procedures [12,16].

2-[(2-Oxo-1-phenylhydrazonepropan-1-yl)amino]benzoic acid (**6a**).

A homogeneous solution of 2-aminobenzoic acid (1.65 g, 12 mmol) in aqueous methanol (70%, 20 ml) and triethylamine (3 ml) was added dropwise to a stirred and cooled (0 °C) solution of 1-phenylhydrazone-1-chloropropanone (**5a**) (2.16 g, 11 mmol) in tetrahydrofuran (30 ml). Additional triethylamine (3 ml) in tetrahydrofuran (5 ml) was then introduced dropwise into the reaction mixture, which was further stirred at 0 °C for 15-20 minutes, and then at room temperature for 2-3 hours. The organic solvents were then removed *in vacuo* from the reaction mixture, and the residual aqueous solution was directly acidified with glacial acetic acid (3 ml). The resulting crude solid product was collected, dried and recrystallized from hot chloroform/methanol (5:1 v/v) as fine yellow needles. Yield of **6a** = 82%; mp 210-211 °C; ir (potassium bromide): 3480 (OH), 3310 (NH), 3246 (NH), 1690 (CO), 1603, 1584, 1516, 1491, 1232 cm^{-1} ; ms: m/z (% rel. int.): 297 (M^+ , 100), 279 (8), 236 (18), 208 (4), 174 (18), 148 (49), 118 (60), 108 (87); hrms: Calcd for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_3$: 297.111316. Found: 297.108719; ^1H nmr (300 MHz, $\text{DMSO}-d_6$): δ 2.46 (s, 3H, COCH_3), 6.10 (dd, 1H, $J = 8.3$ Hz, 0.7 Hz, H-3), 6.78 (m, 1H, H-5), 6.89 (m, 1H, H-4'), 7.22-7.33 (m, 5H, H-2'/H-6', H-3'/H-5' and H-4), 7.88 (dd, 1H, $J = 7.9$ Hz, 1.6 Hz, H-6), 9.25 (s, 1H, $\text{C}_2\text{-NH}$), 10.02 (s, 1H, $\text{C}_1\text{-NH}$), 13.10 (s, 1H, CO_2H); ^{13}C nmr (75 MHz, $\text{DMSO}-d_6$): 25.1 (COCH_3), 114.2 (C-1), 114.7 (C-2'/C-6'), 115.8 (C-3), 118.5 (C-5), 121.9 (C-4'), 129.5 (C-3'/C-5'), 131.9 (C-6), 134.4 (C-4), 134.9 (NH-C=N-), 144.2 (C-1'), 145.7 (C-2), 170.5 (CO_2H), 193.3 ($\text{CH}_3\text{C=O}$).

Anal. Calcd. for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_3$ (297.32): C, 64.64; H, 5.09; N, 14.13. Found: C, 64.35; H, 5.03; N, 14.11.

2-[[2-Oxo-1-(4-methylphenylhydrazonopropan-1-yl)]amino]-benzoic Acid (**6b**).

This compound was prepared from 2-aminobenzoic acid (1.65 g, 12 mmol) and 1-chloro-1-(4'-methylphenylhydrazono)-propanone (**5b**) (2.30 g, 11 mmol) by following the same procedure and experimental conditions described above for obtaining **6a**. The product was recrystallized from chloroform/methanol as pale yellow prisms. Yield of **6b** = 86%; mp 214–215 °C; ir (potassium bromide): 3466 (OH), 3332 (NH), 3305 (NH), 1674 (CO), 1574, 1555, 1524, 1501, 1244 cm⁻¹; ms: m/z (% rel. int.): 311 (M⁺, 84), 293 (13), 250 (7), 222 (4), 189 (26), 174 (13), 148 (26), 146 (28), 132 (100), 106 (74); hrms: Calcd for C₁₇H₁₇N₃O₃: 311.1270. Found: 311.1250; ¹H nmr (300 MHz, DMSO-d₆): δ 2.20 (s, 3H, C₄-CH₃), 2.44 (s, 3H, COCH₃), 6.08 (d, 1H, J = 8.3 Hz, H-3), 6.77 (dd, 1H, J = 7.8 Hz, 7.5 Hz, H-5), 7.06 (d, 2H, J = 8.3 Hz, H-3'/H-5'), 7.22 (d, 2H, J = 8.3 Hz, H-2'/H-6'), 7.29 (m, 1H, H-4), 7.87 (dd, 1H, J = 7.8 Hz, 1.3 Hz, H-6), 9.21 (br s, 1H, C₂-NH), 9.97 (s, 1H, C₁-NH), 13.09 (br s, 1H, CO₂H); ¹³C nmr (75 MHz, DMSO-d₆): δ 20.8 (C₄-CH₃), 25.0 (COCH₃), 114.1 (C-1), 114.8 (C-2'/C-6'), 115.7 (C-3), 118.4 (C-5), 130.0 (C-3'/C-5'), 130.8 (C-6), 131.8 (C-4), 134.3 (C-4'), 134.4 (NH-C=N-), 141.9 (C-1'), 145.8 (C-2), 170.3 (CO₂H), 193.1 (CH₃-C=O).

Anal. Calcd. for C₁₇H₁₇N₃O₃ (311.34): C, 65.58; H, 5.50; N, 13.50. Found: C, 65.44; H, 5.52; N, 13.38.

2-[[2-Oxo-1-(4-chlorophenylhydrazonopropan-1-yl)]amino]-benzoic Acid (**6c**).

This compound was prepared from 2-aminobenzoic acid (1.65 g, 12 mmol) and 1-chloro-1-(4'-chlorophenylhydrazono)-propanone (**5c**) (2.31 g, 11 mmol) by following the same procedure and experimental conditions described above for obtaining **6a**. Recrystallization from chloroform/methanol produced light orange granules. Yield of **6c** = 88%; mp 234–235 °C; ir (potassium bromide): 3450 (OH), 3348 (NH), 3244 (NH), 1695 (CO), 1652, 1586, 1567, 1502, 1490, 1360, 1281, 1240, 1088, 1028 cm⁻¹; ms: m/z (% rel. int.): 331 (M⁺, 100), 313 (30), 270 (11), 235 (13), 174 (28), 148 (95), 126 (97), 125 (88); hrms: Calcd for C₁₆H₁₄ClN₃O₃: 331.0723. Found: 331.0692; ¹H nmr (300 MHz, DMSO-d₆): δ 2.46 (s, 3H, COCH₃), 6.10 (dd, 1H, J = 8.3 Hz, 0.7 Hz, H-3), 6.79 (ddd, 1H, J = 8.1 Hz, 7.2 Hz, 1.0 Hz, H-5), 7.27–7.34 (m, 5H, H-2'/H-6', H-3'/H-5', H-4), 7.89 (dd, 1H, J = 7.2 Hz, 1.5 Hz, H-6), 9.29 (s, 1H, C₂-NH), 10.10 (s, 1H, C₁-NH), 13.09 (br s, 1H, CO₂H); ¹³C nmr (75 MHz, DMSO-d₆): δ 25.11 (COCH₃), 114.3 (C-1), 115.9 (C-3), 116.2 (C-2'/C-6'), 118.6 (C-5), 125.4 (C-4'), 129.4 (C-3'/C-5'), 131.9 (C-6), 134.4 (C-4), 135.3 (NH-C=N-), 143.2 (C-1'), 145.4 (C-2), 170.3 (CO₂H), 193.4 (CH₃-C=O).

Anal. Calcd. for C₁₆H₁₄ClN₃O₃ (331.76): C, 57.93; H, 4.25; Cl, 10.69; N, 12.67. Found: C, 57.68; H, 4.15; Cl, 10.53; N, 12.58.

2-Acetyl-4-phenyl-1,4-dihydro-1*H*-1,3,4-benzotriazepin-5-one (**7a**).

To a mixture of 1,1'-carbonyldiimidazole (2.0 g, 12.5 mmol) and compound **6a** (3.0 g, 10 mmol) was added dry tetrahydrofuran (50 ml), and the resulting solution was stirred at room temperature for 1–2 hours. The solvent was then removed *in vacuo* and the residue was immediately treated with water (40 ml) and extracted with chloroform (2 x 40 ml). The combined organic extracts were dried (Na₂SO₄), the solvent chloroform was evapo-

rated and the residual solid product was recrystallized from CHCl₃/petroleum ether (bp 40–60 °C) forming beige fine crystals. Yield of **7a** = 78 %; mp 152–153 °C; ir (potassium bromide): 3319 (NH), 1711 (CO), 1590, 1500, 1465, 1366, 1294, 1270, 1166, 1011 cm⁻¹; ms: m/z (% rel. int.): 279 (M⁺, 100), 250 (10), 236 (44), 220 (10), 174 (9), 146 (22), 132 (64), 130 (17), 105 (24); hrms: Calcd for C₁₆H₁₃N₃O₂: 279.1008. Found: 279.0975; ¹H nmr (300 MHz, CDCl₃): δ 2.53 s (3H, COCH₃), 6.72 (dd, 2H, J = 8.5 Hz, 1.1 Hz, H-2'/H-6'), 6.92 (m, 1H, H-4'), 7.16 (m, 2H, H-3'/H-5'), 7.46 (br s, 1H, N₁-H), 7.43 (m, 1H, H-7), 7.74 (m, 2H, H-8/H-9), 8.16 (ddd, 1H, J = 7.6 Hz, 1.9 Hz, 1.0 Hz, H-6); ¹³C nmr (75 MHz, CDCl₃): δ 28.7 (COCH₃), 114.5 (C-2'/C-6'), 122.4 (C-5a), 122.8 (C-4'), 127.0 (C-6), 128.1 (C-7), 128.2 (C-9), 129.4 (C-3'/C-5'), 135.2 (C-8), 145.7 (C-1'), 146.6 (C-9a), 153.7 (C-2), 160.6 (O=C₅-N), 194.2 (CH₃-C=O).

Anal. Calcd. for C₁₆H₁₃N₃O₂ (279.3): C, 68.81; H, 4.69; N, 15.04. Found: C, 68.63; H, 4.55; N, 14.86.

2-Acetyl-4-(4-methylphenyl)-1,4-dihydro-1*H*-1,3,4-benzotriazepin-5-one (**7b**).

This compound was prepared from **6b** (3.1 g, 10 mmol) and 1,1'-carbonyldiimidazole (2.0 g, 12.5 mmol) by following the same procedure and experimental conditions described above for **7a**. The product was recrystallized from dichloromethane/*n*-hexane as yellow prisms. Yield of **7b** = 81%; mp 155–156 °C; ir (potassium bromide): 3268 (NH), 1705 (CO), 1626, 1606, 1523, 1510, 1344, 1286, 1234, 1130 cm⁻¹; ms: m/z (% rel. int.): 293 (M⁺, 45), 279 (9), 250 (4), 189 (21), 167 (28), 149 (97), 132 (19), 119 (22), 91 (100); hrms: Calcd for C₁₇H₁₅N₃O₂: 293.1164. Found: 293.1117; ¹H nmr (300 MHz, CDCl₃): δ 2.38 (s, 3H, C₄-CH₃), 2.43 (s, 3H, COCH₃), 6.69 (d, 1H, J = 7.9 Hz, H-8), 6.97 (dd, 1H, J = 7.4 Hz, 7.7 Hz, H-7), 7.23 (d, 2H, J = 8.3 Hz, H-3'/H-5'), 7.36 (d, 2H, J = 8.3 Hz, H-2'/H-6'), 7.33 (m, 1H, H-9, overlapped with H-2'/H-6'), 7.30 (br s, 1H, N₁-H), 7.96 (d, 1H, J = 7.7 Hz, H-6); ¹³C nmr (75 MHz, CDCl₃): δ 21.2 (C₄-CH₃), 24.1 (COCH₃), 118.3 (C-8), 121.9 (C-5a), 122.7 (C-7), 126.0 (C-2'/C-6'), 129.4 (C-3'/C-5'), 134.0 (C-6), 134.6 (C-9), 137.5 (C-4'), 141.5 (C-1'), 143.6 (C-9a), 145.8 (C-2), 165.6 (O=C₅-N), 194.0 (CH₃-C=O).

Anal. Calcd. for C₁₇H₁₅N₃O₂ (293.33): C, 69.61; H, 5.15; N, 14.33. Found: C, 69.37; H, 5.02; N, 14.15.

2-Acetyl-4-(4-chlorophenyl)-1,4-dihydro-1*H*-1,3,4-benzotriazepin-5-one (**7c**).

This compound was prepared from **6c** (3.3 g, 10 mmol) and 1,1'-carbonyldiimidazole (2.0 g, 12.5 mmol) by following the same procedure and experimental conditions described above for **7a**. Recrystallization from dichloromethane/*n*-hexane gave pale yellow granules. Yield of **7c** = 83 %; mp 181–182 °C; ir (potassium bromide): 3313 (NH), 1732 (CO), 1701 (CO), 1608, 1595, 1491, 1467, 1413, 1374, 1168, 1097 cm⁻¹; ms: m/z (% rel. int.): 313 (M⁺, 94), 278 (3), 270 (21), 254 (10), 235 (22), 189 (10), 174 (13), 146 (25), 132 (58), 126 (84), 125 (100), 111 (63); hrms: Calcd for C₁₆H₁₂ClN₃O₂: 313.0618. Found: 313.0592; ¹H nmr (300 MHz, CDCl₃): δ 2.57 (s, 3H, COCH₃), 6.69 (m, 2H, H-2'/H-6'), 7.16 (m, 2H, H-3'/H-5'), 7.20 (br s, 1H, N₁-H), 7.54 (ddd, 1H, J = 8.0 Hz, 6.3 Hz, 1.5 Hz, H-7), 7.79 (m, 2H, H-8/H-9), 8.22 (dd, 1H, J = 8.0 Hz, 1.5 Hz, H-6); ¹³C nmr (75 MHz, CDCl₃): δ 28.6 (COCH₃), 116.0 (C-2'/C-6'), 122.3 (C-5a), 127.2 (C-6), 128.1 (C-4'), 128.3 (C-9), 128.4 (C-7), 129.4 (C-3'/C-5'), 135.3 (C-8), 144.3 (C-1'), 146.4 (C-9a), 152.9 (C-2), 160.5 (O=C₅-N), 193.9 (CH₃-C=O).

Anal. Calcd. for $C_{16}H_{12}ClN_3O_2$ (313.75): C, 61.25; H, 3.86; Cl, 11.30; N, 13.39. Found: C, 61.12; H, 3.81; Cl, 11.18; N, 13.35.

Acknowledgement.

We wish to thank the Advanced Pharmaceutical Industries Co. Ltd., Amman-Jordan for financial support.

REFERENCES AND NOTES

- [*] E-mail: mustelab@ju.edu.jo
- [1a] L. H. Sternbach, in the *Benzodiazepines*, S. Garattini, E. Mussini and L. O. Randall, eds., Raven Press, New York, 1973, pp. 1-25; [b] H. W. Gschwend, in *Anxiolytics*, S. Fielding and H. Lal, eds., Futura Publishing Co., Mt. Kisco, New York, 1979, Chapter 1; [c] L. H. Sternbach, *J. Med. Chem.*, **22**, 1(1979); [d] S. J. Childress, in *Antianxiety Agents in Burger's Medicinal Chemistry*, M. E. Wolff, ed., 4th edition, part III, John Wiley and Sons, New York, 1981, p. 981; [e] D. J. Greenblatt and R. I. Shader, in *Benzodiazepines in Clinical Practice*, Raven Press, New York, 1974, p.17.
- [2a] N. B. Peet, *Synthesis*, 1065 (1984); [b] N. P. Peets and S. Sunder, *J. Heterocyclic Chem.*, **21**, 1807 (1984); [c] P. Scheiner, L. Frank, I. Giusti, S. Arwin, S. A. Pearson, F. Excellent and A. P. Harper, *J. Heterocyclic Chem.*, **21**, 1817 (1984); [d] R. W. Leiby, *J. Heterocyclic Chem.*, **21**, 1825 (1984); [e] J. Y. Mérour, *J. Heterocyclic Chem.*, **19**, 1425 (1982).
- [3a] C. K. Reddy, P. S. N. Reddy and C. V. Ratnam, *Indian J. Chem.*, **24B**, 902 (1985); [b] L. M. Korkodinova, V. L. Gein and M. I Vakhrin, *Zh. Obsch. Khim.*, **63**, 699 (1993); *Chem Abstr.*, **120**, 8575h (1994); the structure of benzotriazepinones, prepared in ref [3b], was presented as the tautomeric 1,4-dihydro-1*H* form of **3f**.
- [4] M. Takahashi, S. Onizawa and T. Satoh, *Bull. Chem. Soc. Jpn.*, **47**, 2724 (1974).
- [5] *Jpn. Kokai Tokkyo Koho JP*, **82 109 772** (1982) (Otsuka Pharmaceutical Co., Ltd, Japan); *Chem. Abstr.*, **97**, 216249p (1982).
- [6a] G. T. Panse and S. K. Kamat, *Indian J. Chem.*, **13**, 834 (1975); [b] C. Podeseva, G. Kohan and K. Vagi, *Can. J. Chem.*, **47**, 489 (1969); [c] C. K. Reddy, P.S.N. Reddy and C. V. Ratnam, *Synthesis*, 842 (1983); [d] A. M. M. E. Omar, F. A. Ashour and J. Bourdais, *J. Heterocyclic Chem.*, **16**, 1435 (1979); [e] J. S. Davidson, "Abstracts of the English International Congress of Heterocyclic Chemistry" p. 214 (1981); [f] A. L. Langis, US 3 542 767 (1970) *Chem. Abstr.*, **74**, 88089x (1971); [g] G. M. Reddy and P. S. N. Reddy, *Indian J. Chem.*, **36B**, 166 (1997).
- [7a] N. P. Peet, S. Sunder and D. L. Trepanier, *Indian. J. Chem.*, **14B**, 701 (1976); [b] M. J. Kornet, *J. Pharm. Sci.*, **62**, 834 (1973); [c] S. Sunder, N. P. Peet and R. J. Barbuch, *J. Heterocyclic Chem.*, **18**, 1601 (1981); [d] E. Tihanyi, M. Jal and P. Dvorstak, *Heterocycles*, **20**, 571 (1983); [e] G. M. Reddy and P. S. N. Reddy, *Indian J. Chem.*, **37B**, 207 (1998).
- [8] S. Sunder, N. P. Peet and D. L. Trepanier, *J. Org. Chem.*, **41**, 2732 (1976).
- [9] R. W. Leiby and N. D. Heindel, *J. Org. Chem.*, **41**, 2736 (1976).
- [10a] R. N. Bulter and F. L. Scott, *Chem.&Ind.* (London), 1216 (1970); [b] H. Ulrich, *The Chemistry of Imidoyl Halides*, Plenum Press, New York, 1968, Chapter 7, pp. 174-192; [c] A. F. Hegarty, J. B. Aylward and F. L. Scott, *J. Chem. Soc. (C)*, 2587 (1967); [d] P. Dalla Corce, P. Del Buttero, E. Licandro and S. Maiorana, *Synthesis*, 299 (1979); [e] G. Heubach, *Liebigs Ann. Chem.*, 1376 (1980); [f] H. M. Hassaneen, H. A. H. Mousa and N. M. Abed, *Heterocycles*, **27**, 695 (1988); [g] T. Benincori and F. Sannicoló, *J. Org. Chem.*, **53**, 1309 (1988).
- [11a] V. A. Galishev, V. N. Chistokletov and A. A. Petrov, *Zh. Obshch. Khim.*, **45**, 1695 (1975); [b] A. S. Shawali and C. Párkányi, *J. Heterocyclic Chem.*, **17**, 833 (1980).
- [12] M. M. El-Abadelah, A. Q. Hussein and B. A. Abu Thaher, *Heterocycles*, **32**, 1879 (1991).
- [13] N. F. Eweiss and A. Osman, *J. Heterocyclic Chem.*, **17**, 1713 (1980).
- [14] R. Fusco and R. Romani, *Gazz. Chim. Ital.*, **76**, 419 (1946).
- [15a] R. R. Phillips, *Org. Reactions*, **10**, 143 (1959); [b] H. C. Yao and P. Resnick, *J. Am. Chem. Soc.*, **84**, 3514 (1962).
- [16] G. C. Barrett, M. M. El-Abadelah and M. K. Hargreaves, *J. Chem. Soc. (C)*, 1986 (1970).