

Published on Web 12/21/2006

Catalytic Enantioselective Allenylation Reactions of Aldehydes with Tethered Bis(8-quinolinolato) (TBOx) Chromium Complex

Guoyao Xia and Hisashi Yamamoto*

Department of Chemistry, The University of Chicago, 5735 South Ellis Avenue, Chicago, Illinois 60637

Received November 12, 2006; E-mail: yamamoto@uchicago.edu

Allenes have proven to be versatile and useful intermediates in organic synthesis due to the existence of the two orthogonal π -bonds.¹ Recently, α -allenic alcohols have drawn much attention because of their unique reactivities and the ease of further conversion into compounds with other functional groups.^{1,2} Practical methods for the synthesis of α -allenic alcohols by coupling propargylic reagents with carbonyl groups have often been limited by lack of chemoselectivity as well as enantioselectivity. This is due to the ambient nature of propargylic carbanions, which generally exist as an equilibrium mixture of allenic and propargylic organometallic derivatives.1a,d,3 Despite these complications, several methods have been developed for the regiospecific construction of α -allenic alcohols in recent years.^{1d,4} Among these enantioselective synthetic methods,1f,5 only a very limited number of catalytic asymmetric allenylation have been reported.^{6,7} The nucleophilic addition of organochromium reagents to aldehydes is a powerful C-C bondforming method.8 The allenylation reactions between aldehydes and propargylic halides catalyzed by chromium complexes are known to be very useful due to excellent chemoselectivity, broad compatibility with different functional groups, and environmentally benign process.⁷ However, there are still some difficulties in terms of (1) the enantioselectivities of α -allenic alcohols, (2) scope of substrates, and (3) the ease of operation with commercially available reagents.

We earlier developed and synthesized a tethered bis(8-quinolinolato) (TBOx) chromium catalyst and applied it to asymmetric pinacol coupling reactions⁹ of aldehydes and asymmetric Nozaki— Hiyama allylation reactions.¹⁰ This successful catalytic redox system was further applied to the asymmetric allenylation reaction between benzaldehyde and commercially available 1-trimethylsilyl-3-bromopropyne.

First, TBOxCr(III)Cl (5 mol %) and Mn powder were mixed in THF under an atmosphere of Ar at room temperature. After 10 min, to the reaction mixture was added 1-trimethylsilyl-3-bromopropyne,¹¹ and the mixture was stirred for 30 min. Then, benzaldehyde and the silyl chloride were added successively and slowly. After 48 h, the isolated crude product was treated with TBAF in THF at room temperature to afford 1-phenylbuta-2,3dien-1-ol as the only product. If the crude product was treated with 0.5 N HCl in THF, TMS-substituted α-allenic alcohol could be obtained in lower yield.¹² As shown in Table 1, by using TMSCl (1 equiv), the α -allenic alcohol was obtained in 77% yield with only 17% ee (entry 1). When TESCI (1 equiv) was used, the enantioselectivity of the product was increased dramatically accompanied by good yield (entry 4). The different reaction rates between TMSCl (entry 2) and TESCl (entry 5) were observed when no catalyst was applied. If more than 1 equiv of silvl chloride was used, the desired product was generated in higher yield but lower ee (entries 3 and 6). Finally, when 1-trimethylsilyl-3-bromopropyne (1.5 equiv) and TESCI (1 equiv) were applied, the desired α -allenic alcohol was produced with 91% yield and 96% ee (entry 8).

Under the optimized reaction conditions, a very wide scope of aldehydes was successfully allenylated in moderate to high yields

Table 1.	Asymmetric	Allenylation	Reactions	of Benzaldehyde
	,,	,		0. 20. 20. 20. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0

PhCHO ·	+ Br → Br TMS 2) TE	OxCr(III)CI(5 mol%) Mn, THF, rt, 48 h ↓AF,THF, rt Ph		N CI Cr < O N N DxCr(III)CI
	Br			ee ^b
entry	TMS	SiCl	yield ^a (%)	(%) (config) ^c
1	1 equiv	1 equiv of TMSCl	77	17 (<i>R</i>)
2^d	1 equiv	1 equiv TMSCl	37	
3	1 equiv	1.2 equiv of TMSCl	83	8 (R)
4	1 equiv	1 equiv of TESC1	71	96 (R)
5^d	1 equiv	1 equiv of TESC1	<5	
6	1 equiv	1.2 equiv of TESCl	74	95 (R)
7	1.25 equiv	1 equiv of TESC1	82	96 (R)
8	1.5 equiv	1 equiv of TESC1	91	96 (R)
9 ^e	1 equiv	1 equiv of TESCl	89	96 (R)

^{*a*} Isolated yield after chromatographic purification. ^{*b*} Enantiomeric excess was determined by chiral HPLC analysis. ^{*c*} Assigned by comparison of the sign of optical rotation with reported value. ^{*d*} No TBOxCr(III)Cl was used. ^{*e*} The reaction was complete in 30 h by using TBOxCr(III)Cl (10 mol %).

with excellent enantioselectivities. As shown in Table 2, *para*substituted aromatic aldehydes with either electron-donating or electron-withdrawing groups gave excellent enantioselectivities (entries 1, 4, 5, and 7). However, a higher catalyst loading, more TESCl, and 1-trimethylsilyl-3-bromopropyne had to be applied to achieve higher yields for the benzaldehydes with an electronwithdrawing group (entries 6 and 8). The *ortho-* and *meta*substituted benzaldehyde could also be allenylated in good yields with good ee (entries 2 and 3). Bulky aryl aldehydes (entries 9 and 10), heterocyclic aldehyde (entry 11), α,β -unsaturated aldehyde (entry12), and aliphatic aldehydes (entries 13–15) proved to be good substrates for this method.

To further explore the substrate scope, the allenylation reactions of aldehydes with terminally alkyl-substituted propargylic bromide, which had never succeeded with high enantioselectivities by Crcatalyzed asymmetric allenylation reactions previously,13 were examined under the optimized reaction conditions. As shown in Table 3, the allenylation reaction between benzaldehyde and commercially available 1-bromo-2-butyne provided 2-methyl-1phenylbuta-2,3-dien-1-ol in 84% yield with 97% ee as the major product with <5% yield of homopropargylic alcohol (entry 1). In the allenylation of other aromatic aldehydes with an electrondonating or electron-withdrawing group (entries 2 and 3), an aliphatic aldehyde (entry 4) also gave excellent enantioselectivities with moderate to good yields when R₂ was the methyl group. Increasing the size of the R₂ group caused the enantioselectivities of the allenylation reactions to go down slightly. When R2 is an ethyl group, benzaldehyde and hydrocinnamylaldehyde were allenylated in good yield with excellent ee (entry 5 compared to entry 1, and entry 6 compared to entry 4). When R_2 is a phenyl group, benzaldehyde was allenylated in 58% yield with 88% ee (entry 7 compared to entries 1 and 5).

Table 2. Asymmetric Allenylation	Reactions of Aldehydes
----------------------------------	------------------------

RCHO	Br	1) TBOxCr(III)Cl(5 mol%) TESCI, Mn, THF, rt, time		он	
Konc	TMS	2) TBAF,THF,	rt	R * *	
entry	RCHO	time (h)	yield ^a (%)	ee ^b (%)	
1	СНО	48	83	90	
2	СНО	60	77	84	
3	MeO	48	85	96	
4	МеО	48	81	95	
5	СНО	60	35	94	
6 ^c		60	51	91	
7	СНО	60	40	93	
80	Br	60	57	92	
9	СНО	60	33	93	
10 ^c		60	45	90	
11	ОСНО	48	79	97	
12	СНО	60	51	90	
13	СНО	48	81	93 ^d	
14	СНО	60	72	90 ^d	
15	СНО	60	75	85 ^d	

^{*a*} Isolated yield after chromatographic purification. ^{*b*} Enantiomeric excess was determined by chiral HPLC or GC analysis. ^{*c*} 1-Trimethylsilyl-3-bromopropyne (2 equiv), TESCI (1.2 equiv), and TBOxCr(III)Cl (7 mol %) were used. ^{*d*} The absolute configuration of the product was determined to be *S*.

Table 3.	Asymmetric	Allenylation	Reactions	of Aldehy	/des with
Terminall	y Alkyl-Subs	tituted Propa	argylic Bron	nide	
		1) TE	BOYCr(III)CI(5)	mol%)	0 4

R₂CH	0 + ^{Br}	TESCI, Mn, THF, rt, time			
R ₂ 1.5 eq.		2) TBAF,THF, rt		$\mathbf{R}_1 + \mathbf{R}_2$	
entry	R₁CHO	R_2	time (h)	yield ^a (%)	ee ^b (%)
1	СНО	Ме	48	84	97
2	СНО	Ме	48	76	97
3	СІСНО	Ме	60	48	90
4	СНО	Ме	48	69	93 ^c
5	СНО	Et	60	87	95
6	СНО	Et	60	72	92 ^c
7	СНО	Ph	60	58	88

^{*a*} Isolated yield after chromatographic purification. ^{*b*} Enantiomeric excess was determined by chiral HPLC analysis. ^{*c*} The absolute configuration of the product was determined to be *S*.

In summary, TBOxCr(III)Cl was shown to efficiently catalyze the asymmetric allenylation reactions of both aromatic and aliphatic aldehydes. Utilizing this methodology, most of the difficulties were overcome: (1) Excellent enantioselectivities (up to 97% ee) for the allenylation reaction of versatile aldehydes were obtained. (2) Easy operation with commercially available terminally substituted propargylic bromide could be achieved. (3) First success for the allenylation reactions of aldehydes with terminally alkyl-substituted propargylic bromides that provided excellent chemoselectivities (α -allenic alcohol as the major product in up to 84% yield) as well as unprecedented enantioselectivities (up to 97% ee). Studies are currently underway to elucidate the mechanism.¹⁴ The applications of TBOxH in other asymmetric catalysis will be reported in due course.

Acknowledgment. Support of this research was provided by the National Science Foundation (NSF) and Merck & Co.

Supporting Information Available: Experimental procedures and spectral data. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (a) Moreau, J. L. In *The Chemistry of Ketenes, Allenes, and Related Compounds*; Patai, S., Ed.; John Wiley & Sons: New York, 1980; Part 1.
 (b) *The Chemistry of Allenes*; Landor, S. R., Ed.; Academic Press: London 1982.
 (c) Schuster, H. F.; Coppola, G. M. *Allenes in Organic Synthesis*; John Wiley & Sons: New York, 1988.
 (d) Yamamoto, H. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: Oxford, 1991.
 (e) *Modern Acetylene Chemistry*; Stang, P. J., Diederich, F., Eds.; VCH: Weinheim, Germany, 1995.
 (f) Marshall, J. A.; Gung, B. W.; Grachan, M. L. In *Modern Allene Chemistry*; Krause, N., Hashmi, A. S. K., Eds.; VCH: Weinheim, Germany, 2004.
- A., Oung, B. W., Orachan, M. L. In *Modern Antene Chemmsols*, Nalase, N., Hashmi, A. S. K., Eds.; VCH: Weinheim, Germany, 2004.
 (2) Some examples: (a) Olsson, L. I.; Claesson, A. Synthesis 1979, 9, 743–745. (b) Nikam, S. S.; Chu, K. H.; Wang, K. K. J. Org. Chem. 1986, 51, 745–747. (c) Friesen, R. W. Tetrahedron. Lett. 1990, 31, 4249–4252. (d) Marshall, J. A.; Pinney, K. G. J. Org. Chem. 1993, 58, 7180–7184. (e) Laurens, M.; Delanghe, P. H. M. J. Am. Chem. Soc. 1994, 116, 8526–8535. (f) Ma, S.; Zhao, S. J. Am. Chem. Soc. 1999, 121, 7943–7944. (g) Hoffmann-Röder, A.; Krause, N. Org. Lett. 2001, 3, 2537–2538. (h) Ma, S.; Zhao, S. J. Am. Chem. Soc. 2001, 123, 5578–5579. (i) Krause, N.; Hoffmann-Röder, A.; Canisius, J. Synthesis 2002, 1759–1774. (j) Alcaide, B.; Almendros, P.; Aragoncillo, C. Chem.—Eur. J. 2002, 8, 1719–1729. (k) Xu, D.; Xu, Y.; Li, L.; Ma, S. Chem.—Eur. J. 2002, 8, 5012–5018. (l) Yoneda, E.; Zhang, S.; Zhou, D.; Onitsuka, K.; Takahashi, S. J. Org. Chem. 2003, 68, 8371–8376. (m) Mushima, S. K.; Blanco, M. C.; Fischer, D.; Bats, J. W. Eur. J. Org. Chem. 2006, 1387–1389.
- (3) Reviews: (a) Epsztein, R. In Comprehensive Carbanion Chemistry; Buncel, E., Durst, T., Eds.; Wiley: New York, 1980. (b) Brandsma, L.; Verkruijsee, H. D. Synthesis of Acetylenes, Allenes and Cumelenes; Elsevier: Amsterdam, 1981.
- (4) (a) Kobayashi, S.; Nishio, K. J. Am. Chem. Soc. 1995, 117, 6392-6393.
 (b) Hamada, T.; Mizojiri, R.; Urabe, H.; Sato, F. J. Am. Chem. Soc. 2000, 122, 7138-7139. (c) Loh, T. P.; Lin, M. J. J. Am. Chem. Soc. 2003, 125, 13042-13043. (d) Krause, N.; Hoffmann-Röder, A. Tetrahedron 2004, 60, 11671-11694.
- (5) (a) Corey, E. J.; Jones, G. B. *Tetrahedron Lett.* **1991**, *32*, 5713–5716.
 (b) Brown, H. C.; Khire, U. R.; Narla, G. J. Org. Chem. **1995**, *60*, 8130–8131. (c) Marshall, A.; Adams, N. D. J. Org. Chem. **1997**, *62*, 8976–8977. (d) Schultz-Fademrecht, C.; Wibbeling, B.; Froehlich, R.; Hoppe, D. Org. Lett. **2001**, *3*, 1221–1224. (e) Denmark, S. E.; Fu, J. Chem. Rev. **2003**, *103*, 2763–2793. (f) Hernandez, E.; Soderquist, J. A. Org. Lett. **2005**, *7*, 5397–5400. (g) Hernandez, E.; Burgos, C. H.; Alicea, E.; Soderquist, J. A. Org. Lett. **2006**, *8*, 4089–4091.
- D. Org. Lett. 2001, 3, 1221-1224. (e) Denmark, S. E.; Fu, J. Chem. Rev. 2003, 103, 2763-2793. (f) Hernandez, E.; Soderquist, J. A. Org. Lett. 2005, 7, 5397-5400. (g) Hernandez, E.; Burgos, C. H.; Alicea, E.; Soderquist, J. A. Org. Lett. 2006, 8, 4089-4091.
 (6) (a) Yu, C. M.; Yoon, S. K.; Baek, K.; Lee, J. Y. Angew. Chem., Int. Ed. 1998, 37, 2392-2395. (b) Iseki, K.; Kuroki, Y.; Kobayashi, Y. Tetrahedron: Asymmetry 1998, 9, 2889-2894. (c) Nakajima, M.; Saito, M.; Hashimoto, S. Tetrahedron: Asymmetry 2002, 13, 2449-2452.
 (7) For a recent excellent report shout Cr a optimized supmatria ellowulation.
- (7) For a recent excellent report about Cr-catalyzed asymmetric allenylation reactions of aldehydes, see: Inoue, M.; Nakada, M. Angew. Chem., Int. Ed. 2006, 45, 252–255.
- (8) Pioneer works and recent reviews: (a) Fürstner, A.; Shi, N. J. Am. Chem. Soc. 1996, 118, 12349–12357. (b) Fürstner, A. Chem.-Eur. J., 1998, 4, 567–570. (c) Fürstner, A. Chem. Rev. 1999, 99, 991–1045. (d) Wess-johann, L. A.; Scheid, G. Synthesis 1999, 1, 1–36. (e) Avalos, M.; Babiano, R.; Cintas, P.; Jimenez, J. L.; Palacios, J. C. Chem. Soc. Rev. 1999, 28, 169–177. (f) Takai, K.; Nozaki, H. Proc. Jpn. Acad., Ser. B 2000, 76B, 123–131.
- (9) Takenaka, N.; Xia, G.; Yamamoto, H. J. Am. Chem. Soc. 2004, 126, 13198-13199.
- (10) Xia, G.; Yamamoto, H. J. Am. Chem. Soc. 2006, 128, 2554-2555.
- (11) The silyl group of the propargylic bromide affected the yields and enantioselectivities. See Scheme 1 in Supporting Information.
- (12) See Scheme 2 in Supporting Information.(13) For example, ref 7 in comparison to entry 1 in Table 3.
- (14) See Scheme 3 in Supporting Information for the proposed catalytic cycle and transition structures.

JA0679578