

Substituent Effects on the Antibacterial Activity of Nitrogen–Carbon-Linked (Azolylphenyl)oxazolidinones with Expanded Activity Against the Fastidious Gram-Negative Organisms *Haemophilus influenzae* and *Moraxella catarrhalis*

Michael J. Genin,^{*,†} Debra A. Allwine,[†] David J. Anderson,[†] Michael R. Barbachyn,[†] D. Edward Emmert,[†] Stuart A. Garmon,[†] David R. Graber,[†] Kevin C. Grega,[†] Jackson B. Hester,[†] Douglas K. Hutchinson,[§] Joel Morris,[†] Robert J. Reischer,[†] Charles W. Ford,[‡] Gary E. Zurenko,[‡] Judith C. Hamel,[‡] Ronda D. Schaadt,[‡] Douglas Stapert,[‡] and Betty H. Yagi[‡]

Pharmacia & Upjohn, Inc., Kalamazoo, Michigan 49001

Received July 19, 1999

A series of new nitrogen–carbon-linked (azolylphenyl)oxazolidinone antibacterial agents has been prepared in an effort to expand the spectrum of activity of this class of antibiotics to include Gram-negative organisms. Pyrrole, pyrazole, imidazole, triazole, and tetrazole moieties have been used to replace the morpholine ring of linezolid (**2**). These changes resulted in the preparation of compounds with good activity against the fastidious Gram-negative organisms *Haemophilus influenzae* and *Moraxella catarrhalis*. The unsubstituted pyrrolyl analogue **3** and the 1*H*-1,2,3-triazolyl analogue **6** have MICs against *H. influenzae* = 4 µg/mL and *M. catarrhalis* = 2 µg/mL. Various substituents were also placed on the azole moieties in order to study their effects on antibacterial activity in vitro and in vivo. Interesting differences in activity were observed for many analogues that cannot be rationalized solely on the basis of sterics and position/number of nitrogen atoms in the azole ring. Differences in activity rely strongly on subtle changes in the electronic character of the overall azole systems. Aldehyde, aldoxime, and cyano azoles generally led to dramatic improvements in activity against both Gram-positive and Gram-negative bacteria relative to unsubstituted counterparts. However, amide, ester, amino, hydroxy, alkoxy, and alkyl substituents resulted in no improvement or a loss in antibacterial activity. The placement of a cyano moiety on the azole often generates analogues with interesting antibacterial activity in vitro and in vivo. In particular, the 3-cyanopyrrole, 4-cyanopyrazole, and 4-cyano-1*H*-1,2,3-triazole congeners **28**, **50**, and **90** had *S. aureus* MICs ≤ 0.5–1 µg/mL and *H. influenzae* and *M. catarrhalis* MICs = 2–4 µg/mL. These analogues are also very effective versus *S. aureus* and *S. pneumoniae* in mouse models of human infection with ED₅₀s in the range of 1.2–1.9 mg/kg versus 2.8–4.0 mg/kg for the eperezolid (**1**) control.

Introduction

Multi-drug-resistant Gram-positive bacterial pathogens^{1–3} including methicillin-resistant *Staphylococcus aureus* (MRSA)⁴ and *Staphylococcus epidermidis* (MRSE),⁴ vancomycin-resistant enterococci (VRE),^{5,6} and penicillin- and cephalosporin-resistant streptococci⁷ have become a serious problem in hospitals and the community. Particularly alarming is the emergence of staphylococcal strains with reduced susceptibility to vancomycin, the so-called vancomycin/glycopeptide intermediate strains (VISA or GISA).^{8–11} More recently, a multinational team reported high rates of resistance among aerobic Gram-negative bacilli in European intensive care units.¹⁴ Thus, the search for novel potent broad-spectrum antibacterial agents is being fervently pursued by pharmaceutical houses worldwide.

The totally synthetic oxazolidinones typified by eperezolid (**1**) and linezolid (**2**) are one such class of antibacterial agents with potent activity against Gram-

positive organisms including MRSA, MRSE, and VRE.¹³ They have been shown to selectively and uniquely bind to the 50S ribosomal subunit and inhibit bacterial translation at the initiation phase of protein synthesis.^{14,15} In addition, single-step selection studies demonstrated that eperezolid- and linezolid-resistant mutants develop with a very low spontaneous mutation frequency of <10^{–9} among selected staphylococcal bacteria.¹⁶ Furthermore, evidence for the rapid development of bacterial resistance to both compounds could not be found via serial passage studies using spiral drug concentration gradient plates.¹⁷ Recently, phase III clinical trials with linezolid (PNU-100766) for the treatment of Gram-positive infections have been successfully completed.


We have been interested in expanding the spectrum of activity of the oxazolidinones to include the fastidious Gram-negative bacteria *Haemophilus influenzae* and *Moraxella catarrhalis*, which are key organisms in cases of community acquired pneumonia (CAP) and otitis media. Linezolid exhibits only modest in vitro activity against these organisms (MIC₉₀ = 16 µg/mL). In our efforts to improve the spectrum of activity, the (azolylphenyl)oxazolidinone subclass has been discovered wherein the morpholine ring of linezolid has been

* To whom correspondence should be addressed. Tel: 616-833-9869 or 4000. Fax: 616-833-2516. E-mail: michael.j.genin@am.pnu.com.

[†] Combinatorial and Medicinal Chemistry Research.


[‡] Infectious Diseases Research.

[§] Current address: Abbott Laboratories, One Abbott Park Rd., Abbott Park, IL 60064.

Scheme 1^a

^a Reagents: (a) pyrrole, NaH, THF, 66%; or pyrazole, K₂CO₃, DMSO, 90 °C, 87%; or imidazole, K₂CO₃, DMSO, 90 °C, 92%; or 1,2,3-triazole, K₂HPO₄, DMSO, 90 °C, 50% 1*H*-isomer + 43% 2*H*-isomer; or 1,2,4-triazole, K₂HPO₄, DMSO, 90 °C, 72% 1*H*-isomer; or tetrazole, NEt₃, CH₃CN, 10% 2*H*-isomer + 4% 1*H*-isomer; (b) H₂, 5% Pt/S/C, THF; or H₂, RaNi, EtOAc/MeOH; or H₂, 10% Pd/C, THF; (c) benzyl chloroformate, NaHCO₃, THF, rt, 37–92% for steps b + c; (d) (i) LiHMDS, THF –78 °C, (ii) *R*-(–)-glycidyl butyrate, –78 °C–rt, 48–85%; (e) MsCl, NEt₃, CH₂Cl₂; (f) NaN₃, DMF, 65 °C; (g) H₂, Pd/C, THF/MeOH; (h) Ac₂O, pyridine, 0 °C–rt, 47–66% for steps e–h.

replaced with various five-membered nitrogen-containing heterocycles (azoles). Some of these analogues have interesting levels of antibacterial activity. In particular, the pyrrole **3** and the 1*H*-1,2,3-triazole **6** analogues have excellent Gram-positive activity (MICs < 0.5–1 µg/mL), vide infra. More interesting, however, was their good activity against *H. influenzae* and *M. catarrhalis* (MICs = 2–4 µg/mL). In an effort to further increase the activity of the azolyphenyl class of oxazolidinones, we have explored the effects of various substituents on the azole rings. Previously we reported the discovery of the 3-cyanopyrrole and 4-cyanopyrazole analogues (PNU-171933 and PNU-172576) which have *S. aureus* MICs ≤ 0.5 µg/mL and *H. influenzae* and *M. catarrhalis* MICs = 2–4 µg/mL.¹⁸ These analogues also exhibit excellent pharmacokinetics and in vivo activity.¹⁸ We now wish to report a more extensive survey of this class in which numerous pyrrole, pyrazole, imidazole, triazole, and tetrazole analogues have been synthesized and tested for antibacterial activity.



Chemistry


Unsubstituted Azole Analogues. The synthesis of the unsubstituted azole analogues **3–8** and **10** is outlined in Scheme 1. The general preparation involved nucleophilic aromatic substitution reaction between the requisite azole and 3,4-difluoronitrobenzene (**12**) to give the 3-fluoro-4-azolylnitrobenzene intermediates **13a–g**. Reduction of the nitro group and Cbz protection of

the resulting aniline gave intermediates **14a–g**. Deprotonation with base followed by treatment with (*R*)-(–)-glycidyl butyrate afforded oxazolidinones **15a–g**.¹⁹ The hydroxymethyl side chain was then elaborated to the final acetamide analogues **3–10** via standard transformations. The synthesis of the pyrrole, pyrazole, imidazole, and 1*H*-1,2,4-triazole derivatives **3–5** and **9** proceeded smoothly. Preparation of the 1*H*-1,2,3-triazole analogue **6** and the 2*H*-1,2,3-triazole analogue **7** was made possible by the formation of both 1*H*- and 2*H*-regioisomers in the reaction between 1,2,3-triazole and 3,4-difluoronitrobenzene.

Also shown in Scheme 1, treatment of an acetonitrile solution of 3,4-difluoronitrobenzene with tetrazole in the presence of triethylamine afforded in low yield a mixture of regioisomers, which were separable. The major 2*H*-isomer **13g** was taken on to the target 2*H*-tetrazole analogue **10**. The minor 1*H*-isomer adduct was not available in sufficient quantity for preparation of the 1*H*-tetrazole analogue **11** via this route. Thus an alternate approach was used to prepare this compound (Scheme 2). This route involved the preparation of the aniline **20**, which served as a key intermediate in the synthesis of **11** and numerous other derivatives. Reaction of **12** with benzylamine followed by bis-protection afforded **17**. Oxazolidinone ring formation and side chain elaboration gave the bis-protected aniline **19**, which was deprotected to afford **20**. Reaction of **20** with triethyl orthoformate and sodium azide afforded the desired tetrazole analogue **11** directly.^{20,21}

The synthesis of the 4*H*-1,2,4-triazole analogue **9** is also shown in Scheme 2. Reaction of **20** with 1,1-thiocarbonyldi-2(1*H*)-pyridone²² gave an excellent yield of the isothiocyanate **21** which was treated with formic hydrazide to give the thiosemicarbazide **22**.²³ Cyclization of **22** in aqueous base gave the 2-mercaptotriazole intermediate **23**.²³ Treatment of this material with nitric acid afforded the desired analogue **9**.²⁴

Substituted Azole Analogues. 3-Substituted Pyrrole Analogues: Substituted pyrrole analogues were synthesized as shown in Scheme 3. The aniline **20** was condensed with dimethoxytetrahydrofuran-carboxaldehyde²⁵ **24** to yield the 3-formylpyrrole **25** which was exploited for the preparation of **26**, **27**, **29**, and **31–35** via standard transformations. Dehydration of the oxime **26** with triphenylphosphine in a mixture of acetonitrile/


Scheme 2^a

^a Reagents: (a) BnNH_2 , DIEA, CH_3CN , 90°C , 84%; (b) H_2 , 5% Pt/C, THF; (c) *N,N*-dimethylaniline, benzyl chloroformate, THF, 0°C –rt, 72% for steps b + c; (d) (i) *n*-BuLi, THF, -78°C , (ii) *R*-(–)-glycidyl butyrate, -78°C –rt, 77%; (e) MsCl , NEt_3 , CH_2Cl_2 ; (f) potassium phthalimide, CH_3CN , 95°C , 93% for steps e + f; (g) hydrazine hydrate, MeOH, 80°C , 99%; (h) Ac_2O , pyridine, 0°C –rt, 89%; (i) H_2 , 10% Pd/C, EtOH, 78%; (j) HC(OEt)_3 , NaN_3 , HOAc, reflux, 47%; (k) 1,1-thiocarbonyldi-2(1*H*)-pyridone, CH_2Cl_2 , 0°C , 94%; (l) formic hydrazide, THF, 70°C , 91%; (m) KOH, H_2O , 98%; (n) 20% HNO_3 , steam bath, NH_4OH , 77%.

CCl_4 yielded the 3-cyano analogue **28**.²⁶ The aldehyde **25** was oxidized to the ester **30** with manganese(IV) oxide according to the method of Corey and co-workers.²⁷ Analogues **38**–**41** were synthesized from 3-acetylpyrrole **36** and **12** as shown.

4-Substituted Pyrazole Analogues: Several 4-substituted pyrazole analogues were synthesized utilizing the ethyl 4-pyrazolecarboxylate analogue **47**. This material was prepared in a regioselective manner as outlined in Scheme 4. The arylhydrazine **42** was synthesized and condensed with the known (ethoxycarbonyl)malondialdehyde (**43**)²⁸ to give the ethyl 4-pyrazolecarboxylate **44**.²⁹ This material was then converted to **47** employing the azide intermediate **46** for the introduction of the acetylaminomethyl side chain. Ester **47** was further converted to **48**–**50**.

Another approach to 4-substituted pyrazoles wherein the 4-iodopyrazole **54** serves as a key intermediate was investigated. Thus, **54** was prepared in several steps as outlined in Scheme 4. Palladium-mediated carboxylation of **54** yielded an intermediate acid, which was converted to **55** via Curtius rearrangement.³⁰ This material was then deprotected and acylated to give **56**. The intermediate **52** was also used to prepare the terminal acetylene derivative **58** as shown.


Scheme 3^a

^a Reagents: (a) **20**, HOAc, reflux, 85%; (b) NH_2OH or NH_2OMe , $\text{MeOH}/\text{CH}_2\text{Cl}_2$, K_2CO_3 , 63–94% *E/Z* mixture; (c) NaBH_4 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 84%; (d) NaCN , MnO_2 , HOAc, MeCN/MeOH , 71%; (e) PPh_3 , $\text{CH}_3\text{CN}/\text{CCl}_4$, 72%; (f) NH_3/MeOH , KCN, 50°C , 20%; (g) (*i*-PrO)₂POCH₂CO₂Et, *t*-BuOK, -78°C –rt, 83%; (h) CuCl, NaBH₄, 0°C –rt, 99%; (i) LiBH₄, THF, 39%; (j) (i) MsCl , NEt_3 , CH_2Cl_2 , (ii) $\text{CH}_3\text{SO}_2\text{NH}_2$, NaH, DMF, 26%; (k) K_2CO_3 , DMSO, 50°C , 98%; (l) same steps as in Scheme 1, 19%; (m) NaBH₄, MeOH, 72%.


3-Substituted Pyrazole Analogues: The (3-aminopyrazolylphenyl)oxazolidinone **66** was targeted as a key intermediate in this series. In addition to the usual N-acylation, N-sulfonylation, and N-alkylation chemistry, we anticipated that the 3-amino derivative would allow access to the 3-cyano analogue **71** via Sandmeyer chemistry.³¹ Thus, the Boc-protected (3-aminopyrazolylphenyl)oxazolidinone **65** was prepared as outlined in Scheme 5. Removal of the Boc group and functional group manipulation gave the analogues **66**–**71**. The trifluoromethyl congener **64** was synthesized in a similar fashion (Scheme 5).

It was anticipated that the arylhydrazine **72** would be a useful intermediate in azole synthesis. This material was prepared from the aniline **20** as shown in Scheme 6. Diazotization followed by reduction of the diazonium species afforded the arylhydrazine **72**.³² Condensation of **72** with acetylacetaldehyde dimethyl acetal proceeded to give the 3- and 5-methylpyrazole analogues **73** and **74** as a 2:1 mixture (Scheme 6).³³ The hydrazine **72** was also utilized in the regiochemical synthesis of the (3-hydroxypyrazolylphenyl)oxazolidinone analogue **77**. The known ethoxycarbonyl chloride (**75**)³⁴ was reacted with **73** to yield the key acylhydrazine intermediate **76**. This material was cyclized in HCl to afford the desired analogue **77**,³⁵ which was converted to the acetoxy and methoxy analogues **78** and **79**.


Imidazole Analogues: The synthesis of the imidazole analogues of interest is outlined in Scheme 7. The 2-hydroxymethylimidazole (**80**) was converted to the oxazolidinone derivative **83** as described above for the preparation of **3**–**8**. Removal of the silyl group yielded

Scheme 4^a


^a Reagents: (a) hydrazine hydrate, K₂CO₃, CH₃CN, 94%; (b) NaOAc, EtOH, reflux, 65%; (c) H₂, 10% Pd/C, MeOH, 96%; (d) benzyl chloroformate, NaHCO₃, THF, 98%; (e) (i) *n*-BuLi, THF, -78 °C, (ii) *R*-(-)-glycidyl butyrate, -78 °C-rt, 79%; (f) MsCl, NEt₃, CH₂Cl₂, 98%; (g) NaN₃, DMF, 65 °C, 93%; (h) (i) PPh₃, THF, (ii) H₂O, 65 °C, 77%; (i) Ac₂O, pyridine, CH₂Cl₂, 99%; (j) NH₃/MeOH, KCN, 70 °C, 34%; or MeNH₂/H₂O/MeOH, 70 °C, 53%; (k) SOCl₂, DMF, 0 °C, 71%; (l) **12**, DMSO, 95 °C, 95%; (m) SnCl₂, EtOH, reflux, 88%; (n) Pd(PPh₃)₂Cl₂, CuI, 85%; (o) (i) PdOAc, dppp, CO, NEt₃, H₂O/DMF, 60%, (ii) DPPA, NEt₃, *t*-BuOH, reflux, 23%; (p) (i) TFA, CH₂Cl₂, (ii) Ac₂O, pyridine, 91%; (q) K₂CO₃, MeOH, 98%.

Scheme 5^a

^a Reagents: (a) **12**, 3-trifluoromethylpyrazole, DMSO, 90 °C, 94%; or 3-aminopyrazole, K₂CO₃, CH₃CN, 43%; (b) Boc₂O, THF, 77%; (c) same steps as in Scheme 6; (d) TFA, CH₂Cl₂, 83%; (e) benzyloxycarbonyl chloride, NEt₃, 97%; (f) Ac₂O, NEt₃, CH₂Cl₂, 78%; (g) MeSO₂Cl, pyridine, 73%; (h) HCl, NaNO₂ then CuCN, KCN, 38%; (i) H₂, 10% Pd/C, 99%.

Scheme 6^a

^a Reagents: (a) NaNO₂, HCl then SnCl₂, 85%; (b) HCl, EtOH, reflux, 80%; (c) NEt₃, THF, reflux; (d) cond HCl, reflux, 65% for steps c + d; (e) Ac₂O, 89%; (f) MeI, K₂CO₃, acetone, reflux, 74%.


Scheme 7^a

^a Reagents: (a) K₂CO₃, DMSO, 90 °C, 47%; (b) TBDMSCl, imidazole, DMF, 64%; (c) same as in Scheme 1; (d) AcOH/THF/MeOH (98%); (e) (i) (PhO)₂PON₃, DBU, (ii) 2,5-dimethyl-3-hexyne-2,5-diol, 10% Pd/C, EtOH, reflux, 25%.

the hydroxymethyl congener **84** which was treated with diphenyl phosphorazidate and DBU according to the method of Mizuno and Shiori generating an intermediate azide.³⁶ The azide was then subjected to palladium-catalyzed decomposition to the nitrile **85** according to the method of Hayashi and co-workers.³⁷

1*H*-1,2,3- and 2*H*-1,2,3-Triazole Analogues: 1*H*-1,2,3- and 2*H*-1,2,3-triazole analogues were synthesized from the aniline derivative **20** (Scheme 8). Formation of the azide **86** followed by 1,3-dipolar cycloaddition in refluxing toluene with various commercially available alkynes gave 1*H*-1,2,3-triazoles **87** and **89–91**.³⁸ These materials were converted to other analogues of interest as shown.


The primary analogue of interest in the 2*H*-1,2,3-triazole series was the cyano derivative **99**. Diazotization of aniline **20** followed by reaction of the diazonium salt with malononitrile afforded the dicyano intermediate **97**.³⁹ Treatment of this material with hydrazine

Scheme 8^a

^a Reagents: (a) (i) NaNO₂, HCl, (ii) NaN₃, NaOAc, >90%; (b) PhCH₃, reflux, 33–56%; (c) NH₂OH·HCl, K₂CO₃, MeOH, 20%; (d) NH₃/MeOH, KCN, 70 °C; or MeNH₂/MeOH 70 °C, 65–67%; (e) MeOH, NaBH₄, 88%; (f) NH₂OH or NH₂OMe, K₂CO₃, MeOH/THF, 72%; (g) (i) HNO₂, (ii) CH₂(CN)₂, 88%; (h) N₂H₄·H₂O, EtOH, reflux, 97%; (i) Cu(OAc)₂, pyridine, 100 °C, 7%; (j) (i) NaNO₂, HCl, (ii) CNCH₂CO₂CH₃, NaHCO₃, 78%; (k) LiBH₄, *i*-PrOH, 78%; (l) Swern, 17%; (m) NH₄OH, CH₃CN, 82%; (n) TFAA, pyridine, 83%.


afforded the diaminopyrazolylhydrazone intermediate **98** which was cyclized with cupric acetate⁴⁰ to give the targeted cyanotriazole **99**.

1H-1,2,4-Triazole Analogues: Several 1H-1,2,4-triazoles were also prepared from the advanced aniline intermediate **20** (Scheme 8). Diazotization and condensation with methyl isocyanoacetate afforded the ester analogue **100**.⁴¹ This material was converted to compounds **101–104** under standard conditions. The 3-methyl-1,2,4-triazole congener was prepared as shown in eq 1. Treatment of ethyl acetimidate with the arylhydra-



zine **72** in ethanol followed by addition of methanolic HCl gave an intermediate imidohydrazone, which upon refluxing in triethyl orthoformate according to the procedure of Westermann and co-workers afforded the methyltriazole **106**.⁴²

Additional 1H-1,2,4-triazole analogues were synthesized as depicted in Scheme 9. The known aminooxaziazole **107**⁴³ was prepared and converted to the iminoformate intermediate **108** with triethyl orthoformate.⁴⁴ Condensation of this ethoxyformylamino derivative **108** with the aniline **20** afforded the formamidine **109** which was then cyclized to the phenylacetylaminotriazole **110**.⁴⁴ This process however cleaved the base-sensitive oxazolidinone ring to the amino alcohol. Reformation of the oxazolidinone with triphosgene yielded the protected analogue **111**. Removal of the phenylacetyl protecting group gave the amino analogue **112** which was converted to the chloro derivative **113** via diazotization and Sandmeyer reaction with copper(I) chloride.⁴⁵


Scheme 9^a

^a Reagents: (a) HC(OEt)₃, reflux, 92%; (b) **20**, THF/EtOH, reflux, 82%; (c) 10% NaOH, EtOH, 89%; (d) triphosgene, CH₂Cl₂, 74%; (e) EtOH, HCl, reflux, 92%; (f) (i) NaNO₂, 2 N HCl, 0 °C, (ii) CuCl, rt, 22%.

2H-Tetrazole Analogues: Once again, the advanced aniline intermediate **20** was used for the preparation of the tetrazole analogues of interest (Scheme 10). Cycloaddition of the phenylsulfonylhydrazone **114**⁴⁶ with the diazonium salt of **20** in pyridine afforded the cinnamyltetrazole **115**.⁴⁶ Ozonolysis with reductive workup gave the aldehyde **116** that was further converted to **117** and **118**.

Results and Discussion

The oxazolidinone analogues prepared above were tested in vitro versus a panel of Gram-positive and Gram-negative bacterial isolates.¹⁹ Minimum inhibitory concentration (MIC) values were determined using standard agar dilution methods and are shown in Tables

Scheme 10^a

^a Reagents: (a) (i) **20**, NaNO₂, HCl, H₂O, EtOH, (ii) **114**, pyridine, 35%; (b) O₃, CH₂Cl₂, -78 °C then DMS, rt, >90%; (c) NH₂OH, K₂CO₃, CH₂Cl₂, 44%; (d) PPh₃, CCl₄, CH₃CN, 69%.

1–3. Many of the analogues tested had excellent in vitro antibacterial activity several times more potent than that of linezolid, eperezolid, and vancomycin. Notably, several analogues were very potent against the fastidious Gram-negative bacteria *H. influenzae* and *M. catarrhalis*. Also, included in Tables 1–3 is in vivo efficacy data for selected analogues tested in a lethal systemic *S. aureus* infection model in mice (po administration).¹⁹

Considering the unsubstituted azole analogues, the pyrrolyl and 1*H*-1,2,3-triazolyl derivatives **3** and **6** emerge as the most interesting followed by the imidazolyl analogue **5**. All three have very good activity against Gram-positive and Gram-negative bacteria. It is interesting to note the profound effect on Gram-negative activity exerted by the number and location of the nitrogens of the azole rings. For example, whereas the pyrrole congener **3** is very active against *H. influenzae* and *M. catarrhalis*, the pyrazole analogue **4**, with a nitrogen in the 2-position, is not. However, moving the 2-nitrogen of the pyrazole to the 3-position to give the imidazole variant **5** results in a significant increase in the Gram-negative activity. Interestingly, replacing the 2-CH of the imidazole with a nitrogen to give the 1*H*-1,2,3-triazole **6** results in one of the most potent analogues. Thus, one can conclude that a nitrogen at the 2-position is not deleterious to activity in and of itself. Likewise, the absence of a nitrogen at the 2-position does not explain the improved activity of the imidazole **5**. It appears that the overall electronics of the azole ring play a subtle, yet important, role in the antibacterial activity of these compounds, especially when one considers the Gram-negative activity.

Several of the parent azole analogues were also tested in vivo versus a lethal systemic *S. aureus* infection in a mouse model of human infection.¹⁹ The analogues were dosed orally and compared to either orally administered eperezolid or linezolid or subcutaneously dosed vancomycin as controls. Of the unsubstituted azoles, the 1*H*-1,2,3-triazole analogue **6** was the most efficacious with an ED₅₀ = 4.7 mg/kg.

With these results in hand an extensive survey of substituted derivatives within each class was undertaken in order to increase the antibacterial activity as well as to attain and/or improve in vivo potency. This discussion will be limited to monosubstituted analogues at the 3- or 4-position of the azole rings since disubstitution generally led to decreases in activity. In addition, substitution at the 2-position (i.e. α to the phenyl ring)

was not extensively explored, again due to significant losses in activity.

The pyrrole analogues were studied initially due to the excellent activity of the lead compound **3** in this series. Several 3-substituted pyrrole analogues were prepared which retain good broad-spectrum activity such as the formyl, oxime, nitrile, and hydroxymethyl derivatives **25**–**28**. Other modifications resulted in a decrease in activity across the panel of isolates. Activity against *H. influenzae* was especially sensitive to structural changes. Upon testing in vivo, the methoxime, cyano, and acetyl congeners **27**, **28**, and **38** all displayed an increase in efficacy vis-a-vis the parent pyrrole, with the cyano analogue **28** being the most efficacious (ED₅₀ = 1.9 mg/kg).





In the pyrazole series substitution at the 3- or 4-position of the ring often led to a decrease in activity. The noted exceptions to this are both the 3- and 4-cyano analogues **50** and **71** which show an increase in activity against all organisms. It is interesting to note the change in activity when moving the cyano group from the 3-position to the 4-position of the pyrazole ring. The 4-cyano congener **50** is 2–4 times more potent than the 3-cyano analogue **71**. In particular, the 4-cyano analogue has very good Gram-negative activity whereas the 3-cyano analogue has only moderate activity versus these bacteria. Furthermore, although the parent pyrazole **4** was devoid of in vivo activity, the 4-cyano-substituted compound **50** is very potent (ED₅₀ = 1.2 mg/kg). The acetylene moiety was incorporated at the 4-position for a comparison to the 4-cyano congener. Although this change resulted in a reasonably active Gram-positive compound **58**, the Gram-negative activity dropped dramatically (**58** vs **50**). Because the cyano moiety imparted such interesting activity, the corresponding analogue was targeted in the imidazole series. Unfortunately, the 4-cyanoimidazole analogue **85** did not show an improvement in antibacterial activity.

Due to the very potent in vitro and in vivo activity of the unsubstituted 1*H*-1,2,3-triazole analogue **6**, a more in-depth survey of substituent effects on the various triazoles was investigated (Table 2). Several analogues in the 1*H*-1,2,3-triazole series have excellent broad-spectrum activity. Analogues **87**, **90**, **91**, and **94** are potent antibacterial agents versus both Gram-positive and Gram-negative organisms. Once again the cyano and formyl derivatives were the most active in this series, and the cyano analogue **90** maintains excellent in vivo efficacy with an ED₅₀ = 1.9 mg/kg.

The 4-cyano analogue **99** was also prepared in the 2*H*-1,2,3-triazole series, and a significant increase in in vitro antibacterial activity was seen. Good broad-spectrum activity was attained, with the exception of *H. influenzae* where an MIC = 8 μg/mL was observed. This analogue is also active in vivo with an ED₅₀ = 3.2 mg/kg.

The 1*H*-1,2,4-triazole series was not as interesting. Although the parent analogue **8** has good in vivo activity, it lacks Gram-negative activity. Unfortunately, the incorporation of the cyano moiety, analogue **104**, in this series did not significantly increase the Gram-negative activity. Furthermore, all other substituents explored were not beneficial. One interesting exception is the chloro derivative **113** that is equipotent with

Table 1. Antibacterial Activity (MIC, $\mu\text{g/mL}$) of (Pyrrolylphenyl)-, (Pyrazolylphenyl)-, and (Imidazolylphenyl)oxazolidinones

<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>3, 25-35, 38-41</p> </div> <div style="text-align: center;">  <p>4, 47-50, 54-58</p> </div> <div style="text-align: center;">  <p>64-66, 68-71, 73, 77-79</p> </div> <div style="text-align: center;">  <p>5, 85</p> </div> </div>										
compd	R	<i>S.a.</i> ^a	<i>S.a.</i> ^b	<i>S.e.</i> ^c	<i>S.p.</i> ^d	<i>E.f.</i> ^e	<i>H.inf.</i> ^f	<i>M.cat.</i> ^g	ED ₅₀ ^h	control ED ₅₀ ⁱ
Pyrrolyl										
3	H	1	1	≤0.5	≤0.5	1	4	2	13.2 (7.9–32.5)	2.7 (ND) ^v
25	CHO	0.25	0.25	0.125	≤0.06	0.25	2	1	>20	1.7 (0.9–2.6) ^v
26	CH=NOH	1	0.5	0.25	≤0.125	0.5	4	1	5.5 (3.5–9.1)	2.6 (2.2–2.9) ^e
27	CH=NOCH ₃	1	1	0.5	0.25	1	16	16	7.3 (4.3–13.3)	3.5 (2.2–6.1) ^e
28	CN	0.5	0.25	≤0.125	≤0.125	0.25	4	1	1.9 (1.1–2.7)	4.0 (0.1–6.6) ^e
29	CH ₂ OH	1	1	0.25	0.25	1	4	2	>20	0.9 (0.5–1.5) ^v
30	CO ₂ CH ₃	2	2	0.5	0.25	1	>16	16	>20	2.9 (1.8–4.4) ^e
31	CONH ₂	16	8	1	0.25	4	>16	>16		
32	CH=CHCO ₂ Et	4	4	2	0.5	2	32	4	>20	2.1 (1.3–3.7) ^e
33	(CH ₂) ₂ CO ₂ Et	4	4	1	0.5	2	>16	2		
34	(CH ₂) ₃ OH	4	4	1	0.5	2	>16	8		
35	(CH ₂) ₃ NHSO ₂ Me	8	4	2	0.5	2	>16	8	>20	1.8 (0.8–2.3) ^e
38	COCH ₃	2	1	0.5	0.25	1	16	4	6.5 (ND)	10 (ND) ^e
39	CH(OH)CH ₃	16	16	4	2	8	>16	>16		
40	C(NO ₂)CH ₃	2	1	0.5	0.25	1	8	2	15 (ND)	10 (ND) ^e
41	C(NOCH ₃)CH ₃	4	2	1	0.5	2	>16	8		
Pyrazolyl										
4	H	2	2	1	1	2	32	16	>20	1.8 (1.1–2.8) ^v
47	CO ₂ Et	4	4	2	1	4	>16	>16		
48	CONH ₂	>16	>16	4	0.5	>16	>16	16		
49	CONHMe	>16	>16	2	0.5	16	>16	16		
50	CN	0.5	0.5	0.25	≤0.125	0.5	4	2	1.2 (0.8–2.0)	3.3 (1.8–7.0) ^e
54	I	2	2	1	0.5	2	>16	8		
55	NHBoc	8	4	2	1	8	>16	>16		
56	NHAc	>16	16	2	1	16	>16	>16		
57	CC-TMS	4	4	2	2	8	>16	>16		
58	CCH	2	2	1	0.5	2	>16	8		
64	CF ₃	4	4	2	1	8	>16	>16		
65	NH–Boc	4	2	2	0.5	2	>16	8		
66	NH ₂	2	4	1	0.5	4	16	16		
68	NHCOCH ₂ OH	16	16	4	1	8	>16	>16		
69	NHAc	8	8	2	0.5	4	>16	>16		
70	NHSO ₂ Me	>16	>16	4	4	>16	>16	>16		
71	CN	1	1	0.5	0.5	2	16	8		
73	CH ₃	4	4	1	1	4	>16	16		
77	OH	8	8	1	1	8	16	8		
78	OAc	8	8	2	2	16	16	16		
79	OCH ₃	4	4	2	2	8	>16	>16		
Imidazolyl										
5	H	2	1	≤0.5	≤0.5	2	8	4	9.5 (5.5–22.1)	2.2 (1.43–3.5) ^v
85	CN	4	4	0.5	0.25	2	8	4		
linezolid		4	2	1	1	4	16	8	5.6 (2.9–8.5)	3.9 (2.5–6.4) ^v
eperezolid		4	1	0.5	0.5	2	16	8	1.9 (1.4–3.8)	3.9 (2.5–6.4) ^v
vancomycin		1	1	2	0.5	4	>32	>32	3.9 (2.5–6.4)	

^a Methicillin-susceptible *S. aureus* UC9213. ^b Methicillin-resistant *S. aureus* UC6685. ^c Methicillin-resistant *S. epidermidis* UC12084. ^d *Str. pneumoniae* UC9912. ^e *E. faecalis* UC9217. ^f *H. influenzae* UC30063. ^g *M. catarrhalis* UC30610. Minimum inhibitory concentration (MIC): lowest concentration of drug ($\mu\text{g/mL}$) that inhibits visible growth of the organism. ^h ED₅₀ is the amount of drug required after oral administration (mg/kg/day) to cure 50% of infected mice subjected to a lethal systemic infection of *S. aureus*. Numbers in parentheses are 95% confidence ranges. Data shown is from one experiment ($n = 36$ mice/drug). ⁱ v = vancomycin, e = eperezolid, l = linezolid as controls. ^j ND = not determined.

linezolid against Gram-positive bacteria and has excellent activity against *M. catharralis*.





In the tetrazole series, the 5-cyano-2*H*-tetrazole derivative **118** has good Gram-positive activity, but it is only moderately active against the Gram-negative organisms and shows no appreciable benefit over linezolid.

Conclusion

In conclusion, an effort to expand the spectrum of antibacterial activity of the oxazolidinones to include Gram-negative organisms has led to the discovery of the

(azolylphenyl)oxazolidinone subclass. Certain members of this class have very potent activity versus both Gram-positive and Gram-negative organisms. Interesting differences in antibacterial activity were seen in many analogues that cannot be rationalized solely on the basis of sterics and position/number of nitrogen atoms in the azole ring. It seems that differences in activity rely strongly on subtle changes in the electronic character of the overall azole system. Electronic differences or changes in dipole may effect such characteristics as partitioning through the bacterial cell wall and/or efflux

Table 2. Antibacterial Activity (MIC, $\mu\text{g/mL}$) of (Triazolylphenyl)oxazolidinones

<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>6, 87-96</p> </div> <div style="text-align: center;">  <p>7, 99</p> </div> <div style="text-align: center;">  <p>8, 100-104, 106, 111-113</p> </div> <div style="text-align: center;">  <p>9</p> </div> </div>										
compd	R	<i>S.a.</i> ^a	<i>S.a.</i> ^b	<i>S.e.</i> ^c	<i>S.p.</i> ^d	<i>E.f.</i> ^e	<i>H.inf.</i> ^f	<i>M.cat.</i> ^g	ED ₅₀ ^h	control ED ₅₀ ⁱ
1H-1,2,3										
6	H	1	1	≤0.5	≤0.5	1	4	2	4.7 (3.1–7.8)	2.0 (ND) ^v
87	COCH ₃	1	1	1	0.5	2	8	8	8.8 (5.7–14.5)	4.4 (2.9–7.1)e
88	C(NO ₂)Me	4	4	2	0.5	8	16	4		
89	CO ₂ CH ₃	4	4	2	1	4	>16	16	>20	15.3 (ND)e
90	CN	1	0.5	0.25	≤0.125	1	2	2	1.9 (1.1–3.3)	2.8 (1.7–4.5)e
91	CHO	0.5	0.25	≤0.125	0.25	0.5	4	2	>20	1.1 (0.5–1.5)e
92	CONH ₂	4	4	1	0.5	8	>16	8		
93	CONHCH ₃	8	4	2	1	8	>16	>16		
94	CH ₂ OH	4	4	0.5	0.25	2	4	2		
95	HC=NOH	8	2	1	0.5	8	>16	8		
96	HC=NOMe	4	2	1	0.5	4	>16	16	8.6 (ND)	3.8 (1.9–6.6)e
2H-1,2,3										
7	H	4	2	1	1	4	>16	>16		
99	CN	0.5	0.5	≤0.125	≤0.125	0.5	8	2	3.2 (2.0–4.9)	3.8 (1.9–6.6)e
1H-1,2,4										
8	H	4	4	1	1	4	32	16	5.7 (3.1–13.1)	2.4 (1.0–3.2)v
100	CO ₂ CH ₃	>16	>16	4	4	>16	>16	>16		
101	CH ₂ OH	16	16	2	1	16	16	8		
102	CHO	4	2	0.5	0.5	4	8	8		
103	CONH ₂	>16	>16	16	2	>16	>16	16		
104	CN	4	2	2	1	4	16	16		
106	CH ₃	8	4	2	1	8	>16	16		
111	NHCOBn	>16	>16	>16	2	>16	>16	>16		
112	NH ₂	8	4	2	1	16	>16	16		
113	Cl	4	2	0.5	≤0.125	2	16	2	9.4 (6.2–16.5)	5.0 (ND)
4H-1,2,4										
9	na	16	8	4	1	8	>16	8	>20	2.0 (1.1–2.8)e
linezolid		4	2	1	1	4	16	8	5.6 (2.9–8.5)	3.9 (2.5–6.4)v
eperezolid		4	1	0.5	0.5	2	16	8	1.9 (1.4–3.8)	3.9 (2.5–6.4)v
vancomycin		1	1	2	0.5	4	>32	>32	3.9 (2.5–6.4)	

^a Methicillin-susceptible *S. aureus* UC9213. ^b Methicillin-resistant *S. aureus* UC6685. ^c Methicillin-resistant *S. epidermidis* UC12084. ^d *Str. pneumoniae* UC9912. ^e *E. faecalis* UC9217. ^f *H. influenzae* UC30063. ^g *M. catarrhalis* UC30610. Minimum inhibitory concentration (MIC): lowest concentration of drug ($\mu\text{g/mL}$) that inhibits visible growth of the organism. ^h ED₅₀ is the amount of drug required after oral administration (mg/kg/day) to cure 50% of infected mice subjected to a lethal systemic infection of *S. aureus*. Numbers in parentheses are 95% confidence ranges. Data shown is from one experiment ($n = 36$ mice/drug). ⁱ v = vancomycin, e = eperezolid, l = linezolid as controls. ^j ND = not determined.

mechanisms which have been shown to be responsible for the lack of activity of the oxazolidinones against Gram-negative bacteria.^{47,48} The placement of a cyano moiety on the azole often leads to a significant increase in antibacterial activity in vitro and in vivo. In particular, the 3-cyanopyrrole, 4-cyanopyrazole, and 4-cyano-1H-1,2,3-triazole congeners **28**, **50**, and **90** had *S. aureus* MICs ≤ 0.5–1 mg/mL and *H. influenzae* and *M. catarrhalis* MICs = 2–4 mg/mL. Furthermore, these analogues are very effective versus *S. aureus* and *S. pneumoniae*¹⁸ in mouse models of human infection.

Experimental Section



General. Melting points were determined on a Fisher-Johns or a Thomas-Hoover apparatus and are uncorrected. ¹H NMR spectra were recorded on either a Bruker AM-300 or ARX-400 spectrometer. Chemical shifts are reported in δ units (ppm) relative to TMS as internal standard. Mass spectra and combustion analyses were obtained by the Structural, Analytical and Medicinal Chemistry Department of Pharmacia & Upjohn, Inc. Unless otherwise indicated all reactions were conducted in commercially available anhydrous solvents under

a nitrogen atmosphere in oven- or flame-dried glassware. Chromatography was carried out on EM Science 230–400 mesh ASTM silica gel. Elemental analyses were within $\pm 0.4\%$ of calculated values. Biological assays were performed as described previously in ref 19.

3-Fluoro-1-nitro-4-(1H-pyrrol-1-yl)benzene (13a). Sodium hydride (630 mg of 60% in oil, 15.75 mmol) was suspended in THF (85 mL) and treated with pyrrole (1.0 g, 15 mmol), followed by warming at 50 °C for 15 min. The solution was then treated with 3,4-difluoronitrobenzene (2.51 g, 15.75 mmol) and refluxed for 18 h. The mixture was cooled and treated with saturated ammonium chloride solution (10 mL). The mixture was diluted with ethyl acetate and washed with water (2×). Drying (Na₂SO₄) and concentration in vacuo afforded a dark brown solid, which was chromatographed over 75 g silica gel, eluting with 30% CH₂Cl₂/hexane. These procedures afforded 2.05 g (66%) of **13a** as a light yellow solid: mp 101–102 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.14 (m, 2 H), 7.56 (t, $J = 8.7$ Hz, 1 H), 7.16 (m, 2 H), 6.44 (m, 2 H); HRMS (EI) calcd for C₁₀H₇FN₂O₂ 206.0492, found 206.0504.

General Procedure for the Preparation of 3-Fluoro-1-nitro-4-(azoly)benzenes 13b–f, 37, 61, 62, and 81. The azole (1 equiv), K₂CO₃ or dibasic potassium phosphate (2 equiv) and 3,4-difluoronitrobenzene (1 equiv) were heated to 90 °C

Table 3. Antibacterial Activity (MIC, $\mu\text{g/mL}$) of (Tetrazolylphenyl)oxazolidinones

										
compd	R	<i>S.a.</i> ^a	<i>S.a.</i> ^b	<i>S.e.</i> ^c	<i>S.p.</i> ^d	<i>E.f.</i> ^e	<i>H.inf.</i> ^f	<i>M.cat.</i> ^g	ED ₅₀ ^h	control ED ₅₀ ⁱ
2H-Tetrazolyl										
10	H	2	2	1	0.5	1	8	8		
116	CHO	8	4	2	1	16	16	16		
117	CH=NOH	4	4	1	0.5	4	16	8		
118	CN	2	1	0.5	0.5	2	16	4		
1H-Tetrazolyl										
11	na	4	2	2	1	2	16	16	>20	7.8 (4.0–26.7)e
linezolid		4	2	1	1	4	16	8	5.6 (2.9–8.5)	3.9 (2.5–6.4)v
eperezolid		4	1	0.5	0.5	2	16	8	1.9 (1.4–3.8)	3.9 (2.5–6.4)v
vancomycin		1	1	2	0.5	4	>32	>32	3.9 (2.5–6.4)	

^a Methicillin-susceptible *S. aureus* UC9213. ^b Methicillin-resistant *S. aureus* UC6685. ^c Methicillin-resistant *S. epidermidis* UC12084. ^d *Str. pneumoniae* UC9912. ^e *E. faecalis* UC9217. ^f *H. influenzae* UC30063. ^g *M. catarrhalis* UC30610. Minimum inhibitory concentration (MIC): lowest concentration of drug ($\mu\text{g/mL}$) that inhibits visible growth of the organism. ^h ED₅₀ is the amount of drug required after oral administration (mg/kg/day) to cure 50% of infected mice subjected to a lethal systemic infection of *S. aureus*. Numbers in parentheses are 95% confidence ranges. Data shown is from one experiment ($n = 36$ mice/drug). ⁱ v = vancomycin, e = eperezolid, l = linezolid as controls.

in DMSO for 18 h. The reaction was cooled diluted with water (100 mL) and extracted with ethyl acetate (2×100 mL). The organic layers were washed with water (5×100 mL) and brine (75 mL). Drying (Na_2SO_4) and concentration in vacuo afforded crude products.

3-Fluoro-1-nitro-4-(1H-pyrazol-1-yl)benzene (13b). Workup afforded an off-white solid which was crystallized from hot hexanes to afford 5.3 g (87%) of an off-white solid: mp 128–129 °C; ¹H NMR (400 MHz, CDCl_3) δ 8.28 (m, 1 H), 8.17 (m, 3 H), 7.82 (d, $J = 1.4$ Hz, 1 H), 6.58 (m, 1 H); MS (EI) m/z 207 ($M + \text{H}$); HRMS (EI) calcd for $\text{C}_9\text{H}_6\text{FN}_3\text{O}_2$ 207.0444, found 207.0454. Anal. Calcd for $\text{C}_9\text{H}_6\text{FN}_3\text{O}_2$: C, 52.18; H, 2.92; N, 20.28. Found: C, 52.01; H, 2.75; N, 20.22.

3-Fluoro-1-nitro-4-(2H-1,2,3,4-tetrazol-1-yl)benzene (13g). Tetrazole (15.83 g, 0.23 mol), 3,4-difluoronitrobenzene (7.19 g, 45.16 mmol) and NEt_3 (31.5 mL, 0.23 mol) in 90 mL acetonitrile was heated at reflux for 18 h, followed by cooling and concentration in vacuo. The residue was dissolved in EtOAc (250 mL) and washed with water (2×100 mL) and brine (100 mL). Drying (Na_2SO_4) and concentration in vacuo afforded a black oil that was chromatographed over 360 g silica gel, eluting with 3% acetone/ CH_2Cl_2 . These procedures afforded 0.93 g (10%) of the less polar 2H-regioisomer **13g** as a yellow solid and 0.38 g (4%) of the more polar 1H-regioisomer as a yellow-brown solid. **13g**: mp 111–115 °C; ¹H NMR (400 MHz, DMSO) δ 9.71 (s, 1 H), 8.57 (dd, $J = 9.6, 1.6$ Hz, 1 H), 8.32 (m, 2 H); HRMS (FAB) calcd for $\text{C}_7\text{H}_4\text{FN}_5\text{O}_2 + \text{H}$ 210.0427, found 210.0424. Anal. Calcd for $\text{C}_7\text{H}_4\text{FN}_5\text{O}_2$: C, 40.20; H, 1.93; N, 33.49. Found: C, 40.28; H, 1.87; N, 33.26.

1H-isomer: ¹H NMR (DMSO) δ 10.04 (s, 1 H), 8.57 (dd, $J = 10.4, 2.0$ Hz, 1 H), 8.34 (d, $J = 8.9$ Hz, 1 H), 8.23 (t, $J = 8.1$ Hz, 1 H).

General Procedure for the Preparation of 3-Fluoro-1-(phenylmethoxycarbonylamino)-4-(azoly)benzenes 14a–g. Compounds **13a–g** (1 equiv) and the indicated catalyst in THF were hydrogenated for 18 h. The mixture was filtered through Celite treated with saturated NaHCO_3 solution and cooled to –20 °C. Benzyl chloroformate (1.2 equiv) was added followed by warming to ambient temperature for 48 h. The mixture concentrated to ca. half the original volume diluted with EtOAc and washed with water ($4 \times$) and brine. Drying (Na_2SO_4) and concentration in vacuo afforded crude products **14a–g** which were purified as indicated.

3-Fluoro-1-(phenylmethoxycarbonylamino)-4-(1H-pyrrol-1-yl)benzene (14a). 5% Pt on sulfide carbon was used as catalyst and workup afforded a brown solid, which was chromatographed over 35 g silica gel, eluting with 1:3:32 to 1:3:16 $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ /hexane gradient to yield 581 mg (77%)

of **14a** as a brown solid: mp 124–126 °C; ¹H NMR (400 MHz, CDCl_3) δ 7.40 (m, 7 H), 7.08 (d, $J = 8.7$ Hz, 1 H), 6.98 (m, 2 H), 6.79 (bs, 1 H), 6.35 (m, 2 H), 5.23 (s, 2 H); HRMS (EI) calcd for $\text{C}_{18}\text{H}_{15}\text{FN}_2\text{O}_2$ 310.1117, found 310.1132. Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{FN}_2\text{O}_2$: C, 69.67; H, 4.87; N, 9.03. Found: C, 69.32; H, 4.89; N, 9.07.

General Procedure for Oxazolidinone Ring Formation from Cbz-Protected Anilines 14a–g To Give Compounds 15a–g and 18. A solution of Cbz derivative **14a–g** (1 equiv) THF at –78 °C was treated with 1.0 M lithium bis(trimethylsilyl)amide (1.1 equiv) followed by stirring at –78 °C for 30 min. The solution was treated with (*R*)-(–)-glycidyl butyrate (1 equiv) followed by gradual warming to ambient temperature for 24 h. The mixture was diluted with EtOAc and washed with H_2O and brine. Drying (Na_2SO_4) and concentration in vacuo afforded crude products **15a–g**.

(S)-[3-[3-Fluoro-4-(1H-pyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methanol (15a). Workup gave a red-brown oil which was subjected to radial chromatography on a 2 mm plate, eluting with 2–5% $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ gradient. These procedures afforded 157 mg (59%) of **15a** as a light tan solid: mp 112–113 °C; $[\alpha]_D^{25} = -46^\circ$ (c 0.84, DMSO); ¹H NMR (400 MHz, CDCl_3) δ 7.63 (dd, $J = 13.2, 2.5$ Hz, 1 H), 7.37 (m, 3 H), 7.00 (m, 2 H), 6.36 (m, 2 H), 4.79 (m, 1 H), 4.05 (m, 3 H), 3.79 (dd, $J = 12.7, 3.7$ Hz, 1 H); HRMS (EI) calcd for $\text{C}_{14}\text{H}_{13}\text{FN}_2\text{O}_3$ 276.0910, found 276.0918. Anal. Calcd for $\text{C}_{14}\text{H}_{13}\text{FN}_2\text{O}_3$: C, 60.87; H, 4.74; N, 10.14. Found: C, 60.97; H, 4.77; N, 9.84.

General Procedure for Conversion of Alcohols 15a–g to Final Acetamide Analogues 3–8 and 10. The alcohols **15a–g** (1 equiv) and NEt_3 (1.75 equiv) were cooled to 0 °C in CH_2Cl_2 and treated with methanesulfonyl chloride (1.25 equiv) followed by stirring at 0 °C for 30 min. The solution was warmed to ambient temperature, diluted with CH_2Cl_2 , and washed with water ($3 \times$) and brine. Drying (Na_2SO_4) and concentration in vacuo afforded intermediate mesylates that were dissolved in DMF and treated with sodium azide (10 equiv) at 60 °C for 18 h. The mixture was cooled, diluted with EtOAc and washed with water ($6 \times$) and brine. Drying (Na_2SO_4) and concentration in vacuo afforded intermediate azides sufficiently pure for use. Then, a solution of the azide (1 equiv) in $\text{CH}_3\text{OH}/\text{THF}$ was treated with an appropriate catalyst and hydrogenated at 1 atm for 24 h. The solution was filtered through Celite and concentrated in vacuo and the residue was dissolved in pyridine and treated with acetic anhydride (1.1 equiv) followed by stirring at ambient temperature for 12 h. The solution was concentrated under high vacuum to yield crude products **3–8** and **10**.

(S)-N-[[3-[3-Fluoro-4-(1H-pyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (3). Workup afforded a brown solid which was chromatographed over 50 g of silica gel, eluting with 1–2% CH₃OH/CH₂Cl₂ gradient to yield 582 mg (56%) of **3** as an off-white solid: mp 198–199.5 °C; [α]_D²⁵ = –26° (c 0.93, DMSO); ¹H NMR (400 MHz, CD₃OD) δ 7.70 (dd, *J* = 18.3, 3.3 Hz, 1 H), 7.47 (t, *J* = 11.7 Hz, 1 H), 7.37 (dd, *J* = 11.8, 3.2 Hz, 1 H), 7.01 (m, 2 H), 6.27 (m, 2 H), 4.84 (m, 1 H), 4.18 (t, *J* = 12.1 Hz, 1 H), 3.85 (dd, *J* = 12.3, 8.5 Hz, 1 H), 3.57 (d, *J* = 6.6 Hz, 2 H), 1.97 (s, 3 H); HRMS (EI) calcd for C₁₆H₁₆FN₃O₃ 317.1176, found 317.1183. Anal. Calcd for C₁₆H₁₆FN₃O₃: C, 60.56; H, 5.08; N, 13.24. Found: C, 60.28; H, 5.10; N, 13.14.

N-(4-Amino-2-fluorophenyl)-N-benzylamine (16). To a solution of 3,4-difluoronitrobenzene (0.7 mL, 6.29 mmol) in acetonitrile (12 mL) was added *N,N*-diisopropylethylamine (1.64 mL, 9.43 mmol) followed by benzylamine (0.82 mL, 7.54 mmol). After heating at 90 °C in an oil bath for 6 h the reaction was allowed to cool and stand at room temperature for 16 h and then concentrated in vacuo. The residue was treated with EtOAc and the suspended solid was removed by filtration and washed with additional EtOAc. The combined filtrates were concentrated in vacuo and the residue was chromatographed over silica gel with 5% EtOAc/hexane to yield after recrystallization from EtOAc/hexane 1.19 g (77%) of intermediate *N*-benzylnitroaniline. A solution of this material (0.50 g, 2.03 mmol) in THF (150 mL) containing of 5% platinum-on-carbon (0.1 g) was shaken with H₂ on a Parr hydrogenator for 6 h. The catalyst was removed by filtration through Celite and the filtrate was concentrated in vacuo to yield 0.54 g of **16** as a dark oil that was used as is in the next step: ¹H NMR (300 MHz, CDCl₃) δ 3.91 (broad s, 3 H), 4.29 (s, 2 H), 6.52 (m, 3 H), 7.35 (m, 5 H).

Benzyl Benzyl(4-[(benzyloxy)carbonyl]amino)-2-fluorophenyl carbamate (17). To a solution of **16** (0.54 g, 4.47 mmol) in THF (36 mL) was added *N,N*-dimethylaniline (0.58 mL, 4.57 mmol) followed by benzyl chloroformate (0.64 mL, 4.47 mmol). After 40 min the ice bath was removed and then after stirring at room temperature for 17 h the reaction mixture was concentrated in vacuo. A solution of the residue in CH₂Cl₂ (100 mL) was washed in turn with cold 1 N HCl (3×), water, and aqueous NaHCO₃, dried over MgSO₄ and concentrated in vacuo. The residue was chromatographed over silica gel with 10–15% EtOAc/hexane gradient to yield 0.74 g (72%) of **17**: mp 105–106.5 °C. An analytical sample was recrystallized from methyl *tert*-butyl ether: mp 107–108 °C; ¹H NMR (300 MHz, CHCl₃) δ 4.77 (broad s, 2 H), 5.16 (m, 4 H), 6.85–7.36 (m, 19 H); MS *m/z* 484 (M + H). Anal. Calcd for C₂₉H₂₅FN₂O₄: C, 71.89; H, 5.20; N, 5.78. Found: C, 71.86; H, 5.27; N, 5.74.

Benzyl 4-[(5S)-5-[(Acetyl amino)methyl]-2-oxo-1,3-oxazolidin-3-yl]-2-fluorophenyl(benzyl)carbamate (19). Methanesulfonyl chloride (0.16 mL, 2.05 mmol) was added dropwise to a solution of **18** (0.71 g, 1.58 mmol) and NEt₃ (0.48 mL, 3.47 mmol) in CH₂Cl₂ (25 mL) at 0 °C. After stirring in the cold for 45 min the reaction was washed with water, aqueous NaHCO₃, and brine, dried over MgSO₄ and concentrated in vacuo to yield 0.86 g of the mesylate as a gummy residue. This material was placed in acetonitrile (25 mL) and potassium phthalimide (0.875 g, 4.73 mmol) was added. The reaction was heated at 95 °C for 48 h and allowed to cool. A suspended solid was collected on a filter and washed well with acetonitrile. The combined filtrates were concentrated in vacuo. The residue was chromatographed over silica gel with 0.5% MeOH/CH₂Cl₂ to yield 0.85 g (93%, two steps) of the phthalimide. This phthalimide (0.85 g, 1.47 mmol) was suspended in MeOH and hydrazine monohydrate (0.075 g, 0.073 mL) was added. The mixture was heated at 80 °C for 5 h and then allowed to cool and stand at room temperature. After standing for 16 h the reaction mixture was poured into 3% aqueous sodium carbonate (25 mL) and extracted with EtOAc. The combined extracts were dried over MgSO₄ and concentrated in vacuo to give 0.65 g of the free amine as a gummy residue which was placed in pyridine (8.5 mL) at 0 °C. Acetic

anhydride (2.8 mL) was added dropwise and the reaction was stirred for 18 h at room temperature and partitioned between water and EtOAc. The mixture was extracted with EtOAc (3×) and the combined extracts were dried over MgSO₄ and concentrated in vacuo. The residue was chromatographed over silica gel with 1% MeOH/0.1% NH₄OH/CH₂Cl₂ followed by 2% MeOH/0.2% NH₄OH/CH₂Cl₂ to yield 0.36 g of pure product **19** followed by 0.25 g of product mixed with a small amount of polar impurity: mp 64–66 °C; [α]_D –13° (c 0.93, EtOH); ¹H NMR (300 MHz, CDCl₃) δ 2.00 (s, 3 H), 3.64 (m, 2 H), 3.71 (m, 1 H), 3.98 (t, 1 H), 4.74 (m, 1 H), 4.81 (s, 2 H), 5.15, 5.23 (s, s, 2 H), 6.15 (broad s, 1 H), 6.95–7.52 (m, 13 H); HRMS calcd for C₂₇H₂₆FN₃O₅ 491.1856, found 491.1860. Anal. Calcd for C₂₇H₂₆FN₃O₅: C, 65.98; H, 5.33; N, 8.55. Found: C, 65.29; H, 5.41; N, 8.44.

N-[(5S)-3-(4-Amino-3-fluorophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (20). To a solution of **19** (2.381 g, 4.84 mmol) in absolute ethanol (150 mL) was added 0.5 g of 10% Pd/C catalyst. The mixture was placed on a Parr hydrogenator for 7.5 h. There was then added an additional 0.1 g of 10% Pd/C catalyst and the reaction was allowed to continue. After an additional 15.5 h the catalyst was removed by filtration through Celite. The filtrate was concentrated in vacuo and the residue was recrystallized from MeOH to yield **20**: 1.0 g (77%); mp 168–169 °C; [α]_D –24° (c 0.70, DMSO); ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.83 (s, 3 H), 3.38 (m, 2 H), 3.63 (d, d, 1 H), 4.01 (t, 1 H), 4.65 (m, 1 H), 5.04 (s, 2 H), 6.76 (t, 1 H), 6.96 (d, d, 1 H), 7.30 (dd, 1 H), 8.25 (t, 1 H); MS (EI) *m/z* 267 (M + H). Anal. Calcd for C₁₂H₁₄N₃F₃O₃: C, 53.93; H, 5.28; N, 15.72. Found: C, 53.90; H, 5.36; N, 15.66.

(S)-N-[[3-[3-Fluoro-4-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (11). A slurry of **20** (500 mg, 1.87 mmol), sodium azide (182 mg, 2.81 mmol), and triethyl orthoformate (450 mg, 3.02 mmol) in acetic acid (10 mL) was refluxed for 4 h. The mixture was cooled and added to ice water (20 mL). After setting at ambient temperature for 48 h, the precipitated product was collected by filtration and washed with cold CH₃OH to yield 172 mg (29%) of **11**: mp 167–169 °C; ¹H NMR (400 MHz, DMSO) δ 9.96 (s, 1 H), 8.25 (t, *J* = 5.76 Hz, 1 H), 7.89 (t, *J* = 8.7 Hz, 1 H), 7.85 (dd, *J* = 13.2, 2.4 Hz, 1 H), 4.79 (m, 1 H), 4.20 (t, *J* = 9.1 Hz, 1 H), 3.81 (dd, *J* = 9.2, 6.5 Hz, 1 H), 3.44 (t, *J* = 5.4 Hz, 1 H); HRMS (FAB) calcd for C₁₃H₁₃FN₆O₃ + H 321.1111, found 321.1118. Anal. Calcd for C₁₃H₁₃FN₆O₃·1/3CH₃OH: C, 48.39; H, 4.37; N, 25.39. Found: C, 48.72; H, 4.29; N, 24.71.

N-[(5S)-3-(3-Fluoro-4-isothiocyanatophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (21). To a solution of 1,1-thiocarbonyldi-2(1*H*)-pyridone (0.96 g, 4.12 mmol) in CH₂Cl₂ (30 mL) at 0 °C was added **20** (1.0 g, 3.74 mmol) in one portion. Over the course of 1 h **20** slowly went into solution as a second material came out. After stirring in the cold for 2.5 h and then at room temperature for 2 h, the reaction mixture was concentrated in vacuo. The residue was treated with water (15 mL) and the solid was finely divided with a spatula and collected on a filter, washed with a small amount of water and dried overnight under vacuum at 55 °C to yield 1.09 g (94%) of **21**: mp 173–176 °C; MS *m/z* 309 (M + H).

N-[(5S)-3-(3-Fluoro-4-[(2-formylhydrazino)carbothioyl]amino)phenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (22). Formic hydrazide (0.23 g) was added to a mechanically stirred mixture of **21** (1.11 g, 3.6 mmol) in THF (25 mL). The mixture was heated at 70 °C and over 3.5 h a heavy precipitate came out of solution. The mixture was then allowed to cool and stir at room temperature for 1 h. The suspended solid was collected and washed well with EtOAc. The resulting solid was dried under vacuum at 60 °C to yield 1.2 g (91%) of **22** that was carried on to the next reaction without further purification.

(S)-N-[[3-[4-[3-Mercapto-4-[1,2,4]triazolyl]-3-fluorophenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (23). To a mechanically stirred suspension of **22** (1.10 g, 3.0 mmol) in water (22 mL) was added 1 N KOH (3.1 mL). Over 30 min the solid gradually became a finely divided suspension. After stirring for 1 h at room temperature the reaction mixture was

acidified with 1 N HCl (3 mL). The resulting precipitate was collected, washed with a little 0.1 N HCl and dried under vacuum at 60 °C to yield 1.03 g (98%) of **23**: mp 266.5–268 °C. An analytical sample was recrystallized from DMF:H₂O: mp 272–273 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.84 (s, 3 H), 3.44 (m, 2 H), 3.78 (m, 1 H), 4.19 (t, 1 H), 4.78 (m, 1 H), 7.48 (d, 1 H), 7.66 (t, 1 H), 7.75 (d, 1 H), 8.29 (t, 1 H), 8.70 (s, 1 H), 14.02 (broad s, 1 H); MS *m/z* 351 (M + H). Anal. Calcd for C₁₄H₁₄FN₅O₃S: C, 47.86; H, 4.02; N, 19.93. Found: C, 48.01; H, 4.29; N, 19.70.

(S)-N-[[3-[4-[4-[1,2,4]Triazolyl]-3-fluorophenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (9). A mixture of **23** (0.33 g, 0.94 mmol) and 20% HNO₃ (0.6 mL) were heated on a steam bath in a test tube with vigorous effervescing. After 1 min the sides of the tube were rinsed down with an additional 0.6 mL of 20% HNO₃. Heating was continued for 7 min until the solid dissolved and then the reaction was cooled in an ice bath. The pH was adjusted to 9.5 with 1 N NH₄OH (5.7 mL) and the solution was concentrated in vacuo. The residue was chromatographed over silica gel with 5% MeOH/0.5% NH₄OH/CH₂-Cl₂ to yield 0.23 g (77%) of **9**: mp 204–206 °C; [α]_D²⁵ = –26° (c 0.92, DMSO); ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.82 (s, 3 H), 3.42 (t, 2 H), 3.78 (m, 1 H), 4.16 (t, 1 H), 4.75 (m, 1 H), 7.46 (dd, 1 H), 7.72 (t, 1 H), 7.77 (dd, 1 H), 8.24 (t, 1 H), 8.90 (s, 1 H), 8.91 (s, 1 H); HRMS (FAB) calcd for C₁₄H₁₄FN₅O₃ + H₁ 320.1159, found 320.1159. Anal. Calcd for C₁₄H₁₄FN₅O₃: C, 52.67; H, 4.42; N, 21.93. Found: C, 52.42; H, 4.56; N, 21.68.

(S)-N-[[3-[3-Fluoro-4-(1H-pyrrol-1-yl)-3-carboxaldehyde]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (25). A solution of **20** (2.0 g, 7.5 mmol) and 2,5-dimethoxy-3-tetrahydrofuran-2-carboxaldehyde (1.68 g, 10.5 mmol) in acetic acid (55 mL) was refluxed for 2 h. The solution was cooled and the solvent removed under high vacuum, azeotroping the residue with toluene to remove the last traces of acetic acid. The residue was chromatographed over 300 g of silica gel with 0–3% CH₃OH/CH₂Cl₂ to yield 2.21 g (85%) of **25** as a light yellow amorphous solid: mp 165–168 °C; [α]_D²⁴ = –24° (c 0.38, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 9.86 (s, 1 H), 7.70 (dd, *J* = 13.0, 2.4 Hz, 1 H), 7.59 (q, *J* = 1.9 Hz, 1 H), 7.41 (t, *J* = 8.7 Hz, 1 H), 7.27 (d, *J* = 6.9 Hz, 1 H), 7.0 (m, 1 H), 6.80 (dd, *J* = 3.1, 1.6 Hz, 1 H), 6.07 (m, 1 H), 4.83 (m, 1 H), 4.10 (t, *J* = 9.0 Hz, 1 H), 3.85 (dd, *J* = 9.1, 6.8 Hz, 1 H), 3.70 (m, 2 H), 2.04 (s, 3 H); HRMS (EI) calcd for C₁₇H₁₆FN₃O₄ 345.1125, found 345.1129. Anal. Calcd for C₁₇H₁₆FN₃O₄·0.1H₂O: C, 58.82; H, 4.70; N, 12.10. Found: C, 58.71; H, 4.77; N, 12.01.

General Procedure for the Preparation of Oxyimino Analogues 26, 27, 40, 41, 88, 95, 96, and 117. The aldehyde or ketone (1 equiv), hydroxyl- or methoxyamine hydrochloride (1.5 equiv) and K₂CO₃ (1 equiv) were stirred in CH₃OH/CH₂Cl₂ (1:1). Water and EtOAc were added, the phases were separated, and the aqueous portion was extracted with EtOAc and CH₂Cl₂. The combined organics were dried (MgSO₄) and concentrated to give crude materials which were purified as indicated.

(S)-N-[[3-[3-Fluoro-4-(3-((hydroxyimino)methyl)-1H-pyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (26). The residue was purified on a 2.3 × 28 cm medium-pressure silica column with 5% MeOH/CH₂Cl₂ to give 0.116 g (72%) of **26** as a mixture of *E* and *Z* isomers: mp 216–218 °C; [α]_D²³ = –23° (c 0.47, DMSO); ¹H NMR (400 MHz, DMSO) δ 11.08 (s, 1 H), 10.56 (s, 1 H), 8.24 (m, 1 H), 8.01 (s, 1 H), 7.75 (m, 1 H), 7.70 (m, 1 H), 7.62 (m, 1 H), 7.40 (m, 1 H), 7.30 (m, 1 H), 7.14 (bs, 1 H), 6.63 (m, 1 H), 6.49 (m, 1 H), 4.76 (m, 1 H), 4.16 (t, *J* = 9.1 Hz, 1 H), 3.78 (dd, *J* = 9.0, 6.6 Hz, 1 H), 3.42 (t, *J* = 5.5 Hz, 2 H), 1.83 (s, 3 H); MS (EI) *m/z* 360 (M + H); HRMS (EI) calcd for C₁₇H₁₇FN₄O₄ 360.1234, found 360.1237. Anal. Calcd for C₁₇H₁₇FN₄O₄: C, 56.66; H, 4.75; N, 15.55. Found: C, 56.26; H, 4.85; N, 15.17.

(S)-[[3-[3-Fluoro-4-(3-cyanopyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (28). An acetonitrile (15 mL) solution of **26** (0.068 g, 0.19 mmol), triphenylphosphine (0.198 g, 0.75 mmol), and CHCl₄ (0.04 mL, 0.41 mmol) was stirred overnight. More reaction tetrachloride (2.5 mL, 25.9 mmol) was added, the reaction mixture was stirred for 4 h

and partitioned between water and CH₂Cl₂. The combined organic layer was washed (H₂O), dried (MgSO₄), and concentrated in vacuo to give 0.254 g of crude material. This material was purified on a 2.5 × 23 cm medium-pressure silica column with 5% MeOH/CH₂Cl₂ to give 0.047 g (72%) of **28**: mp 167–168 °C; [α]_D²⁵ = –26° (c 0.49, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 7.68 (dd, *J* = 13.2 Hz, *J'* = 2.4 Hz, 1 H), 7.41 (d, *J* = 1.6 Hz, 1 H), 7.35 (t, *J* = 8.4 Hz, 1 H), 7.28 (dd, *J* = 9.6 Hz, *J'* = 2.4 Hz, 1 H), 6.92 (t, *J* = 3.2 Hz, 1 H), 6.58 (dd, *J* = 2.8 Hz, *J'* = 1.6 Hz, 1 H), 6.40 (t, *J* = 6.0 Hz, 1 H), 4.84 (m, 8 lines, 1 H), 4.09 (t, *J* = 8.8 Hz, 1 H), 3.87 (dd, *J* = 8.8 Hz, *J'* = 6.8 Hz, 1 H), 3.69 (dd, *J* = 6.0 Hz, *J'* = 4.8 Hz, 2 H), 2.03 (s, 3 H); HRMS calcd for C₁₇H₁₅FN₄O₃ 342.1128, found 342.1140. Anal. Calcd for C₁₇H₁₅FN₄O₃: C, 59.65; H, 4.42; N, 16.37. Found: C, 59.58; H, 4.57; N, 16.07.

N-[[[(5S)-3-{3-Fluoro-4-[3-(hydroxymethyl)-1H-pyrrol-1-yl]phenyl]-2-oxo-1,3-oxazolan-5-yl]methyl]acetamide (29). A solution of aldehyde **25** (125 mg, 0.36 mmol) in 2:1 CH₃OH/CH₂Cl₂ (6 mL) at 0 °C was treated with sodium borohydride (7 mg, 0.18 mmol) followed by warming to ambient temperature for 4 h. The solution was diluted with CH₂Cl₂, treated with saturated ammonium chloride solution, and washed. Drying (Na₂SO₄) and concentration in vacuo afforded a white solid which was crystallized from boiling EtOAc/hexane (with a few drops of methanol) to afford **29** as a white solid 106 mg (84%): mp 155–157 °C.

(S)-N-[[3-[3-Fluoro-4-(3-carbomethoxy-1H-pyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (30). A solution of **25** (220 mg, 0.64 mmol) in 1:1 acetonitrile–CH₃OH (10 mL) was treated with NaCN (164 mg, 0.64 mmol), activated manganese dioxide (1.15 g, 13.2 mmol), and HOAc (58 μL, 1.0 mmol). The mixture was stirred at ambient temperature for 36 h, at which point additional activated manganese dioxide (550 mg, 6.32 mmol) and HOAc (58 μL, 1.0 mmol) were added. The solution was stirred for an additional 24 h, and was filtered through Celite. The filtrate was concentrated in vacuo, diluted with EtOAc and washed with water. The organic layer was dried (Na₂SO₄) and concentrated in vacuo to afford a pink solid. This material was subjected to radial chromatography on a 2 mm plate, eluting with a 0–2% CH₃OH/CH₂Cl₂ gradient to afford 170 mg (71%) of **30** as a white solid: mp 134–135 °C.

General Procedure for the Amonolysis of Esters 30, 47, 89, and 100 To Give Amides 31, 48, 49, 92, and 93. A solution of the ester (1 equiv) and KCN (0.25 equiv) in saturated methanolic ammonia or 2 N MeNH₂/CH₃OH was heated in a sealed tube at 50 °C. The solvent was evaporated and the residue was purified as indicated.

(S)-N-[[3-[3-Fluoro-4-(3-amidopyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (31). The residue was purified on a 2.5 × 25 cm medium-pressure silica column with 5–10% MeOH/CH₂Cl₂ to give 0.029 g (20%) of **31** as a light yellow solid: mp 119–121 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.61 (dd, *J* = 13.2 Hz, *J'* = 2.4 Hz, 1 H), 7.51 (s, 1 H), 7.35 (t, *J* = 8.4 Hz, 1 H), 7.22 (d, *J* = 9.2 Hz, 1 H), 6.90 (m, 1 H), 6.57 (m, 1 H), 4.75 (m, 1 H), 4.04 (t, *J* = 9.2 Hz, 1 H), 3.77 (dd, *J* = 9.2 Hz, *J'* = 6.8 Hz, 1 H), 3.58 (m, 1 H), 3.56 (m, 1 H), 3.01 (s, 3 H), 1.96 (s, 3 H); HRMS calcd for C₁₇H₁₇FN₄O₄ + H 361.1312, found 361.1314. Anal. Calcd for C₁₇H₁₇FN₄O₄·1.6H₂O: C, 52.47; H, 5.23; N, 14.39. Found: C, 52.78; H, 5.22; N, 13.69.

(S)-N-[[3-[3-Fluoro-4-(3-(2-carbomethoxyvinyl)-1H-pyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (32). A solution of diisopropyl ethoxycarbonylmethylphosphonate (805 mg, 3.19 mmol) in THF (7 mL) at 0 °C was treated with potassium *tert*-butoxide (3.19 mL 1.0 M/THF, 3.19 mmol) and warmed to ambient temperature for 1 h. The solution was cooled to –78 °C and treated via cannula with a solution of **25** (500 mg, 1.45 mmol) in THF (4 mL). The reaction was stirred at –78 °C for 30 min, followed by warming to ambient temperature for 2 h. The solution was quenched by addition of 1 mL saturated ammonium chloride solution, diluted with EtOAc and washed with water. Drying (Na₂SO₄) and concentration in vacuo afforded a yellow solid, which was chromatographed

graphed over 100 g of silica gel with 1–2% CH₃OH/CH₂Cl₂ to yield 497 mg (83%) of **32** as a light yellow solid: mp 161–163 °C.

(S)-N-[[3-[3-Fluoro-4-[3-(2-carbethoxyethyl)-1H-pyrrol-1-yl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (33). A solution of **32** (150 mg, 0.36 mmol) and copper(I) chloride (54 mg, 0.54 mmol) in 1:1 CH₃OH–THF (15 mL) at 0 °C was treated with NaBH₄ (136 mg, 3.6 mmol). The solution was stirred at 0 °C for 30 min and then warmed to ambient temperature for 1 h. The solution was treated with 2 mL saturated ammonium chloride solution and filtered through Celite. The filtrate was concentrated in vacuo and the residue was diluted with EtOAc and washed with water. Drying (Na₂SO₄) and concentration in vacuo afforded **33** 150 mg (99%) as a white solid.

(S)-N-[[3-[3-Fluoro-4-[3-(3-hydroxypropyl)-1H-pyrrol-1-yl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (34). A solution of the ester **33** (275 mg, 0.66 mmol) in THF (15 mL) was treated with lithium borohydride (287 mg, 13.2 mmol) and stirred at ambient temperature for 18 h. The solution was treated with 1 mL of saturated ammonium chloride solution, diluted with EtOAc, and washed with water. Drying (Na₂SO₄) and concentration in vacuo afforded an oil, which was chromatographed over 20 g silica gel eluting with 2–4% CH₃OH/CH₂Cl₂ to yield 96 mg (39%) of **34** as a white solid.

(S)-N-[[3-[3-Fluoro-4-[3-(3-methanesulfonylamino)propyl]-1H-pyrrol-1-yl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (35). A solution of the alcohol **34** (100 mg, 0.266 mmol) and NEt₃ (0.12 mL, 0.87 mmol) CH₂Cl₂ (2 mL) at 0 °C was treated with methanesulfonyl chloride (58 mL, 0.73 mmol) for 30 min. The solution was diluted with CH₂Cl₂ and washed with water and saturated sodium bicarbonate solution. Drying (Na₂SO₄) and concentration in vacuo afforded the mesylate as a solid. This was dissolved in DMF (2 mL) and added via cannula to a 0 °C solution of methanesulfonamide (38 mg, 0.40 mmol) in DMF (2 mL) that had been previously treated with NaH (17 mg of 60% in oil, 0.43 mmol). The reaction was then warmed at 60 °C for 18 h and cooled and the DMF removed in vacuo. The residue was dissolved in EtOAc and washed with water. Drying (Na₂SO₄) and concentration in vacuo afforded a solid which was subjected to radial chromatography on a 2 mm plate, eluting with 1–4% CH₃OH/CH₂Cl₂ to yield 31 mg (26%) of **35** as a white solid.

(S)-[[3-[3-Fluoro-4-(3-acetylpyrrol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (38). The same procedures were used as described for the preparation of **3–8** above. Starting with **37** (11.612 g, 42.38 mmol) and making noncritical variations **38**, 1.12 g (19%), was isolated as a pink solid: mp 145–148 °C.

(S)-[[3-[3-Fluoro-4-[3-(1-hydroxyethyl)pyrrol-1-yl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (39). Sodium borohydride (0.046 g, 1.21 mmol) was added to a CH₂Cl₂ (2.5 mL) and CH₃OH (2.5 mL) solution of **38** (0.101 g, 0.28 mmol). After stirring overnight, the reaction was quenched with 1 mL saturated ammonium chloride and the mixture was partitioned between water and CH₂Cl₂. The aqueous portion was extracted with CH₂Cl₂. The combined organic portions were dried (MgSO₄) and concentrated in vacuo to give of crude material that was purified on a preparative TLC plate (1000 μm, eluted with 5% MeOH/CH₂Cl₂ 2×) to give 0.073 g (72%) of the **39** as a white solid: mp 169–171 °C.

3-Fluoro-4-hydrazinonitrobenzene (42). K₂CO₃ (35 g, 251 mmol) and 3,4-difluoronitrobenzene (20 g, 126 mmol) were stirred in CH₃CN (200 mL) overnight. The solvent was removed in vacuo and the residue was partitioned between EtOAc/H₂O. The organic layer was washed with brine and dried (Na₂SO₄). Removal of solvent in vacuo gave pure **42** as a yellow solid: 20.2 g (94%); mp 144–145 °C; ¹H NMR (400 MHz, CDCl₃) δ 3.75 (br s, 2 H), 5.97 (br s, 1 H), 7.25 (t, *J* = 8.7 Hz), 7.89 (dd, *J* = 2.3, 9.4 Hz, 1 H), 8.05 (d, *J* = 9.1 Hz, 1 H); MS (+ESI) *m/z* 172 (*M* + H). Anal. Calcd for C₆H₆FN₃O₂: C, 42.11; H, 3.53; N, 24.55. Found: C, 42.49; H, 3.52; N, 24.13.

3-Fluoro-1-nitro-4-(1H-4-carbethoxypyrazol-1-yl)benzene (44). The dialdehyde **43**²⁸ (6.5 g, 45 mmol), NaOAc (3.7

g, 45 mmol) and **42** (8.5 g, 49.6 mmol) were placed in absolute EtOH and stirred at room temperature for 30 min. The reaction was then heated to reflux until complete by TLC, cooled and the solvent removed in vacuo. The residue was partitioned between EtOAc/H₂O. The organic layer was washed with brine and dried (Na₂SO₄). Removal of solvent gave **44** as a brown solid which was crystallized from EtOAc/hexane as orange needles 11.5 g (65%); mp 127–127 °C.

(S)-[3-[3-Fluoro-4-(1H-4-carbethoxypyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methanol (45). The nitro compound **44** (10 g, 35.8 mmol) was hydrogenated in MeOH with 10% Pd/C (1 g) at 20 psi on a Parr shaker for 3 h. The reaction was filtered through Celite and the solvent removed in vacuo to give pure aniline as an orangish solid 8.63 g (97%). This aniline (8.36 g, 33.5 mmol) and NaHCO₃ (5.6 g, 67 mmol) were placed in dry THF (200 mL) and cooled to 0 °C. Benzyl chloroformate (5.3 mL, 36.9 mmol) was added and the reaction was allowed to warm to room temperature. It was partitioned between EtOAc/H₂O and the organic layer was washed with brine and dried (Na₂SO₄). Removal of solvent gave pure Cbz-aniline as an orange solid. 12.7 g (98%). The Cbz-aniline (1.23 g, 3.2 mmol) was placed in dry THF at 78 °C under N₂. BuLi was added dropwise and the reaction was stirred 45 min. The *R*-(–)-glycidyl butyrate (0.48 mL, 3.4 mmol) was added dropwise and stirring was continued at –78 °C for 1 h then at room temperature overnight. The mixture was quenched with saturated NaHCO₃ and partitioned between H₂O and EtOAc. The organic layer was washed with brine and dried (Na₂SO₄). Removal of solvent gave an orange residue which was passed through a 4 × 5 cm silica gel plug with 50%EtOAc/hexane to 100% EtOAc gradient. Product **45** was isolated as a white solid: 0.885 g (79%); mp 133–134 °C; [α]_D –39° (*c* 0.93, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 1.38 (t, *J* = 7.1 Hz, 3 H), 2.35–2.45 (br m, 1 H), 3.77–3.83 (m, 1 H), 4.02–4.10 (m, 3 H), 4.34 (q, *J* = 7.1 Hz, 2 H), 4.78–4.82 (m, 1 H), 7.28 (d, *J* = 10.7 Hz, 1 H), 7.77 (dd, *J* = 2.4, 11.4 Hz, 1 H), 7.88 (t, *J* = 8.8 Hz, 1 H), 8.11 (s, 1 H), 8.43 (d, *J* = 2.4 Hz, 1 H); HRMS calcd for C₁₆H₁₆FN₃O₅ + H 350.1152, found 350.1153. Anal. Calcd for C₁₆H₁₆FN₃O₅: C, 55.01; H, 4.62; N, 12.03. Found: C, 55.68; H, 4.74; N, 11.66.

(S)-[3-[3-Fluoro-4-(1H-4-carbethoxypyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl Azide (46). The alcohol **45** (0.76 g, 2.2 mmol) and NEt₃ (0.46 mL, 3.3 mmol) were placed in dry CH₂Cl₂ at 0 °C under N₂. Methanesulfonyl (0.185 mL, 2.4 mmol) was added dropwise. The reaction was allowed to stir at room temperature for 4 h at which time it was partitioned between saturated NaHCO₃ and CH₂Cl₂. The organic layer was washed with brine and dried (Na₂SO₄). Removal of solvent gave the mesylate as an off white solid which was pure by TLC and NMR: 0.925 g (98%). The mesylate (0.74 g, 1.73 mmol) and NaN₃ (0.84 g, 13 mmol) were placed in dry DMF and heated to 65 °C overnight. The reaction was cooled, diluted with EtOAc, washed with water (2×) and brine, and dried (Na₂SO₄). Removal of solvent gave **46** as an off white solid: 0.6 g (93%); mp 128–130 °C; [α]_D –105° (*c* 0.96, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 1.38 (t, *J* = 7.1 Hz, 3 H), 3.63 (dd, *J* = 4.2, 13.3 Hz, 1 H), 3.76 (dd, *J* = 4.4, 13.2 Hz, 1 H), 3.91 (dd, *J* = 6.2, 8.9 Hz, 1 H), 4.13 (t, *J* = 8.9 Hz, 1 H), 4.34 (q, *J* = 7.1 Hz, 2 H), 4.82–4.87 (m, 1 H), 7.28 (d, *J* = 7.3 Hz, 1 H), 7.79 (dd, *J* = 2.4, 13.7 Hz, 1 H), 7.91 (t, *J* = 8.8 Hz, 1 H), 8.12 (s, 1 H), 8.45 (d, *J* = 2.4 Hz, 1 H); HRMS calcd for C₁₆H₁₅FN₆O₄ 374.1139, found 374.1143.

(S)-N-[[3-[3-Fluoro-4-(1H-4-carbethoxypyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (47). The azide **46** (0.59 g, 1.58 mmol) was placed in dry THF and a solution of triphenylphosphine (0.5 g, 1.89 mmol) in THF was added dropwise. The reaction was stirred at room temperature for 1.5 h at which time water (3 mL) was added and the temperature raised to 65 °C overnight. The reaction was cooled and the solvent removed in vacuo. The residue was partitioned between CH₂Cl₂ and water. The aqueous was extracted with CH₂Cl₂ and the combined extracts were dried (Na₂SO₄). Removal of solvent in vacuo gave a residue which was chromatographed on a 2 × 30 cm column with 5–10% MeOH/

CHCl₃ gradient. The intermediate amine was isolated as a white solid: 0.43 g (77%). The amine (0.35 g, 1 mmol) and pyridine (0.243 mL, 3 mmol) were placed in CH₂Cl₂. Acetic anhydride (0.132 mL, 1.4 mmol) was added and the reaction was stirred at room temperature until complete as determined by TLC. The reaction was then diluted with CH₂Cl₂, washed with 1 N HCl (2×), saturated NaHCO₃, and brine, and dried (Na₂SO₄). Removal of solvent gave a residue which was chromatographed on a 2 × 35 cm column with 5% MeOH/CH₂Cl₂. Product **47** was isolated as a white solid: 0.39 g (99%); mp 160–161 °C; [α]_D²⁰ –23° (c 0.47, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 1.39 (t, *J* = 7.1 Hz, 3 H), 2.04 (s, 3 H), 3.61–3.69 (m, 1 H), 3.70–3.79 (m, 1 H), 3.84 (dd, *J* = 6.8, 15.8 Hz, 1 H), 4.10 (t, *J* = 9.0 Hz, 1 H), 4.35 (q, *J* = 7.1 Hz, 2 H), 4.81–4.90 (m, 1 H), 5.95 (br t, 1 H), 7.28 (d, *J* = 8.8 Hz, 1 H), 7.75 (dd, *J* = 2.4, 13.7 Hz, 1 H), 7.91 (t, *J* = 8.9 Hz, 1 H), 8.12 (s, 1 H), 8.45 (d, *J* = 2.4 Hz, 1 H); HRMS calcd for C₁₈H₁₉FN₄O₅ + H 391.1418, found 391.1421. Anal. Calcd for C₁₈H₁₉FN₄O₅·0.25H₂O: C, 54.75; H, 4.98; N, 14.19. Found: C, 54.67; H, 4.98; N, 14.11.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-4-cyanopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**50**). The carboxamide **48** (0.23 g, 0.64 mmol) was placed in dry DMF (7.5 mL) and cooled to 0 °C under N₂. SOCl₂ (0.07 mL, 0.95 mmol) was added dropwise and the reaction was stirred at room temperature for 30 min. More SOCl₂ (0.07 mL, 0.95 mmol) was added and stirring was continued another 30 min. Saturated NaHCO₃ was added and the reaction was extracted with CH₂Cl₂. The extracts were dried (Na₂SO₄) and the solvent removed. The residue was chromatographed on a 2 × 30 cm column with 5% MeOH/CH₂Cl₂. Product **50** was isolated as a white solid: 155 mg (71%); mp 194–196 °C; [α]_D²⁰ –27° (c 0.57, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 2.04 (s, 3 H), 3.65–3.76 (m, 2 H), 3.86 (dd, *J* = 6.9, 8.9 Hz, 1 H), 4.10 (t, *J* = 8.9 Hz, 1 H), 4.81–4.90 (m, 1 H), 6.07 (br t, 1 H), 7.29 (d, *J* = 7.6 Hz, 1 H), 7.78 (dd, *J* = 1.9, 13.8 Hz, 1 H), 7.89 (t, *J* = 8.8 Hz, 1 H), 8.00 (s, 1 H), 8.38 (s, 1 H); HRMS calcd for C₁₆H₁₄FN₅O₃ + H 344.1159, found 344.1153. Anal. Calcd for C₁₆H₁₄FN₅O₃·0.5H₂O: C, 54.55; H, 4.29; N, 19.87. Found: C, 54.66; H, 4.24; N, 19.46.

3-Fluoro-1-(phenylmethoxycarbonylamino)-4-(1*H*-4-iodopyrazol-1-yl)benzene (52). 4-Iodopyrazole (10 g, 52 mmol), 3,4-difluoronitrobenzene (8.3 g, 52 mmol) and K₂CO₃ (7.2 g, 52 mmol) were placed in dry DMSO (100 mL) and heated to 90 °C under N₂ for 24 h. The reaction was cooled and partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc and the combined extracts were dried (Na₂SO₄). Removal of solvent gave a residue which was taken up in Et₂O, washed with H₂O (3×) and dried (Na₂SO₄). Removal of solvent again gave pure pyrazolynitrobenzene intermediate as a yellow solid, 16.5 g (95%). This pyrazolynitrobenzene (2.6 g, 7.8 mmol) and SnCl₂ (8.8 g, 39 mmol) were placed in absolute EtOH and heated to 75 °C for 1 h. The reaction was cooled and poured into ice. The mixture was made basic with saturated NaHCO₃ and filtered. The filtrate was extracted with EtOAc (3×). The combined extracts were washed with brine and dried (Na₂SO₄). Removal of solvent gave the aniline as a yellow oil 2.1 g (88%). This aniline (9.41 g, 31.05 mmol) and NaHCO₃ (5.2 g, 62 mmol) were placed in dry THF and benzyl chloroformate was added. The reaction was stirred at room temperature overnight, diluted with water and extracted with EtOAc (3×). The extracts were dried (Na₂SO₄) and the solvent was removed in vacuo to give **52** as a yellow solid: 13.72 g (99% from the aniline).

3-Fluoro-1-(phenylmethoxycarbonylamino)-4-(1*H*-4-trimethylsilylacetylenylpyrazol-1-yl)benzene (53). The Cbz intermediate **52** (2.0 g, 4.6 mmol), TMS-acetylene (0.78 mL, 5.5 mmol) and diethylamine (20 mL) were placed under N₂. Pd(PPh₃)₂Cl₂ (0.1 mmol, 70 mg) and CuI (0.005 g) were added. The reaction was stirred at room temperature under N₂ for 8 h. The reaction was evaporated to dryness and the residue was purified on a 2 × 40 cm column. Product **53** was isolated as a solid 1.6 g (85%); mp 133–136 °C.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-4-iodopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**54**). The Cbz-pro-

TECTED ANILINE **52** (3.17 g, 7.25 mmol) was reacted as above for the preparation of compound **47** making noncritical variations. The procedures afforded **54** as a white solid: 0.88 g (27%); mp 199–201 °C.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-4-*N*-(*tert*-butoxycarbonylamino)pyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**55**). The aryl iodide **54** (0.265 g, 0.6 mmol), PdOAc₂ (13.3 mg, 0.06 mmol), dppp (29.7 mg, 0.072 mmol), and Et₃N (0.1 mL, 0.72 mmol) were heated in 1:10 H₂O/DMF (10 mL) at 100 °C an atmosphere of CO with vigorous stirring for 2 days. The reaction was cooled, filtered and the solvent removed. The residue was triturated in hot acetone and the solid acid filtered off to give 0.2 g (92%); HRMS (FAB) calcd for C₁₆H₁₅FN₄O₅ 362.1104, found 362.1265. This acid (0.15 g, 0.4 mmol), DPPA (0.09 mL, 0.4 mmol), and NEt₃ (0.057 mL, 0.4 mmol) were refluxed in *t*-BuOH for 3 days. The solvent was removed and the residue was chromatographed on a 2 × 30 cm column with 5% MeOH/CH₂Cl₂ to give pure **55** as an ivory solid: 110 mg (58%); mp 177–179 °C.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-4-*N*-(acetylamino)pyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**56**). The Boc-amine **55** (20 mg, 0.05 mmol) was deprotected in 1:1 TFA/CH₂Cl₂ (4 mL) at room temperature under N₂ for 4 h. The solvent and excess TFA were removed in vacuo and the residue was triturated in Et₂O. The solvent was removed again and the solid product was placed in CH₂Cl₂ (5 mL) and NEt₃ (0.5 mL) was added followed by Ac₂O (0.5 mL). The reaction was stirred at room temperature for 2 h and evaporated to dryness in vacuo. The residue was taken up in CH₂Cl₂ and washed with saturated NaHCO₃. The organic layer was dried (Na₂SO₄) and the solvent removed. The product **56** was purified on a 2 × 25 cm column with 10% MeOH/CH₂Cl₂ and isolated as a white solid: 15 mg (80%); mp 229–230 °C dec.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-4-trimethylsilylacetylenylpyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**57**). The acetylene derivative **53** (1.2 g, 3 mmol) was reacted as described above for the preparation of compound **47** making noncritical variations. The procedures afforded **57** as a white solid: 0.35 g (28%); mp 197–199 °C.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-4-acetylenylpyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**58**). The TMS-acetylide **57** (0.31 g, 0.75 mmol) was treated with K₂CO₃ (0.1 g) in MeOH (20 mL) at room temperature for 1 h. The solvent was removed in vacuo and the residue was partitioned between CH₂Cl₂ and water. The organic layer was dried (Na₂SO₄) and the solvent removed to give pure product as a white solid: 0.25 g (98%); mp 180–182 °C.

3-Fluoro-1-nitro-4-(1*H*-3-*tert*-butoxycarbonylamino-pyrazol-1-yl)benzene (63). The aminopyrazole derivative **62** (3 g, 13.5 mmol), di-*tert*-butyl dicarbonate (2.95 g, 13.5 mmol), NEt₃ (1.88 mL, 13.5 mmol) and (dimethylamino)pyridine (20 mg) were placed in dry DMF and stirred under nitrogen overnight. The reaction was partitioned between water and Et₂O and the organic layer was washed with water, brine and dried (Na₂SO₄). Removal of solvent gave a residue that was purified on a 3 × 30 cm column to give **63**: 2.2 g (51%); ¹H NMR (400 MHz, CDCl₃) δ 1.55 (s, 9 H), 5.97 (d, *J* = 2.6 Hz, 1 H), 7.12 (br s, 1 H), 8.07–8.14 (m, 4 H); MS (+ESI) *m/z* 322 (M + H); MS (–ESI) *m/z* 321 (M – H).

(*S*)-[[3-[3-Fluoro-4-(1*H*-3-trifluoromethylpyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (**64**). The nitro compound **61** (6.5 g, 23.6 mmol) was reacted as described above for the preparation of compound **47** making noncritical variations. The procedures afforded **64** as a white solid: 1.32 g (14%); mp 180–181 °C; [α]_D²⁵ –23° (c 0.95, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 2.04 (s, 3 H), 3.65–3.78 (m, 2 H), 3.85 (dd, *J* = 6.9, 9.0 Hz, 1 H), 4.10 (t, *J* = 8.9 Hz, 1 H), 4.79–4.85 (m, 1 H), 6.09 (br t, 1 H), 6.73 (d, *J* = 2.4 Hz, 1 H), 7.24 (d, *J* = 10.4 Hz, 1 H), 7.77 (dd, *J* = 2.5, 13.7 Hz, 1 H), 7.90 (t, *J* = 8.8 Hz, 1 H), 8.00 (s, 1 H); MS (+ESI) *m/z* 387 (M + H). Anal. Calcd for C₁₆H₁₄F₄N₄O₃: C, 49.75; H, 3.65; N, 14.50. Found: C, 50.01; H, 3.70; N, 14.10.

(*S*)-*N*-[[3-[3-Fluoro-4-(1*H*-3-*tert*-butoxycarbonylamino-pyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]ac-

etamide (65). The Boc intermediate **63** (1.8 g, 5.6 mmol) was reacted as described above for the preparation of compound **47**. The main variation was that 2 equiv of *n*-BuLi were used in the oxazolidinone ring forming reaction. The procedures afforded **65** as a pale yellow solid: 300 mg (12%); mp 193–194 °C.

(S)-N-[[3-[3-Fluoro-4-(1H3-aminopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (66). The Boc-protected material **65** (0.25 g, 0.58 mmol) was dissolved in CH₂Cl₂ (7 mL) and cooled to 0 °C under nitrogen. Trifluoroacetic acid (5 mL) was added and the reaction was allowed to warm to room temperature and stirred 3 h. The solvent and excess TFA were removed in vacuo and the residue was partitioned between 1 M NaHCO₃ and CH₂Cl₂. The aqueous layer was extracted with 5% MeOH/CH₂Cl₂ and the extracts were combined and dried (Na₂SO₄). Removal of solvent gave pure **66** as a pale yellow solid: 0.16 g (83%).

(S)-N-[[3-[3-Fluoro-4-(1H3-benzyloxymethylacetylaminopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (67). The arylamine **66** (25 mg, 0.075 mmol) and NEt₃ (0.012 mL, 0.082 mmol) were placed in dry methylene chloride. Benzyloxyacetyl chloride (0.013 mL, 0.082 mmol) was added and the reaction was stirred under nitrogen for 4 h and then partitioned between saturated NaHCO₃ solution and CH₂Cl₂. The organic layer was dried (Na₂SO₄) and the solvent removed to give a residue which was chromatographed on a 2 × 30 cm column with 7% MeOH/CHCl₃. The product **67** was isolated as a tan solid: 35 mg (97%).

(S)-N-[[3-[3-Fluoro-4-(1H3-hydroxymethylacetylaminopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (68). The intermediate **67** (30 mg, 0.062 mmol) was hydrogenated in MeOH on a Parr shaker at 30 psi with 10% Pd/C for 6 h. The reaction was filtered through Celite and evaporated to dryness. The residue was purified on a 2 × 35 cm column with 10% MeOH/CHCl₃ to give the product as a tan solid: 22 mg (99%); mp 129–132 °C.

(S)-N-[[3-[3-Fluoro-4-(1H3-acetylaminopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (69). Intermediate **66** (26 mg, 0.078 mmol) and NEt₃ (0.012 mL, 0.085 mmol) were placed in dry CH₂Cl₂. Acetic anhydride (2 mL) was added and the reaction was stirred at room temperature for 2 h and evaporated to dryness. The residue was purified on a 2 × 30 cm column with 10% MeOH/CHCl₃ to give product as a white solid: 25 mg (78%); mp 212–214 °C.

(S)-N-[[3-[3-Fluoro-4-(1H3-methanesulfonylaminopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (70). Intermediate **66** (20 mg, 0.06 mmol) was placed in pyridine (3 mL) and cooled to 0 °C. Methanesulfonyl chloride (0.005 mL, 0.066 mmol) was added and the reaction was stirred at room temperature overnight. The reaction was diluted with CH₂Cl₂, washed with saturated NaHCO₃ and dried (Na₂SO₄). Removal of solvent in vacuo gave the product as a pale yellow foam: 18 mg (73%); mp 145–146 °C.

(S)-N-[[3-[3-Fluoro-4-(1H3-cyanopyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (71). The amino derivative **67** (36 mg, 0.108 mmol) was placed in 2 N HCl (3 mL) at 0 °C. A solution of NaNO₂ in water (1 mL) was added and the mixture was stirred 30 min at 0 °C before being neutralized with solid NaHCO₃. In a separate flask CuCN (12.6 mg, 0.14 mmol), KCN (10.5 mg, 0.162 mmol), water (2 mL) and EtOAc (3 mL) were cooled to 0 °C. The diazonium salt was added to this mixture with a pipet. The reaction was stirred at 0 °C for 30 min and then at room temperature for 2 h. It was partitioned between saturated NaHCO₃ and CHCl₃ and extracted with CHCl₃. The extracts were dried (Na₂SO₄) and evaporated to dryness. The residue obtained was purified on a 2 × 35 cm column with 5% MeOH/CHCl₃. Product was isolated as a yellow free flowing powder: 14 mg (38%); mp 190–192 °C dec; ¹H NMR (400 MHz, CDCl₃) δ 2.04 (s, 3 H), 3.62–3.73 (m, 2 H), 3.86 (dd, *J* = 6.7, 9.1 Hz, 1 H), 4.11 (t, *J* = 8.9 Hz, 1 H), 4.80–4.88 (m, 1 H), 5.98 (br t, 1 H), 6.88 (d, *J* = 2.6 Hz, 1 H), 7.27 (d, *J* = 6.1 Hz, 1 H), 7.80 (dd, *J* = 2.4, 13.8 Hz, 1 H), 7.89 (t, *J* = 8.8 Hz, 1 H), 8.04 (t, *J* = 2.5 Hz, 1 H); HRMS calcd for C₁₆H₁₄FN₅O₃ 343.1081, found 343.1078.

Anal. Calcd for C₁₆H₁₄FN₅O₃·1.25CH₃OH: C, 54.04; H, 4.99; N, 18.27. Found: C, 53.65; H, 4.35; N, 18.10.

N-[[[(5S)-3-(3-Fluoro-4-hydrazinophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (72). A solution of the aniline **20** (5.34 g, 20.0 mmol) in 8 mL of methanol, 20 mL of water plus 6 mL of 37% HCl at –10 °C was treated dropwise with a solution of NaNO₂ (1.40 g, 20.2 mmol) in 4 mL of water over a 2 min period. After stirring for 8 min at –10 °C, the contents were transferred to a dropping funnel and added over a 2 min period to a vigorously stirred solution of SnCl₂·2H₂O (10.80 g, 47.8 mmol) in 52 mL of 37% HCl at –10 °C. After stirring for an additional 40 min, the pH of the reaction mixture was adjusted to 7–8 by the dropwise addition of 10 N NaOH (80 mL) with cooling in an ice/acetone bath. The milky white aqueous suspension was concentrated and the gelatinous residue triturated (5 × 150 mL) with chloroform–methanol (9:1). The combined organic extracts were dried with anhydrous Na₂SO₄ and concentrated in vacuo to obtain 5.2 g of crude product. Recrystallization from a methanol, methylene chloride, diethyl ether mixture afforded 4.26 g (75.5%) of analytically pure **72** as a white solid: mp 173–175 °C; [α]_D²⁵ –26° (*c* 0.79, DMSO); ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.22 (t, *J* = 5.77 Hz, 1 H), 7.32 (dd, *J* = 2.34, 14.14 Hz, 1 H), 7.11 (q, *J* = 8.95 Hz, 1 H), 7.04 (dd, *J* = 2.27, 8.87 Hz, 1 H), 6.53 (s, 1 H), 4.64 (m, 1 H), 4.01 (m, 3 H), 3.63 (m, 1 H), 3.37 (t, *J* = 5.54 Hz, 2 H), 1.81 (s, 3 H); MS (EI) *m/z* 282 (M + H). Anal. Calcd for C₁₂H₁₅FN₄O₃: C, 51.06; H, 5.36; N, 19.85. Found: C, 51.07; H, 5.34; N, 19.76.

(S)-N-[[3-[3-Fluoro-4-(1H3-methylpyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (73). Acetaldehyde dimethyl acetal (0.105 mL, 0.71 mmol) and the arylhydrazine **72** (0.2 g, 0.71 mmol) were heated at reflux in ethanol for 2 h. The solvent was removed in vacuo and the residue was partitioned between CH₂Cl₂ and saturated NaHCO₃. The organic layer was dried (Na₂SO₄) and the solvent removed in vacuo. The residue was purified on a 2 × 35 cm column with 5% MeOH/CH₂Cl₂ to give the desired product **73** (high *R*_f) as a solid 110 mg (47%); mp 129–131 °C; [α]_D²⁵ = –29° (*c* 0.74, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 2.01 (s, 3 H), 2.35 (s, 3 H), 3.66 (dd, *J* = 4.6, 6.0 Hz, 2 H), 3.81 (dd, *J* = 6.7, 9.1 Hz, 1 H), 4.06 (t, *J* = 9.0 Hz, 1 H), 4.76–4.83 (m, 1 H), 6.23 (d, *J* = 2.4 Hz, 1 H), 6.71 (br t, *J* = 6.0 Hz, 1 H), 7.17 (dd, *J* = 1.7, 8.9 Hz, 1 H), 7.63 (dd, *J* = 2.4, 13.9 Hz, 1 H), 7.82 (d, *J* = 1.7 Hz, 1 H); HRMS (FAB) calcd for C₁₆H₁₇FN₄O₃ + H⁺ 333.1363, found 333.1349. Anal. Calcd for C₁₆H₁₇FN₄O₃·0.25H₂O: C, 57.05; H, 5.24; N, 16.63. Found: C, 57.05; H, 5.10; N, 16.44.

(S)-N-[[3-[3-Fluoro-4-(1H5-methylpyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (74). In addition to the above material a lower *R*_f product was isolated which was found to be the 5-regioisomer by 2D NMR: 60 mg (25%); mp 139–141 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.01 (s, 3 H), 2.20 (s, 3 H), 3.59–3.71 (m, 2 H), 3.84 (dd, *J* = 6.8, 9.1 Hz, 1 H), 4.08 (t, *J* = 9.0 Hz, 1 H), 4.76–4.83 (m, 1 H), 6.19 (s, 1 H), 6.63 (br t, *J* = 6.0 Hz, 1 H), 7.27 (dd, *J* = 1.8, 8.7 Hz, 1 H), 7.44 (t, *J* = 8.5 Hz, 1 H), 7.60 (d, *J* = 1.5 Hz, 1 H), 7.67 (dd, *J* = 2.4, 12.2 Hz, 1 H); HRMS (FAB) calcd for C₁₆H₁₇FN₄O₃ + H⁺ 333.1363, found 333.1349. Anal. Calcd for C₁₆H₁₇FN₄O₃·0.75H₂O: C, 55.57; H, 5.39; N, 16.20. Found: C, 55.71; H, 5.10; N, 16.00.

N-[[[(5S)-3-(4-{2-[(E)-3-ethoxy-2-propenoyl]hydrazino}-3-fluorophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (76). The arylhydrazine **72** (1.3 g, 4.6 mmol), the acryloyl chloride **75**³⁴ (0.42 g, 3 mmol) and triethylamine (0.42 mL, 3 mmol) were heated to reflux in dry THF for 5 h. The reaction was cooled and diluted with a large volume of Et₂O. The precipitated product was isolated as an orange solid and used without further purification: ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.0 Hz, 3 H), 1.82 (s, 3 H), 3.38 (t, 2 H), 3.65–3.70 (m, 1 H), 3.91 (q, *J* = 7.0 Hz, 2 H), 4.03 (t, 1 H), 4.62–4.70 (m, 1 H), 5.41 (d, 1 H), 6.72 (t, 1 H), 7.05 (d, 1 H), 7.39 (d, 1 H), 7.42 (br s, 1 H), 7.53 (br s, 1 H), 8.21 (br t, 1 H); MS (+ESI) *m/z* 381 (M + H); MS (–ESI) *m/z* 379 (M – H).

(S)-[[3-[3-Fluoro-4-(1H3-hydroxypyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (77). The acylhy-

drazine **76** was stirred in concentrated HCl for 4 h at room temperature. An equal volume of water was added and the reaction was evaporated to dryness under Hi-vac. The residue was chromatographed on a 2 × 30 cm column with 5%MeOH/CH₂Cl₂. The product **77** was isolated as a yellow solid: 0.65 g (65% for 2 steps); mp 234–235 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.83 (s, 3 H), 3.42 (t, *J* = 5.5 Hz, 2 H), 3.76 (dd, *J* = 6.5, 9.1 Hz, 1 H), 4.12 (t, *J* = 9.0 Hz, 1 H), 4.70–4.78 (m, 1 H), 5.82 (d, *J* = 2.5 Hz, 1 H), 7.39 (d, *J* = 9.0 Hz, 1 H), 7.65 (dd, *J* = 2.4, 13.2 Hz, 1 H), 7.70 (t, *J* = 9.1 Hz, 1 H), 7.86 (t, *J* = 2.6 Hz, 1 H), 8.23 (br t, 1 H); MS (+ESI) *m/z* 335 (M + H); MS (–ESI) *m/z* 333 (M – H). Anal. Calcd for C₁₅H₁₅FN₄O₄·1/3H₂O: C, 53.20; H, 4.64; N, 16.46. Found: C, 53.20; H, 4.73; N, 16.02.

(S)-[3-[3-Fluoro-4-(1*H*-3-acetoxypyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methylacetamide (**78**). The alcohol **77** (20 mg, 0.06 mmol) was placed in acetic anhydride (3 mL) and pyridine (0.03 mL) was added. The reaction was stirred at room temperature for 3 h and then evaporated to dryness. The residue was purified on a 2 × 30 cm column with 3% MeOH/CH₂Cl₂ to give the product as an oil. This was crystallized from CH₂Cl₂/hexane to give 20 mg (89%); mp 125–128 °C.

(S)-[3-[3-Fluoro-4-(1*H*-3-methoxypyrazol-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methylacetamide (**79**). The alcohol **77** (0.11 g, 0.33 mmol), methyl iodide (3.3 mmol) and K₂CO₃ were heated to reflux in acetone for 3 h. The reaction was cooled, filtered and evaporated to dryness. The residue was purified on a 2 × 30 cm column with 5% MeOH/CH₂Cl₂ to give the product as a gum which was solidified in CH₂Cl₂/hexane to yield a tan solid: 85 mg (74%); mp 132–133 °C.

4-([*tert*-Butyldimethylsilyloxy]methyl)-1-(2-fluoro-4-nitrophenyl)-1*H*-imidazole (**82**). Compound **81** (5.8 g, 24.5 mmol) and imidazole (3.32 g, 48.8 mmol) were dissolved in dry DMF (40 mL) and cooled to 0 °C. TBDMSCl (5.5 g, 36.5 mmol) was added and the reaction was stirred at room temperature overnight. The solvent was removed in vacuo and the residue was chromatographed with a 30–50% EtOAc/hexane gradient. Product was isolated as a yellow solid: 5.53 g (64%); ¹H NMR (400 MHz, CDCl₃) δ 0.15 (s, 6 H), 0.94 (s, 9 H), 4.78 (s, 2 H), 7.26 (d, *J* = 3 Hz, 1 H), 7.34 (t, *J* = 8 Hz, 1 H), 7.96 (s, 1 H), 8.17 (s, 1 H), 8.20 (s, 1 H); MS (ESI+) *m/z* 352 (M + H); MS (ESI–) 350 (M – H).

N-[(5*S*)-3-[4-[4-([*tert*-Butyldimethylsilyloxy]methyl)-1*H*-imidazol-1-yl]-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**83**). Compound **82** (5.53 g, 15.6 mmol) was reacted as described above for the preparation of compounds **3–8** making noncritical variations. The procedures afforded **83** as an amber solid: 1.67 g (23%).

N-[(5*S*)-3-[4-[4-(Hydroxymethylmethyl)-1*H*-imidazol-1-yl]-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**84**). Compound **83** (1.67 g, 3.6 mmol) was dissolved in 3:1:1 AcOH/THF/H₂O (50 mL) and stirred at room temperature for 16 h. The solvent was removed in vacuo and the residue was dissolved in CH₂Cl₂ and diluted with Et₂O. The resulting precipitate was collected as a white solid and used as is: 1.27 g.

N-[(5*S*)-3-[4-(4-Cyano-1*H*-imidazol-1-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**85**). A slurry of **84** (1.27 g, 3.65 mmol) in CH₂Cl₂ (10 mL) was treated with DBU (1.33 mL, 8.9 mmol) followed by diphenyl phosphorazidate (1.8 mL, 8.4 mmol). The mixture was stirred at room temperature for 2 days. The solvent was removed in vacuo and the residue was purified on silica gel with 0.5–2%MeOH/CH₂Cl₂ gradient. The intermediate azide was isolated as an amber syrup: 1.17 g (86%). The azide (1.17 g, 3.14 mmol) was suspended in EtOH (80 mL). 2,5-dimethyl-3-hexyne-2,5-diol (0.58 g, 4.12 mmol) and 10% Pd/C (0.58 g) were added. The mixture was heated to reflux for 3 h. The reaction was filtered and the solvent removed in vacuo. The residue was purified on silica gel with 0.5–2%MeOH/CH₂Cl₂ gradient to give **85** as a white solid: 308 mg (28%); mp 204–205 °C; [α]_D²⁵ = –25° (c 0.79, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 2.04 (s, 3 H), 3.65–3.78 (m, 2 H), 3.88 (dd, *J* = 7, 9 Hz, 1 H), 4.10 (t, *J* = 9

Hz, 1 H), 4.84 (br m, 1 H), 6.01 (br t, 1 H), 7.34–7.43 (m, 2 H), 7.75–7.80 (m, 3 H); MS (ESI+) *m/z* 344 (M + H); MS (ESI–) *m/z* 342 (M – H). Anal. Calcd for C₁₆H₁₄FN₅O₃: C, 55.98; H, 4.11; N, 20.40. Found: C, 55.86; H, 3.99; N, 20.31.

N-[(5*S*)-3-(4-Azido-3-fluorophenyl)-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**86**). The aniline **20** (2 g, 7.5 mmol) was dissolved in concentrated HCl (10 mL) and water (10 mL) and cooled to 0 °C. Sodium nitrite was added and the yellow solution was stirred at 0 °C for 2 h. A solution of NaN₃ (0.97 g, 15 mmol) and NaOAc (12.3 g, 150 mmol) was added dropwise. The mixture was extracted with EtOAc and the combined extracts were washed with brine and dried (Na₂SO₄). Removal of solvent gave product as a tan solid, 2.1 g (96%), that was used without further purification.

General Procedure for the Preparation of 1,2,3-triazoles **87** and **89–91** from Azide **86** and Alkynes. The azide **86** (1 equiv) was suspended in benzene and the alkyne (3 equiv) was added and the mixture was refluxed. The reaction was cooled and the precipitated products were isolated.

N-[(5*S*)-3-[4-(4-Acetyl-1*H*-1,2,3-triazol-1-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**87**): yield 0.41 g (66%); mp 183–186 °C; [α]_D²⁵ = –29° (c 0.73, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 2.05 (s, 3 H), 2.77 (s, 3 H), 3.67–3.80 (m, 2 H), 3.88 (dd, *J* = 9, 7 Hz, 1 H), 4.12 (t, *J* = 9 Hz, 1 H), 4.81–4.90 (m, 1 H), 6.05 (br t, 1 H), 7.35 (d, *J* = 11 Hz, 1 H), 7.82 (dd, *J* = 2, 13 Hz, 1 H), 7.97 (t, *J* = 9 Hz, 1 H), 8.55 (d, *J* = 2 Hz, 1 H); MS (EI) *m/z* 361 (M); HRMS (EI) calcd for C₁₆H₁₆FN₅O₄ 361.1186, found 361.1183. Anal. Calcd for C₁₆H₁₆FN₅O₄: C, 53.19; H, 4.46; N, 19.38. Found: C, 53.03; H, 4.54; N, 19.13.

N-[(5*S*)-3-[3-Fluoro-4-[4-(hydroxymethyl)-1*H*-1,2,3-triazol-1-yl]phenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**94**). The aldehyde **91** (100 mg, 0.29 mmol) was suspended in MeOH, NaBH₄ (11 mg, 0.29 mmol) was added and the mixture became homogeneous. The reaction was stirred at room temperature overnight, and a few drops of water was added and the solvent removed in vacuo. The residue was purified on silica gel with 10% MeOH/CH₂Cl₂. Product was isolated as a white solid: mp 186–188 °C.

N-[(5*S*)-3-[4-[2-(1-Cyano-2-nitriloethylidene)hydrazino]-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**97**). A mixture of concentrated HCl (100 mL) and water (100 mL) was stirred at 25 °C and aniline **20** (20.0 g, 74.9 mmol) was added. The solution was cooled in ice and sodium nitrite (5.5 g, 79.7 mmol) added over 5 min. After 1 h of vigorous mechanical stirring, a solution of sodium acetate (12.5 g, 1.52 mole) in water (200 mL) was added. The mixture became very thick and a solution of malononitrile (5.0 g, 75.8 mmol) in ethanol (50 mL) was added. The reaction was allowed to come to room temperature then filtered. The precipitate was washed with NaHCO₃ solution (150 mL) and water (200 mL) and dried. The filter cake was ground up and dried in vacuo, 25 °C for 3 days, to afford a bright yellow solid (22.7 g, 88%); mp 235 °C dec; ¹H NMR δ (DMSO-*d*₆) δ 1.81 (s, 3 H), 3.40 (t, *J* = 5.5 Hz, 2 H), 3.73 (dd, *J* = 6.5, 9.1 Hz, 1 H), 4.11 (t, *J* = 9.0 Hz, 1 H), 4.66–4.78 (m, 1 H), 7.34 (d, *J* = 9.0 Hz, 1 H), 7.50 (t, *J* = 9.0 Hz, 1 H), 7.60 (dd, *J* = 2.3, 13.8 Hz, 1 H), 8.22 (t, *J* = 5.8 Hz, NH); MS (ESI+) *m/z* 345 (M + H). Anal. Calcd for C₁₅H₁₃FN₆O₃: C, 52.33; H, 3.81; N, 24.41. Found: C, 52.32; H, 3.84; N, 24.41.

N-[(5*S*)-3-[4-[2-(3,5-Diamino-4*H*-pyrazol-4-ylidene)hydrazino]-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methylacetamide (**98**). To the dinitrile **97** (1.0 g, 2.90 mmol) suspended in ethanol (15 mL) was added hydrazine hydrate (1.0 mL). A yellow solution resulted which was heated to reflux for 2 h. After cooling, filtration afforded **98** as a caked golden solid (1.065 g, 97%); mp 256 °C dec; ¹H NMR δ (400 MHz, DMSO) δ 1.82 (s, 3 H), 3.40 (t, *J* = 5.1 Hz, 2 H), 3.75 (t, *J* = 7.7 Hz, 1 H), 4.14 (t, *J* = 9.0 Hz, 1 H), 4.64–4.78 (m, 1 H), 6.13 (bd, 3 H), 7.28 (d, *J* = 9.0 Hz, 1 H), 7.55 (d, *J* = 13.6 Hz, 1 H), 7.81 (t, *J* = 9.0 Hz, 1 H), 8.23 (t, *J* = 5.4 Hz, NH); MS (ESI+) *m/z* 375 (M + H). Anal. Calcd for C₁₅H₁₇FN₈O₃: C, 47.87; H, 4.56; N, 29.77. Found: C, 47.49; H, 4.61; N, 29.52.

***N*-((5*S*)-3-[4-(4-Cyano-2*H*-1,2,3-triazol-2-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl)methylacetamide (99).** Cupric acetate monohydrate (4.0 g, 20 mmol) was stirred in pyridine (20 mL) to give a brilliant blue mixture. Diaminopyrazole **98** (1.88 g, 5.0 mmol) was added and the mixture heated at 100 °C for 1 h. After cooling, the solvent was removed at 40 °C/1 mm and the dark residue partitioned between chloroform (150 mL) and 1 N HCl (100 mL). The acid layer was washed with chloroform (2 × 100 mL) and the combined organics dried (MgSO₄), filtered and evaporated. The chloroform was evaporated and the residue chromatographed over silica gel (50 g) eluting with a 0–3% methanol–chloroform gradient. The product was obtained as a creamy yellow powder (128 mg, 7%): mp 164–5 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.04 (s, 3 H), 3.60–3.78 (m, 2 H), 3.87 (dd, *J* = 6.8, 9.1 Hz, 1 H), 4.11 (t, *J* = 9.0 Hz, 1 H), 4.80–4.90 (m, 1 H), 5.99 (t, *J* = 6 Hz, NH), 7.37 (d, *J* = 9.0 Hz, 1 H), 7.76 (dd, *J* = 2.4, 12.9 Hz, 1 H), 7.83 (t, *J* = 8.5 Hz, 1 H), 8.20 (s, 1 H); HRMS (EI) calcd for C₁₅H₁₃FN₆O₃ 344.1033, found 344.1022. Anal. Calcd for C₁₅H₁₃FN₆O₃·0.2H₂O: C, 51.79; H, 3.88; N, 24.15. Found: C, 51.87; H, 3.99; N, 23.78.

Methyl 1-(4-((5*S*)-5-[(Acetylamino)methyl]-2-oxo-1,3-oxazolidin-3-yl)-2-fluorophenyl)-1*H*-1,2,4-triazole-3-carboxylate (100). A mixture of **20** (5.00 g, 18.73 mmol) and 3 N HCl (26.25 mL, 78.75 mmol) at 0 °C was treated dropwise with a solution of NaNO₂ (1.31 g, 18.91 mmol) in 6 mL of water over a 4 min period. After stirring for 8 min, the mixture was added to an ice cooled, mechanically stirred solution of NaHCO₃ (20.45 g, 243.5 mmol) in 200 mL of water over a 1 min period. After stirring for an additional 5 min, a solution of methyl isocyanacetate (2.04 g, 20.60 mmol) in 25 mL of CH₃OH was added dropwise over a 5 min period at 0–5 °C. The reaction was stirred at 0–5 °C for 5 h. The mixture was diluted with CHCl₃–CH₃OH (9:1) and 200 mL of water, and the aqueous layer was extracted six times with CHCl₃–CH₃OH (9:1). The combined organics were dried with Na₂SO₄ and concentrated with 26 g of silica gel. The material was chromatographed over 350 g of silica gel, packed and eluted with EtOAc–CH₃OH (95:5) with increasing polarity gradient to 8% CH₃OH to obtain 5.50 g as an off-white solid. Trituration with 200 mL of Et₂O at room temperature gave 5.50 g (78%) of **100** as a white solid: mp 165–166.5 °C; [α]_D²⁵ –27° (*c* 0.27, methanol); ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.15 (s, 1 H), 7.84 (t, *J* = 9 Hz, 1 H), 7.79 (dd, *J* = 2, 13 Hz, 1 H), 7.53 (dd, *J* = 2, 11 Hz, 1 H), 4.77 (m, 1 H), 4.17 (t, *J* = 9 Hz, 1 H), 3.88 (s, 3 H), 3.79 (m, 1 H), 3.43 (t, *J* = 5 Hz, 2 H), 1.82 (s, 3 H); MS (EI) *m/z* 377 (M + H). Anal. Calcd for C₁₆H₁₆FN₅O₅: C, 50.93; H, 4.27; N, 18.56. Found: C, 50.86; H, 4.34; N, 18.34.

***N*-[(5*S*)-3-[3-Fluoro-4-[3-(hydroxymethyl)-1*H*-1,2,4-triazol-1-yl]phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl]acetamide (101).** A mixture of lithium borohydride (0.880 g, 95% assay, 0.040 mol) in 160 mL of 2-propanol under N₂ was treated with **100** (7.54 g, 0.020 mol) at room temperature. The reaction was stirred at room temperature overnight. The mixture was treated with 30 mL of water and stirred to decompose the excess LiBH₄. The residuals were suspended in CH₃OH/H₂O and adsorbed onto 35 g of silica gel at reduced pressure. The material was chromatographed with 400 g of silica gel packed with CHCl₃–CH₃OH (92:8) and eluted with increasing CH₃OH gradient to 12% to obtain 5.07 g (73%) of product. Recrystallization from EtOAc, CH₃OH and Et₂O mixture gave analytically pure **101** as white crystals: mp 180–183 °C.

***N*-((5*S*)-3-[3-Fluoro-4-(3-formyl-1*H*-1,2,4-triazol-1-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methylacetamide (102).** A mixture of **101** (0.349 g, 1.00 mmol), 4-methylmorpholine *N*-oxide (0.176 g, 1.50 mmol), tetrapropylammonium perruthenate (TPAP) (0.018 g, 0.05 mmol) and 0.50 g of flame-dried 4 Å molecular sieves in 8 mL of acetonitrile was heated at 60–65 °C for 10 h. The mixture was diluted with EtOAc and washed with 50% saturated brine (40 mL). The aqueous layer was backed extracted two times with EtOAc and the combined organics were dried over Na₂SO₄ and concentrated in vacuo. Chromatography over 50 g of silica gel, packed and eluted with

acetone–methylene chloride (2:1) afforded 0.058 g (17%) of **102** as a white solid: mp 186.5–188.5 °C.

1-(4-((5*S*)-5-[(Acetylamino)methyl]-2-oxo-1,3-oxazolidin-3-yl)-2-fluorophenyl)-1*H*-1,2,4-triazole-3-carboxamide (103). To a solution of 20 mL of 29% NH₄OH in 40 mL of CH₃CN at 50 °C was added **100** (0.500 g, 0.133 mmol). After 3.5 h, the reaction mixture was cooled to 0–5 °C for 25 min, and the white solid was collected and dried in a vacuum oven at 50 °C overnight to yield 375 mg (78%) of **103**: mp 269–270 °C dec.

***N*-((5*S*)-3-[4-(3-Cyano-1*H*-1,2,4-triazol-1-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl)methylacetamide (104).** A suspension of **103** (1.20 g, 3.31 mmol) in 45 mL of CH₂Cl₂ at 0–5 °C in an ice bath was treated with pyridine (0.916 g, 11.60 mmol) followed by the dropwise addition of trifluoroacetic anhydride (2.44 g, 11.60 mmol) in 5 mL of CH₂Cl₂ over a 3 min period. The cooling bath was removed after 20 min and the contents allowed to stir overnight at room temperature. The mixture was poured into 110 mL of saturated NaHCO₃ and stirred to decompose the excess reagent. The contents were extracted three times with CH₂Cl₂, dried with anhydrous Na₂SO₄ and concentrated in vacuo with 6 g of silica gel. Chromatography over 55 g of silica gel, packed and eluted with acetone–methylene chloride (1:3), afforded 0.952 g (83.5%) of **104** as a white solid: mp 202–203 °C; [α]_D²⁵ –26 (*c* 0.51, DMSO); ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.39 (s, 1 H), 8.23 (t, *J* = 6 Hz, 1 H), 7.85 (t, *J* = 9 Hz, 1 H), 7.80 (dd, *J* = 2, 13 Hz, 1 H), 7.54 (dd, *J* = 2, 9 Hz, 1 H), 4.77 (m, 1 H), 4.18 (t, *J* = 9 Hz, 1 H), 3.79 (m, 1 H), 3.43 (t, *J* = 5 Hz, 2 H), 1.82 (s, 3 H); HRMS (FAB) calcd for C₁₅H₁₃FN₆O₃ + H 345.1111, found 345.1117. Anal. Calcd for C₁₅H₁₃FN₆O₃: C, 52.33; H, 3.81; N, 24.41. Found: C, 52.22; H, 3.89; N, 24.35.

***N*-((5*S*)-3-[3-Fluoro-4-(3-methyl-1*H*-1,2,4-triazol-1-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methylacetamide (106).** To a stirred suspension of ethyl acetimidate hydrochloride (**105**) (0.492 g, 4.00 mmol) in 10 mL of absolute ethanol at room temperature under N₂ was added triethylamine (0.424 g, 4.20 mmol) and the mixture stirred for 13 min. A sample of hydrazine **72** (1.13 g, 4.00 mmol) was added with 2 mL of an additional absolute ethanol rinse. In a separate flask acetyl chloride (0.343 g, 4.40 mmol) was added to 3 mL of CH₃OH to prepare HCl in situ. After cooling to 0 °C, the methanolic HCl was added at once to the reaction mixture at 0 °C. The cooling bath was removed after addition, and the contents were stirred at ambient temperature for 10 min and concentrated in vacuo. The dry solid was treated with 15 mL of triethyl orthoformate and heated for 25 min at 100 °C, allowing the ethanol to distill off. Absolute ethanol (3 mL) was added to dissolve some insoluble material and the mixture was heated at 85 °C. The mixture was diluted with CHCl₃–CH₃OH (95:5) and washed successively with saturated NaHCO₃ (25 mL) and 50% saturated brine (30 mL) and the combined aqueous washings were back-extracted with CHCl₃–CH₃OH (95:5) (4 × 30 mL). The combined organics were dried with Na₂SO₄, concentrated at reduced pressure with 10 g of silica gel and chromatographed over 100 g of silica gel packed and eluted with CHCl₃–CH₃OH (97:3 to 96:4) to obtain 478 mg (36%) of **106** as a white solid: mp 206–208 °C; [α]_D²⁵ –27 (*c* 0.41, methanol); ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (brs, 1 H), 8.25 (m, 1 H), 7.76 (m, 2 H), 7.46 (d, *J* = 9 Hz, 1 H), 4.75 (m, 1 H), 4.16 (t, *J* = 9 Hz, 1 H), 3.77 (t, *J* = 9 Hz, 1 H), 3.42 (t, *J* = 5 Hz, 2 H), 2.34 (s, 3 H), 1.82 (s, 3 H); MS (EI) *m/z* 333 (M + H). Anal. Calcd for C₁₅H₁₆FN₅O₃: C, 54.05; H, 4.84; N, 21.01. Found: C, 54.01; H, 4.81; N, 20.92.

Ethyl 5-Benzyl-1,2,4-oxadiazol-3-yliminoformate (108). 3-Amino-5-benzyl-1,2,4-oxadiazol (**107**)⁴³ (12 g, 68.5 mmol) was suspended in triethyl orthoformate and refluxed overnight. The reaction was evaporated to dryness under Hi-vac to give **108** as a yellow oil which was used as is: 14.51 g (92%); ¹H NMR (400 MHz, CDCl₃) δ 1.38 (t, *J* = 7 Hz, 3 H), 4.17 (s, 2 H), 4.43 (q, *J* = 7 Hz, 2 H), 7.28–7.37 (m, 5 H), 8.48 (s, 1 H).

***N*-((5*S*)-3-[4-((5-Benzyl-1,2,4-oxadiazol-3-yl)imino)methyl]amino)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl)methylacetamide (109).** Compound **108** (13.9 g, 60

mmol) and **20** (16 g, 60 mmol) were placed in THF (100 mL) and absolute EtOH (50 mL) and refluxed overnight. The reaction was cooled to room temperature and the precipitated product **109** was isolated as a white solid: 22 g (82%); mp 233–235 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.83 (s, 3 H), 3.43 (t, *J* = 5 Hz, 1 H), 3.68 (br s, 1 H), 3.78 (dd, *J* = 7, 9 Hz, 1 H), 4.17 (t, *J* = 9 Hz, 1 H), 4.99 (br m, 1 H), 7.23–7.32 (m, 6 H), 7.48 (dd, *J* = 2, 9 Hz, 1 H), 7.72–7.78 (m, 2 H), 8.23 (t, *J* = 6 Hz, 1 H), 8.81 (d, *J* = 2 Hz, 1 H), 10.8 (br s, 1 H).

N-[1-(4-[(2*R*)-3-(Acetylamino)-2-hydroxypropyl]amino)-2-fluorophenyl]-1*H*-1,2,4-triazol-3-yl]-2-phenylacetamide (110**). The intermediate **109** (5 g, 11 mmol) was dissolved in EtOH (80 mL) and 10% NaOH (12 mL) and stirred at room temperature overnight. AcOH (11 mL) was added and the reaction was concentrated in vacuo at which time the solid product formed and was isolated, washed with cold EtOH and dried: 4.2 g (89%); mp 191–193 °C; HRMS (FAB) calcd for C₂₁H₂₃FN₆O₃ + H 427.1894, found 427.1886. Anal. Calcd for C₂₁H₂₃FN₆O₃: C, 59.15; H, 5.44; N, 19.71. Found: C, 56.13; H, 5.58; N, 18.38.**

N-[1-(4-[(5*S*)-5-[(Acetylamino)methyl]-2-oxo-1,3-oxazolidin-3-yl]-2-fluorophenyl)-1*H*-1,2,4-triazol-3-yl]-2-phenylacetamide (111**). The amino alcohol **110** (2.3 g, 5.4 mmol) was suspended in CH₂Cl₂ (125 mL) and triphosgene (2.4 g, 8.1 mmol) was added. The reaction was stirred overnight at room temperature and concentrated and saturated NaHCO₃ was added. The aqueous was decanted to leave a sticky gum that was dissolved in acetone. Precipitated starting material was filtered off and the filtrate was evaporated. The residue was dissolved in MeOH and the desired product precipitated as a white solid: 0.81 g (59%); mp 204–206 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.82 (s, 3 H), 3.42 (t, *J* = 5 Hz, 1 H), 3.68 (br s, 1 H), 3.79 (dd, *J* = 7, 9 Hz, 1 H), 4.17 (t, *J* = 9 Hz, 1 H), 4.77 (br m, 1 H), 7.25 (br m, 1 H), 7.3–7.33 (m, 5 H), 7.48 (d, *J* = 11 Hz, 1 H), 7.72–7.78 (m, 2 H), 8.23 (br t, 1 H), 8.81 (d, *J* = 2 Hz, 1 H), 10.85 (br s, 1 H). Anal. Calcd for C₂₂H₂₁FN₆O₄: C, 58.40; H, 4.68; N, 18.58. Found C, 58.09; H, 4.77; N, 18.38.**

N-[(5*S*)-3-[4-(3-Amino-1*H*-1,2,4-triazol-1-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (112**). The benzoyl derivative **111** (1 g, 2.2 mmol) was suspended in EtOH and concentrated HCl (20 drops) was added. The mixture was refluxed for 3 h and the solvent was removed in vacuo. Saturated NaHCO₃ was added to the residue and the mixture was diluted with water. The precipitated product was isolated and dried to give a white solid: 0.68 g (92%); mp 222–224 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.83 (s, 3 H), 3.42 (t, *J* = 5 Hz, 2 H), 3.77 (dd, *J* = 7, 9 Hz, 1 H), 4.15 (t, *J* = 9 Hz, 1 H), 4.75 (br m, 1 H), 5.69 (s, 2 H), 7.43 (d, *J* = 9 Hz, 1 H), 7.68–7.73 (m, 2 H), 8.23 (br t, 1 H), 8.44 (d, *J* = 2 Hz, 1 H); HRMS (EI) calcd for C₁₄H₁₅FN₆O₃ 334.1190, found 334.1192. Anal. Calcd for C₁₄H₁₅FN₆O₃·1/3H₂O: C, 49.41; H, 4.64; N, 24.69. Found: C, 49.62; H, 4.76; N, 24.37.**

N-[(5*S*)-3-[4-(3-Chloro-1*H*-1,2,4-triazol-1-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl]acetamide (113**). The amino analogue **112** (1 g, 3 mmol) was placed in 2 N HCl (30 mL) at 0 °C and NaNO₂ (0.23 g, 3.2 mmol) in water was added. The yellow mixture was stirred at 0 °C for 30 min and CuCl (34 mg, 3.4 mmol) was added. The reaction was warmed to room temperature and stirred overnight. Saturated NH₄Cl was added and the mixture was extracted with EtOAc and dried (Na₂SO₄). Removal of solvent gave a residue that was purified on a silica gel column with 2% MeOH/CH₂Cl₂ to give 245 mg (22%); mp 182–183 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.04 (s, 3 H), 3.67–3.74 (m, 2 H), 3.86 (dd, *J* = 7, 9 Hz, 1 H), 4.11 (t, *J* = 9 Hz, 1 H), 4.84 (br m, 1 H), 6.07 (br t, 1 H), 7.28 (d, *J* = 9 Hz, 1 H), 7.81 (dd, *J* = 2, 14 Hz, 1 H), 7.85 (t, *J* = 9 Hz, 1 H), 8.52 (d, *J* = 2 Hz, 1 H). Anal. Calcd for C₁₄H₁₃ClFN₅O₃: C, 47.54; H, 3.70; N, 19.80. Found: C, 47.46; H, 3.63; N, 19.63.**

Cinnamaldehyde Tosylhydrazone (114**)**. Cinnamaldehyde (3.4 mL, 26.96 mmol) was added to an ethanol (50 mL) solution of *p*-toluenesulfonyl hydrazide (5.008 g, 26.89 mmol). The reaction mixture was stirred for 1 h, then was cooled to 0

°C. The resulting yellow solid was filtered and recrystallized from boiling ethanol and water to give 7.101 g (88%) of the desired material as white needles: mp 162–163 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.62 (bs, 1H), 2.43 (s, 3 H), 6.81–6.88 (m, 2 H), 7.30–7.37 (m, 5 H), 7.40–7.42 (m, 2 H), 7.57 (d, *J* = 8 Hz, 1 H), 7.88 (d, *J* = 14 Hz, 2 H); HRMS (EI) calcd for C₁₆H₁₆N₂O₂S 300.0932, found 300.0938. Anal. Calcd for C₁₆H₁₆N₂O₂S: C, 63.98; H, 5.37; N, 9.33. Found: C, 63.91; H, 5.42; N, 9.32.

N-[(5*S*)-3-(3-Fluoro-4-{5-[(*E*)-2-phenylethenyl]-2*H*-1,2,3,4-tetraazol-2-yl}phenyl)-2-oxo-1,3-oxazolan-5-yl]methyl]acetamide (115**). A precooled (0 °C) aqueous (0.7 mL) solution of sodium nitrite (0.071 g, 1.03 mmol) was added to a solution of **20** (0.274 g, 1.03 mmol) in water (1 mL), hydrochloric acid (0.37 mL), and ethanol (1 mL). This reaction mixture was stirred at 0 °C for 10 min. Then it was added, via cannula, over 30 min, to a pyridine (8 mL) solution of **114** (0.303 g, 1.01 mmol) which had been cooled in an ice/salt bath. After stirring 2 h, chloroform and water were added to the reaction mixture. The phases were separated, and the organic portion was washed with water and brine, dried with MgSO₄, and evaporated. The residue was chromatographed on a 2.3 × 25 cm medium-pressure silica column with 2% MeOH/CH₂Cl₂ to give **115** (0.150 g, 35%) as a brown solid: mp 180–182 °C; ¹H NMR (400 MHz, CDCl₃ + MeOD) δ 1.99 (s, 3 H), 3.60–3.64 (m, 2 H), 3.84 (dd, *J* = 9, 7 Hz, 1 H), 4.10 (t, *J* = 9 Hz, 1 H), 4.77–4.84 (m, 1 H), 7.19 (d, *J* = 16 Hz, 1 H), 7.31–7.39 (m, 5 H), 7.56 (d, *J* = 7 Hz, 2 H), 7.76 (dd, *J* = 13, 2 Hz, 1 H), 7.81 (d, *J* = 17 Hz, 1 H), 7.83 (dd, *J* = 12, 9 Hz, 1 H); HRMS (FAB) calcd for C₂₁H₁₉FN₆O₃ + H 423.1581, found 423.1585. Anal. Calcd for C₂₁H₁₉FN₆O₃: C, 59.71; H, 4.53; N, 19.90. Found: C, 59.45; H, 4.56; N, 19.77.**

N-[(5*S*)-3-[3-Fluoro-4-(5-formyl-2*H*-1,2,3,4-tetraazol-2-yl)phenyl]-2-oxo-1,3-oxazolan-5-yl]methyl]acetamide (116**). A suspension of **115** (1.355 g, 3.21 mmol) in CH₂Cl₂ (100 mL) was cooled to –78 °C. Ozone was bubbled through the solution until it turned blue. After stirring for 20 min, nitrogen was then bubbled through the reaction mixture for about 5 min. Then dimethyl sulfide (6 mL) was added, and the mixture was stirred at –78 °C for 5 min then warmed to room temperature. The solvent was evaporated, and the crude material, a cream solid, was used in the next reaction as is: mp 230 °C dec.**

N-[(5*S*)-3-[4-(5-Cyano-2*H*-1,2,3,4-tetraazol-2-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolan-5-yl]methyl]acetamide (118**). Acetonitrile (7.5 mL) and carbon tetrachloride (1 mL) were added to a flask containing **117** (0.052 g, 0.14 mmol) and triphenylphosphine (0.070 g, 0.27 mmol). The reaction mixture was heated at 85 °C overnight under nitrogen atmosphere. Water and CH₂Cl₂ were added to the reaction mixture. The phases were separated, and the aqueous portion was extracted with CH₂Cl₂, the combined organic portions were dried (MgSO₄) and evaporated. The crude material was chromatographed on a 2.3 × 26 cm medium-pressure silica column with 2–5% MeOH/CH₂Cl₂ to give 0.034 g (69%) of the desired compound **118** as a white solid: mp 199–200 °C; [α]_D²⁵ = –18° (c 0.93, DMSO); ¹H NMR (400 MHz, CDCl₃) δ 1.98 (s, 3 H), 3.63 (d, *J* = 4 Hz, 2 H), 3.86 (dd, *J* = 9, 7 Hz, 1 H), 4.11 (t, *J* = 9 Hz, 1 H), 4.79–4.85 (m, 1 H), 7.20 (bt, 1 H), 7.42 (dt, *J* = 9, 1 Hz, 1 H), 7.82 (m, 1 H), 7.84 (t, *J* = 9 Hz, 1 H); HRMS (FAB) calcd for C₁₄H₁₂FN₇O₃ + H 346.1064, found 346.1078. Anal. Calcd for C₁₄H₁₂FN₇O₃: C, 48.70; H, 3.50; N, 28.40. Found: C, 48.68; H, 3.54; N, 28.05.**

Acknowledgment. We thank the Structural, Analytical and Medicinal Chemistry Department of Pharmacia & Upjohn for mass spectral and combustion analysis data and the Oxazolidinone Program Team for many helpful discussions.

Supporting Information Available: Full experimental descriptions and analytical data for compounds **13c,d–f**, **14b–g**, **15b–g**, **18**, **4–8**, **10**, **27**, **37**, **40**, **41**, **48**, **49**, **61**, **62**, **81**, **88–93**, **95**, **96**, and **117** and complete analytical data for com-

pounds **29**, **30**, **32–35**, **38**, **39**, **44**, **52–58**, **65–70**, **78**, **79**, **83**, **94**, **101–103**, and **116**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Swartz, M. N. Hospital-acquired Infections: Diseases with Increasingly Limited Therapies. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 2420–2427.
- Tomasz, A. Multiple-Antibiotic-Resistant Pathogenic Bacteria. *N. Engl. J. Med.* **1994**, *330*, 1247–1251.
- Fekete, T. Bacterial Genetics, Antibiotic Usage, and Public Policy: The Crucial Interplay in Emerging Resistance. *Perspect. Biol. Med.* **1995**, *38*, 363–382.
- Brumfitt, W.; Hamilton-Miller, J. M. T. The Challenge of Methicillin-Resistant *Staphylococcus aureus*. *Drugs Exp. Clin. Res.* **1994**, *XX*, 215–224.
- Low, D. E.; Willey, B. M.; McGeer, A. J. Multidrug-resistant Enterococci: a Threat to the Surgical Patient. *Am. J. Surg.* **1995**, *54* (Suppl.), 8S–12S.
- Spera, R. V., Jr.; Farber, B. F. Multidrug-Resistant *Enterococcus faecium*: An Untreatable Nosocomial Pathogen. *Drugs* **1994**, *48*, 678–688.
- Appelbaum, P. C. Antimicrobial Resistance in *Streptococcus pneumoniae*. *Clin. Infect. Dis.* **1992**, *15*, 77–83.
- Hiramatsu, K.; Hanaki, H.; Ino, T.; Yabuta, K.; Oguri, T.; Tenover, F. C. Methicillin-resistant *Staphylococcus aureus* Clinical Strain with Reduced Vancomycin Susceptibility. *J. Antimicrob. Chemother.* **1997**, *40*, 135–136.
- Smith, T. L.; Pearson, M. L.; Wilcox, K. R.; Cruz, C.; Lancaster, M. V.; Robinson-Dunn, B.; Tenover, F. C.; Zervos, M. J.; Band, J. D.; White, E.; Jarvis, W. R. Emergence of Vancomycin Resistance in *Staphylococcus aureus*. *N. Engl. J. Med.* **1999**, *340*, 493–501.
- Sieradzki, K.; Roberts, R. B.; Haber, S. W.; Thomasz, A. The Development of Vancomycin Resistance in a Patient with Methicillin-Resistant *Staphylococcus aureus* Infection. *N. E. J. Med.* **1999**, *340*, 517–523.
- Waldvogel, F. A. New Resistance in Resistant *Staphylococcus aureus*. *N. Engl. J. Med.* **1999**, *340*, 556–557.
- Hanberger, H.; Garcia-Rodriguez, J.-A.; Gobernado, M.; Goossens, H.; Nilsson, L. E.; Struelens, M. J.; Antibiotic Susceptibility Among Aerobic Gram-negative Bacilli in Intensive Care Units in 5 European Countries. *J. Am. Med. Assoc.* **1999**, *281*, 67–71.
- Brickner, S. J. Oxazolidinone Antibacterial Agents. *Curr. Pharm. Des.* **1996**, *2*, 175–194.
- Lin, A. H.; Murray, R. W.; Vidmar, T. J.; Marotti, K. R. The Oxazolidinone Eperezolid Binds to the 50S Ribosomal Subunit and Competes with Binding of Chloramphenicol and Lincomycin. *Antimicrob. Agents Chemother.* **1997**, *41*, 2127–2131.
- Shinabarger, D. L.; Marotti, K. R.; Murray, R. W.; Lin, A. H.; Melchior, E. P.; Swaney, S. M.; Dunyak, D. S.; Demyan, W. F.; Buysse, J. M. Mechanism of Action of Oxazolidinones: Effects of Linezolid and Eperezolid on Translation Reactions. *Antimicrob. Agents Chemother.* **1997**, *41*, 2132–2136.
- Zurenko, G. E.; Yagi, B. H.; Schaadt, R. D.; Allison, J. W.; Kilburn, J. O.; Glickman, S. E.; Hutchinson, D. K.; Barbachyn, M. R.; Brickner, S. J. In Vitro Activities of U-100592 and U-100766, Novel Oxazolidinone Antimicrobial Agents. *Antimicrob. Agents Chemother.* **1996**, *39*, 839–845.
- Kaatz, G. W.; Seo, S. M. In Vitro Activities of Oxazolidinone Compounds U-100592 and U-100766 against *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Antimicrob. Agents Chemother.* **1996**, *39*, 799–801.
- Genin, M. J.; Hutchinson, D. K.; Allwine, D. A.; Hester, J. B.; Emmert, D. E.; Garmon, S. A.; Ford, C. W.; Zurenko, G. E.; Hamel, J. C.; Schaadt, R. D.; Stapert, D.; Yagi, B. H.; Friis, J. M.; Shobe, E. M.; Adams, W. J. Nitrogen–Carbon-Linked (Azolylphenyl)oxazolidinones with Potent Antibacterial Activity Against the Fastidious Gram-Negative Organisms *Haemophilus influenzae* and *Moraxella catarrhalis*. *J. Med. Chem.* **1998**, *41*, 5144–5147.
- Brickner, S. J.; Hutchinson, D. K.; Barbachyn, M. R.; Manninen, P. R.; Ulanowicz, D. A.; Garmon, S. A.; Grega, K. C.; Hendges, S. K.; Toops, D. S.; Ford, C. W.; Zurenko, G. E. Synthesis and antibacterial activity of U-100592 and U-100766, two oxazolidinone antibacterial agents for the potential treatment of multidrug-resistant gram-positive bacterial infections. *J. Med. Chem.* **1996**, *39*, 673–679.
- Hutchinson, D. K. Preparation of N-(heteroarylphenyl)oxazolidin-2-ones as bactericides. WO 9623788 (CAN 125:247827).
- Litkei, G.; Patonay, T.; Patonay-Peli, E.; Khilya, V. P. A simple synthesis of 3-(1-tetrazolyl)benzopyranoids. *Pharmazie* **1989**, *44*, 791–792.
- Kim, S. Novel synthetic reactions with 2-pyridyl related reagents. *Org. Prep. Proc. Int.* **1988**, *20*, 147–172.
- Wikel, J. H.; Paget, C. J. Synthesis of s-triazolo[4,3-b]benzothiazoles. *J. Org. Chem.* **1974**, *39*, 3506–3508.
- Aulaskari, P.; Ahlgren, M.; Rouvinen, J.; Vainiotalo, P. Preparation and Structure Determination of 1-Benzyl-, 1-Methyl- and 1H-5-[(2-Nitro-2-phenyl)ethenyl]imidazoles. *J. Heterocycl. Chem.* **1996**, *33*, 1345–1354.
- Vega, S.; Gil, M. S. Synthesis of 5,6-Dihydro-4H-pyrrolo[1,2-a]thieno[2,3-f][1,4]diazepines[1,2]. *J. Heterocycl. Chem.* **1991**, *28*, 945–950.
- Kim, J. N.; Chung, K. H.; Ryu, E. K. Improved Dehydration Method of Aldoximes to Nitriles. Use of Acetonitrile in Triphenylphosphine/Carbon Tetrachloride System. *Synth. Commun.* **1990**, *20*, 2785–2788.
- Corey, E. J.; Gilman, N. W.; Ganem, B. E. New Methods for the Oxidation of Aldehydes to Carboxylic Acids and Esters. *J. Am. Chem. Soc.* **1968**, *90*, 5616–5617.
- Bertz, S. H.; Dabbagh, G.; Cotte, P. New Preparations of Ethyl 3,3-Diethoxypropionate and (Ethoxycarbonyl)malondialdehyde. Cu(II)-Catalyzed Acetal Formation from a Conjugated Triple Bond. *J. Org. Chem.* **1982**, *47*, 2216–2217.
- Holzer, W.; Seiringer, G. N-1 Substituted Ethyl 4-Pyrazolecarboxylates: Synthesis and Spectroscopic Investigations. *J. Heterocycl. Chem.* **1993**, *30*, 865–872.
- Ninomiya, K.; Shioiri, T.; Yamada, Y. Phosphorous in Organic Synthesis-VII. Diphenyl Phosphorazidate (DPPA). A New Convenient Reagent for a Modified Curtius Reaction. *Tetrahedron* **1974**, *30*, 2151–2157.
- Clarke, H. T.; Read, R. R. *o*-Tolunitrile and *p*-Tolunitrile. *Org. Synth.* **1941**, *1*, 514.
- Barclay, B. M.; Campbell, N. Dehydrogenation of Tetrahydrocarbazoles by Chloranil. *J. Chem. Soc.* **1945**, 530–532.
- Burness, D. M. β -Keto Acetals. I. Synthesis of Pyrazoles and Pyrimidines and the Steric Inhibition of Resonance in 5-Alkyl-1-*p*-nitrophenylpyrazoles. *J. Org. Chem.* **1956**, *21*, 97.
- Tietze, L. F.; Schneider, C.; Pretor, M. Improved Synthesis of (E)-3-Alkoxy- and (E)-3-Phenoxyacryloyl Chlorides. *Synthesis* **1993**, 1079.
- Effenberger, F.; Hartmann, W. Synthesen mit β -Aldoxyacrylsäurechloriden. *Chem. Ber.* **1969**, *102*, 3260.
- Mizuno, M.; Shiori, T. Efficient method for the one-pot azidation of alcohols using bis(*p*-nitrophenyl)phosphorazidate. *J. Chem. Soc., Chem. Commun.* **1997**, *22*, 2165–2166.
- Hayashi, H.; Ohno, A.; Oka, S. Palladium-catalyzed Decomposition of Azides. I. Formation of Nitriles. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 506.
- Palacios, F.; Ochoa de Retana, A. M.; Pagalday, J.; Sanchez, J. M. Cycloadditions of Azidoalkylcarboxylates to Acetylenes and Enamines. Regioselective Synthesis of Substituted Triazoles. *Org. Prep. Proc. Int.* **1995**, *27*, 603–612.
- Elnagdi, M. H.; Allah, S. O. A. Reactions with the Arylhydrazones of Some α -Cyanoketones. *J. Prakt. Chem.* **1973**, *315*, 1009.
- Goldberg, M. W.; Rachlin, A. I. U.S. 2,658,889; *Chem. Abstr.* **49**, 1103g.
- Matsumoto, K.; Suzuki, M.; Tomie, M.; Yoneda, W.; Miyoshi, M. A New Synthesis of 1-Aryl-1H-1,2,4-triazole-3-carboxylic Acid Esters. *Synthesis* **1975**, 609.
- Westermann, P.; Paul, H.; Hilgetag, G. Über die Umsetzung des Benzyloxycarbonylamino-acetamidessäure-athylesters mit Alkyliden- und Arylhydrazinen. *Chem. Ber.* **1966**, *99*, 1111.
- Buscemi, S.; Vivona, N.; Caronna, T. A generalized and Efficient Synthesis of 3-Amino, 3-(N-N-Dialkylamino)-5-alkyl-1,2,4-oxadiazoles by Irradiation of 3-Alkanoylamino-4-phenyl-1,2,5-oxadiazoles (Furazans). *Synthesis* **1995**, *8*, 917–919.
- Ruccia, M.; Vivona, N.; Cusmano, G. Rearrangements in the 1,2,4-Oxadiazole Series. IV. Conversion of N-(1,2,4-Oxadiazol-3-yl)-N'-arylformamidines into 3-Acylamino-1-aryl-1,2,4-triazoles. *J. Heterocycl. Chem.* **1971**, *8*, 137.
- Suzuki, N.; Kaneko, Y.; Nomoto, T.; Izawa, Y. Synthetic Reactions in Polyethylene Glycol. Diazotization and Sandmeyer Reactions of Anilines in Polyethylene Glycol-Methylene Dichloride. *J. Chem. Soc., Chem. Commun.* **1984**, *22*, 1523.
- Ito, S.; Tanaka, Y.; Kakehi, A.; Kondo, K. A Facile Synthesis of 2,5-Disubstituted Tetrazoles by the Reaction of Phenylsulfonylhydrazones with Arenediazonium Salts. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 1920–1923.
- Ford, C.; Hamel, J.; Stapert, D.; Moerman, J.; Hutchinson, D. K.; Barbachyn, M. R.; Zurenko, G. Oxazolidinones: A New Class of Antimicrobials. *Infect. Med.* **1999**, 435–445.
- Buyse, J. M.; Demyan, W. F.; Dunyak, D. S.; Stapert, D.; Hamel, J. C.; Ford, C. W. Mutation of the *acrAB* antibiotic efflux pump in *Escherichia coli* confers susceptibility to oxazolidinone antibiotics. Program and Abstracts of the 36th Interscience Conference on Antimicrobial Agents and Chemotherapy, 1996: Abstr. 62.